

SOME BIOCHMICAL CHANGES IN SERUM OF FEMALE ALBINO RATS ADMINISTERED AQUEOUS EXTRACT OF *PIPER GUINEENSE* SEEDS

Abstract

This study examined the biochemical changes in serum of female albino rats administered aqueous extract of *Piper guineense* seeds extract. The test animals groups 2, 3 and 4 received 25mg/kg, 50mg/kg and 75mg/kg of the extract respectively for 21 days, while group 1 served as control and received a placebo of 0.5% physiological saline which is the vehicle for administration. On the 22nd day the test animals which have been fasted overnight were sacrificed with chloroform anesthesia. Blood was collected through cardiac puncture. Results show that serum protein, albumin and packed cell volume concentration significantly ($p \leq 0.05$) increased. Creatinine and urea levels decreased ($p \leq 0.05$) significantly, while cholesterol profile decreased ($p \leq 0.05$) significantly, except HDL. Superoxide dismutase, glutathione-s-transferase, increased ($p \leq 0.05$) significantly, while lipid peroxidation decreased. This study has shown that nursing mothers in South East of Nigeria who use the *Piper guineense* seeds may derive immense benefit from its effect on the biochemical parameters assayed in this study.

Key words: Spices, Lipid, Antioxidants, Protein, Creatinine, Urea

Introduction

Spices are aromatic vegetable substances that are used in small quantities for seasoning of foods and also as preservatives or flavoring [1]. Spices enhance the taste of food, beverage and drug [2]. Spices contain phytochemicals which gives them their characteristic odour flavor and other properties [3].

Piper guineense (Utazi or black paper) is a spice, a West African tropical plant that is widely consumed and is considered medicinal by some people [4]. The aroma, flavor and preservative properties of *Piper guineense* are probably due to the nutritional and non-nutritional factors of the plant. It is rich in protein, fat, fiber and carbohydrate. The vitamin content is also high, especially vitamin C, which is an antioxidant. Peroxide value and free fatty acid level are generally low. The essential oils content is fairly high and range from 0.1 to 5.2%. The fruits of *Piper guineense* (which is the part of the plant that is traditionally used) are rich in a wide range of natural products, including volatile oils, lignans, amides, alkaloids, flavonoids and polyphenols [5]. Juliani *et al.*; [6] reported that *Piper guineense* is an important plant used in

traditional medicine and as a spice. As a medicinal plant, *Piper guineense* seeds are used to relieve discomfort in the stomach caused by excess gas, while the leaves are used to treat respiratory infections, rheumatism and syphilis [7]. The leaves and seeds of the plant are used to flavor soup, stew and food generally [8]. The presence of an alkaloid – piperine has also been reported. The piperine contains strong anticonvulsant called anti- epileparine [N- 3-4- methylene dioxycynamyl piperidine] which has been synthesized for the treatment of epilepsy (Abila *et al.*; [9], Oleifa *et al.*; [10], Amosan *et al.*; [11]. Udoh *et al.*; [12] and Echo *et al.*; [5] have also reported that the plant has therapeutic effect for the management of convulsion, leprosy, inflammation and rheumatoid pains. Bacteriostatic and bactericidal properties of *Piper guineense* has also been reported Zollo *et al.*; [13], Okigbo and Igwe [14], while Echo *et al.*; [5] reported the molluscidal properties of the methanolic extract of the plant, and suggested it could be used for the treatment of Bilharzia. Mbongwe *et al.*; [15] reported that the leaves of *Piper guineense* are used for female infertility, while the seeds are used as an aphrodisiac. Hence the Igbos of the South Eastern part of Nigeria use the seeds to prepare soups for nursing mothers from the first day of delivery, to prevent post partum contractions and to aid the fast return of the uterine muscles of the uterus to their original shape. It is also believed that it aids the increase in breast milk production and flow.

The present study is therefore aimed at investigating the biochemical changes that may occur in female albino rats due the administration of aqueous extract of *Piper guineense* seeds.

Materials and Methods

Piper guineense seeds

The *Piper guineense* seeds were bought from the Eke Okigwe market as sold. The seeds were identified and authenticated at the Department of Plant Science and Biotechnology, Abia State University, Uturu Nigeria and voucher specimens deposited at the departmental herbarium.

Preparation of extract

The *Piper guineense* seeds were cleaned and sorted for healthy seeds. The seeds were sun dried for five days, and milled into fine powder using a milling machine. 40g of the powder was dissolved in 300ml of distilled water and left to stand for 48hrs with occasional shaking. The mixture was then filtered using a muslin bag. 263ml of the filtrate was recovered. The concentration of the extract was determined to be 10.15mg/ml.

Experimental Animals and Design

Forty female albino rats, 7 weeks old and weighing between 128g-142g were used for the study. The animals were randomly placed into four groups of 10 animals each. Group 1 served as control, Group 2, 3 and 4 served as test animals. The animals were allowed feed and water *ad libitum*. Standard laboratory protocols for animal studies were maintained. Approval for animal

studies was obtained from the Animal Ethics Committee of the College of Medicine and Health Sciences, Abia State University, Uturu, Nigeria.

Treatment of Animals

The animals were starved overnight prior to the commencement of the administration of the *Piper guineense* seeds extract. The administration of the *Piper guineense* seeds extract was by oral intubation once daily at 25mg/kg body weight, 50mg/kg body weight and 75mg/kg body weight for group 2, 3 and 4 animals respectively for 21 days.

Collection of blood samples and serum preparation

At the 22nd day the animals were sacrificed after overnight starving. Incisions were made into their thoracic cavity. Blood samples were collected by heart aorta puncture using a 10mL hypodermic syringe and allowed to clot in sample vials. The samples were centrifuged at 3000 rpm for 5 min. using the Bran Scientific and Instrument Company England centrifuge. The supernatant (serum) was harvested by simple aspiration with Pasteur pipette and stored in clean tubes at - 4°C until analysis.

Estimation of parameters

Determination of Total Cholesterol

Total cholesterol was determined by the enzymatic colorimetric cho-PAP method, High density lipoprotein (HDL) Very low density lipoprotein (VLDL), and Low density lipoprotein (LDL) were determined as described by Burstein *et al.*; [16]. Triacylglycerol and cholesterol were assayed as described by Fletcher [17], Richmond [18] and Barlett [19], using Biosystem kits. The measurements were performed according to the manufacturers' instruction.

Determination of Total protein

Total protein test kits produced by Randox laboratories were used. This test involves formation of colored complex between cupric ions in alkaline medium with peptide bonds. The intensity of the color formed is proportional to the concentration of the protein.

Determination of albumin

Albumin test kit produced by Biosystem kits were used to estimate albumin. The principle of this test is based on albumin quantitatively binding to bromocresol green. The complex of albumin-BCG so formed absorbs maximally at 578nm, and the absorbance is directly proportional to the concentration of albumin in the sample.

Determination of lipid peroxidation

Thiobarbituric acid reactive substances (TBARS) method as described by Buege and Aust [20] was used to determine lipid peroxidation,

Determination of Antioxidant enzymes

Glutathione-s- transferase (GST) activity was determined according to the method described by Habig *et al.*; [21], while Superoxide Dismutase was measured as described by Kakkar *et al.*; [22].

Hematological parameters

Commercial kits (Biosystem kits) were used to estimate packed cell volume, creatinine and urea. Packed cell volume and creatinine were assayed as described by Dacie and Lewis [23].

Statistical Analysis:

Values were represented as Mean $\pm$ SD. Data obtained were subjected to one way Analysis of Variance (ANOVA) and group means were compared using Duncan's new multiple range tests. Differences were considered to be significant at ($p < 0.05$).

Results

Table 1: Effect of *Piper guineense* seed extract on animal weight (g)

	Group 1 (Control)	Group 2	Group 3	Group 4
Body weight	92.56 $\pm$ 0.65*	98.74 $\pm$ 0.35	101.46 $\pm$ 0.86 *	102.15 $\pm$ 0.63*

*Values are mean $\pm$ SD of triplicate determinations (n=10)

Values with * are statistically significant ($p < 0.05$)

Table 2: Effect of *Piper guineense* seed extract on fatty acid levels (mg/dL)

Parameter	Group 1 (Control)	Group 2	Group 3	Group 4
Triacylglycerol	73.46 $\pm$ 3.28*	72.95 $\pm$ 4.72	70.54 $\pm$ 2.35*	68.42 $\pm$ 3.82*
Total cholesterol	1.43 $\pm$ 0.18	1.37 $\pm$ 0.09	1.26 $\pm$ 0.10	1.20 $\pm$ 0.12*
HDL	35.56 $\pm$ 0.86*	36.43 $\pm$ 0.16*	37.76 $\pm$ 0.24*	38.57 $\pm$ 0.11*
LDL	32.68 $\pm$ 2.14	31.82 $\pm$ 2.06	30.32 $\pm$ 2.21*	28.54 $\pm$ 2.16
VLDL	11.56 $\pm$ 0.85*	11.14 $\pm$ 0.14*	10.76 $\pm$ 0.56*	10.25 $\pm$ 0.43*

*Values are mean $\pm$ SD of triplicate determinations (n=10)

Values with * are statistically significant ($p < 0.05$)

Table 3: Effect of *Piper guineense* seed extract on Lipid peroxidation levels and antioxidant enzymes

Parameter	Group 1 (Control)	Group 2	Group 3	Group 4
Lipid peroxidation (Mol/L)	4.69 $\pm$ 0.65*	3.28 $\pm$ 0.16	3.11 $\pm$ 0.72	3.01 $\pm$ 0.52*
SOD (IU/L)	68.76 $\pm$ 6.74	104.57 $\pm$ 5.64*	108.26 $\pm$ 5.96*	112.39 $\pm$ 6.86*

GST (IU/L)	6.86 ± 1.64	7.56 ± 1.22	8.73 ± 1.45*	9.28 ± 1.32*
------------	-------------	-------------	--------------	--------------

*Values are mean ±SD of triplicate determinations (n=10)

Values with * are statistically significant (p<0.05)

Table 4: Effect of *Piper guineense* seed extract on some macromolecules

Parameter	Group 1 (Control)	Group 2	Group 3	Group 4
Serum protein(mg/dl)	246.45 ± 12.25	252.25 ± 10.24	265.53 ± 9.65*	278.48 ± 9.82*
Albumin (mg/dl)	2.16 ± 0.13*	2.37 ± 0.17	2.52 ± 0.15*	2.58 ± 0.22*
PCV (L/L)%	30.25 ± 0.56*	33.14 ± 1.05	34.68 ± 1.27*	34.94 ± 1.21*
Creatinine (mg/dl)	0.65 ± 0.03	0.52 ± 0.02	0.46 ± 0.06	0.41 ± 0.04
Urea (mg/dl)	38.74 ± 1.43*	38.14 ± 1.69	36.45 ± 1.36	36.16 ± 1.76*

*Values are mean ±SD of triplicate determinations (n=10)

Values with * are statistically significant (p<0.05)

Discussion

Body weight of experimental animal increased (p≤0.05) significantly (p≤0.05) compared to control (Table 1). Animals treated with 75mg/kg body weight of extract showed significant increase in body weight. The effect of the extract on body weight seems to be dose dependent, as shown on table 1. The increase in weight may be due to the high content of protein in the extract and the phytochemicals which could elicit increase in protein synthesis.

Triacylglycerol, total cholesterol, Low density and very low density cholesterol decreased (p≤0.05) non-significantly compared to control; while high density cholesterol increased (p≤0.05) significantly (Table 2). Cholesterol is a precursor of other steroids such as sex hormones. It is also an important constituent of cell membranes and bile acids [24, 25]. Low density and high density cholesterol are involved in the transport of cholesterol to and from the liver. The reduction of these fatty acids may show that the *Piper guineense* seed extract may have hypocholesterolemic effect. The increase in High density cholesterol could be of immense benefit to the animals, since HDL is also known as the 'good' cholesterol and is involved in transport of other lipids to the liver for disposal in bile [26].

Lipid peroxidation decreased (p≤0.05) significantly (Table 3) compared to control. The significant decrease in lipid peroxidation values in the experimental animals may be as a result of the extract aiding in the reduction of lipid oxidation since the *Piper guineense* seeds are rich in vitamin C which is an antioxidant. Lipid peroxidation is due to the oxidation of lipids especially polyunsaturated lipids to generate peroxides and aldehydes which have been associated with the development of diseases [27, 28]. The aldehydes and peroxides are long-lived and can easily spread to distant sites through blood circulation to cause further peroxidation of cells, hence they have been reported to be more dangerous than reactive oxygen species (ROS) [27]. Lipid peroxidation causes cytotoxicity and injury that leads to cell damage due to its ability to form malondialdehyde (MDA) [29].

The antioxidant enzymes superoxide dismutase (SOD) and glutathione s-transferase (GST) increased (p≤0.05) significantly (Table 3) compared to control. The increase in SOD and GST

could be attributed to the presence of phytochemicals in the *Piper guineense* seeds extract especially flavonoids. The seeds of *Piper guineense* are rich in flavonoids, which may have induced detoxification enzymes through up-regulation of their genes by interacting with antioxidant response elements [30, 31, 32]. Prestera *et al.*; [33] reported that substances like flavonoids that possess electrophilic centers are able to react with sulfhydryl groups through oxido-reduction or alkylation. Pantuck *et al.*; [34] also reported that an increase in GST activity translates to increased capacity to conjugate and excrete toxic intermediates that can cause diseases such as diabetes. The significant increase in antioxidant enzymes further confirms the high antioxidant capacities of the seeds of *Piper guineense*.

In humans the damaging effects of oxidants and lipid peroxidation can be checked by an organized antioxidant defense mechanism. Several enzymatic and non-enzymatic processes are involved that protect cells against oxidative damage. Farombi *et al.*; [35] reported that the major enzymatic anti-oxidant systems include superoxide dismutase (SOD), glutathione peroxidase (GSH), glutathione – s- transferase (GST) and catalase, while the non – enzymatic antioxidants include glutathione and dietary constituents such as vitamins, flavonoids and carotenoids. The antioxidants protect cells from the damaging effects of free radicals and lipid peroxidation.

Serum protein concentration increased ($p \leq 0.05$) significantly, while albumin increased non-significantly (Table 4). This could be attributed to the high protein content of the *Piper guineense* seeds [5]. However, the non-significant increase in albumin content could infer a possible preservation of the stability of the liver. Packed cell volume increased ($p \leq 0.05$) significantly, while creatinine and urea decreased ($p \leq 0.05$) significantly (Table 4). The increase in PCV may be contributed by the rich phytochemical content of *Piper guineense* seeds which increased the rate of hemoglobin synthesis in the experimental animals, while increasing the excretion of creatinine and urea [36].

The kidney is a major organ for excretion and its stability is estimated by the creatinine clearance, which is a function of the glomerular filtration rate.

Conclusion

This study has been able to show that the *Piper guineense* seeds increased the body weight of the experimental animals, hence may elicit and increase protein synthesis. Total cholesterol and triacylglycerol were significantly reduced, hence the seeds may also have the ability to control cholesterol levels. Lipid peroxidation levels were reduced while antioxidant enzymes increased; albumin, pack cell volume, creatinine and urea were all reduced. The reduction in urea, creatinine and cholesterol levels as observed in this study, shows that *Piper guineense* seeds can be used to improve kidney function and for the management of hypercholesterolemia. These observations have lend credence to the claims of alternative medicine practitioners that nursing mothers who use this spices may benefit from its effect on some biochemical indices as reported in this study.

Consent

It is not applicable

References

1. Hafner UC, John DA and Mickey CB. Nutrition for Developing countries. *Journal of Nutrition*. 2003; **6**:18-25
2. Marcel KK, Akhanovna MJ, Yves-Alain B, Gabin DB, Jérôme ZB . *In vitro* antioxidant activities of total flavonoids extracts from leaves and stems of *Adenia lobata* (Jacq.) Engl. (Passifloraceae). *Journal Pharmacog Phytother*. 2010; **3** (1):8-12.
3. Olusegun Ekundayo, Intolaakso Rita Marie, Babajide Oguntimein and Abayomi Sofowora Essential oil constituents of Ashanti pepper (*Piper guineense*) fruits. *J. Agric. Food Chem*. 2000; **26**(5):880-882
4. Iwu M, Hand Book of African medicinal plants. CRC Press, Boca Raton, Flip, 1993; pp: 183-184.
5. Echo IA, Osuagwu AN, Agbor RB, Okpako EC, Ekanem, B.E. Phytochemical Composition of *Aframomum melegueta* and *Piper guineense* Seeds *World Journal of Applied Environmental Chemistry*. 2012; **2** (1): 17-21
6. Juliani Rodolfo H, Koroch AR, Giordano L, Amekuse L, Koffa SJ, Asante-Dartey SE , Simon JE . *Piper guineense* (Piperaceae): Chemistry, Traditional Uses and Functional Properties of West African Black Pepper. African Natural Plant Products. Volume II: Discoveries and Challenges in Chemistry, Health and Nutrition. ACS Symposium series, 2013; Vol.1127 pp 33-48
7. Bep EO. Medicinal Plants in Nigeria. *J. of Agric and Food Sci*. 1999; **7**: 218-225
8. Agoha AE . Uses of *Piper guineense* fruits. *Journal of Food Sci*. 2002; **10** (3): 231-238
9. Abila B, Richens A, Davies JA. Anticonvulsant effects of extracts of the West African black pepper, *Piper guineense*. *Journal of Ethnopharmacology* 1993; **39** (2):113-117
10. Olaifa JI, Eehum WO, Akingbohunge EA (2002). Insecticidal activity of some Nigerian Plants *Insect Science Application* **8**:221-224
11. Amosan CA , Okerie TG and Mike BC. The use of *Piper guineense* fruit oil as a protectant of dried fish against dermatis and maculates. *Global J. Pure Sci*. 2002; **8** (2): 197-201
12. Udo FVI, Lot TY, Braide VB. Effects of extracts of seed and leaf of *Piper guineense* on skeletal muscle activity in rat and frog. *Phytother Res*. 1999; **13** (2):106-10.
13. Zollo, PHA., Biyiti, L., Tchoumboungang F., Menut, C., Lamaty, G. and Bouchet, P. Aromatic plants of Tropical Central. Part XXXII. Chemical composition and antifungal activity of thirteen essential oils from aromatic plants of Cameroon. *Flav. Fragr. J*. 1998; **13**: 107-114.
14. Okigbo RN and Igwe DI. Antimicrobial effects of *Piper guineense* ‘Uziza’, and

- 279 *Phyllanthus amarus* 'Ebe-Benizo' on *Candida albicans* and *Streptococcus faecalis*.
280 *Acta Microbiologica et Immunologica Hungarica*. 2007; **54**(4): 353-366
- 281 15. Mbongue F G, Kamtchouing P, Essame O J, Yewah P M, Dimo T, Lontsi D. Effect
282 of the aqueous extract of dry fruits of *Piper guineense* on the reproductive function of
283 adult male rats. *Indian Journal Pharmacol*. 2005; **37**:30-32
- 284 16. Burstein M., Scholnick H.R., Morfin R. Rapid method for isolation of lipoproteins
285 from human serum by precipitation with polyanions. *J Lipid Res*, 1970; **11**:583- 595
- 286 17. Fletcher MJ. A colorimetric method for estimation of Serum triglycerides.
287 *Clin. Chem. Acta*,1968; **22**: 393-397
- 288 18. Richmond N. Determination of Cholesterol in Serum.
289 *Clin. Chem.* **19**: 1973; 1350-1396
- 290 19. Barlett GR. Colorimetric assay methods for free and phosphorylated glyceric acids.
291 *Journal Biol. Chem*, 1959; **234**: 469-471
- 292 20. Buege J and Aust S. Microsomal lipid peroxidation. In: Methods in Ezymology,
293 Biomembranes. Part C, volume 52. (Packer L and Fleischer S. eds)
294 Academic Press Inc. 1978; pp 302-310
- 295 21. Habig WH, Pabst MJ, Jakoby WB. Glutathione S-transferases:
296 The first enzymatic step in mercapturic acid formation. *J. Biol. Chem.* 1974;
297 **249**:7130-7139
- 298 .
- 299 22. Kakkar P, Das B and Viswanathan PN. A modified spectrophotometric assay of
300 Superoxide dismutase. *Indian J. Biochem. Biol.* 1984; **21**:130-132
- 301 23. Dacie, J.V. and Lewis, S.M. Practical Hematology.
302 Churchill Livingstone New York 1990; 34
- 303 24. Murray IK, Mayer PA, Granner OK and Rodwel V. Cholesterol biosynthesis,
304 Degradation and transport. In: Harper's Biochemistry. 2nd Edn. Prentice Hall
305 International Inc. 1990; pp 240-254
- 306 25. Boles, SJ, Bradley D, Force RA and Brown BR. The new dietary Fats in Health and
307 Diseases. *J. Am. Diabetic Assoc.* 1994; **97**: 280-286.
- 308 26. Vance IS. Metabolism of cholesterol. In: Biochemistry of Lipids and Lipoprotein and
309 Membranes. Amsterdam Elsevier Science Publishers, Amsterdam. 1990
- 310 27. Esterbauer H, Gebicki J, Puhl H and Jurgens G . The role of Lipid peroxidation and
311 antioxidants in oxidation modification of LDL. *Free Radic. Biol. Med.* 1992; **13**: 341-390
- 312 28. Steinberg D. Thematic review series The pathogenesis of atherosclerosis. An
313 interpretive history of cholesterol controversy, part v. The discovery of the statins and
314 the end of the controversy. *J. Lipid Res.* 2006; **47**: 1339-1351
- 315 29. Del RD, Stewart AJ and. Pellegrini N. A review of recent studies on
316 malondialdehyde as toxic molecule and biological marker of oxidative stress.
317 *Nutr. Metab Cardiovasc. Dis.* 2005; **15** (4): 316-328

- 318 30. Ferguson LR . Role of plant polyphenols in genomic stability.
319 *Mutat. Res.* 2001; **475**:89-111.
- 320 31. Ross JA, Kasum CM . Dietary flavonoids: bioavailability,
321 metabolic effects, and safety. *Annu. Rev. Nutr.* 2002; **22**:19–34.
- 322 32. Ren W, Qiao Z, Wang H, Zhu L, Zhang L. Flavonoids: promising
323 anticancer agents. *Med. Res. Rev.* 2003; **23**(4):519-534
- 324 33. Prestera T, Holtzclaw WD, Zhang Y, Talalay P. Chemical and molecular regulation
325 of enzymes that detoxify carcinogens. *Proc. Natl. Acad. Sci. USA* 1993; **90**:2965 - 2969.
- 326 34. Pantuck EJ, Pantuck CB, Anderson KE, Wattenberg LW, Conney AH, Kappas A.
327 Effect of Brussels sprouts and Cabbage on drug conjugation. *Clin. Pharmacol. Ther.*
328 1984; **35**: 161-169
- 329 35. Farombi EO. African Indigenous plants with chemotherapeutic potentials and
330 biotechnological approach to the production of bioactive prophylactic agents.
331 *Afr. J. Biotechnol* 2003; **2**(12):662-671
- 332 36. Adepoju GKA and Odubena OO. Effects of *Mucuna pruriens* on some hematological
333 and biochemical parameters in Albino rats. *Journal of Medicinal Plants Research.*
334 2009; **3**(2):73-76
335
336
337
338