

A REVIEW ON EFFECTS OF HEAVY METAL TOXICITY ON HEALTH OF AUTISTIC POPULATION

Abstract

Autism is a neurodevelopmental disorder that predominantly affects the younger generation. The etiology which contributes to the occurrence of autism is not well defined. However, apart from genetic factors, environmental factors such as heavy metal exposure have been controversial from the last decade. Contamination of several heavy metals was proposed to be responsible for oxidative stress production, mitochondria dysfunction and immune abnormalities which lead to characteristics of autism in children. Objective of review is to analyze the relationship between the most studied toxic heavy metals namely mercury, zinc, cadmium and zinc and autism. Based on the findings, heavy metal toxicity due to lead and mercury are strongly correlated with autism, whereas arsenic and cadmium show a weak correlation with autism based on the research articles throughout the years. In addition, lack of essential minerals in autistic children who were exposed to heavy metals has also precipitated the autistic disorder. However, high quality epidemiological studies with minimal biases should be conducted to support the correlation of heavy metal with autism.

Keywords: Autism, Heavy metals, toxicity

INTRODUCTION

Autistic spectrum disorder (ASD) is a neurodevelopmental disorder that predominantly established in early childhood and is identified by impairments in social interactions, marked abnormal communication skills as well as restricted, repetitive and stereotyped behavior. ^[1] According to Autism and Developmental Disabilities Monitoring Network (ADDM), among 14 sites in United States, the prevalence per 1000 children was around 6 from 2000 to 2002. However, the number of children affected by autism increased tremendously since 2004 and reached the peak value in 2008 in which the range of prevalence over 1000 children was from 4.8 to 21.2, whereas about 1 in 150 children affected by autism from 2000 to 2002 and almost 2 children suffered from autism disorder among 176 people. ^[1] Researchers have conducted a survey based on different number of reported sites. Boys were found to have higher risk

factor than girls. Centers for Disease Control and Prevention (CDC) had done a government survey on autism prevalence rate in 2009, it shows that 1 in 110 of 8 years old children among 14 communities in United State is affected by autism and similarly, boys carry four to five times higher risk than girls.^[1]




Figure 1: Prevalence of autism spectrum disorder from 2000 to 2012 in United States.^[1]

The pathophysiological etiology, which contributes to autism, is currently not well-defined, non-specific and controversial but genetic and environmental factors have been implicated. Autism is linked to genetic susceptibility such as fragile X syndrome which is a hereditary factor that lead to intelligence problem and tuberous sclerosis.

^[3]Environmental factor is also an inevitable cause of autism. Contamination of several heavy metals like mercury, lead, arsenic and cadmium are believed to be responsible for oxidative stress production, mitochondria dysfunction, immune abnormalities and lead to characteristics of autism in children.^[4] It is suspected that children with autism may have liver or renal function impairment, hence their bodies accumulate large amount of heavy metals and eventually, causing neurodegeneration.^[4]

Heavy metals are widespread environmental elements and acute or chronic exposure to these toxins will lead to developmental problems including endocrine disruption, behavioral disturbance and lower intelligence quotient (IQ).

^[5] Windham et al, found that the amount of airborne pollutant especially mercury is correlated to the increase risk in autism in an epidemiological study. ^[6] James et al explained that infants are at risk of metal toxicity due to low serum level of glutathione as toxic metals are commonly being removed by glutathione conjugation through bile excretion. In addition, children with predominantly all-milk food in everyday diet will indirectly decrease their heavy metal excretion abilities by a factor of three in an animal study. ^[7]

Heavy metal and the correlation with autism

The aim of this study is to explore the relationship between toxic heavy metal and autism among the children. Therefore, based on the analysis of several research articles, we have deduced that there were few toxic heavy metals that were identified in hair and nail sample of autistic children.

Researches showed that 18 children in Dallas, USA aged between 1 to 4 year old, we have identified that the common heavy toxic metal were mercury, chromium, tin, manganese, lead, arsenic, cadmium, cobalt and uranium. However, the most common heavy metal have been identified as lead which is having mean value of 0.63 where the minimum level should be 0.18 μ g of metal for every 1g of hair. Hair mineral analysis indicates the mineral composition accumulated over a long time span, which is proportional to the level of elements in the body.

Research conducted in Egypt, among 44 children aged 3 to 9 years, it were noted that hair concentration of aluminum, arsenic, cadmium, mercury, nickel, antimony and lead were elevated in autistic as compared to non-autistic children. The mean levels of hair toxic metal concentration have been mentioned in Figure 2.




Figure 2: Mean value of toxic heavy metal in autistic children in 44 children.

In this review article, we are mainly discussing about the toxic heavy metals that are commonly related to autism which are mercury, lead, cadmium, and arsenic.

Mercury

Mercury is a toxic heavy metal that is usually related to autism. According to the WHO factsheet titled 'Mercury and Health', even a small amount of mercury exposure may result in serious health problems. ^[8] This is supported by Charles Lee, who states that mercury exposure can result in serious health consequences as it damages the renal and nervous system besides interfering the brain development in unborn and very young children. ^[9]

To understand better about the relationship between mercury and health problems, mercury cycle should be revised. Various forms of mercury are present and these include the metallic mercury, organic mercury and inorganic mercury. Metallic mercury is also known as elemental mercury and it exists as liquid in room temperature and evaporates easily to form mercury vapours which are readily absorbed and transported to the brain. ^[10] Metallic mercury can accumulate in the brain as it passes through the blood brain barrier. ^[10] Metallic mercury is commonly used in amalgam, thermometer and switches.

Dental amalgam is a mercury-based filling which made up of 50% metallic mercury.^[11] While chewing, metallic mercury from the filling is released as mercury vapour and then inhaled. According to the US Food and Drug Administration, the use of mercury in dental amalgam may cause neurotoxic effects on the nervous system of developing foetus and children.^[12] This is in line with a prospective study conducted by Geier and colleagues in 2009, where children whose mother received more than six dental amalgams filling during pregnancy were 3.2 folds more likely to be diagnosed with autism.^[13]

A study carried out by Palkovica et al, also reported that dental amalgam fillings should be used cautiously in women during the childbearing age in order to avoid prenatal mercury exposure.^[14] This is due to the increased mercury level found in the cord blood as a result of greater number of maternal amalgam fillings. The adverse health effects of mercury from dental fillings are however not studied in this research. Due to the possible side effects of mercury to human, dental amalgam were even banned from being use in country like Norway (2008), Denmark (2008) and Swedish (2009).^[12]

In contrast, the American Dental Association (ADA) has considered the use of mercury in amalgam as safe and what's more, it does not leak from the amalgam.^[12] A systematic review conducted by Rasines G. in 2008 finds out that the mercury released from amalgam fillings are not associated with neurotoxicity in children owing to insignificant difference in neurobehavioral and neuropsychological score that were reported in children receiving dental amalgam fillings or composite fillings.^[15] Another more recent survey in 2009, conducted by Ye X and colleagues on 198 children who had amalgam filing and 205 children with no history of amalgam fillings, also shows no differences in renal function, neurobehavioural, neuropsychological or intelligence tests between the groups.^[16]

Combination of mercury with elements such as oxygen, chlorine or sulphur results in the formation of inorganic mercury which is also called the mercury salts.^[17] It is usually found in the cosmetic products and enters into the body through skin absorption.^[10] Unlike the metallic and organic mercury, inorganic mercury does not easily pass through the blood brain barrier and this explains why the neurological damage caused by inorganic mercury is unlikely to occur and less severe.^[10]

Organomercury is the most toxic mercury among the three types of mercury exposure where it is formed when mercury is combined with carbon.^[17] Methylmercury, ethylmercury and phenyl mercury are examples of organomercury. Similar to metallic mercury, methylmercury can pass through the blood brain barrier and its accumulation in the central nervous system leads to neurological disorder.^[10] Prenatal exposure of mothers to methylmercury also affects the foetus despite the function of placenta as a barrier to mercury. The condition is even worse, with the fact that neurotoxicity is more expressive in the developing brain as compared to the fully mature brain.^[10] Methylmercury poisoning is aroused from the contaminated fish and shellfish consumption as this mercury bioaccumulates in the ascending food chain of the marine life.^[10] Higher concentration of methylmercury is usually found in larger fishes that feed on other smaller fishes due to the longer life-span.^[18] Swordfish, sharks and king mackerels that usually contain a mercury level of greater than 1 part per million (ppm), are the highest mercury-containing fishes.^[18]

According to study by Shea and colleagues in 2004, declined in infants' and children performances in the neurobehavioural tests are found to be due to the prenatal exposure to methylmercury which can be easily passed to the new-born through the placenta and breast milk.^[18] Young children are more susceptible to mercury toxicity and their exposure to mercury in the first year of life may cause their brain to be more affected. This explained why these children are unable to perform in the attention, language skills, visual-spatial abilities and memory test.^[18] However, conflict of interest has been reported in study carried out by van Wijngaarden et al., which states that no association is found regarding the prenatal exposure of methylmercury with ASD phenotypic behaviours in the study of high fish consumption cohort.^[19]

Other type of organomercury is called as ethylmercury which is formed from the dissociation of thiomersal which functions as a preservative in vaccines.^[10] Its use as preservative was first introduced by Eli Lilly in 1931.^[20] In the United State (US), 12 times increase in autism cases were reported, after three additional thiomersal-containing vaccines were introduced to newborns in the early 1990s.^[20] This is much higher as compared to country like Germany and Denmark which has reduced the use of thiomersal in vaccines. Though autism cases are much lower in these countries, increase in autism is reported despite the removal of thiomersal-containing vaccines since 1992.^[10]

A cross sectional study conducted by Gallagher and Goodman in 2010, male newborns who receive thiomersal-containing hepatitis B vaccination in the first month of life are 3 times more likely to be diagnosed with autism.^[21] This association is also reported in study done by Geier and Geier in 2004, where children receiving thiomersal-containing diphtheria-tetanus-acellular pertussis (DTaP) vaccines are associated with a greater risk of neurodevelopmental disorder adverse events as compared to children receiving thiomersal-free DTaP vaccines.^[22] Autism, mental retardation, speech disorder, thinking abnormalities, infantile spasms and ataxia are the neurodevelopmental disorder adverse events studied.^[23] Geier and Geier further prove the validity of their study in 2007, where ASD infants who had normal development in their first year of life, undergo regression into autistic disorder after a significant exposure to medicinal sources of mercury.^[24]

Debate has been going on among researchers on the correlation between thiomersal-exposure and autism in children. Besides the supporting evidence proposed, there are also researches that found no correlation between thiomersal-exposure and autism in children. A study in Canada, by Fombonne and colleagues in 2006, found that the increase in pervasive developmental disorder (PDDs) is not related to thiomersal exposure as the incidence of PDDs are greater in the thiomersal-free birth groups rather than the thiomersal-exposed groups.^[25] Another study done by Price and colleagues in 2010, also found no evidence to conclude that increasing ethylmercury exposure through thiomersal-containing injection was in line with a greater risk of autisms.^[26]

A critical review proposed by Parker and colleagues in 2005, further showed invalid correlation between thiomersal-containing vaccines with ASD. Parker and colleagues reviewed a total of 12 articles published between the years of 1966 to 2004 before they came to the conclusion that the incidence of ASD is not associated with the use of thiomersal-containing vaccines.^[27] Ethylmercury is eliminated out of the body efficiently, owing to its 7 days, short half-life. The very low concentration of ethylmercury used in vaccines also cause harm results from thiomersal-containing vaccines to be less likely.^[28]

Autistic children are associated with reduced cysteine and glutathione synthesis which are crucial for natural mercury detoxification.^[29] This is proven in study conducted by James and colleagues in 2004 where 19% lower plasma cysteine and 46% lower plasma glutathione were identified in autistic children.^[29] As the ability of autistic

children in mercury detoxification and excretion were impaired, higher mercury concentrations can be found in tissues of the nervous system. The longer half-life of mercury in autistic children also increases their susceptibility to mercury exposure. As there is a delay in mercury detoxification, phospholipid methylation process that is responsible for normal attention is impaired in autistic children^[20]

The impaired ability of autistic children in mercury excretion is proven in a study conducted by Holmes and colleagues in 2003. This study was conducted to measure the mercury level in the first baby haircuts of autistic children. A significant reduction in mercury levels, which is about 8-folds lower, is found in autistic children as compared to normal children.^[30] This is in accordance to the fact that autistic children have impaired ability in excreting mercury. A more recent study conducted by Adams et al in 2009 also reported that children with lower mercury level in the first babies' haircut had 2.5-fold higher chance of getting ASD when compared to those with higher mercury level.^[31] Hair mercury analysis can be used to test for autism as children with more severe autism have a lower mercury level in their hair.^[30]

Lead

One of the most common causes of autism is childhood lead poisoning.^[32] Harmful effects on the development of widespread brain areas including those implicated in mental, communication, and social functioning have been identified in lead poisoning in the childhood.^[32] As early as 1976, a study conducted by Cohen et al has noticed elevated blood lead levels in children with autism.^[33] The toxic effects of lead are usually attributed to multiple mechanisms and one of the key factors is the ability of lead to mimic calcium action in the body.^[32] Lead is capable of competing with or substituting for calcium in activating both protein kinase C (PKC) and calmodulin. It was reported that inhibition of brain microvascular formation and function can happen with lead concentrations that are effective in activating PKC. This will impair the development of nervous system in children who are exposed to considered amount of lead. Besides that, improper activation of calmodulin may alter the activity of secondary messenger pathways, which play crucial roles long-term potentiation in synapses, spatial learning, and memory processes.^[34]

Several studies have been made to determine the correlations between lead exposure and autism. One study has been conducted to measure the levels of 10 toxic metals in hair samples of autistic children as compared to non-autistic children through hair mineral analysis, and correlate the levels of these toxic metals to severity of autism. A total of 44 autistic test group (aged 3 to 9 years) were compared to a similar control group, consisting of a total of 106 non-autistic children. ^[35] As a result, there was a statistically significant positive correlation between lead and verbal communication ($p=0.020$), and between lead and general impression ($p=0.008$). ^[35] Brockel et al, claimed that children with autism are commonly found to have problems with communication and learning skills. They are typically associated with symptoms, such as retarded childhood development, decreased cognitive ability, learning and behavioral disabilities, attention deficit hyperactivity disorder (ADHD), impetuosity, and responding incoherently. ^[35] Researchers found out that there is sufficient evidence to state that a biomarker related to heavy metal toxicity especially lead has relationship with the severity of autism where elevations in urinary porphyrins which have relationship with lead were significantly associated with Childhood Autism Rating Scales. ^[35] Present investigation shows that increasing order of toxic metals lead concentration in the hair and nail samples and their correlation with degree of severity. In several cases, they have noted temporal association between elevated blood lead levels and the emergence of autistic symptoms. ^[32]

Evidence has been stated by Abdullah et al, in his article on Toxic trace elements in the hair of children of autism that children with autism had significantly ($p < 0.001$) higher in hair concentration level of lead compared to other heavy metals such as antimony, arsenic, beryllium, cadmium and aluminium. ^[36] Wasserman et al, have stated that inorganic lead poisoning have been implicated with autism and reported an association between increased exposure to the inorganic lead exposure and teacher parents' ratings of children behavior especially in distractibility, impulsive behavior, poor concentration and non -persistence.

Cadmium

Commonly studied heavy metal in cases of autism is cadmium. Cadmium is one of the metals that known to affect the development of infant brain particularly during the first three months of growth. Therefore, several studies have been conducted to determine the relationship between heavy metals and autism.

Children with ASD might have inabilities in eliminating the toxins from the body. Thus, toxins build up in body and result in increased free radical body which eventually affects the nervous system. However, detoxification and excretion of heavy metals depends on their conjugation to glutathione, whose levels are significantly reduced in autistic subjects.^[37] Low level of glutathione can be part of the reason that reduces the ability to remove toxic metals. This is further supported by the urine analysis of Yorbik et al which compare the urine levels of heavy metals such as cadmium, chromium and lead in 30 autistic children with 20 healthy subjects using atomic absorption spectrometry. The results have shown that the cadmium level in urine of autistic children is significantly decreased as compared to that of healthy subjects.^[38]

However, there are a few study indicates there is no correlation between autism and heavy metal. A study has been conducted to understand the toxicological status of children with autism compared to neurotypical children where the levels of toxic metals are measured in total blood, red blood cell and urine. Cadmium is detectable for whole blood and urine but hardly spotted in RBC as the level was below the detection limit which is 0.001mcg/g. Children with autism has significantly lower level of cadmium in whole blood except urinary level is being the most sensitive. However, compared to other metals, cadmium levels are considered low in blood and urine because cadmium is primarily excreted in bile. Thus, this suggests that there is not much difference of cadmium body burden in children with autism and neurotypical children.^[39] In a case study by Abdullah et al, in 40 autistic children who were tested for heavy metal toxicity in their hair sample, the p value were not significant, hence does not contribute to autism. Hence, more studies need to be done to confirm the relationship between cadmium and autism in children.^[36]

Arsenic

It has been believed that autism spectrum disorders are in part caused by environmental exposure to heavy metals such as arsenic. Arsenic is a sulfhydryl-reactive metal found to be toxic to the nervous system even at low exposure levels. The more toxic form, inorganic arsenic, is found to be carcinogenic and affects neurodevelopment. Inorganic arsenic exists in soil and rocks. It is highly water soluble thus readily mobile in the environment. Organic arsenic, though less toxic compared to the inorganic counterpart, can be harmful to neurodevelopment too.^[40] Human can be exposed to inorganic arsenic in many ways including:

- Arsenate which widely contaminates drinking water source
- Consuming crops grown in arsenic-contaminated soil
- Bioaccumulation in the human body from high frequency consumption of seafood (eg shellfish) over a long period.^[40]

Arsenic exposure in human is assessed using the blood concentrations. Concentrations of the metal in urine samples indicate more recent exposures (for past several days). Arsenic levels in hair or fingernails would represent exposure from the past several months.^[40] As mentioned earlier, the exact causes of autism spectrum disorders have remained obscure; among them one postulation suggested is the epigenetics theory. Autism is among the monogenetic brain disorders associated with DNA methylation and histone modification defects.^[41] The epigenetic dysregulation of the gene expression is considered to be one of the pathophysiological factors contributing to neurodevelopmental disorders such as autism.^[42] Arsenic has been viewed to induce epigenetic alterations of the gene expression. For example, arsenite exposure is reported to induce DNA hypo and hypermethylation.^[43] Studies have been conducted to investigate the possible correlation between arsenic exposure and autistic spectrum disorder. Rahbar et al in a Jamaican age and sex-matched case control study did not find a significant association between ASDs and total blood arsenic concentrations. This was the first study to take into account the confounding factor diet such as consumption of fruits and vegetables, drinking water source and seafood consumption.^[40] In another study, Fido and Al-saad used 40 older children with autism as subjects. They found that there was no difference of arsenic levels in haircuts in between the group of autistic children and the control group of 40 normal controls.^[44]

Geier et al indicated that there is no significant association between arsenic and ASD severity. In the study, autistic subjects of 1 to 6 years old had their autism severity measured by the Childhood Autism Rating Scale (CARS) score. The arsenic concentrations in their haircuts were tested. The strength of the study was that it included only children with moderate to severe autism. However the sample size was small, though it was statistically adequate enough to establish an association between heavy metal concentrations and autism severity.^[45] However, some studies obtained conflicting results. The study done by Obrenovich et al indicated significantly higher levels of arsenic in the hair of children with an ASD compared to the controls ($p < 0.0001$). The source of arsenic was not

known.^[46] Eleonor et al found raised hair concentrations of arsenic in autistic children compared to those non-autistic.

^[47] Al-ayadhi evaluated 77 Riyadh autistic children and significantly higher levels of arsenic were found in the hair of the autistic group. ^[48]It would be interesting to source more studies investigating exposure of arsenic in the prenatal/fetal period. The studies looked at above only measure levels of arsenic in postnatal period (children). There have been evidences that environmental chemicals can cause insults *in utero*, leading to dysregulation of the fetal epigenome which would then possibly exert an effect on the neurodevelopment of the child. ^[49] In fact, arsenic was shown to be able to easily cross the placenta. ^[40]

DISCUSSION

The exposure of environmental toxin and the influence for the neurodevelopmental disorders such as autism have been studied in many cases. In a PUBMED search, 58 empirical reports of autism and heavy metal toxicity were reported where 43 suggest the correlation and 13 reports deny the correlation between heavy metal toxicity and autism. However, the tendencies for null results were not reported and it cannot be said that there is no evidence for a link between heavy metal toxicity and autism. ^[50]However, there are several controversies and conflict of interest in the correlation between toxic heavy metal and autism. A very recent study in 2014, by Rossignol et al, suggested that controversies in the research of heavy metal associated with autism may exist due to the significant limitations in the studies, retrospective design, publication biases, limited sample size, and the utilization of non-standard tool to diagnose autism. ^[51]

In several cases such as the thiomersal in vaccines biases in publication causes the unclear relationship between the heavy metal toxin and autism. Wakefield et al, published in the medical journal of The Lancet that autism were associated with administration of vaccines. However, there were conflict of interest and it was found that standard ethical practices were violated when the research is carried out due to false findings because it was identified that the vaccines used did not contain thiomersal in the vaccine. In a separate scenario, it has been studied that it has pharmaceutical companies offer the significant monetary support to the researchers whose findings do not support link between toxic heavy metals in vaccine. Therefore, the findings in the research are significantly affected due to the

professional ethics of the researcher and pharmaceutical companies.^[50] Besides that, other factor that precipitates autism with heavy metals is polymorphism in genes that increases the susceptibility of autism.^[50] In addition, toxic metals affects the absorption of the trace element, interacts with the essential elements and affects the threshold level besides the toxicity effect which may contribute to the neurodevelopmental condition such as autism.^[50] For instance, toxic metals namely cadmium, lead and mercury interacts with nutritionally essential metals.^[50] Lead may interact with calcium and causes the impairment in the cognitive development in children.^[50] Besides that, it has been observed that children's presented with autistic disorder have very low level of essential metals such as calcium, iron, iodine, magnesium, manganese, zinc, and selenium. However, the level of copper have exceeded the reference range and it is confirmed in a finding by Faber et al, who stated that the incidence of zinc deficiency and copper intoxication is high in children with Autism Spectrum Disorder.^[50] This is further supported by Al- Ayadhi who stated that compared to the hair sample of normal children, hair sample of children with Autistic Spectrum Disorder shows significant lower concentration of essential trace elements such as calcium, manganese, iron and chromium.^[52]

It has been identified that selenium protects from mercury and methylmercury toxicity. It has been established that mercury toxicity is prominent in the children of autism. This is further supported by Jory and Woody et al in their study where lower level of selenium is identified in autistic children.^[52] Apparently, the ability to adapt to the environment is affected by low level of selenium concentration in the children. Besides that, Fagala and Wigg mentioned that selenium is a vital component of glutathione peroxidase which appears to protect from toxicity from metal such as lead, mercury and cadmium.^[52] Therefore, we can identify that, autistic children with selenium deficiency are highly susceptible to the heavy metal toxicity which causes them to become regressive, unable to concentrate, and display some learning disorders.^[52] In addition, toxic heavy metal exposure will affect the absorption and utilization of the nutrient elements in our body. For example, lead substitutes zinc on hem enzymes and cadmium causes replacement of zinc n metallothionin.^[53] Hence, low level of zinc have been associated to fear, verbal communication impairment, and nervousness which may enhance the effect of lead which supports the positive correlation between lead and verbal communication ($p=0.020$).^[54] Therefore, toxic heavy metal toxicity in addition to nutritional inadequacies potentially heightens the toxicity of heavy metals. Nutritive elements such as trace metals are deficient among the autistic children which further contribute to the autism spectrum disorder.^[55]

CONCLUSION

Heavy metal toxicity due to lead and mercury are strongly correlated to autism, however arsenic and cadmium shows a weak correlation on autism based on the research articles throughout the year. A strong correlation of mercury and few other toxins have been shown in the pathogenesis of autism, similarly the lack of excess essential minerals have been identified to precipitate the incidence of autism spectrum disorder among children. Increase environmental exposure at children developmental stage plays a significant role in the etiology of autistic disorder. However, further research is needed on a wider scale particularly in heavy metals such as cadmium and arsenic because they have significant role in the autistic spectrum disorder. Therefore, high quality epidemiological studies with minimal biases should be conduct to support the correlation of heavy metal with autism.

REFERENCES

1. Jon Baio. Prevalence of autism spectrum disorders. Autism and Developmental Disabilities Monitoring Network Surveillance. Report number: 61(SS03); 1-19, 2012.
2. What is autism spectrum disorder : National Institute of Mental Health; 2011
3. Parris M. Kidd, Autism, An Extreme Challenge to Integrative Medicine, Part 1: The Knowledge Base, *Alternative Medicine Review*, 7(4), 2002
4. Mark E. Obrenovich, Raymond J. Shamberger, Derrick Lonsdale. Altered heavy metals and transketolase found in autistic spectrum disorder. *Biological Trace Element Research*. 2011 July 14; 144: 475–486.
5. Paula E. Goines, Paul Ashwood. Cytokine dysregulation in autism spectrum disorders (ASD): possible role of the environment. *Neurotoxicology and Teratology*. 2012 August 17; 36: 67-81.
6. G. C. Windham, L. Zhang, R. Gunier, L. A. Croen, J. K. Grether, "Autism spectrum disorders in relation to distribution of hazardous air pollutants in the San FranciscoBay area," *Environmental Health Perspectives*, vol. 114, no. pp. 1438–1444, 2006.
7. James B Adams, Matthew Baral, Elizabeth Geis, Jessica Mitchell, Julie Ingram, Andrea Hensley, Irene Zappia, Sanford Newmark, Eva Gehn, Robert A Rubin, Ken Mitchell, Jeff Bradstreet and Jane El-Dahr. 'Safety and efficacy of oral DMSA therapy for children with autism spectrum disorders: Part A - Medical results', *BMC Clinical Pharmacology*. 2009 October 23. 9:16.
8. Preventing Disease Through Healthy Environments: Action is needed on chemicals of major public health concern; World Health Organization; 2010
9. Mercury Poisoning Linked to Skin Products. 2012 March 6
10. Kwok KN, Chan CH, Soo MT, ShingYI. Low-level chronic mercury exposure in children and adolescence: Meta-analysis. *Pediatrics International*. 2007; 49:80-87
11. Agency for Toxic Substances and Disease Registry (ATSDR). Public Health Statement Mercury.

12. Oken E, Bellinger DC. Fish Consumption, methylmercury and child neurodevelopment. NIH Public Access Author Manuscript. 2008 April; 20(2):178-183.
13. Edlich RF, Rhoads SK, Cantrell HS, Sabrina M. Azavedo, Anthony T. Newkirk. Banning Mercury Almagam. 9-16.
14. Palkovicova L, Ursinyova M, Masanova V, Yu Zw, Irva Hertz-picciotto. Maternal amalgam dental fillings as the source of mercury exposure in developing fetus and newborn. Journal of Exposure Science and Environment Epidemiology. 2008; 18:326-331.
15. Geier M, Kern JK, King PG, Sykes L, David A. Geier. Mercury Induced Autism. 2014; 1411-1432.
16. Rasines G, Mercury released from amalgam restorations does not give rise to toxic effects on the nervous system of children. Evidence-Based Dentistry. 2009; 9:25-27.
17. Ye X, Qian H, Xu P, Lin Z, Matthew P. Longnecker, Hua F. Nephrotoxicity, Neurotoxicity, and Mercury Exposure among Children with and without Dental Amalgam Fillings. NIH Public Access Author Manuscript. 2009 July; 212(4):1-15.
18. Griesbauer L. Methylmercury Contamination in Fish and Shellfish. 2007.
19. Van Wijngaarden E, Davidson PW, Smith TH, Evans K, Yost K, Love T, Thurston SW, Watson GE, Zareba G, Burns CM, Shamlaye CF, Myers GJ. Autism spectrum disorder phenotypes and prenatal exposure to methylmercury. Epidemiology. 2013 Sep; 24(5):651-9.
20. Mutter J, Naumann J, Schneider R, Walach H, Haley B. Mercury and autism: Accelerating Evidence? Neuroendocrinology Letters. 2005 October 5; 26(5):439-446.
21. Gallagher CM, Goodman MS. Hepatitis b vaccination of male neonates and autism diagnosis, NHIS 1997–2002. Journal of Toxicology and Environmental Health, Part A. 2010; 73:1665-1677.
22. Geier DA, Geier MR. Neurodevelopmental Disorders Following Thiomersal-Containing Immunization: A Follow-Up Analysis. International Journal of Toxicology. 2004; 23:369-376.
23. Geier DA, Geier MR. An evaluation of the effects of thiomersal on neurodevelopmental disorders reported following DTP and Hib vaccines in comparison to DTPH vaccine in the United States. Journal of Toxicology and Environmental Health, Part A. 2006; 69:1481-1495.
24. Geier DA, Geier MR. A Prospective Study of Mercury Toxicity Biomarkers in Autistic Spectrum Disorders. Journal of Toxicology and Environmental Health, Part A. 2007; 70:1723-1730.
25. Fombonne E, Zakarian R, Bennett A, Meng L, Mclean-heywood D. Pervasive Developmental Disorders in Montreal, Quebec, Canada: Prevalence and Links With Immunizations. PEDIATRICS. 2006 July; 118(1):139-150.
26. Price CS, Thompson WW, Goodson B, et al. Prenatal and Infant Exposure to Thiomersal from Vaccines and Immunoglobulins and Risk of Autism. PEDIATRICS. 2010 October; 126(4):656-664.
27. Parker SK, Schwartz B, Todd J, Pickering Lk. Thiomersal-Containing Vaccines and Autistic Spectrum Disorder: A Critical Review of Published Original Data. PEDIATRICS. 2004 Sept; 114(3):793-804.
28. Tomljenovic L, Dorea JS, Shaw CA. Commentary: A Link Between mercury exposure, Autism Spectrum Disorder, and other neurodevelopmental Disorders? Implications for thiomersal-containing Vaccines. Journal on Developmental Disabilities. 2012 Nov; 18(1):34-42.
29. Zhang J, Wheeler JJ. Mercury and Autism: A Review. Education and Training in Autism and Developmental Disabilities. 2010; 45(1):107-115

30. Adams JB, Holloway CE, George F, Quig D. Analyses of Toxic Metals and Essential Minerals in the Hair of Arizona Children with Autism and Associated Conditions, and Their Mothers. *Biological Trace Element Research*. 2006; 110:193-209.
31. Adams JB, Romdalvik G, Levine KE, Hu LW. Mercury in first cut baby hair of children with autism vs. typically-developing children. *Environ Toxicol Chem*. 2008; 90: 739-53.
32. Lidsky TI, Schneider JS. Lead neurotoxicity in children: basic mechanisms and clinical correlates. *Brain*. 2003;126:5-19
33. Phasing Out Leaded Gasoline. Blacksmith Institute.2014 Feb.
34. Toxicological Profile for Lead. U.S. Agency for Toxic Substances and Disease Registry. 2007 August. pg. 214-216.
35. Blaurock-Busch E et al. Toxic Metals and Essential Elements in Hair and Severity of Symptoms among Children with Autism. *Maedica, A Journal of Clinical Medicine*.2012; 7(1): 38-48.
36. Abdullah Fido et al, Toxic trace elements in the hair of children with autism, Sage Publication, The National Autistic Centre 2005.
37. Yahya M. Al-Farsi, Mostafa I. Waly. Levels of Heavy Metals and Essential Minerals in Hair Samples of Children with Autism in Oman: a Case–Control Study. *Biological Trace Element Research*. 2012 November 28; 51:181–186.
38. Yorbik O, KurtI, HasimiA, Ozturk O. Chromium, Cadmium, and Lead Levels in Urine of Children with Autism and Typically Developing Controls. *Biological Trace Element Research*. 2009 August 18; 135:10-15.
39. James B. Adams, TapanAudhya, Sharon McDonough. Toxicological Status of Children with Autism vs. Neurotypical Children and the Association with Autism Severity. *Biological Trace Element Research*. 2012 August 3.
40. Rahbar MH, Samms-Vaughan M, Ardjomand-Hessabi M, Loveland KA, Dickerson AS, Chen Z et al. The role of drinking water sources, consumption of vegetables and seafood in relation to blood arsenic concentrations of Jamaican children with and without Autism Spectrum Disorders. *Sci Total Environ*. 2012 September 1; 433C: 362–370.
41. Arita A, Costa M. Epigenetics in metal carcinogenesis: Nickel, Arsenic, Chromium and Cadmium. *Metallomics*.2009; 1: 222–228.
42. Jakovcevski M, Akbarian S. Epigenetic mechanisms in neurodevelopmental and neurodegenerative disease. *Nat Med*. 2012 August; 18(8): 1194–1204.
43. Baccarelli A, Bollati V. Epigenetics and environmental chemicals. *CurrOpinPediatr*. 2009 April; 21(2): 243–251.
44. Fido A, Al-Saad S. Toxic trace elements in the hair of children with autism. *SAGE Publications and the National Autistic Society*. 2005; 9(3): 290-298.
45. Geier DA, Kern JK, King PG, Sykes LK, Geier MR. Hair Toxic Metal Concentrations and Autism Spectrum Disorder Severity in Young Children. *Int. J. Environ. Res. Public Health*.2012; 9: 4486-4497.
46. Obrenovich ME, SHamberger RJ, Lonsdale D. Altered Heavy Metals and Transketolase Found in Autistic Spectrum Disorder. *Biol Trace Elem Res*. 2011 June.
47. EleonorBlaurock-busch; Omnia R. AMIN; Hani H. DESSOKI; Thanaarabah.Toxic Metals and Essential Elements in Hair and Severity of Symptoms among Children with Autism.*Maedica A Journal of Clinical Medicine*. 2012; 7(1): 38-48.

48. Al-Ayadhi LY. Heavy metals and trace elements in hair samples of autistic children in central Saudi Arabia. *Neurosciences (Riyadh)*. 2005 Jul; 10(3):213-8.
49. Perera F, Herbstman J. Prenatal Environment Exposures, Epigenetics, and Disease. *Reprod Toxicol*. 2011 April; 31(3): 363–373.
50. Mary Catherine De soto, Sorting out the spinning of autism: Heavy metals and the question of incidence, *Acta Neurobiol Exp* 2010, 70: 165-176
51. Rossignol D A, et al, Environmental toxicants and autism spectrum disorders: a systematic review, *Translational Psychiatry*. 2014, 4 e360
52. Goyer RA , Toxic and Essential Metal Interactions. *Ann Rev of Nutr*. 1997; 17:37-50.
53. AL-Ayadhi L, Heavy Metals and Trace Elements in Hair Samples of Autistic and Normal Children in central Saudi Arabia. *Neurosci*. 2005; 10:213-218.
54. Black MM, Zinc Deficiency and child development. *Am J Clin Nutrition*. 1998; 68:464-469.
55. Jory J, Woody R, Red-Cell Trace Minerals in Children with Autism McGinnis. *Am J Biochem Biotech* 2008; 4:101-104.