

Original Research Article

Inhibitory Activity of Ethanolic Extract of Horseradish (*Moringa oleifera*) (Lam.) on Oxidative Stress in some organs of Dexamethazone Induced Hypertensive Wistar Albino Rats.

ABSTRACT

Local communities, throughout the world over prevent and treat indigenous diseases using local herbal

remedies. *Moringa oleifera* (horseradish) is one of the many plant species used in herbal preparations in West Africa

for treating numerous human ailments such as malnutrition, cardiovascular diseases and uterine conditions among others. Despite its wide use by communities around the world, there is inadequate scientific information available on the actual pharmacological effects of ethanolic extract of *Moringa oleifera* leaves on the heart and kidney dysfunction. The aim of this study ~~however~~ was to determine the inhibitory activity of ethanolic extract of horseradish on oxidative stress in the heart and kidney of dexamethazone induced hypertensive wistar albino rats. Air dried *Moringa oleifera* leaves were pulverized and an ethanolic crude extract was prepared. The

animals were grouped into four groups of five animals each. Group A animals served as control and fed

with only water and feed ~~only~~, while Groups B and C animals were administered with 0.5mL of 40% ethanolic extract of *Moringa oleifera* leaf and 0.5 ml of 1.67 mg/kg body weight dexamethasone, respectively. Group D animals were treated with 0.5 ml of 1.67 mg/kg body

weight dexamethasone and 0.5mL of 40% ethanolic extract of *Moringa oleifera* leaf intermittently for ~~twenty-one~~21 days. The results obtained from the study showed that the ethanolic

extract of *Moringa oleifera* leaf possessed the ability to inhibit and reverse the dexamethasone

mediated tissues oxidative damage of both organs as seen in cholesterol, reduced glutathione and malondialdehyde concentrations. The same protective trend was also observed in activities of alkaline phosphatase, aspartate transaminase, alanine transaminase and catalase enzymes in both organs of the hypertensive rats. Hence, ethanolic extract of *Moringa oleifera* leaf reduced the extent of antioxidant loss and restoration of organ dysfunction caused by dexamethasone in the

28 rats.

29 Keyword: Moringa oleifera, oxidative stress, dexamethazone, chemoprevention, antioxidant.

Introduction

Herbal medicine is the oldest form of health care known to mankind and most cultures have long folk medicine histories that include the use of plants. The WHO recognizes herbal medicines as a valuable and readily available resource for Primary Health Care, and it has endorsed their safe and effective use (Tilburt and Kaptchuk, 2008). It, however, recommends that many herbal remedies still need to be studied scientifically while recognizing the experience obtained from their prolonged safe use over the years in the treatment of various conditions in both humans and animals, such as cardiovascular diseases (CVDs) and obstetric conditions.

In recent decades, cardiovascular diseases have emerged as major causes of morbidity and mortality in many countries, with hypertension being the commonest (Akinkugbe, 1990). Hypertension is generally defined as an abnormal persistently raised blood pressure greater than 140/90 mmHg (Massie, 2004) and not due to physical exercise. It is estimated that a sixth of the world population is afflicted by sustained high blood pressure (Kearney et al., 2005). Hypertension is a chronic disease and its management requires prolonged treatment, which is very expensive and often not accessible to poor individuals; hence people resort to the use of herbal medicines.

Obstetric conditions are also a common problem especially in developing countries and more so in the rural areas. Obstetric conditions are those maternal conditions that are as a result of a mother being pregnant or arise in the process of giving birth. These conditions can be direct or indirect (Johanna et al., 2003). Direct conditions occur before or within 6 weeks of delivery (such as haemorrhage, abortion, eclampsia and sepsis), while indirect conditions include anaemia, malaria, hepatitis and cardiovascular disease. There are also psychological stressors like puerperal psychoses, post-partum depression (baby blues), suicide, and strong fear of pregnancy and childbirth resulting from obstetric complications, interventions or cultural practices (AbouZahr, 2003). The management of these conditions often involves the use of non-pharmaceutical approaches like exercise and nutrition, but also to a large extent allopathic medicine. Despite their proven efficacy, allopathic medicines are more expensive and are also presumed to be associated with a lot of side effects, and they are moreover not readily accessible to the majority of the people who need them.

On the other hand, herbal remedies are seen as less expensive and less toxic (Hughes and Bradly, 2004). In Nigeria, rural people often have to travel long distances to reach the nearest

health centre, and here the ‘modern/western’ type of medicine is often out of stock or may be unaffordable. These hardships force people to resort to their local remedies which are plentiful, available, and affordable, and are already known to be efficacious in managing certain illnesses. People are thus increasingly willing to manage their medical needs by using complementary and alternative medicines like herbs (Hughes and Bradly, 2004). Among the medicinal plants that are commonly used in the management of various conditions is *Moringa oleifera*. This herb has been reported in the management of cardiovascular disease such as hypertension (Nath et al., 1992). Substances in *Moringa oleifera*, which are similar to those derived from other herbs (for example glycosides), have become a basis for some of the commercial medications used today for the treatment of heart disease, high blood pressure, pain, obstetric, and other medical problems such as diabetes, cancer, and AIDS (Peltzer et al., 2008). Traditionally women use herbal preparations for the process of labour. Herbs are also prescribed by traditional birth attendants and healers for various reasons including increasing chances of having twins, correcting a mal-positioned fetus, anaemia in pregnancy among others (Azriani et al., 2007). In Nigeria the practice of using herbs has been passed on for generations and it is used not only by rural but also by many ‘modern’ women who continue using the herbal adjuncts in addition to their regular antenatal and postnatal prescriptions. *Moringa oleifera* has been reported to be used by women for contraception and as an abortifacient (Nath et al., 1992). Herbs such as *Moringa oleifera* are reportedly used for treatment of painful uterus, inducing uterine contractions, management of retained placenta and post partum bleeding among other conditions and have thus been found useful in obstetric care (Kamatenesi-Mugisha, 2002).

Materials and Methods

Sample collection and preparation

Fresh leaves of *Moringa oleifera* were obtained from a the Federal University of Technology

satellite Estate in Akure, Ondo State, Nigeria. The leaf identification was carried out in the Department of Plant Science, Ekiti State University, Ado Ekiti, Nigeria by plant taxonomist.

Fresh leaves of *Moringa oleifera* were collected and air dried under shade and ground into powder with a Marlex Excella laboratory blender. The An ethanolic extract of *Moringa oleifera*

leaves was made. 200g Two Hundred grams of the powder was weighed and mixed in 500 mL of ethanol in a coked

bottle, shaken and allowed to stand overnight with constant shaking on a mechanical shaker.

91 Experimental protocol

92 The study was performed on ~~twenty~~ 20 male wistar albino rats housed in ventilated cages in the
 93 Animal House of Biochemistry Department, Ekiti State University, Ado-Ekiti, Nigeria. They
 94 were acclimatized for two weeks before administration of the drugs. Animals were divided into
 95 four groups of five rats each. Group A served as the positive control and received distilled water
 96 and rat feed. Group B animal were administered with 0.5 mL of 40% ethanolic extract of
 97 Moringa olifera. Group C animal served as the negative control and were orally administered
 98 with 0.5mL single dose of dexamethasone (1.67 mg/kg body weight) while Group D animals
 99 were orally given 0.5mL dexamethasone (1.67 mg/kg body weight) followed by 0.5 mL of 40%
 100 ethanolic extract of Moringa oleifera. Animals were kept at optimum temperature with a 12 hour
 101 light/dark cycle, and given rat feed and water. The period of induction and treatment lasted for
 102 three weeks.

103 Preparation of organ homogenate and plasma collection

104 After three weeks of induction and treatment of the rats, they were sacrificed anesthetically using
 105 chloroform. Blood was collected using syringes and needles from the heart into anti-coagulant
 106 bottles. The organs (kidney and heart) of interest were also collected which were homogenized in
 107 6.7mM potassium phosphate buffer and the supernatants were collected after spinning at 10,000
 108 rpm in a centrifuge.

109 Biochemical Assay

110 Standard Randox kits were used to determine Cholesterol, Total protein, Alkaline
 111 Phosphatase (ALP), Aspartate Transaminase (AST) and Alkaline Transaminase (ALT).

112 ALT

113
$$\alpha\text{-oxoglutarate} + \text{L-alanine} \xrightarrow{\text{GPT}} \text{L-glutamate} + \text{pyruvate}$$

114 Alanine Aminotransferase was measured by monitoring the concentration of pyruvate
 115 hydrazone formed with 2, 4-dinitrophenylhydrazine.

116 AST

117
$$\alpha\text{-oxoglutarate} + \text{L-aspartate} \xrightarrow{\text{GOT}} \text{L-glutamate} + \text{oxaloacetate}$$

118 AST was measured by monitoring the concentration of oxaloacetate hydrazone formed
 119 with 2, 4-dinitrophenylhydrazine.

120 CHOLESTEROL

121 The cholesterol was determined after enzymatic hydrolysis and oxidation. The indicator
 122 quinoneimine is formed from hydrogen peroxide and 4-aminoantipyrine in the presence of
 123 phenol and peroxidase.

124

125 Cholesterol esterase

126
$$\text{Cholesterol ester} + \text{H}_2\text{O} \rightarrow \text{Cholesterol} + \text{Fatty acids}$$

Cholesterol oxidase



Peroxidase



Determination of plasma Malondialdehyde (MDA)

Lipid peroxidation was determined by measuring the formation of thiobarbituric acid reactive substances (TBARS) according to the method of Varshney and Kale (1990). An aliquot of 0.4mL of the plasma or other organ homogenates was mixed with 1.6mL of Tris-KCl buffer to which 0.5mL of 30% trichloroacetic acid (TCA) was added. Then 0.5mL of 0.75% TBA was added and placed in a water bath for 45 minutes at 80°C. This was then cooled on ice and centrifuged at 3000g. The clear supernatant was collected and absorbance measured against a reference blank of distilled water at 532nm. The MDA level was calculated according to the method of Adam-Vizi and Seregi (1982). Lipid peroxidation in units/mg protein or gram tissue was computed with a molar extinction coefficient of $1.56 \times 10^5 \text{ M}^{-1}\text{Cm}^{-1}$.

Calculation

$$\text{MDA (units/mg protein)} = \frac{\text{Absorbance} \times \text{volume of mixture}}{\text{E}_{532\text{nm}} \times \text{volume of sample} \times \text{mg protein}}$$

Determination of reduced glutathione

The method of Jollow et al. (1974) was followed in estimating the level of reduced glutathione (GSH). 0.2mL of sample was added to 1.8mL of distilled water and 3mL of the precipitating solution was mixed with sample. The mixture was then allowed to stand for approximately 10minutes and then centrifuged at 3000g for 5 minutes. 0.5mL of the supernatant was added to 4mL of 0.1M phosphate buffer. Finally 0.5mL of the Ellman's reagent was added. The absorbance of the reaction mixture was read within 30 minutes of colour development at 412nm against a reagent blank.

Determination of catalase activity

This experiment was carried out using the method described by Sinha (1972). 0.2ml of sample was mixed with 0.8ml distilled H₂O to give 1 in 5 dilution of the sample. The assay mixture contained 2mL of solution (800μmol) and 2.5mL of phosphate buffer in a 10mL flat bottom flask. 0.5mL of properly diluted enzyme preparation was rapidly mixed with the reaction mixture by a gentle swirling motion. The reaction was run at room temperature. A 1mL portion of the reaction mixture was withdrawn and blown into 1mL dichromate/acetic acid reagent at 60

seconds intervals. The hydrogen peroxide content of the withdrawn sample was determined by the method described above. The mononuclear velocity constant, K, for the decomposition of H₂O₂ by catalase was determined by using the equation for a first-order reaction: $K = 1/t \log S_0/S$, where S₀ is the initial concentration of H₂O₂ and S is the concentration of the peroxide at t min. The values of the K are plotted against time in minutes and the velocity constant of catalase K₍₀₎ at 0 min determined by extrapolation. The catalase contents of the enzyme preparation were expressed in terms of Katalase feiahigkeit or 'Katf' according to von Euler and Josephson (1927).

$$\text{Kat. f} = \frac{K_{(0)}}{\text{mg protein/ml}}$$

Determination of Superoxide Dismutase (SOD)

The level of SOD activity was determined by the method of Misra and Fridovich (1972). 1mL of sample was diluted in 9mL of distilled water to make a 1 in 10 dilution. An aliquot of the diluted sample was added to 2.5ml of 0.05M carbonate buffer (pH 10.2) to equilibrate in the spectrophotometer and the reaction started by the addition of 0.3ml of freshly prepared 0.3mM adrenaline to the mixture which was quickly mixed by inversion. The reference cuvette contained 2.5ml buffer, 0.3ml of substrate (adrenaline) and 0.2ml of water. The increase in absorbance at 480nm was monitored every 30 seconds for 150 seconds.

Calculation

$$\text{Increase in absorbance per minute} = \frac{A_3 - A_0}{2.5}$$

where A₀ = absorbance at 0 second

A₃ = absorbance at 150 seconds

$$\% \text{ inhibition} = \frac{\text{increase in absorbance for substrate}}{\text{increase in absorbance of blank}} \times 100$$

1 unit of SOD activity was given as the amount of SOD necessary to cause 50% inhibition of the oxidation of adrenaline to adrenochrome during 1 minute.

Statistical Analysis: The experimental results of the analyses were obtained in triplicates.

Various formulae were used to calculate the individual parameter and enzyme activities. The means and standard deviations of the triplicates results were determined using Microsoft Office Excel 2007

RESULTS AND DISCUSSIONS

Table 1: Specific activity of alkaline phosphatase (ALP) in selected tissues of hypertension induced rats treated with ethanolic extract of *Moringa oleifera* leaf.

Groups	Specific activity (IU/mg protein)		
	Heart	Kidney	Plasma
A	41.40 ±1.06	5.52 ±0.42	16.56 ±2.10
B	39.78 ±0.56	11.04 ±0.22	19.32 ±1.64
C	13.80 ±0.43	30.36 ±0.12	32.76 ±0.06
D	25.52 ±0.23	8.28 ±0.76	11.04 ±1.26

Mean of triplicate results ±SEM

Table 1 showed the activity of alkaline phosphatase in the heart, kidney and serum of hypertensive induced rats treated with ethanolic extract of *Moringa oleifera* leaves. We observed about 67% decrease in ALP activity in the heart of the rats treated with dexamethazone when compared with the control (group A) while the ethanolic extract caused about 28% reversal in the activity of ALP in the heart when compared with control. There was a contrary observation in the kidney as the drug showed increased ALP activity when compared with the control though treatment with the extract caused a reversal to the control as seen in group A. These observations showed the toxic effect of the drug and the ability of the ethanolic extract of the plant to reverse the toxic effect on both organs of the animals. The activity of the enzyme was also monitored in the plasma. Normally, enzymes will not always be found in the serum except when there is damage to one or more organs of the body. Therefore, enzymes from diseased organs may become manifested in the serum resulting in increased activity since they must have leaked from the diseased organ. The results obtained showed that the ethanolic extract of the plant in groups B and D in both organs and plasma caused a reversal in the ALP activity, adjusting their values toward those recorded in the respective controls. The observations in this work showed a resemblance to trends observed in the effect of aqueous extract of nutmeg in bromate induced renal toxicity in rats (Oseni et al., 2015). Increase in plasma level of ALP is due to increased synthesis, in presence of increasing organ dysfunction, as measurement of ALP activity is of clinical and toxicological importance as changes in its activity is indicative of kidney damage by toxicants or in diseased conditions (Singh et al., 2001).

Table 2: Specific activity of aspartate transaminase (AST) in selected tissues of hypertension induced rats treated with ethanolic extract of *Moringa oleifera* leaf.

Groups	Specific activity (IU/mg protein)		
	Heart	Kidney	Plasma
A	163.00 ±5.98	82.00±3.39	29.50±2.88
B	161.00 ±6.34	76.00±7.23	24.50±7.89
C	130.00 ± 4.86	115.50±8.80	123.00±3.25
D	140.00 ± 2.86	68.50±5.54	67.00±2.78

Mean of triplicate results ±SEM

Figures-Tables 2 and 3 showed the AST and ALT activities in the tissues of hypertension induced rats treated with ethanolic extract of *Moringa oleifera* leaves, the results obtained showed

that the extract reversed the effects of the drug in both organs. These tissues participate in a variety of metabolic activities and contain a host of enzymes. The increased plasma AST and ALP activities of the hypertension induced animals showed that the drug resulted in membrane disruption, cellular leakage, and loss of functional integrity of cell membrane the organs of the animals in the group C, while the extract performed corrective roles as observed in the treated animals in group D, while the results of animals in group B showed that the extract maintained the organs integrity.

Table 3: Specific activity of alanine transaminase (ALT) in selected tissues of hypertension induced rats treated with ethanolic extract of *Moringa oleifera* leaves.

Groups	Specific activity (IU/mg protein)		
	Heart	Kidney	Plasma
A	5.40 ±1.65	3.00 ±1.21	67.00 ±2.67
B	6.25 ±0.80	5.60 ±1.10	69.58 ±2.43
C	168.20 ±5.34	20.80 ±3.07	125.00 ±2.30
D	6.80 ±0.75	7.60 ±1.98	57.40 ±1.64

Mean of triplicate results ±SEM

The elevated activities of AST and ALT in plasma are indicative of cellular leakage and loss of the functional integrity of cell membranes (Drotman and Lawhorn, 1978). Administration of dexamethasone significantly raises the plasma level of AST and ALT activities in rats as

observed in our results, which indicated that apart from the hypertensive effect of the drug, it can also cause damage to other organs of the body, measurement AST and ALT activities in rats as observed in our results is of clinical and toxicological importance as changes in their activities are indicative of tissues damage by toxicants or in diseased conditions (Singh et al., 2001). The high level of activities of AST, ALT and ALP in the kidney and heart tissues of the rats and their corresponding decreased activities in the plasma is indicative of non toxicity of the ethanolic extract in these organs.

AST in plasma was increased in dexamethasone-induced rats. A marked elevation in the concentration of the enzyme was also recorded in the kidney and the heart after the injection of the drug. This observation was supported by Fleischer and Schwartz (1971) who reported that any damage done to the cell membrane may lead to leakage of AST and ALP, which is a marker enzyme in the plasma membrane, into extracellular fluid. The study is also corroborated by the work Rajesh and Latha, (2004) that recorded the same trend in CCl₄-intoxicated rats treated with Kamilari. Oral administration of ethanolic extract of *Moringa oleifera* ~~leaf~~-leaves caused a decrease in the activity of the above enzymes, which may be a consequence of the stabilization of plasma membrane as well as repair of damage caused to the organ by dexamethasone. This, ~~however~~-however, indicate that the ethanolic extract displayed a high degree of protection in the organs and plasma.

Table 4: Reduced GSH in selected tissues of hypertensive induced rats treated with ethanolic extract of *Moringa oleifera* leaf.

Groups	(µg/mL)		
	Heart	Kidney	Plasma
A	0.769 ±0.43	0.114 ±0.04	0.641 ±0.42
B	0.841 ±0.03	0.126 ± 0.02	0.763 ±0.04
C	0.280 ±0.04	0.053 ± 0.10	0. 201 ±0.06
D	0.652 ±0.08	0.123 ± 0.03	0. 428 ±0.05

Mean of triplicate results ±SEM

Glutathione is a small tripeptide protein synthesized in the liver (Kaplowitz, et ~~al.~~, 1985). It is a potent antioxidant with high redox potential and it also serves as a co-factor for several oxidative stress detoxifying enzymes (glutathione peroxidase and glutathione transferase) (Valko et al., 2007; Parris, 1997). Glutathione also helps in the regeneration of some important antioxidant vitamins such as C and E. Depletion of GSH has been reported in apoptosis and many degenerative conditions (Parris, 1997). The results of the present study showed that the level of

glutathione (GSH) was significantly decreased in the plasma, heart and kidney homogenates of the animals fed with dexamethazone when compared with the control group A animals. This depletion in blood and organs glutathione is attributable primarily to the dexamethasone, a xenobiotic and an inducer of hypertension. The effects of ethanolic extract of horseradish (*Moringa oleifera*) leaf treatments on the organs and plasma concentration of reduced Glutathione (GSH) showed a significant reversal to the control levels in the tissues and plasma as seen in Table 4. The observations in this study were, however, similar to what was obtained by Ologundudu et al., (2010) in their work on the effect of Hibiscus sabdariffa anthocyanins on 2, 4-dinitrophenylhydrazine-induced tissue damage in rabbits. The reduction in glutathione to the level that was observed in this study could lead to a devastatingly significant decrease in the total antioxidant status of the animals because glutathione helps in recycling cellular antioxidants, inhibits free radical damage, and plays a key role in the detoxification of harmful compounds (Robert et al., 2000). This agrees with earlier works carried out by (Unt et al., 2008) and ; Dominquez et al., (1998) who reported reduced total plasma antioxidant capacity in uncontrolled hypertension. However, the plant extract intake by the hypertensive rats increased their glutathione status to near the control level. ~~T and~~ this is remarkable as this implies that Moringa ethanolic extract could have an ameliorating effect on the altered antioxidant status of the hypertensive rats. It has been reported that decreased hepatic and serum GSH has been consistent with exposure to toxic and oxidative agents, which may also lead to more free ROS (write out the 1st time) leading to excessive oxidative damage (Vijaya et al., 2011; Ahmed et al., 2008; Videla et al., 1991). The results obtained in this work revealed that dexamethazone treated group showed reduced glutathione in both heart, kidney and plasma when compared with the control, which, however, corroborate the results of other previous investigators.

Table 5 revealed the effects of ethanolic extracts of horseradish (*Moringa oleifera*) leaf treatments on the catalase activity of dexamethasone induced hypertensive rat. Catalase is an enzyme which functions in the detoxification of hydrogen peroxide. The reduction in catalase activity after injection with dexamethasone in the heart, kidney and plasma respectively is another significant finding in this study. This decreased concentration in catalase activity of organs and plasma is attributable in part to the reduced synthesis of this antioxidant enzyme whose concentrations would have fallen as a result of the dexamethasone that was injected into

the animals. This study is in agreement with earlier reports by (Erukainure et al. (2012), Udoh et al. (2007) and Tagami et al. (1992) who reported a decreased red blood cell catalase activity in diabetic conditions, while treatment with aqueous extracts of *Moringa oleifera* leaves caused a slight increase in catalase activity.

Table 5: Specific activity catalase in selected tissues of hypertension induced rats treated with ethanolic extract of *Moringa oleifera* leaf.

	($\mu\text{g/ml}$)		Groups
	Heart	Kidney	Plasma
A	18.40 $\pm$ 1.20	10.3 $\pm$ 0.20	11.42 $\pm$ 0.20
B	22.56 $\pm$ 2.50	9.62 $\pm$ 0.01	10.72 $\pm$ 0.04
C	4.25 $\pm$ 0.20	1.66 $\pm$ 0.01	3.42 $\pm$ 0.01
D	16.50 $\pm$ 0.75	9.18 $\pm$ 0.01	8.97 $\pm$ 0.01

Mean of triplicate results $\pm$ SEM

Table 6 revealed the total cholesterol concentration in heart, kidney and plasma of hypertension induced rats that were treated with ethanolic extract of *Moringa oleifera* leaves. The observations in this study revealed that dexamethazone caused hypercholesterolemia, which was subsequently reversed by the ethanolic extract of *Moringa oleifera*. This was however contrary to observation by Oseni et al. (2015) where total cholesterol concentration was depleted as a result of rats fed with bromate. However, the reverse of the total cholesterol concentration to almost control level was observed by the ethanolic extract of *Moringa oleifera*. The ethanolic extract also showed its protective ability against hypercholesterolemia on the normal control animals fed with the extract as seen in Table 6 group B animals

Table 6: Total cholesterol concentration in selected tissues of hypertension induced rats treated with ethanolic extract of *Moringa oleifera* leaves.

	mMol/dL		
Groups	Heart	Kidney	Plasma
A	21.419 $\pm$ 2.09	11.990 $\pm$ 3.11	60.477 $\pm$ 6.34
B	19.138 $\pm$ 3.23	8.349 $\pm$ 1.87	53.547 $\pm$ 4.23
C	77.486 $\pm$ 5.76	16.379 $\pm$ 2.98	101.430 $\pm$ 6.98
D	26.930 $\pm$ 3.87	13.859 $\pm$ 5.87	67.107 $\pm$ 3.47

Mean of triplicate results $\pm$ SEM

352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385

Table 7 showed the effects of ethanolic extract of horseradish (~~Moringa oleifera~~) leaf treatment on the malondialdehyde concentration in dexamethasone-induced hypertensive rats. The concentration of plasma Malondialdehyde (MDA) was shown to be significantly increased in the heart, kidney and plasma of hypertensive rats when compared with the control. However, the hypertensive rats, when placed on Moringa oleifera ethanolic extracts had a remarkable decrease in their cardiac, renal and plasma malondialdehyde levels when compared with the control, thus indicating the extract ability to protect against cellular lipid peroxidation caused by the drug and free radical scavenging ability of the ethanolic extract of Moringa oleifera ~~leaf-leaves~~ in hypertensive rats. The ethanolic extract also showed its protective ability against cellular lipid peroxidation on the normal animals fed with the extract as seen in Table 7 group B animals.

The ability of the ethanolic extract to inhibit the process of lipid peroxidation in vivo may be due to the free radical scavenging activities of its phytochemical components. The observations in this study, however, allude to previous work (Oseni et al., 2015) on effects of aqueous extract of nutmeg on potassium bromated induced renal toxicity, and ~~Olorunnisola et al., 2012~~ on protective effect of T. ~~(write out genus name)~~ violacea rhizome extract against hypercholesterolemia-induced oxidative stress in Wistar rats (~~Olorunnisola et al., 2012~~)

Table 7: Malondialdehyde concentration in selected tissues of hypertension induced rats treated with ethanolic extract of Moringa oleifera leaf.

Groups	(Units/ mg)		
	Heart	Kidney	Plasma
A	0.22 ± 0.01	0.12±0.002	0.50±0.002
B	0.12±0.001	0.08±0.003	0.32±0.004
C	0.31±0.001	0.35±0.010	1.10±0.030
D	0.15±0.001	0.29±0.001	0.30±0.002

Mean of triplicate results ±SEM

CONCLUSSION AND RECOMMENDATION

Moringa oleifera leaf extract has chemopreventive and protective benefits on Dexamethasone mediated tissues oxidative damage in rat as it significantly reduced the extent of antioxidant loss and restoration of organ dysfunction caused by Dexamethasone in rat.

The results of this study show preventive and protective effect on tissues functions ~~which-that~~ might be due to the presence of some bioactive compounds in plant ethanolic extract. On the

basis of our study, however, it can be concluded that *Moringa oleifera* leaf ethanolic extract has plasma, cardiac and renal antitoxic and protective effects as well as antioxidant abilities. Further investigations may therefore be necessary to elucidate the active components of *Moringa oleifera* leaf and establish their chemical nature.

REFERENCES

- AbouZahr C. Global burden of maternal death and disability. *British Medical Bulletin* 2003;67:1-11.
- Adam-vizi V, Seregi A. Receptor independent stimulatory effect of noradrenaline on Naq-Kq ATPase in rat brain homogenate. *Biochem Pharmacol.* 1982;31:2231]6.
- Ahmed RS, Suke SG, Seth V, Chakraborti A, Tripathi AK, Banerjee BD. (2008). Protective effects of dietary ginger (*Zingiber officinales* Rosc) on lindane induced oxidative stress in rats. *Phytother Res*; 22:902-6.
- Akinkugbe O. Epidemiology of cardiovascular disease in developing countries. *Journal of Hypertension* 1990;8:233-238.
- Azriani R et al. The use of herbal medicines during pregnancy and perinatal mortality in Tumpat District, Malaysia. *Southeast Asian J Trop Med* 2007;38(6):1150-7.
- Dominquez, C., E. Ruiz, M. Gussinye and A.C. Carrascosa, 1998. Oxidative stress at onset and early stages of type1 diabetes in children and adolescents. *Diabetes Care*, 21: 1736-1742.
- Drotman, R.B., Lawhorn, G.T., 1978. Serum enzymes as indicators of chemical-induced liver damage. *Drug and Chemical Toxicology* 1, 163–171.
- Erukainure, O.L., F.O. Adeboyejo, G.N. Elemo and O.A. Ebuehi, 2012. Modulation of lipid peroxidation, hypolipidemic and antioxidant activities in brain tissues of diabetic rats by fibre-Enriched biscuits. *J. Acute Dis.*, 1: 42-46.
- Hughes E and Bradly J. Complementary and Alternative medicine. In: Tierney L et al, editor. *Current medical Diagnosis and treatment*. 43rd ed; 2004.
- Jedi W. *Moringa oleifera*: A review of the medical evidence for its nutritional, therapeutic and prophylactic properties. *Trees of Life Journal* 2005;1(1):5.
- Johanna W et al. Critical care obstetrics. In: *Current obstetric and gynaecologic diagnosis and treatment*. 9th ed; 2003.

416 Jollow DJ, Michell JR, Zampaglionic, Gillete JR. 1974. Bromoibenzene-induced Liver necrosis:
 417 Protective role of glutathione and evidence for 3,4- Bromobenzene oxide as hepatotoxic
 418 metabolite. Pharmacology. 11:151-169.

419 Kamatenesi-Mugisha. The Socio-Cultural Aspects in Utilisation of Medicinal Plants in
 420 Reproductive Health Care in Western Uganda; 2002.

421 Kaplowitz, N.; Aw, T.Y.; Ookhtens, M. (1985) The regulation of hepatic glutathione. Ann.
 422 Rev. Pharmacol. Toxicol. 25, 715–744.

423 Katzung B. Cardiac Glycosides and other drugs used in Congestive heart failure. In: Basic and
 424 Clinical Pharmacology. 8th ed; 2001. p. 200-217.

425 Katzung B. Drugs with important actions on smooth muscle. In: Basic and Clinical
 426 Pharmacology. 8th ed; 2001. p. 265-291,

427 Katzung B. The Eicosanoids. In: Basic and Clinical Pharmacology. 8th ed; 2001. p. 311-322.

428 Kearney P et al. Global burden of hypertension: Analysis of worldwide data. Lancet
 429 2005;365(9455):217-23.

430 Massie B. Systemic Hypertension. In: Tierney L et al, editor. Current Medical Diagnosis and
 431 Treatment. 43rd ed: Lange; 2004. p. 401-427.

432 Misra HP, Fridovich I. 1972. The role of superoxide anion in the autoxidation of epinephrine and
 433 a simple assay for superoxide dismutase. J Biol Chem. 247:3170-5.

434 Nath D et al. Commonly used Indian abortifacient plants with special reference to their
 435 teratological effects in rats. J Ethnopharmacol 1992;36(2):147-154.

436 Ologundudu, A., Ologundudu, A. O. Oluba, O. M. Omotuyi, I. O. and Obi, F. O. (2010) Effect
 437 of Hibiscus sabdariffa anthocyanins on 2, 4-dinitrophenylhydrazine-induced tissue
 438 damage in rabbits. Journal of Toxicology and Environmental Health Sciences Vol. 2(1),
 439 pp. 1-6

440 Oseni, O. A., Olagboye, S. A. and Idowu, A. S. K. (2015) Potassium Bromate Induced Renal
 441 Toxicity in Wistar Albino Rats: Effects of Aqueous Extract of Nutmeg (*Myristica*
 442 *fragrans* Houtt). British Journal of Medicine & Medical Research 5(12): 1547-1556

- 443 Parris, M.K. (1997). Glutathione systemic protectant against oxidative and free radical damage.
444 Altern. Med. Rev. 2, 155–176.
- 445 Peltzer K et al. Use of traditional, complementary and alternative medicine for HIV patients in
446 KwaZulu-Natal, South Africa. BMC public Health 2008(8):255.
- 447 Robert, K.M., K.G. Darry and A.M. Peter, 2000. Harpers Biochemistry. 23rd Edn., Appleton and
448 Lange, USA., pp: 150-151.
- 449 Rajesh, M.G. and Latha, M.S. (2004). Preliminary evaluation of the antihepatotoxic activity of
450 Kamilari, a polyherbal formulation. Journal of Ethinopharmacology, 91; 99-104.
- 451 Sinha, AK. 1972. Colorimetric assay of catalase. Anal Biochem. 47:389-94.
- 452 Tagami, S., T. Kondo, K. Yoshidfa, J. Hirokaw, Y. Ohsuka and Y. Kaweasmi, 1992. Effect of
453 insulin on impaired antioxidant activities on aortic endothelial cells from diabetic
454 rabbits. Metabolism, 41: 1053-1058.
- 455 Tilburt J and Kaptchuk T. Herbal medicine research and global health: an ethical analysis.
456 Bulletin of the World Health Organisation 2008 August 2008:577-656.
- 457 Udoh, A.E., I. Ntu, O. Essien and M. Ndon, 2007. Red cell catalase activity in diabetics. Pak. J.
458 Nutr., 6: 511-515.
- 459 Unt, E., C. Kairane, I. Vaher and M. Zilmer, 2008. Red blood cell and whole blood glutathione
460 redox status in endurance-trained men following a ski marathon. J. Sports Sci. Med.,
461 7: 344-349.
- 462 Valko, M.; Leibfritz, D.; Moncol, J.; Cronin, M.T.D.; Mazur, M.; Telser, J. (2007) Free radicals
463 and antioxidants in normal physiological functions and human disease. Int. J. Biochem.
464 Cell Biol. 39, 44–48
- 465 Varshney R, Kale RK. Effects of Calmodulin Antagoniss. Int. J Rad. Biol. 1990;58:733-743.
- 466 Videla LA, Simizu K, Barros SB, Junqueira VB. (1991). Mechanisms of lindane-induced
467 hepatotoxicity: alterations of respiratory activity and sinusoidal glutathione efflux in the
468 isolated perfused rat liver. Xenobiotica; 21:1023-32.
- 469 Vijaya Padma V, Sowmya P, Arun FT, Baskaran R, Poornima P. (2011)Protective effect of
470 gallic acid against lindane induced toxicity in experimental rats. Food Chem Toxicol; 49:

471 991-8.

472 von Euler HV, Josephson K. 1927. European Journal of Organic Chemistry - EUR J ORG

473 CHEM.; 452(1):158-181.

474 Wells, F.E., 1988. Tests in liver and biliary tract disease. In: Gowenlock, H.A. (Ed.), Varley's

475 Practical Clinical Biochemistry. CRC Press, Florida.

476 Zimmerman, H.J., Seef, L.B., 1970. Enzymes in hepatic disease. In: Goodly, E.I. (Ed.),

477 Diagnostic Enzymology. Lea and Febiger, Philadelphia.

478

479