

Original Research Article
**IN VITRO AND IN VIVO EFFECT OF QUERCETIN
ON HEPATOCYTE TRANSMEMBRANE Na^+ , K^+ -
ATPase ACTIVITY IN RATS**

ABSTRACT

Aims: This study investigated *in vitro* and *in vivo* effect of quercetin on transmembrane hepatocyte Na^+ , K^+ -ATPase activity in normal Wistar rats.

Study design: One factor experimental design.

Place and Duration of Study: Department of Biochemistry, Federal University of Technology, Akure, Ondo State, Nigeria, between April and September 2010.

Methodology: *In vitro* effect of quercetin on transmembrane hepatocyte Na^+ , K^+ -ATPase activity was investigated through enzyme assay by measuring the released inorganic phosphate (Pi) through the method of Fiske and Subbarow, while in *in vivo* experiment, three (3) main groups of male Wistar rats weighing 200-250g (n=5) were randomly selected. Group 1 (IP; Intraperitoneal), group 2 (IV; Intravenous) and 3 (OR; Oral) group. All three groups were administered with quercetin 50 mg/kg body weight (once/day) for 5 days through respective route. The control for each group received equal volume of vehicle through respective route of administration. Hepatocyte Na^+ , K^+ -ATPase activity for each group was assayed after days of quercetin administration.

Results: *In vitro* results indicated that quercetin, with IC_{50} of 5.75 μM showed high inhibitory activity on Na^+ , K^+ -ATPase, while *in vivo* result showed inhibitory activity on hepatic Na^+ K^+ -ATPase irrespective of the routes of administration. Quercetin through the oral route showed significant ($F_{5, 12} = 265.8$; $P = 0.05$) inhibitory effect on hepatic Na^+ K^+ -ATPase compared to control. Administration of quercetin through the parenteral routes (intraperitoneal and intravenous) showed significant ($F_{5, 12} = 265.8$; $P < 0.001$) inhibitory effect compared to respective control. The parenteral routes of administration of quercetin have significant ($F_{5, 12} = 265.8$; $P < 0.001$) inhibitory effect on hepatic Na^+ K^+ -ATPase compared to the oral route.

Conclusion: Quercetin showed high inhibitory effect *in vitro* on hepatocyte Na^+ , K^+ -ATPase activity in normal rats. *In vivo* administration of quercetin showed significant inhibitory effect on hepatocyte Na^+ , K^+ -ATPase activity in normal rats, with the parenteral routes of administration exerting significant inhibitory effect compared to the oral route.

Keywords: Quercetin, hepatocyte Na^+ , K^+ -ATPase, *In vitro*, *In vivo*.

1. INTRODUCTION

Ubiquitous transmembrane enzyme, Na^+ , K^+ -ATPase, also known as sodium pump transports Na^+ and K^+ across the plasma membrane by hydrolyzing ATP [1]. Like the H^+ , K^+ -ATPase, Na^+ , K^+ -ATPase belongs to the P-type ATPases. It is composed of majorly two subunits α and β . The catalytic α subunit is a large membrane protein approximately between 110,000 – 113,000 Da, and it contains the binding sites for ATP, sodium, potassium, phosphate and cardiac glycoside [2, 3], while the β subunit with approximately 35,000 to 55,000 Da is assumed to be necessary for the activity of the complex [4]. Of the four (4) α isoforms that have been identified in mammals with tissue-specific pattern of expression, the $\alpha 1$ isoform is however ubiquitous and has been identified in hepatocytes in the basolateral and bile canalicular membranes [5, 6].

The Na^+ , K^+ -ATPase plays an important role in the control of the ionic intracellular milieu, this process is needed for the maintenance of the characteristic intracellular concentration of sodium and potassium, and to determine the transmembrane electrochemical gradients for these ions. It is also needed for regulation of metabolism, proliferation, differentiation and most importantly cell volume. It is a common belief that the electrochemical sodium gradient provides the driving force for secondary active sodium-coupled transport of a variety of solutes, including certain amino acids and bile, and also drives the activity of many other transporters like Na^+ , H^+ exchanger and Na^+ - K^+ - 2Cl^- cotransporter involved in the regulation of intracellular pH and cell volume respectively [7]. Knowing that apoptotic cells lose K^+ , and gain Na^+ in a process that involves the Na^+ , K^+ -ATPase underpins the significance of this ubiquitous membrane protein in both physiologic and pathologic states [8, 9]. It has been hypothesized that variations in hepatocyte ATP might be transduced into neural signaling by modulating Na^+ , K^+ -ATPase activity, in that, decrease in hepatocyte ATP which reduce the activity of Na^+ , K^+ -ATPase, thus depolarizing the cell which in turn would affect hepatic neuronal activity through close coupling or release of a neuromodulator. This is evidenced by the observation that feeding behaviour in rats is stimulated by intraperitoneal or hepatic portal administration of ouabain, an inhibitor of the Na^+ , K^+ -ATPase [10 - 12].

Quercetin (3, 3', 4', 5, 7-pentahydroxyflavone) is one of several naturally-occurring dietary flavonoids compounds, ubiquitously found in fruits, vegetables, herbs, tea, wine, etc. Quercetin belongs to the group of flavonoids, and generally, flavonoids are characterized by a phenyl benzo (γ) pyronederived structure consisting of two benzene rings (A and B) linked by a heterocyclic pyran or pyrone ring [13]. In plants, the flavonol aglycone is most commonly present conjugated at the 3- position of the unsaturated C-ring with a sugar moiety, forming O- β -glycosides such as quercitrin or rutin [14]. Quercetin can be obtained from plants via extraction of the quercetin glycosides followed by hydrolysis to release the aglycone and subsequent purification. Flavonols exhibit numerous biological and pharmacological effects, including anti-oxidant, chelation, anti-carcinogenic, cardioprotective, bacteriostatic, and secretory properties [13]. The biological action of quercetin is connected with its antioxidant properties which are mainly due to: (1) its ability to scavenge free radicals and reactive oxygen species (ROS) (superoxide anion, hydroxyl-radical) [15], (2) to form complexes with metal ions, thus preventing oxidation of the metals with oxygen yielding ROS (Haber-Weiss and Fenton reactions) [16, 17]; and (3) reduction of Fe^{3+} to Fe^{2+} [18]. These mechanisms explain the inhibition of the lipid peroxidation reactions by quercetin [19-20] and its ability to prevent deoxyribose degradation [18]. The mechanism(s) contributing to the health beneficial effects of quercetin involve the quenching of free radicals, elevating antioxidant status, and membrane modulating effects [21]. Quercetin has shown a potent effect at membrane level modulating the activity of membrane transport processes of erythrocyte's Na^+ , K^+ -ATPase and Na^+ , H^+ antiport in diabetic patients [22].

Our previous researches showed that quercetin elevated antioxidant status and modulated Na^+ , K^+ -ATPase in rat brain [23], and also potentiated hepatoprotective and antioxidant response to intraperitoneal, intravenous, subcutaneous and oral administration in Wistar rats [24]. Meanwhile, a recent work by Kairane *et. al.* [25] showed that genistein, an exogenous antioxidant like quercetin, has an inhibitory effect on Na^+ , K^+ -ATPase in normal brain, but improve the activity of this pump in Alzheimer's disease brain. Since our prior works showed inhibitory effect of quercetin on cerebral Na^+ , K^+ -ATPase, but antioxidant effect in both the brain and liver in rats, we have taken to investigating the effect of quercetin on hepatic Na^+ , K^+ -ATPase in normal rats.

2. MATERIAL AND METHODS

2.1. Chemicals

Quercetin was obtained from Sigma-Aldrich (Germany). All other chemicals used were of analytical grade and obtained from FLUKA, BDH (Germany) and other standard commercial suppliers.

2.2 Animals

Male adult Wistar rats (200–250 g) obtained from the animal facility of the Department of Biochemistry of the Federal University of Technology, Akure, Nigeria were used in this study. Animals were maintained with food and water ad libitum and under a 12-h light/12-h dark cycle. The “principle of laboratory animal care” (National Institute of Health-NIH publication No. 85-23) guidelines and procedures were followed in this study (NIH publication revised, 5). The Ethical Committee of the Faculty Postgraduate Committee, Faculty of Science, Federal University of Technology, approved the research work.

2.3 *In vitro* Effect of Quercetin on Na⁺, K⁺-ATPase Activity

Immediately after the animals were euthanized, the liver was quickly removed and the homogenate was prepared in 0.05 M Tris–HCl, pH 7.4. The homogenate was centrifuged at 4,000 rpm at 4 °C for 7 min and the supernatant was used for the assay of Na⁺, K⁺-ATPase. The reaction mixture for Mg²⁺-dependent- Na⁺, K⁺-ATPase activity assay contained 3 mM MgCl₂, 125 mM NaCl, 20 mM KCl, 200mM sodium azide and 50 mM Tris–HCl, pH 7.4 and 100–120 µg of protein, in a final volume of 500 µl. The reaction was initiated by addition of ATP to a final concentration of 3.0 mM. Controls were carried out under the same conditions with the addition of 0.1 mM ouabain [26, 27]. Na⁺, K⁺-ATPase activity was calculated by the difference between the two assays. Released inorganic phosphorous (Pi) was measured by the method of Fiske and Subbarow [28].

2.4 *In vivo* Experiment

The animals were divided randomly into the following groups (n=5 per group): group 1 (IP, intraperitoneal); group 2 (IV, intravenous); group, and 3 (OR, oral) group. Groups 1, 2 and 3 were administered quercetin at a dose of 50 mg/kg (once/day) for 5 days. Quercetin was pre-dissolved in ethanol. Control for each group received equal volume of vehicle through respective route of administration.

2.4.1 Tissue preparation for *in- vivo* experiment

Animals were anesthetized with ether and euthanized by decapitation and whole liver were quickly removed, placed on ice and homogenized in cold 50 mM Tris–HCl pH 7.4. The homogenate was centrifuged at 4,000 g for 10 min to yield the low-speed supernatant (S2). For all analyses, protein content was determined by the method of Lowry *et al.* [29], using bovine serum albumin as the standard.

2.4.1.1 In-vivo effect of quercetin on Na⁺, K⁺-ATPase activity

The assay was carried out as described earlier for in-vitro assay. Released inorganic phosphorous (Pi) was measured by the method of Fiske and Subbarow [28].

2.5 Statistical analysis

The results of the *in vivo* experiment are expressed as mean $\pm$ SEM and statistical difference was determined by Student Newman Keul's Test after a one-way analysis of variance (ANOVA) was done. The level of significant was set at 5% ($P < 0.05$) for all treatment carried out. For the *in vitro* experiment, IC_{50} was calculated with the aid of Adobe® Page Maker® 7.0 from graphs plotted with the aid of Graph pad PRISM® version 3.00.

3. RESULTS AND DISCUSSION

3.1 *In vitro* Effect of Quercetin on Na^+ , K^+ -ATPase Activity

The *in vitro* result in this study showed that quercetin, like ouabain a potent inhibitor of Na^+ , K^+ -ATPase, has an inhibitory effect on the hepatocyte Na^+ , K^+ -ATPase activity (see figure 1). While the IC_{50} for ouabain was $3.63 \mu\text{M}$, the IC_{50} for quercetin was $5.75 \mu\text{M}$. These indicate, as noticed in our previous research [23], that quercetin showed inhibitory activity in both normal liver and brain Na^+ , K^+ -ATPase. However, using Rukunga and Simon's classification of activity of some natural products [30], both ouabain and quercetin (with IC_{50} $3.63 \mu\text{M}$ and $5.75 \mu\text{M}$ respectively) have high inhibitory activity on hepatic Na^+ , K^+ -ATPase activity. Both ouabain and quercetin showed a concentration-dependent inhibitory activity on hepatic Na^+ , K^+ -ATPase activity. Genistein, an exogenous antioxidant like quercetin, has been shown to have inhibitory effect on Na^+ , K^+ -ATPase activity in normal brain, but improved the activity of Na^+ , K^+ -ATPase in Alzheimer disease (AD) brain [25]. The speculation to genistein affecting Na^+ , K^+ -ATPase activity in control and AD brains in opposite directions was said to be that Na^+ , K^+ -ATPase is redox sensitive. That whereas shifts both into a more reduced and oxidized states cause the suppression of Na^+ , K^+ -ATPase activity in control brain, the antioxidant activity of genistein may change redox balance back towards the reduced state and thus helping to improve Na^+ , K^+ -ATPase activity in the AD brain. We have shown in our previous work [24] that quercetin showed hepatoprotective and antioxidant effects in normal rat liver. Thus quercetin, an antioxidant like genistein, showing inhibitory effect on normal hepatocyte Na^+ , K^+ -ATPase might achieved this, like genistein, by shifting both into a more reduced and oxidized state to cause inhibitory activity on normal hepatocyte Na^+ , K^+ -ATPase. We used normal hepatocyte Na^+ , K^+ -ATPase in this study, a further work will be necessary to investigate the effect of quercetin on Na^+ , K^+ -ATPase in a diseased liver.





Fig. 1. Na⁺, K⁺-ATPase inhibitory activity of ouabain compared with quercetin in rat liver.

3.2 *In vivo* Effect of Quercetin on Na⁺, K⁺-ATPase Activity

This study also investigated the effect of *in vivo* (oral, intraperitoneal and intravenous) administration of quercetin on hepatic Na⁺ K⁺-ATPase activity. The results indicated that quercetin showed inhibitory activity on hepatic Na⁺ K⁺-ATPase *in vivo* irrespective of the routes of administration. Quercetin through the oral route showed significant ($F_{5, 12} = 265.8$; $P = 0.05$) inhibitory effect on hepatic Na⁺ K⁺-ATPase activity compared to control. Administration of quercetin through the parenteral routes (intraperitoneal and intravenous) showed significant ($F_{5, 12} = 265.8$; $P < 0.001$) inhibitory effect compared to respective control. The parenteral routes of quercetin administration have significant ($F_{5, 12} = 265.8$; $P < 0.001$) inhibitory effect on hepatic Na⁺ K⁺-ATPase activity compared to the oral route (see figure 2). The pattern of inhibition exhibited in this study between the oral and parenteral routes, has also been seen in the hepatoprotective and antioxidant effects of quercetin in normal rats [24]. This has been attributed to poor absorption of quercetin in the small intestine when administered through the oral route [30-32].



179

180 **Fig. 2. Effect of quercetin on hepatocyte Na⁺, K⁺- ATPase activity; comparing different**
 181 **routes of administration.**

182 Each bar is expressed as Mean $\pm$ SEM of released inorganic phosphate (Pi) (n=5 per group). ^a P =
 183 0.05 compared to oral control, ^b P < 0.001 compared to oral and ^c P < 0.001 compared to
 184 intraperitoneal.

185

186 4. Conclusion

187 Quercetin showed high inhibitory effect *in vitro* on hepatocyte Na⁺, K⁺- ATPase activity in
 188 normal rats. In vivo administration of quercetin showed significant inhibitory effect on
 189 hepatocyte Na⁺, K⁺- ATPase activity in normal rats, with the parenteral routes of
 190 administration exerting significant inhibitory effect compared to the oral route.

ETHICAL APPROVAL

The “principle of laboratory animal care” (National Institute of Health-NIH publication No. 85-23) guidelines and procedures were followed in this study (NIH publication revised, 5). The Postgraduate Ethical Committee of the School of Science, Federal University of Technology, approved the research work.

REFERENCES

1. Kaplan JH, Biochemistry of Na, K-ATPase. *Anna. Rev. Biochem.* 2002;71:511-535.
2. Dostanic-Larson I, Van-Huyse JW, Lorenz JN, Lingrel JB. The highly conserved cardiac glycoside binding site of Na,K-ATPase plays a role in blood pressure regulation. *Proc. Natl. Acad. Sci. USA* 2005;102:15845-15850.
3. Vasilets LA, Schwarz W. Structure-function relationship of cation binding in the Na⁺/K⁺-ATPase *Biochimica et Biophysica Acta* 1993;1154(2):201-222.
4. Kirley TL. Determination of three disulfide bonds and one free sulfhydryl in the beta subunit of (Na,K)-ATPase. *J. Biol. Chem.* 1994;269:19659.
5. Lingrel JB Kuntzweiler T. Na⁺ K⁺-ATPase. *J. Biol. Chem.* 1994;269:19659.
6. Brisse J, Sastre B, Bongrand P, Chamlian A. Immunocytochemical study of Na⁺ K⁺-ATPase alpha 1 and beta 1 subunits in human and rat normal hepatocytes using confocal microscopy *Cell Mol. Biol.* 1995;41:499.
7. Erlinger S. Does Na⁺-K⁺ ATPase have any role in bile secretion? *Am. J. Physiol. Gastrointest. Liver Physiol.* 1982;243:G243.
8. Yu SF. Na⁺, K⁺-ATPase: the new face of old player in pathogenesis and apoptotic/hybrid cell death. *Biochem. Pharmacol.* 2003;6:1601.
9. Panayiotidis MI, Bortrar CD, Cidlowski JA. On the mechanism of ionic regulation of apoptosis: would the Na⁺/K⁺ ATPase please stand up? *Acta Physiol.* 2006;186:203.
10. Langhans W. Role of the liver in the metabolic control of eating: what we know- and what we do not know. *Neurosci. Biobehav. Rev.* 1996;20:145-153.
11. Langhans W., Schharrer E. Metabolic control of eating, in: S.A.P. [Ed.] *Metabolic Control of Eating, Energy Expenditure and the Bioenergetics of Obesity*. World Review of Nutrition and Dietetics, S. Karger, Basel. 1993, pp. 1-67.
12. Langhans W. Scharrer E. Evidence of a role of the sodium pump of hepatocytes in the control of food intake. *J. Auton. Nerv. Syst.* 1987;20:199-205.
13. Harwood M, Danielewska-Nikiel B, Borzelleca JF, Flamm GW, Williams GM, Lines TC. A critical review of the data related to the safety of quercetin and lack of evidence of in vivo toxicity, including lack of genotoxic/carcinogenic properties. *Food Chem. Toxicol.* 2007;45:2179-2205.
14. Merck. Quercetin. The Merck Index: An Encyclopedia of Chemicals, Drugs and Biologicals, ed 13, Merck & Co., Inc. Whitehouse Station, New Jersey 2001, 1438.
15. Javanovic SV, Steenken S, Tosic M, Majanovic A, Simic MG. Flavonoids as antioxidants. *J. Am. Chem. Soc.* 1994;116:4846-4851.
16. Morel I, Lescoat G, Gillard P, Gillard J. Role of flavonoids and iron chelation in antioxidant action. *Methods Enzymol.* 1994;234:443.
17. Miral L, Fernandez MT, Santos M, Rocha R, Florencio MH, Jennings KR, Interactions of flavonoids with iron and copper ions: a mechanism for their antioxidant activity. *Free Radical. Res.* 2002;3:1199-1208.
18. Omololu PA, Rocha JBT, Kade IJ. Attachment of rhamnosyl glucoside on quercetin confers protein iron-chelating ability on its antioxidant properties. *Exp. Toxicol. Pathol.* 2011;63(3):249-255.
19. Hayek T, Fuhrman B, Vaya J, Rosenblat M, Belinky P, Coleman R. Reduced Progression of atherosclerosis in apolipoprotein E- deficient mice following consumption of red wine, or its polyphenols quercetin or catechin in association with reduced

244 susceptibility of LDL to oxidation and aggregation. *Arterioscler. Thromb. Vasc Biol.*
 245 1997;17:2744-2752.

246 20. Zhao Y, Gao Z, Li H, Xu H. Hemin/nitrite/H₂O₂ induces brain homogenate oxidation and
 247 nitration: effects of some flavonoids. *Biochim. Biophys. Acta* 2004;1675:105-112.

248 21. Renugadevi J, Prabu SM, Quercetin protect against oxidative stress related renal
 249 dysfunction by cadmium in rats *Exp. Toxicol. Pathol.* 2010;62:471-481.

250 22. Mishra N, Rizvi SI, Quercetin Modulates Na⁺, K⁺ ATPase and Sodium Hydrogen
 251 Exchanger in Type 2 Diabetic Erythrocytes. *Cell Mol. Biol.* 2012;58:148-152.

252 23. Ogundajo A, Imoru J, Kade I, Olawoye T. Quercetin elevates antioxidant status and
 253 Modulates ouabain-sensitive transmembrane sodium pump in rat brain. *TPI Journal.*
 254 2014;3(10):69-75.

255 24. Ogundajo AT, Imoru JO, Asaolu FM. Quercetin potentiates hepatoprotective and
 256 antioxidant response to intraperitoneal, intravenous, subcutaneous and oral
 257 administration in Wistar rats. *Asian J. Biomed. Pharma. Sci.* 2014;04(38):57-61.

258 25. Kairane C, Mahlapuu R, Ehrlich K, Zilmer M, Soomets U. The effects of different
 259 antioxidants on the activity of cerebrocortical MnSOD and Na, K-ATPase from post
 260 mortem Alzheimer's disease and age-matched normal brains. *Current Alzheimer Res.*
 261 2014;11(1):79-85.

262 26. Kade IJ, Paixão MW, Rodrigues OED, Barbosa NBV, Broga AL, Avila DS, Nogveira
 263 CW, Rocha JBT. Comparative Studies on Dicholesteryl Diselenide and Diphenyl
 264 Diselenide as Antioxidant Agents and their effect on the Activities of Na⁺/K⁺ ATPase
 265 and δ-Aminolevulinic acid Dehydratase in Rat Brain. *Neurochem. Res.* 2008;33:167-
 266 178.

267 27. Tsakiris S., Marinou K., Schulpis KH. The Effect of Galactose Metabolic Disorders on
 268 Rat Brain Na⁺, K⁺-ATPase Activity. *Z. Naturforsch.* 2002;57c:939-943.

269 28. Fiske CH, Subbarow TJ. The colorimetric determination of phosphorus. *Journal. Biol.*
 270 *Chem.* 1925;66:375-378.

271 29. Lowry OH, Rosenbrough NJ, Farr AL, Randall RJ. Protein measurement with Folin-
 272 Phenol reagent. *Journal Biol. Chem.* 1951;193:265-275.

273 30. Brown JP. A review of the genetic effects of naturally occurring flavonoids,
 274 anthraquinones and related compounds. *Mutat. Res.* 1980;75:243-277.

275 31. Tamura G, Gold C, Ferro-luzzi A, Ames BN. Fecalase. A model for activation of dietary
 276 glycosides to mutagens by intestinal flora. *Roc. Natl. Acad. Sci. USA.* 1980;77:451-455.

277 32. Bokkenheuser VD, Shackleton CHL, Winter J. Hydrolysis of dietary flavonoid glycosides
 278 by strains of intestinal bacteroides from humans. *Biochem. J.* 1987;248:953-956.

279