

EXTERNAL AUDITORY CANAL FRACTURE AFTER POSTERIOR DISLOCATION OF AN INTACT CONDYLAR HEAD.

Karabouta Irene 1, Kalimeras Eleftherios², Georgiou A. Christos³*

(Karabouta Irene, +30 2320 72922, +30 23920 72468, syomo@otenet.gr)

1. *DDS, MD, PhD , European Interbalkan Medical Center, 10 Asclepiou Str., Pylaia
Thessaloniki, Greece*
2. *DDS, MD, European Interbalkan Medical Center, 10 Asclepiou Str., Pylaia
Thessaloniki, Greece*
3. *DDS, MD, Theageneio - Anticancer Hospital of Thessaloniki, Al. Simeonidi 2,
Thessaloniki, Greece*
syomo@otenet.gr, syomo@otenet.gr

ABSTRACT

Aims: The aim of this article is the kind of applied treatment for fracture of the external auditory canal, following traumatic facial injuries.

Presentation of Case: The case report of the patient described in this article, is evaluated for the posterior dislocation of an intact mandibular condyle after facial injuries she had due to a car accident. The physical and x-ray examination of the External Auditory Canal (EAC) revealed facial fractures, TMJ dysfunctional symptoms with severe disk displacement, as well as fracture of the anterior wall of the EAC. Atraumatic reduction of the fracture was accomplished during the TMJ disc reduction and joint exploration, resulting in a satisfactory outcome with no complications and with improvement of hearing impairment.

Discussion: The anterior osseous wall of the external auditory canal, which represents part of the tympanic portion of the temporal bone, defines the posterior limit of the glenoid fossa, and is situated close to the condyle of the mandible. Due to this intimate anatomical relationship, herniation of the TMJ apparatus into the external auditory canal (EAC) occurs spontaneously or secondary to neoplasia, inflammation, developmental problems and especially trauma. Direct high-energy impact into the chin displaces the condyle posteriorly and the result may be a fracture of the condyle or posterior dislocation of intact condyle without fracture, a dislocation of condyle into middle cranial fossa or temporal fossa, or a fracture on the anterior wall of the EAC.

Conclusion: In summary, a direct blow to the mandible can result in a TMJ apparatus injury. Due to the close relationship between TMJ and EAC, an atypical injury, such as a fracture of the anterior wall of the EAC can occur. An oral & maxillofacial surgeon, when called to examine and diagnose TMJ injury disorders, has the challenging responsibility to take into account the potential concomitant temporal bone fractures or intracranial complications in cooperation with radiologists, ENT doctors and neurosurgeons.

Keywords: fracture, auditory, canal, condylar dislocation.

INTRODUCTION

Temporal bone fractures are, by definition, fractures of the skull base, and are often associated with injuries to other areas of the craniomaxillofacial skeleton[1]. They represent roughly 20% of all skull fractures, while up to 75% of patients with a skull base fracture have a temporal bone fracture as a component of the injury[1]. The most common causes include motor vehicle accidents, falls, bicycle accidents, athletic injuries, assaults and penetrating trauma[2]. Risk factors include younger or older age, male gender.

48 The temporal bone includes the squamous, petrous, mastoid, and tympanic portions, as well as the styloid process[1].
49 Classically, petrous temporal bone fractures are classified as longitudinal (parallel to the long axis of the petrous bone) or
50 transverse (perpendicular to the long axis) depending on the orientation of the fracture line, or mixed when the fracture line
51 extends in any direction across the basal portion of the skull[1].
52 Longitudinal fractures begin at the squamous portion of the temporal bone, run through the EAC and turn anteriorly
53 toward the foramen lacerum. They account for about 80% of cases, whereas transverse fractures are far less common
54 than longitudinal fractures and are frequently caused by a severe blow to the occipital portion of the calvaria or by a
55 direct frontal blow[1]. Transverse fractures extend directly across the petrous pyramid, fracturing the otic capsule, and
56 then extend anteriorly along the eustachian tube and geniculate ganglion.
57 The EAC is divided into an outer cartilaginous, one-third, and an inner osseous, two-thirds, which represent the tympanic
58 portion of the temporal bone[3]. The canal extends from the conchal cartilage to the tympanic membrane and is
59 approximately 25mm long and slightly S shaped. Within the anterior and inferior portions of the cartilaginous ear canal,
60 there are small fenestrations through the cartilage called the "fissures of Santorini". The anterior and inferior walls and the
61 lower portion of the posterior walls of the osseous canal are developed from the tympanic ring. The posterior wall is
62 closed to the mastoid cells and the descending portion of the facial canal. The inferior wall is composed of dense bone
63 and the anterior wall, which defines the posterior limit of the glenoid fossa, is close to the condyle of the mandible. The
64 blood supply of the EAC originates from the external carotid artery (posterior auricular and superficial temporal branches)
65 and from the maxillary artery (deep auricular branch and anterior tympanic artery)[3, 4]. This rich blood supply explains the
66 bleeding from the EAC in patients with maxillofacial fractures.
67 In this article we present a case of a patient with a fracture of the left external auditory canal due to a force applied in the
68 right side of the face.

70

71

72 PRESENTATION OF CASE

73 A 29 year-old female arrived to our clinic with injuries to the left side of her face, which were caused by a car accident 20
74 days before. At the time of the accident the woman was moved to the Emergency Department of a local Hospital, where
75 first aid was given. According to the history, bleeding in the left ear and hearing disturbances were found and evaluated at
76 first by an ENT doctor.

77 After the primary care, the physical and x-ray examination revealed facial injuries, hearing discomfort in the left ear, as
78 well as TMJ dysfunctional symptoms, such as restricted mouth opening, deviation and "locking" of mandible, especially in
79 the morning session due to disk derangements following the accident (Fig.1). She also complained of preauricular pain in
80 the left side. There were no signs of brain injury or neurological deficit. From the history of the accident, she reported a
81 severe blow to the chin and to the lateral craniofacial region of the left side. So bleeding of the external auditory canal
82 was seen as possibly being caused by transversal movement of the condylar head of the lower jaw.

83 When the patient came to our clinic, panoramic radiograph and computed tomography (CT) showed an incomplete
84 parasymphseal fracture of the right mandible, a left two points zygoma fracture, as well as a fracture of the anterior wall
85 of the external auditory canal on the left. The left condyle showed to be in an anterior place during centric occlusion. (Fig.
86 2, Fig. 3, Fig.4)

87 With the patient being under general anaesthesia the parasymphysis and zygoma fractures were reduced and the left
88 TMJ was exposed by the preauricular incision. After the disk reduction using absorbable mini anchor (polylactic acid) with
89 ethibond non-absorbable suture and during the exploration of the joint, a gap was revealed in the posterior wall, which
90 corresponded to the fracture of the anterior wall of the EAC. With careful entry of a blunt elevator inside the EAC and
91 following the canal outline, reduction of the fracture was achieved, with release of characteristic sound.

92 The postoperative course was uneventful. Examination of the patient by an ENT specialist in the following days, showed a
93 considerable improvement in the hearing impairment. Two years after the operation a computed tomograph showed an
94 EAC breadth within normal limits. (Fig. 5)

95

96

97

98 DISCUSSION

99 Hearing loss or impairment, nausea, vomiting and vertigo, TMJ discomfort or dysfunction (TMJ trismus, inability to chew,
00 and localized pain) are symptoms of patients with temporal bone and external auditory canal fractures, as were found in

our case. They consider as pathognomonic signs even though other clinical findings may include ecchymosis, particularly in the periorbital region ("raccoon eyes" sign)[1] or in the postauricular region due to the bleeding from the mastoid veins (Battle's sign)[5]. Physical examination reveals an external auditory canal laceration[6] or haemorrhage[9-10], an a hemotympanum and a cerebrospinal fluid (CSF) an otorrhea or rhinorrhea, which occurs in 20% of temporal bone fractures[1]. Particularly, concerning the association of craniofacial fractures with external auditory canal bleeding (EACB), Lu et al[4] found EACB with an overall frequency of 7.5% (43 of 573 craniofacial fractures) and investigated the presence of EACB between 4 fracture types (skull base, midface, mandibular with and without involvement of the condyle). Statistical analysis showed that skull base and mandibular intracapsular condylar fractures are the two main causes of EACB, while midface and mandibular fracture cases not involving the condyle are quite rare. Facial nerve paralysis exists in most cases, it is noted immediately and remains permanent unless corrected surgically[9]. Hearing loss, as a result of injury to the tympanic membrane, the middle ear ossicles or the presence of a haematoma, is also a common finding, and can be sensorineural or conductive[3, 10].

The TMJ apparatus is located anterior to the external auditory canal and has two articulating bony components ; the mandibular condyle and the articular eminence and glenoid fossa of the temporal bone. The glenoid fossa is limited posteriorly by the petrotympanic fissure. The posterior part of the glenoid fossa is formed by the anterior tympanic plate, which is thin and weak.

Due to this intimate anatomical relationship, herniation of the TMJ into the EAC occurs spontaneously[11] or secondary to neoplasia, inflammation, developmental problems and especially trauma [11-14]. Direct impact into the chin displaces the condyle posteriorly until the movement is stopped by the articular fossa and the ligaments. The result is a fracture of the condyle or posterior dislocation of intact condyle without fracture[15], a dislocation of condyle into middle cranial fossa[16, 17] or temporal fossa[16] or an injury at the anterior wall of the EAC [5]. By contrast, longitudinal temporal bone fractures are usually associated with posteriosuperior quadrant of the EAC near the tympanosquamous suture[3].

The direction, degree, magnitude and precise point of application on the face and the state of dentition and the occlusal position of the mandible, determine the type of the injury in the temporomandibular region. Also, whether the mouth is open, abnormalities of condylar morphology or the presence of a particularly thin roof of the glenoid fossa affect the condylar dislocation. In our patient, the high-energy nature of the causative force applied in the right parasymphysis region caused a fracture in the anterior wall of the left EAC, without a condylar fracture. The result was hearing loss from the left ear due to EAC obstruction.

Computed tomography is the diagnostic modality of choice for mandibular condylar fractures and injuries in the temporal region[11, 18]. It is important to recognize the potential association between mandibular condylar trauma and temporal bone fractures. In our patient, a CT- examination showed a fracture in the anterior wall of left EAC and a decrease in EAC breadth. This case report emphasizes the need to scrutinize the temporal bone particularly the petrous portion of EAC on CT for fractures in a patient with TMJ injury.

Approaches to the joint include preauricular, postauricular, endaural, rhytidectomal and submandibular. We chose the preauricular approach for TMJ reconstruction of internal derangement and easier exploration and confirmation of the EAC fracture.

The reduction of the fracture was atraumatic and simple by the use of an elevator which was inserted inside the EAC. Primary stability was achieved by wedging of adjacent fracture sites. The postoperative period was uneventful. In the next days after the operation, the hearing loss was restored.

Treatment modalities for EAC, tympanic plate and temporal bone fractures from TMJ herniation, in general include either open or closed reduction of the associated mandibular fractures[8, 19], reconstruction of the anterior canal wall, pain control, management of occlusal discrepancies, and physiotherapy to prevent decreased mandibular range of motion.

In contrast to above mentioned interventional surgical method of treatment, in our case report atraumatic reconstruction of the anterior wall of the EAC and fracture reduction was achieved during TMJ disc reduction and joint exploration, with careful entry of a blunt elevator inside the EAC following the canal outline.

CONCLUSION

In summary, a direct blow to the mandible can result in a TMJ apparatus injury especially on disk . Due to the close relationship between TMJ and EAC, an atypical injury such as a fracture on the anterior wall of EAC can occur. An oral & maxillofacial surgeon, when called to examine and diagnose TMJ injury disorders, has the challenging responsibility to take account of potential concomitant temporal bone fractures or intracranial complications in cooperation with radiologists, ENT doctors and neurosurgeons[20]. In such cases soft atraumatic manipulations at the fractured otic region seems to be the treatment of choice. The successful outcome resulting from our case supports the above mentioned opinion.

160

161

162

163

164

165

166

167 **REFERENCES**


- 168 1. Gladwell M and Viozzi. C. Temporal Bone Fractures: A Review for the Oral and
169 Maxillofacial Surgeon. *Journal of Oral and Maxillofacial Surgery* 2008,66(3):513-522.
- 170 2. Johnson F, Semaan MT, Megerian CA . Temporal bone fracture: evaluation and
171 management in the modern era. *Otolaryngol Clin North Am.* 2008,41(3):597-618
- 172 3. Chong VF and Fan YF . External Auditory Canal Fracture Secondary to Mandibular
173 Trauma: Technical Report. *Clin Radiol.* 2000,55(9):714-716.
- 174 4. Lu C , He D, Yang C . Which craniofacial fractures are associated with external
175 auditory canal bleeding? *J Oral Maxillofac Surg.* 2014,72(1):121-6.
- 176 5. Loh FC, Tan KB, Tan KK . Auditory canal haemorrhage following mandibular condylar
177 fracture *Br J Oral Maxillofac Surg.* 1991,29(1):12-13.
- 178 6. Martis C, Karakasis D. Bleeding from the ear in maxillofacial injuries. *J Maxillofac*
179 *Surg.* 1974,2(2-3):126-8.
- 180 7. Dang D. Bilateral mandibular condylar fractures with associated external auditory
181 canal fractures and otorrhagia. *Radiology case Reports.* 2007,2(1):24-29.
- 182 8. Tao KK, Schwartz DT, Rosh A. . Fracture of the external auditory canal mimicking
183 basilar skull fracture. *J Emerg Med.* 2012,42(2):e39-40
- 184 9. Ulug T and Arif Ulubil S. Management of facial paralysis in temporal bone fractures: a
185 prospective study analyzing 11 operated fractures. *Am J Otolaryngol.*
186 2005,26(4):230-8.
- 187 10. Chujo K, Nakagawa T, Komune S. Temporal bone fracture with ossicular dislocation
188 caused by a blow to the opposite side of the head. *Auris Nasus Larynx.*
189 2008,35(2):273-275.
- 190 11. Moriyama M, Kodama S, Suzuki M. Spontaneous temporomandibular joint herniation
191 into the external auditory canal: a case report and review of the literature.
192 *Laryngoscope.* 2005,115(12):2174-7.
- 193 12. Cillo JE, Sinn DP, Ellis E 3rd. Traumatic dislocation of the mandibular condyle into
194 the middle cranial fossa treated with immediate reconstruction: a case report. *J Oral*
195 *Maxillofac Surg.* 2005,63(6):859-65.
- 196 13. Psillas G and Guyot JP. Hernie spontanée de l'articulation temporomandibulaire dans
197 le conduit auditif externe. *Ann Otolaryngol Chir Cervicofac.* 2007,124(6):305-8.



223

224 **Figure 2: Conventional panoramic view. Note the parasymphyseal fracture on the right**
 225 **and the condylar position on the left (arrows).**

226



227

228 **Figure 3 : Two points zygoma fracture (left).**

229

230




231

232 **Figure 4: The fracture of the left EAC**

233

234



235

236 **Figure 5 : Postoperative view, 2 years after reduction (arrows)**

237

238

239