

SPONTANEOUS PLANTS USED IN THE TRADITIONAL SOAP MAKING IN COTE D'IVOIRE

¹COULIBALY Siendou, ¹OUATTARA Djakalia, ¹KONKON N'dri Gilles,
¹KAGOYIRE Kagohiré, ²*KOUAKOU Tanoh Hilaire

¹Université de Cocody, UFR Sciences Biosciences, Laboratoire de Botanique, 22 BP 1414 Abidjan 22,
Côte d'Ivoire ;

²Université d'Abobo Adjamé, UFR Sciences de la Nature, Laboratoire de Biologie et Amélioration des
Productions Végétales, 02 BP 801 Abidjan 02, Côte d'Ivoire.

*Corresponding author: tanohilaire@yahoo.fr; Tel: +225 03350323; Fax: +225 20378118.

COULIBALY Siendou: Université de Cocody, UFR Sciences Biosciences, Laboratoire de
Botanique, 22 BP 1414 Abidjan 22, Côte d'Ivoire (coulibalysindou457@yahoo.fr)

OUATTARA Djakalia: Université de Cocody, UFR Sciences Biosciences, Laboratoire de
Botanique, 22 BP 1414 Abidjan 22, Côte d'Ivoire (odjakal@yahoo.com)

KONKON N'dri Gilles : Université de Cocody, UFR Sciences Biosciences, Laboratoire de
Botanique, 22 BP 1414 Abidjan 22, Côte d'Ivoire (ndri.gilles@yahoo.com)

KAGOYIRE Kagohiré: Université de Cocody, UFR Sciences Biosciences, Laboratoire de
Botanique, 22 BP 1414 Abidjan 22, Côte d'Ivoire (kagore1962@yahoo.fr)

KOUAKOU Tanoh Hilaire : Université d'Abobo Adjamé, UFR Sciences de la Nature, Laboratoire
de Biologie et Amélioration des Productions Végétales, 02 BP 801 Abidjan 02, Côte d'Ivoire
(tanohilaire@yahoo.fr)

ABSTRACT

The aim of the present study is to create the directory of the main species used in traditional soapmaking deals within the Sudan savanna zone in Center-North of Côte d'Ivoire. From an ethnobotanical approach based on direct structured or semi-structured individual interviews, thirty three plants were recorded, among which seven plants are used for the extraction of the fat (oil, butter) and twenty eight for potash manufacture. The intensive use of the stem (trunk and branches) of these species is a real pressure on the resources itself and may lead to the extinction of the most vulnerable one. Plants inventoried were ranked according to their importance for the soap makers. Several types of plants were identified, from the most known, commonly used and abundant in the study site (*Carapa procera*, *Ceiba pentandra*, *Cussonia arborea*) to the least known, little used and scarce. The valorization of these resources can be beneficial to concerned population, it is urgent to adopt a sustainable management approach for the preservation of the used species.

Keywords: Côte d'Ivoire, Ethnobotany, Plants, Soap, Sudan Savanna.

1. INTRODUCTION

Traditionally, the preparation of soap is made from crude palm oil and potash extracted from wood ashes (Caubergs, 2006). In Europe, before the development of the soap factory, the Gauls and Romans were also making soap at home (Bella, 2005). In Africa, despite progress in the soap industry, people are still using soap made traditionally, which remains cheap and have specific virtues. In Ivory Coast, the manufacture of traditional soaps

is widespread, based on the use of plant species which are mostly spontaneous. It therefore contributes to the vulnerability, the scarcity or even extinction of the concerned species. This practice can impact the local ecosystems and may lead to the extinction of the species (Ambé, 2001).

Presently, few are known about the involved species. The present study aims at creating a directory of the main species exploited in traditional soap making in the Center-North of Côte d'Ivoire (regions of Katiola and Dabakala) in order to insure an economical and ecological sustainable use of these materials.


2. MATERIAL AND METHODS

The study site (regions of Katiola and Dabakala) is located in Center-North of Côte d'Ivoire. The climate is Sudanian with 1053.10 mm of rain per annum.

2.1. Ethnobotanical investigation

The study material is represented by an ethnobotanical investigation form developed after a pre-fieldwork. Fourteen localities were selected (Fig. 1) and the ethnobotanical investigation conducted among 193 people, was done through structured or semi-structured individual direct talks. During the investigation, the informants were asked to answer questions from the survey form. Specimens of plants mentioned were collected with the help of informants for confirmation of the vernacular name and constitution of a reference collection. The determination of species (scientific name) was done at the National Center of Floristic (NCF) of the University of Cocody (Abidjan). The scientific names were also checked using the work of Lebrun and Stork (1991, 1992, 1995, 1997).

The information collected on the survey forms were transferred into a database, processed and analyzed using Microsoft Office Excel 2003.



105

2.2. Estimated frequency of use and level of plants abundance

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

Frequency of use (or local preference) species allows us to assess their level of knowledge and exploitation by the people. It was estimated by the method of the open list ("free-listing") of Cotton (1996). This approach, based on the spontaneous citations, rests on the principle that the most significant species are mentioned by several informants and therefore obtain a high ranking. Frequency of use of each species has been evaluated by its citation frequency spontaneous, according to the following formula:

$$\text{Frequency use of a plant} = \frac{\text{Number of people identifying the plant}}{\text{Number of questioned people}} \times 100$$

The abundance level was estimated for each plant species inventoried by the ratio between the number of people recognizing its abundance in the study site and the total number of people identifying the plant:

$$\text{Level of a plant abundance} = \frac{\text{Number of people recognizing the plant abundance in nature}}{\text{Number of people identifying the plant}} \times 100$$

2.3. Frequency of plant type

Following the same principle of Cotton (1996), the frequency of the origin of each plant species (plant type) was calculated, according to whether the species of plant is cultivated, spontaneous or that it comes from a different region of our study site.

3. RESULTS

3.1. Inventoried plants



At the end of the ethnobotanical investigation, thirty-three plants were recorded. These species were left again in thirty-one genera and twenty-one families (Table 1). According to reports, the traditional soapmaking requires two essential plant raw materials: the fat content (oil, butter) and potash, each coming from specific plant species.

3.2. Plants and bodies used for fat extraction

Seven plants were used for the extraction of the fat content (oil or butter). They belong to five families and five genera (Table 2). The Arecaceae and Euphorbiaceae are the most families, each represented with two plant species. The organ used on these plants is always part of the fruit. It is the seed, almond, endosperm or fruit pulp. Two types of organ are used at *Elaeis guineensis*: pulp and almond.

155
156
157
158

Table 1: Inventory of plants harvested in the regions of Katiola and Dabakala (Center-North of Côte d'Ivoire)

Scientific names	Vernacular names			
	Tagbana	Djimini	Malinké (Djamala)	Baoulé
<i>Acacia polyacantha</i>	Gronhon			
<i>Albizia ferruginea</i>			Samandêrê	
<i>Albizia zygia</i>	Dimakanhan			
<i>Amaranthus spinosus</i>	Noumrinhin			
<i>Anogeissus leiocarpus</i>	Nglaha, Guindihi	Nglaga	Krêkêtê, Tchêgbêlêyiri	
<i>Berlinia grandiflora</i>	Kakpaha	Yaurfougo Krougbêguê	Wôngbê Gbohi, Gbohoul, Kobi, Gbogo, Gbowi	Kohndou
<i>Carapa procera</i>			Banda, Bada	
<i>Ceiba pentandra</i>	Sédihi, Kplaba	Séligué, Cheligué, Segbêguê		Nyin
<i>Cocos nucifera</i>	Kpaco	Kpaco	Kpaco	
<i>Coffea spp.</i>	Café	Café	Café	
<i>Cussonia arborea</i>	Fitio, Foutchouhou, Gbotoho	Yémbetché, Dédégué, Tédégué	Borokourou	
<i>Elaeis guineensis</i>	Hèhètihi	Sindigué	Téhi	Mmé
<i>Ficus capensis</i>		Ndassaga	Toro, tro	
<i>Ficus exasperata</i>	Waha			
<i>Gardenia ternifolia</i>			Blé	
<i>Jatropha curcas</i> 	Kapaha, Kahadja	Kapara	Pôropôro Mangandawa-gbê	
<i>Khaya senegalensis</i>		Wêdigué	Djala	
<i>Manihot esculenta</i>		Gbéndé	Gbéndé	
<i>Musa spp.</i>		Blahnda	Barada	
<i>Parkia biglobosa</i>	Nindihi	Nindigué	Nèrè	
<i>Pentadesma butyracea</i>	Djréhé	Ndjéligué	Gbêlèn	
<i>Piliostigma thonningii</i>	Yoganhan, Yéganhan	Yéwangan	Niaman	
<i>Pterocarpus erinaceus</i>	Nagnranhan	Gbatèlgué	Gbin	
<i>Pupalia lappacea</i>		Sibagniguiré		
<i>Ricinus communis</i> 		Tombotigui	Tombotigui	
<i>Securidaca longepedunculata</i>		Félibé	Djoro, Djohi	
<i>Spondias mombin</i>	Tônanhan	Monlimon		
<i>Sorghum spp.</i>	Gbodioho			
<i>Terminalia glaucescens</i>	Kohotihi, Kohoun	Koman		
<i>Theobroma cacao</i>	Cacao	Cacao	Cacao	
<i>Vitellaria paradoxa</i>	Lodihi, Lohodjal, Lokatchlé	Létigué	Kôrô, Kwèi, Sihé	
<i>Vitex doniana</i>	Hangonhon	Sandjo wôô	Koto	
<i>Zanthoxylum zanthoxyloides</i>		Nganhan	Ngôndô	

Tagbana, djimini, malinké and baoulé are local languages.

159
160
161
162
163

Table 2. Plants used for fat extraction in regions of Katiola and Dabakala (Center-North of Côte d'Ivoire)

Family	Species	Organ of the plant
Arecaceae	<i>Cocos nucifera</i> Linn.	Coconut endosperm)
	<i>Elaeis guineensis</i> Jacq.	Pulp, almond
Clusiaceae	<i>Pentadesma butyracea</i> Sab.	Seeds
Euphorbiaceae	<i>Jatropha curcas</i> Linn.	Almond
	<i>Ricinus communis</i> Linn.	Almond
Meliaceae	<i>Carapa procera</i> CD.	Seeds
Sapotaceae	<i>Vitellaria paradoxa</i> Gaertn.	Almond


164
165 **3.3. Plants and organs used for potash manufacture**
166

167 Table 3 presents the twenty eight plants species for the manufacture of potash. They
168 are divided into nineteen families and twenty six genera. Mimosaceae is the most exploited
169 family with four species. Amaranthaceae, Caesalpiniaceae, Combretaceae, Euphorbiaceae,
170 Moraceae and Rubiaceae families are each represented by two species plants. Among the
171 organs harvested from these plants, the stems (trunk and branches) were largely requested. At
172 the kapok (*Ceiba pentandra*), inside the trunk, the foothills and dried fruits are exploited by
173 the population. Fruits and derivatives (pod, pod, pulp, skin) are also used. In *Amaranthus*
174 *spinosus* and *Pupalia lappacea*, the entire aerial part of the plant (stems, leaves, fruit) is
175 exploited.
176

177 **3.4. Types of species used**
178

179 Among the thirty five plants species inventoried, twenty eight plants are spontaneous
180 (79 %), four plants come from other parts of Côté d'Ivoire (12 %) and only three plants are
181 cultivated in the region (9 %), **Table 3**.

182 The spontaneous plants were cited by 84 % of respondents, while 15% cited only
183 plants from other regions of Côte d'Ivoire. Finally, 1 % of respondents know the
184 involvement of crops in the region in the traditional soapmaking (**Fig. 2**).
185



205
206
207
208
209 **Figure 2. Distribution of frequency citation of plant species according on the plant**
210 **type in regions of Katiola and Dabakala (Center-North of Côte d'Ivoire)**
211
212
213

Table 3. Plants used for the potash manufacture in the regions of Katiola and Dabakala (Center-North of Côte d'Ivoire)

Family	Species	Organ of the plant
Amaranthaceae	<i>Amaranthus spinosus</i> Linn.	Aerial part
	<i>Pupalia lappacea</i> (L.) A.Juss.	Aerial part
Anacardiaceae	<i>Spondias mombin</i> Linn.	Stems
Araliaceae	<i>Cussonia arborea</i> Hochst.ex A. Rich.	Stems
Arecaceae	<i>Cocos nucifera</i> Linn.	Fleshy fruit pulp
Bombacaceae	<i>Ceiba pentandra</i> (L.) Gaertn.	Trunk (buttress), cockles
Caesalpiniaceae	<i>Berlinia grandiflora</i> (Vahl) Hutch. et Dalz.	Pods (cockles)
	<i>Piliostigma thonningii</i> Milne Red.	Stem, fruits
Combretaceae	<i>Anogeissus leiocarpus</i> (DC.) Guill. et Perr.	Stems
	<i>Terminalia glaucescens</i> Planch. ex Benth.	Stems
Euphorbiaceae	<i>Jatropha curcas</i> Linn.	Stems
	<i>Manihot esculenta</i> Crantz.	Tuber peel
Fabaceae	<i>Pterocarpus erinaceus</i> Poir.	Stems
Meliaceae	<i>Khaya senegalensis</i> (Desr.) A.Juss.	Stems
Mimosaceae	<i>Acacia polyacantha</i> var. <i>campylacantha</i> (Hochst.) Roberty	Stems
	<i>Albizia ferruginea</i> (Guill. et Perr.) Benth.	Stems
	<i>Albizia zygia</i> (CD.) J.F.Macbr.	Stems
	<i>Parkia biglobosa</i> (Jacq.) R.Br. ex G.Don	Stems
	<i>Ficus capensis</i> Thunb.	Stems
Moraceae	<i>Ficus exasperata</i> Vahl	Stems
Musaceae	<i>Musa</i> spp.	Tuber peel
Poaceae	<i>Sorghum</i> spp.	Stems
Polygalaceae	<i>Securidaca longepedunculata</i> Fres.	Stems
Rubiaceae	<i>Coffea</i> spp.	Fruit pulp
	<i>Gardenia ternifolia</i> Schum. et Thonn.	Stems
Rutaceae	<i>Zanthoxylum zanthoxyloides</i> (Lam.) Zep. et Tim.	Stems
Sterculiaceae	<i>Theobroma cacao</i> Linn.	Cockles (+ rachis)
Verbenaceae	<i>Vitex doniana</i> Sweet.	Stems

3.5. Frequency of use and abundance level of plants used for fat extraction

Figure 3 shows the frequency of use of oilseeds and their abundance level in the place. It appears that plants *Carapa procera* is the most used plant, with 79.48 % of the citations. The shea tree (*Vitellaria paradoxa*) and palm oil (*Elaeis guineensis*), with quotations




Figure 3. Frequency of use and level of abundance of the oleaginous plants used for the soapmaking in the regions of Katiola and Dabakala (Center-North of Côte d'Ivoire)

oilseed species are very well represented in the site, according to informants. However, three trends emerged. First, plants that are very abundant with over 90 % level of abundance; it is *Jatropha curcas* (100 %), *Ricinus communis* (100 %) and *Vitellaria paradoxa* (90.27 %). Then, the fairly abundant plants (60-90 %) which are *E. guineensis* (73.84 %), *Carapa procera* (68.38 %) and *Cocos nucifera* (66.66 %). Only *Pentadesma butyracea* is not very abundant in the area with 33.33 % level of abundance.

3.6. Frequency of use and abundance level of plants species used for potash manufacture

Among the twenty eight plants identified for the manufacture of potash (**Table 4**), only two plants are distinguished (**Fig. 4**). These are *Ceiba pentandra* and *Cussonia arborea* with 51.28 % of all citations. The others twenty six plant species are very slightly mentioned (frequency <9%). Their abundance level, meanwhile, is very high (> 90 %) or high (50-90 %) for almost all inventoried plants. However, both species plants are characterized by very low presence or absence in the region: *Theobroma cacao* (13.33 %) and *Coffea* spp. (0 %).

3.7. Classification of species plant recorded

To present the results, all taxa inventoried were grouped according to the combination with the frequency criteria of use - abundance level (**Table 5**). Abundant species are numerous (22), but only *Carapa procera* was requested much. *Ceiba pentandra*, *Cussonia arborea*, *Elaeis guineensis* and *Vitellaria paradoxa* were moderately used.




Figure 4. Frequency of use and abundance level of used plants for potash manufacture in soapmaking in the regions of Katiola and Dabakala (Center-North of Côte d'Ivoire)

Table 4. Inventoried plants according to their origin in the regions of Katiola and Dabakala (Center-North of Côte d'Ivoire)

Spontaneous plants	Cultivated plants	Plants from other regions
<i>Acacia polyacantha</i>	<i>Manihot esculenta</i>	<i>Cocos nucifera</i>
<i>Albizia ferruginea</i>	<i>Musa spp.</i>	<i>Coffea spp.</i>
<i>Albizia zygia</i>	<i>Sorghum spp.</i>	<i>Elaeis guineensis</i>
<i>Amaranthus spinosus</i>		<i>Theobroma cacao</i>
<i>Anogeissus leiocarpus</i>		
<i>Berlinia grandiflora</i>		
<i>Carapa procera</i>		
<i>Ceiba pentandra</i>		
<i>Cussonia arborea</i>		
<i>Ficus capensis</i>		
<i>Ficus exasperata</i>		
<i>Gardenia ternifolia</i>		
<i>Jatropha curcas</i>		
<i>Khaya senegalensis</i>		
<i>Parkia biglobosa</i>		
<i>Pentadesma butyracea</i>		
<i>Piliostigma thonningii</i>		
<i>Pterocarpus erinaceus</i>		
<i>Pupalia lappacea</i>		
<i>Ricinus communis</i>		
<i>Securidaca longepedunculata</i>		
<i>Spondias mombin</i>		
<i>Terminalia glaucescens</i>		
<i>Vitellaria paradoxa</i>		
<i>Vitex doniana</i>		
<i>Zanthoxylum zanthoxyloides</i>		

Table 5. Inventoried plants for the potash manufacture in the regions of Katiola and Dabakala (Center-North of Côte d'Ivoire)

Plants used in potash manufacture		
1- <i>Ceiba pentandra</i>	11- <i>Terminalia glaucescens</i>	21- <i>Gardenia ternifolia</i>
2- <i>Cussonia arborea</i>	12- <i>Acacia polyacantha</i> var.	22- <i>Pupalia lappacea</i>
3- <i>Anogeissus leiocarpus</i>	13- <i>Parkia biglobosa</i>	23- <i>Securidaca L.</i>
4- <i>Theobroma cacao</i>	14- <i>Pterocarpus erinaceus</i>	24- <i>Ficus capensis</i>
5- <i>Jatropha curcas</i>	15- <i>Albizia zygia</i>	25- <i>Ficus exasperata</i>
6- <i>Piliostigma thonningii</i>	16- <i>Amaranthus spinosus</i>	26- <i>Khaya senegalensis</i>
7- <i>Vitex doniana</i>	17- <i>Cocos nucifera</i> (potasse)	27- <i>Manihot esculenta</i>
8- <i>Coffea</i> spp.	18- <i>Zanthoxylum zanthoxyloides</i>	28- <i>Sorghum</i> spp.
9- <i>Musa</i> spp.	19- <i>Berlinia grandiflora</i>	
10- <i>Spondias mombin</i>	20- <i>Albizia ferruginea</i>	

4. DISCUSSION

Evaluation of plants devoted to traditional soapmaking in Sudanese region (Katiola and Dabakala) of Côte d'Ivoire showed a limited number of plants involved. Ethnobotanical studies conducted with methods similar to ours have an inventory of sixty fifteen wild edible fruit in the region of Seguela (Ambe, 2001), and forty two medicinal plants used by rural residents of (Mehdioui and Kahouadji, 2007). The low number of plants species in this study (33) could be explained either by the restriction of our study area (traditional soap making), the tendency of people (mostly young) to shift away from traditional practice in benefits of modernity, and the reduction of plant resources in the study zone.

Regarding plant organs, the seeds and almonds used for the fat extraction seems to be satisfactory, because these organs are renewable following a regular physiological cycle. Their harvest does not affect the physical integrity of the plants. On the other hand, the almost exclusive use of stem (trunk and branches) for potash manufacture is particularly worrying. Indeed, many plants are often completely slaughtered and incinerated to obtain a good amount of ash, thus potash. This method of harvest puts in danger the survival of exploited plants species. According to Cunningham (1996), there is a clear relationship between the traveled portion of the used plant and the effects of harvesting on its existence. The use of buttresses of *Ceiba pentandra* in potash manufacture was also reported by Tra Bi (1997) in populations living to the neighbourhoods of the classified forests of Haut-Sassandra and Scio (West of Côte d'Ivoire). It was noted that the exploited species are primarily spontaneous, from the savannah. Mehdioui and Kahouadji (2007) explain that the extensive use of local species is due to the proximity of these plants. The limited number of plants used as source of fat (7 of 33 species) might suggest a limitation of oleaginous species in the environment, their ignorance by the people. The difficulties of extracting vegetable oil could also justify the neglect of women with respect to the oleiferous species. This would explain the fact that many oleiferous species quoted in the literature and in the region have not been reported by informants. It is obvious that the valuation of these unused resources can contribute to reduce the present pressure on species already exploited, but also to help

320 traditional soap makers to preserve this ancestral practice. Among the oil crops in the region,
321 *Carapa procera* is the most preferred and used in soapmaking. The best quality of soaps
322 made from oil of this plant and its non-use of this oil for food (because of its bitterness) as
323 well as its therapeutic uses seem to support this preference.

324 Although moderately represented in the region, *C. procera* is known as an ecological
325 restriction. The plant is essentially restricted to gallery forests. The exploitation of lowland
326 rivers and rice is a serious threat to the survival of these species. It is the same for
327 *Pentadesma butyracea* which remains very poorly represented in the study zone because
328 limited to only a few shallows. The unavailability of the last species of plant and its oil very
329 appreciated in the food, seem to justify its weak use in the traditional soap manufacture.

330 These two species plants (*C. procera* and *P. butyracea*) therefore deserve special
331 attention, and can allow their backup or upgrading their local populations. According to
332 Ambe (2001), particular attention should be given to relatively few products consumed
333 today, the latter being for a loss with more or less rapid destruction of their habitat. In
334 addition, Miralles (1983) reported the high content of oleic acid seeds of *Carapa procera*,
335 which could justify its valuation. Thus, the domestication of these plants or their retention in
336 agriculture could be considered as alternatives. Some plants that are well known and used,
337 termed sub-spontaneous, such as *Vitellaria paradoxa*, *Parkia biglobosa* (Ambé, 2001) and
338 *Elaeis guineensis* are already protected by the local population. The threat on *Carapa procera*
339 and *Pentadesma butyracea* is exclusively linked to their habitat (threatened and restricted)
340 and not their exploitation. This explains the mention of *C. procera*, in addition to *P.*
341 *biglobosa* and *Spondias mombin* among the underutilized plant species from Côte d'Ivoire
342 and other parts of Africa (Ahoussou et al., 1995).

343 *Ceiba pentandra* and *Cussonia arborea* were the most representative plants for potash
344 manufacture. If the first specie is very abundant in West Africa (Siepel et al., 2004) as well
345 as in the study site and does not run of risk of imminent extinction, it is the opposite for the
346 second species plant. Indeed, *C. arborea* is dominant in the landscape, i.e. the open savanna
347 vegetation. However, savannahs ecosystems are the most destroyed by agricultural clearing.
348 At this threat, should be add the systematic demolition of the plant for potash manufacture
349 used in traditional soaps making. On the field investigation, we noticed that the plants species
350 is present in the savannah far away from settlements. It is also used for medicinal purposes
351 against conjunctivitis, leprosy, diarrhea and dysentery (Ahoussou et al., 1995). The
352 preferential use of *Ceiba pentandra* and *Cussonia arborea* for making potash could be related
353 to the quality and efficiency of potash compared to other plants. This is even true that many
354 plants are abundant in the site or are very little used for potash manufacture. Therefore, face
355 to the threats on plant species in the region, it is urgent to adopt a sustainable management
356 approach for the safeguarding and preservation of the most exploited plants species. Thus, in
357 addition to the domestication suggested for some plants, the identification of gallery forests
358 portion as biological and ecological interest sites, may be an interesting approach. We note
359 that none of the plants surveyed were among the endemic, rare or endangered in Côte d'Ivoire
360 and West Africa (Ahoussou et al., 1995, Poorter et al., 2004).

362 5. CONCLUSION

364 This study has established a list of wild species used in the traditional soap making in
365 Sudanese zone of Côte d'Ivoire, and evaluated the relative importance of these plants. Thirty
366 three plant species have been recorded among which seven plants are used for the fat
367 extraction and twenty eight plants for potash. Two plants species are being used at a time for
368 fat and potash. The relative importance of each plant species permitted to identify different
369 categories of plants. The most exploited plants are *Carapa procera*, *Vitellaria paradoxa* and

Elaeis guineensis for their fat content and *Ceiba pentandra* and *Cussonia arborea* for potash manufacture. Otherwise, the great majority of plants recorded are spontaneous and abundant in the study site, except for *Pentadesma butyracea* and plants from other regions. Harvesting the trees by stamping out is the most practiced method of collecting. Maintaining all these resources being illusory, identification of the products is more or less easy to value their useful approach. Thus, taking into account their frequency of use by people, their abundance level, three wild species were proposed for possible revalorization. These are *C. procera*, *P. butyracea* and *C. arborea*.

REFERENCES

- Ahoussou, N., Koffi, G., Sangaré, A. (1995).** Côte d'Ivoire: Rapport de pays pour la Conférence technique internationale de la FAO sur les ressources phytogénétiques. Leipzig, Allemagne, 17-23 juin 1995. 75p.
- Ambé, G.A. (2001).** Les fruits sauvages comestibles des savanes guinéennes de Côte d'Ivoire : état de la connaissance par une population locale, les Malinké. Biotechnol. Agron. Soc. Environ., 5, 43-58.
- Bella, O.M. (2005).** Comment fabriquer son savon à domicile? <http://saponification.afrikblog.com/>.
- Caubergs, L. (2006).** La fabrication du savon : Aspects techniques, économiques et sociaux. ATOL, Leuven, Belgique, 97p.
- Cotton, CM. 1996.** Ethnobotany: Principles and Applications. Chichester, UK: Wiley.
- Cunningham, AB. 1996.** Peuples, parc et plantes. Recommandations pour les zones à usages multiples et les alternatives de développement autour du parc national de Bwindi Impénétrable, Ouganda. Documents de travail Peuples et Plantes n°4. UNESCO, Paris, 66p.
- Lebrun, J.P., Stork, A.L. (1991).** Enumération des plantes à fleurs d'Afrique Tropicale : Annonaceae à Pandaceae. CJB, Genève (Suisse), 249p.
- Lebrun, J.P., Stork, A.L. (1992).** Enumération des plantes à fleurs d'Afrique Tropicale: Chrysobalanaceae à Apiaceae. CJB, Genève (Suisse), 257p.
- Lebrun, J.P., Stork, A.L. (1995).** Enumération des plantes à fleurs d'Afrique Tropicale. Monocotylédones : Limnocharitaceae à Poaceae. CJB, Genève (Suisse), 314p.
- Lebrun, J.P., Stork, A.L. (1997).** Enumération des plantes à fleurs d'Afrique Tropicale. Gamopétales: Ericaceae à Lamiaceae. CJB, Genève (Suisse), 712p.
- Mehdioui, R., Kahouadji, A. (2007).** Etude ethnobotanique auprès de la population riveraine de la forêt d'Amsittène : cas de la Commune d'Imi n'Tlit (Province d'Essaouira). Bull. Sci. Vie, 29, 11-20.
- Miralles, J. (1983).** Recherche de nouvelles ressources en huiles végétales. Oléagineux, 38, 665-667.
- Poorter, L., Bongers, F., Kouamé, F.N., Hawthorne, W.D. (2004).** Biodiversity of West African Forests. An Ecological Atlas of Woody Plant Species. CABI Publishing.
- Siepel, A., Poorter, L., Hawthorne, W.D. (2004).** Ecological profiles of large timber species, in: Poorter, L., Bongers, F., Kouamé, F.N., Hawthorne, W.D. (Eds), Biodiversity of

419 West African Forests. An Ecological Atlas of Woody Plant Species, CABI Publishing, pp.
420 391-446.

421 **Tra Bi, F.H. (1997).** Utilisations des plantes, par l'homme, dans les forêts classées du Haut-
422 Sassandra et de Scio, en Côte d'Ivoire. Thèse de doctorat de troisième cycle, UFR
423 Biosciences, Université de Cocody-Abidjan, Côte d'Ivoire, 215p.

424

425

426

427

UNDER PEER REVIEW