

2 **DETERMINATION OF MINERALIZATION RATE OF ORGANIC**
3 **MATERIALS USING CARBON DIOXIDE EVOLUTION AS AN INDEX**

4
5 **¹Makinde E.A. and ²L. S. Ayeni**

6 ¹Department of Botany, Lagos State University, Ojo Lagos

7 ²Department of Agricultural Science, Adeyemi College of Education, Ondo, Ondo State Nigeria

8 esthermak2005@yahoo.com

9 Author for correspondence: ²*leye_sam@yahoo.com

10
11
12
13
14 **ABSTRACT**

15
16
17
18
19
20
21
22
23
24
25
A study was conducted on a sandy loam soil to determine the rate of CO₂ release by Kola Pod Husk (KPH) and Pacesetter Grade B (PGB) (sorted city waste plus cow dung) in southwest Nigeria. Each of KPH and PGB was applied at 0.25 g to 50 g soil; and control without treatment application was incubated for 16 weeks. The treatments were replicated four times on a completely randomized design. Evolution of CO₂ by all the treatments increased as the period of incubation increased from the first week to the 5th week of the experiment. After the 6th week, PGB decreased CO₂ at 7th and 8th week and increased it between 9th and 11th week and thereafter finally decreased it as incubation period progressed. KPH decreased CO₂ between 7th and 8th week and then increased it from 9th - 11th week before the CO₂ finally declined till the termination of the experiment. Compared with control, KPH and PGB significantly ($P < 0.05$) increased CO₂ evolution. The rate of mineralization in the first 1-7 weeks of incubation was in the order of PGB > KPH > control, while the last 12-16 weeks of incubation was in the order of KPH > PGB > control. Pacesetter Grade B reached its peak of CO₂ evolution at 9th week of incubation while KPH reached its peak at 13th week of incubation. **Grade B pacesetter had the highest CO₂ emission.**

16
17 *Keywords:* Kola pod husk, Pacesetter Grade B Fertilizer, organic carbon, soil nutrients

18
19
20
21 **1. INTRODUCTION**

22 Most Nigerian soils are low in native nutrient content and soil organic matter. They
23 are however high in clay content of Kaolinitic (1:1) type (Ogunwale *et al.*, 2002). The soils are
24 hence low in cation exchange capacity and are not able to retain adequate amount of
25 nutrients, they therefore require split application of fertilizers to be able to support good crop

26 growth (Agboola *et al.*, 1981). The procurement costs of the fertilizers are higher than what
27 an average Nigerian farmer can afford (Agbede and Kalu, 1995). The few rich farmers that
28 could afford the procurement of the fertilizer could not get adequate quantity from the
29 market. Most of the raw materials needed for local production are imported at exorbitant cost
30 (Fagbenro and Agboola, 1983). In addition, there has been recent clamour for organic foods
31 and agricultural products in the world market (Pawar *et al.*, 2003). The need to increase the
32 soil organic matter content for sustainable Nigerian agricultural soils (Lombin, 1981;
33 Ogunwale *et al.*, 2002) coupled with the problems above has called for a shift from the use
34 of inorganic fertilizer to the use of organic fertilizers where organic fertilizers are in abundant.

35 Organic materials are capable of promoting crop growth and increasing yield by
36 way of improving soil physical, chemical and biological properties (Titiloye *et al.*, 1985;
37 Wallace, 1994). Organic fertilizers improve the physical properties of soils; maintain the
38 soils in better tilth with; increases water holding capacity (Agboola and Omuetti, 1982; Lal,
39 1986; Ogunwale *et al.*, 2002) and supplies both major and minor plant nutrients (Ayeni,
40 2011, Ayeni *et al.*, 2008). The supplied nutrients can substitute for appreciable amounts of
41 inorganic fertilizer (Tollesa, 1999).

42 Over the years, various organic materials have extensively been used as fertilizers
43 and their beneficial effects documented. They have subsequently been recommended as
44 sources of nutrient supply to farm crops. It is observed that since the introduction of
45 inorganic fertilizers into the Nigerian agriculture, in-depth research into the use of organic
46 fertilizer as sources of plant nutrients for *i.e.*, the rate of decomposition of organic materials
47 has not been adequately determined. Although some information on the mineralization of
48 compost is available as reported by Van De Kerkove (1990) and cited by Jedidi *et al* (1993),
49 but there is scanty information on the rate of Kola pod husk and Pace setter organic fertilizer
50 in south western Nigeria. This study was therefore set up to compare the rate of
51 decomposition of industrially manufactured organic fertilizer called Pacesetter Grade B and
52 Kola Pad Husk, using CO₂ evolution as an index.

53

54 **2. MATERIAL AND METHODS**

55

56 Kola Pod Husks (KPH) and Pacesetter Grade B fertilizer (non fortified sorted city
57 wastes plus cow dung, PGB) were used for the conduct of the experiment. The KPH was
58 obtained from the Kola processing unit of Cocoa Research Institute of Nigeria (CRIN) and
59 PGB fertilizer was obtained from the Pacesetter Fertilizer Plant at Bodija, Ibadan Nigeria.
60 Kola Pod Husk (KPH) was sun dried to constant weight and milled to pass through 2 mm
61 sieve before analysis.

62

63 The CO₂ evolution as index method was used to compare the mineralization
 64 potential of the various amendments. The carbon dioxide evolution study was carried out
 65 according to the procedure described by Moorhead *et al.* (1999) and adopted by Ipinmoroti
 66 *et al.* (1997). Fifty grams of 2 mm sieved soil were weighed into each of the 12 incubation
 67 flasks, with four flasks representing each of the two organic materials and the control without
 68 organic material addition. Each of the milled organic materials was weighed and mixed with
 69 soil in the flasks at the rate equivalent to 10 tonnes ha⁻¹ (i.e. 0.25 g per 50 g soil). Each
 70 treatment was replicated four times. The treated soil in each flask was moistened to 70 %
 71 field capacity of the soil and incubated in the laboratory at temperature of 28°C. Absorbent
 72 was used to cover the mouth of each flask in order to reduce evaporation or gaseous
 73 escape. The moisture content was adjusted fortnightly with de-ionized water. The carbon
 74 dioxide evolved from the flasks was collected in a bottle containing 25 ml of 0.1 M Ca (OH)₂.
 75 The amount of carbon dioxide evolved was determined by titration with 0.05 M HCl, using
 76 phenolphthalein as indicator.

77 The amount of carbonate evolved was calculated using the following equation:

$$\begin{aligned}
 78 \text{ meq of CO}_2 &= 0.2727(25 - (\text{titre} \times f) \times 0.027 \\
 79 f &= \frac{\text{volume of Ca(OH)}_2}{\text{Blank titre}} \\
 80 & \\
 81 \text{ meq of C} &= \text{equivalent weight of C} \times \text{Molarity of Ca (OH)}_2 \\
 82 \text{ Where } 0.2727 &= \text{ratio of carbon in carbon dioxide} \\
 83 \text{ 25ml} &= \text{Volume of Ca (OH)}_2 \text{ in the flask/bottle}
 \end{aligned}$$

85 The soil pH was determined in 1:2.5 soil/water ratio and read with pH meter. Total N
 86 was determined by the normal Microckjedahl method soil OC was determined by wet
 87 dichromate oxidation method (Jackson, 1958). Available P was determined by Bray -1-
 88 method. Exchangeable K, Ca, Na and Mg were extracted with 1N ammonium acetate at pH
 89 7. Exchangeable K was determined by flame photometer while Ca and Mg were determined
 90 by Atomic Absorption Spectrophotometer (AOAC, 1990) Exchangeable acidity was extracted
 91 by 0.1M KCl before titrated with 0.1M HCl. The micronutrients (Mn, Fe, Cu and Zn) were
 92 extracted with HCl and determined by AAS. The ECEC was determined by the summation of
 93 the cations. Ground KPH was ashed in muffle furnace before digested with mixture of nitric-
 94 sulphuric- perchloric acid for the extraction of P, K, Ca and Mg (A O A C, 1990). Nitrogen
 95 was determined by the normal microckjedahl method

96
 97 Data obtained were analysed using ANOVA. Least Significant Difference (LSD) was
 98 employed to separate the differences among the treatments at P < 0.05.

99 **3. RESULTS AND DISCUSSION**

100

101 Kola Pod Husks (KPH) and Pacesetter Grade B fertilizer (non fortified sorted city
102 wastes plus cow dung, PGB) were used for the conduct of the experiment. The KPH was
103 obtained from the Kola processing unit of Cocoa Research Institute of Nigeria (CRIN) and
104 PGB fertilizer was obtained from the Pacesetter Fertilizer Plant at Bodija, Ibadan Nigeria.
105 Kola Pod Husk was sun dried to constant weight and milled to pass through 2 mm sieve
106 before analysis.

107

108 The CO₂ evolution as index method was used to compare the mineralization
109 potential of the various amendments. The carbon dioxide evolution study was carried out
110 according to the procedure described by Ipinmoroti *et al* (1997). Fifty grams of 2 mm sieved
111 soil were weighed into each of the 12 incubation flasks, with four flasks representing each of
112 the two organic materials and the control without organic material addition. Each of the
113 milled organic materials was weighed and mixed with soil in the flasks at the rate equivalent
114 to 10 tonnes ha⁻¹ (i.e. 0. 25 g per 50 g soil). Each treatment was replicated four times. The
115 treated soil in each flask was moistened to 70 % field capacity of the soil and incubated in
116 the laboratory at temperature of 28°C. Absorbent was used to cover the mouth of each flask
117 in order to reduce evaporation or gaseous escape. The moisture content was adjusted
118 fortnightly with de-ionized water. The carbon dioxide evolved from the flasks was collected in
119 a bottle containing 25 ml of 0.1 M Ca (OH)₂. The amount of carbon dioxide evolved was
120 determined by titration with 0.05 M HCl, using phenolphthalein as indicator.

121 The amount of carbonate evolved was calculated using the following equation:

122 meq of CO₂ = 0.2727(25 – (titre x f) 0.027
123 f = $\frac{\text{volume of Ca(OH)}_2}{\text{Blank titre}}$
124 Blank titre
125 meq of C = equivalent weight of C Molarity of Ca (OH)₂
126 Where 0.2727 = ratio of carbon in carbon dioxide
127 25ml = Volume of Ca (OH)₂ in the flask/bottle

128

129 The soil pH was determined in 1:2.5 soil/water ratio and read with pH meter. Total N
130 was determined by the normal Microckjedahl method soil OC was determined by wet
131 dichromate oxidation method (Jackson, 1958). Available P was determined by Bray -1-
132 method. Exchangeable K, Ca, Na and Mg were extracted with 1N ammonium acetate at pH
133 7. Exchangeable K was determined by flame photometer while Ca and Mg were determined
134 by Atomic Absorption Spectrophotometer (AOAC, 1990) Exchangeable acidity was extracted

135 by 0.1M KCl before titrated with 0.1M HCl. The micronutrients (Mn , Fe, Cu and Zn) were
 136 extracted with HCl and determined by AAS. The ECEC was determined by the summation of
 137 the cations. Ground KPH was ashed in muffle furnace before digested with mixture of nitric-
 138 sulphuric- perchloric acid for the extraction of P, K, Ca and Mg (A O A C, 1990). Nitrogen
 139 was determined by the normal microkjedahl method

140

141 Data obtained were analysed using ANOVA. Least Significant Difference (LSD) was
 142 employed to separate the differences among the treatments at $P < 0.05$.

143 **RESULT AND DISCUSSION**

144 The initial properties of the soil used for the incubation study (Table 1) indicated that
 145 the soil was slightly acidic, low in total N, C/N ratio, K, Ca and Mg (Agboola *et al.*, 1980).
 146 The soil was adequate in available P, Fe and Mn. This indicates that the soil is poor in plant
 147 nutrients, thus; the soil needs fertilization

148 **Table 1: Some physical and chemical characteristics of soil used in the study**

Parameters	Soil
pH (H ₂ O)	5.3
Total N (g kg ⁻¹)	1.3
Organic Carbon (g kg ⁻¹)	10.5
C/N ratio	7.7
Available P (mg kg ⁻¹)	8.3
Exchangeable bases (c mol kg⁻¹)	
K	0.2
Ca	0.2
Mg	0.2
ECEC	0.7
H + Al	0.1
% Base Saturation	84.5
Micronutrients (mg kg⁻¹)	
Mn	
Fe	21.0
Cu	11.0
Zn	1.5
Particle size analysis (g kg⁻¹)	
Sand	
Silt	
Clay	912
Texture Class	54
	34
	Sandy Loam

149

150 **Table 2: Chemical properties of PGB and KPH**

Nutrients	Pacesetter Grade B (PGB)	Kola Pod Husk (KPH)
pH (H ₂ O)	6.5	6.8
C:N	13.19	24.8

%		
N	1.46	1.06
C	19.55	26.05
P	0.92	1.11
K	5.83	7.65
Ca	0.33	0.38
Na	0.15	0.2
Mg	0.28	0.28
mgkg ⁻¹		
Zn	10.4	11.0
Cu	1.9	2.0
Mn	30.0	31.0
Fe	11.0	11.1

151

152 In figure 1, evolution of CO₂ by the control experiment, KPH and PGB increased as
 153 the period of incubation increased from the first week up to the fifth week of the experiment.
 154 The control experiment decreased the volume of CO₂ produced as from the sixth week to the
 155 16th week when the experiment was terminated.

156 There was reduction in the volume of CO₂ produced by the PGB Fertilizer at 7th and
 157 8th week of incubation when compared with the rate at which CO₂ was released between 1st
 158 and 6th week of incubation. There were increases in the evolution of CO₂ at 9, 10 and 11th
 159 week of incubation in the soil samples treated with PGB but at gradual rate when compared
 160 to the rate CO₂ evolution at the 8th week of incubation. Also, PGB gradually decreased the
 161 volume of CO₂ evolved as from 9th week till the termination of the experiment. Kola Pod
 162 Husk exhibited slight different characteristics in releasing CO₂ to the soil compared with
 163 control experiment and PGB. Kola Pod Husk increased the volume of CO₂ released as from
 164 the 1st week of incubation to the 6th week, decreased CO₂ evolution between 7th and 8th
 165 week and then increased CO₂ evolution from the 9th week of incubation to the 11th week
 166 before the CO₂ evolved finally declined till the termination of the experiment.

167 Compared with control, PGB significantly increased ($P < 0.05$) CO₂ evolution from the
 168 1st week of the experiment to the 11th week (except 8th week) of incubation. Also, compared
 169 with control, KPH significantly ($P < 0.05$) increased CO₂ evolution from 10th week till the
 170 termination of the experiment (Fig. 1)

171 This work shows that the rate at which PGB and KPH released CO₂ was different
 172 from each other. The percentage change in the volume of CO₂ evolved during the incubation
 173 process by PGB and KPH showed that PGB had higher increase in CO₂ at the earlier stage
 174 of the experiment (1-6 weeks) while KPH had higher increase in CO₂ at the later end of the
 175 experiment. This shows that PGB tended to increase the rate of carbon mineralization than
 176 KPH at early stage of incubation.

177 Through out the period of the incubation, the values of CO₂ in the control experiment
178 were lower than the soil samples amended with PGB and KPH. This shows that the
179 treatments that were applied to the soil samples actually influenced the evolution of CO₂ in
180 the experiment.

181 The results of this experiment showed that the incorporation of the amendments into
182 the soil might have significantly increased the biological activities which could be categorized
183 into three stages (Ayeni, 2011). A stage of intense activity at the beginning of incubation (1-
184 6 weeks) caused by rewetting of the soil plus amendment mixtures, corresponding to the use
185 of the easily metabolized C present in the PGB, KPH and the native soil. Secondly, a stage
186 of reduced activity (7-8 weeks) characterized by a drop in CO₂ as a result of the decreased
187 amount of easily biodegradable organic matter and lastly, a stage of moderate stable activity
188 between 9-16 weeks showing that the decomposition has reached advance stage.

189 The higher CO₂ released by KPH and PGB over the control indicated higher microbial
190 activities in these materials as reported by Ayeni (2011) and Leslie (2002).

191




Figure: Time course of CO₂ evolution (mg kg⁻¹) by kola pod husk (KPH) and pacesetter organomineral fertilizer (PGB)

192

193 The reduction in CO₂ values after the initial upsurge might be caused by exhaustion
194 of the readily oxidized labile contents of the various media which include sugar, starch and
195 cellulose as reported by Obatolu (1991). However, the decomposition of the high molecule

196 carbohydrates and lignin thereafter must have resulted in the second phase of increase in
197 CO₂ values thereafter. This supports the report by Olayinka and Adebayo (1987) that CO₂
198 release increases over a period of time with organic materials. The relative decrease and the
199 subsequent stable values across the various media at the latter weeks (week 12 - 16)
200 confirmed the report of Ayeni (2011) that decomposition of the organic materials had
201 reached advanced stage and their nutrient contents could be made available for plant use.

202 From the results of this study, it could be deduced that mineralization of organic
203 materials in the soil depends upon the type of organic material. Also, the rate of
204 mineralization is faster in the mixture involving city waste materials (PGB) with lower C/N
205 ratio and than in KPH, while the reverse was the case at the latter stage of the incubation.
206 The comparison of the two amendments using CO₂ evolution as an index method showed
207 that PGB was more stable than KPH at the initial stage. Arable crops such as leaf
208 vegetables that could complete their life cycles within eight weeks would benefit more from
209 PGB than KPH due to their nutrient releases patterns, while crops with longer life cycles
210 would benefit more from the use of KPH as fertilizer.

211 **4. CONCLUSION**

212
213 Experiment conducted to show the rate of CO₂ release by kola pod husk and organic
214 fertilizer called Pacesetter Grade B showed that both treatments increased CO₂ at different
215 rates.

216 **COMPETING INTERESTS**

217 Authors have declared that no competing interests exist

218

219 **AUTHORS' CONTRIBUTIONS**

220

221 Dr (Mrs) E.A Makinde designed the study and carried out the incubation study while Dr L.S.
222 Ayeni analyzed the data and prepared the manuscript

223 **REFERENCES**

- 224 [1] Agboola, A A. J.A.I. Omueti and O. Titiloye. 1981. Chemical composition of
225 industrial and agricultural waste products containing water sources. *Proceeding of*
226 *2nd National Conference on Water Pollution*: pp. 198-220
- 227 [2] Agbede, O. O. and B. A. Kalu. 1995. Constraints of small- scale farmers in
228 increasing crop yields: farm size and fertilizer supply. *Nigeria Journal of Soil*
229 *Science*. 11: 139-159.
- 230 [3] Fagbenro, J. A and A A. Agboola. 1983. Effect of different levels of humic acid on
231 the growth and nutrient uptake of Teak seedlings. *Journal of Plant Nutrition*. Vol.
232 16: Pp. 1465-1483.
- 233 [4] Pawar, V.M., M.K.V. Parbhani, S.N. Puri and M.O., K.V. Rahuri. 2003. Organic
234 farming. In *Organic Production*. Pp. 1 – 12.

- 235 [5] Lombin, L. G. 1981. Approximating the potassium fertilization requirement of cotton
236 on some representative semi-arid tropical savannah soils of Nigeria *Canadian*
237 *Journal of Soil Science*, 61: 507-516.
- 238 [6] Ogunwale, J.A, J.O. Olaniyan and M.O. Aduloju. 2002. Morphological, physio-
239 chemical ;and clay mineralogical properties of soils overlaying basement complex
240 rocks in Ilorin East, Nigeria. *Moor Journal of Agricultural Research.*, 3.2: 147 - 154.
- 241 [7] Titiloye, E.O., A.A., Agboola and E.O. Lucas. 1985. Evaluation of fertilizer values of
242 organic waste materials in southwestern Nigeria. *biological Agriculture and*
243 *Horticulture* 3:25-37.
- 244
- 245 [8] Wallace, A. 1994. Soil Organic Matter as Essential Material; in Solving
246 Environmental Problems Com. *Soil Science and Plant Analysis* 25: 1-2.
- 247 [9] Agboola A.A and J. A.I., Omueti. 1982. Soil fertility problems and its management
248 in tropical Africa. In: International conference on land clearing and development.
249 *Proceedings 2: IITA, Ibadan, Nigeria.*
- 250 [10] Lal, R 1986. A Soil surface management in the Topics for Intensive Land use and
251 Sustained Production. *In: Advance in soil science*, 5: 1-108.
- 252 [11] Ayeni, L.S., (2011). Cumulative effect of combined cocoa pod ash, poultry manure,
253 NPK 20:10:10 fertilizer on major cations release for crop production in southwestern
254 Nigeria. *Int. J. Agric. Soil Sci.*, 1(7): 248-253
- 255 [12] Ayeni, L.S., M.T. Adetunji, S.O. Ojeniyi, S.B. Ewulo and A.J. Adeyemo, (2008).
256 Comparative and cumulative effect of cocoa pod husk ash and poultry manure on
257 soil and nutrient contents of maize yield. *American – Eurasian Journal of Sustain.*
258 *Agric.* 2(1): 92-97.
- 259 [13] Tollesa, D. 1999. Effect of organic and inorganic fertilizers on maize grain yield in
260 western Ethiopia. *African Crop Science Conference Proceedings*, 4:229-232.
- 261 [14] Leslie Cooperband. 2002. The Art and science of composting. A resource for
262 farmers and compost producers. University of Wisconsin-isconsin- Madison centre
263 for Integrated Agricultural systems, March 29,2002. Pp. 1-17.
- 264 [15] Jedidi N., O. Vann Cleamput and AM' HIRI 1993. Mineralisation of organic
265 amendments in a Tunishian soil organic matter Dynamics and Sustainability of
266 Tropical Agriculture (Ed) Mutoagoyk and R Merr (eds) HTA/K.U Leuven. Pp 163-
267 169
- 268 [16] Ipinmoroti., R. R., G.O. Adeoye and M.K.C Sridhar. 1997. Potassium extraction
269 from farm wastes. *Proceedings of Soil Science Society of Nigeria Conference.*
270 Benin. 21st - 25th Nov. 1999. Pp. 87 - 90.
- 271 [17] Jackson, M.L. 1958. *Soil chemical analysis, Constable and Co. Ltd.*
272 *London.*

- 273 [18] AOAC. 1990. Methods of Analysis/Association of Analytical Chemist , Washington
274 S.C 71, no2, 247pp
- 275 [19] Moorhead ,D.L; Currie, W.S, Rastetter, E.B; Parton, W.J and Harmon, M.E. 1999.
276 Climatic and litter quality controls on decomposition; An analysis of modelling
277 approaches. Global Biogeochemical Cycles. 13: 575 - 589
- 278 [20] Obatolu C.R 1991. Growth and nutrient uptake of coffee (*coffee spp*) seedlings
279 grown on different organic materials. PhD. Thesis. Dept. of Agronomy. University of
280 Ibadan.xiv + 289pp.