

1  
2 **Factors involved in the early events of spore germination and host**  
3 **colonization by *Botrytis cinerea*.**

4 Salem Nassr and Radwan Barakat <sup>1</sup>

5 <sup>1</sup> Department of Plant Production and Protection, Faculty of Agriculture

6 Hebron University, P.O.Box 40, Hebron, Palestine

7 Email: [RadwanB@hebron.edu](mailto:RadwanB@hebron.edu)

8 **Abstract**

9 *Botrytis cinerea* is a necrotrophic fungal plant pathogen distributed worldwide. The early  
10 stages of epidemiology namely spore germination is a topic of great interest among  
11 researchers. In the current study, the effect of various physical, chemical and nutritional  
12 factors on germination of *B. cinerea* conidia were studied *in vitro*. Results showed that  
13 there was no particular influence of spore age (5-14 days) on germination in 10 millimolar  
14 fructose. In addition, germination-self-inhibition was found to be associated with increased  
15 spore concentrations (above  $4.5 \times 10^5$  conidia/ml) without significant differences between  
16 fungal isolates. When setting different pH values in the medium, conidial germination of  
17 *Botrytis cinerea* was impaired by pH values below 6 and above 8. However, germination of  
18 *Botrytis cinerea* was strongly enhanced (>90% after 24 hours) in the presence of sugars (i.e.  
19 Fructose, Sucrose and Glucose) at concentrations above 100 millimolar, whilst the cations  
20 ( $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{K}^+$ , and  $\text{Fe}^{2+}$ ) had no visible influence on conidial germination at a wide range  
21 of concentrations (0.001-1millimolar). With other additives and in the presence of  
22 inorganic nitrogen forms (i.e.  $\text{NH}_4$  and  $\text{NO}_3$ ), conidial germination responded similarly  
23 with no particular influence on germination, whilst germ tube growth and elongation  
24 increased progressively with increasing concentrations of both N-forms.

25 **Key Words:** *Botrytis cinerea*, conidial germination, early event, germ tube

26 **Introduction**

27 *Botrytis cinerea* Pers. ex. Fr. is the causal agent of gray mold. The name of the sexual stage  
28 or teleomorph is *Botryotinia fuckeliana* (de Bary) Whetzel, but the ascocarps are rarely  
29 observed under field conditions (Polach and Abawi 1975 ). *B. cinerea* is a filamentous,  
30 heterothallic Ascomycete exhibiting great variability in mycelial growth rate, conidial  
31 germination, pathogenicity, incidence of sporulation structures, production of sclerotia, and  
32 resistance to anti-*Botrytis* chemicals (Grindle 1979; Lorbeer 1980; Di Lenna et al. 1981;  
33 Kerssies et al. 1997). The early events of plants infection by plant pathogenic fungi are  
34 essential for disease initiation and progress. Such early events (adhesion, conidial  
35 germination, and formation of external infection structures) were intensively studied lately  
36 on *B. cinerea* throughout several studies (Doehlemann et al. 2006; Klimple et al. 2002;  
37 Schumacher et al. 2008).

38 Conidial germination of *B. cinerea* is induced by different physical and chemical signals,  
39 including the presence and quality of nutrients in particular sugars such as fructose (Kosuge  
40 and Hewitt 1964; Blakeman 1975). Conidial germination in most filamentous fungi  
41 requires the presence of low-molecular-mass nutrients such as sugars, amino acids and  
42 inorganic salts (Carlile and Watkinson 1994). Along with germination and after conidial  
43 adhesion, different mucilages are secreted and assist in anchoring of the germ tube and  
44 appressoria to the host surface. Several groups of proteins have been suggested to assist in  
45 germ tube and appressorium attachment and to mediate the exchange of early signalling  
46 between the fungus and the plant (Prins et al. 2000).

47 Conidia of *B. cinerea* are typically nutrient-dependent; they do not readily germinate in  
48 sterile water, and they usually require an exogenous input of nutrients for germination. In  
49 addition, it has been proposed that conidia of nutrient-dependent phytopathogenic fungi

50 may use germination-stimulating compounds from a host plant as an alternative chemical  
51 cue when nutrient concentrations are too low for conidial germination (Filonow, 2002). In  
52 addition, diverse carbon sources (mono- and disaccharides, acetate) are effective at low  
53 concentrations (10 mM) to induce germination in *B. cinerea*. Rich media such as malt  
54 extract induced rapid germination and early germ tube branching. Induction of conidial  
55 germination by nutrients, in particular sugars, is well known in saprotrophic fungi (Oshero  
56 and May, 2000). The mechanism of nutrient sensing by *B. cinerea* conidia is unknown. As  
57 diverse sugars and acetate induce germination with similar efficiency, it appears unlikely  
58 that nutrient sensing occurs by plasma membrane proteins (Forsberg and Ljungdahl, 2001).  
59 Conidia are also able to germinate on inert artificial surfaces; various amino acids plus  
60 sugars efficiently induced germination of conidia, while mineral salts such as ammonium  
61 and phosphate were effective only in the presence of low concentrations of sugars  
62 (Blakeman, 1975). On cuticular surfaces, however, dry-inoculated conidia can germinate at  
63 high humidity in the absence of liquid water (Prins *et al.*, 2000). Surface hydrophobicity,  
64 together with surface hardness, is well known to induce germination of *B. cinerea* conidia in  
65 the absence of nutrients (Oshero and May, 2000). The current study has illustrated the  
66 effect of such several physical and chemical factors on germination of *B. cinerea* conidia.

## 67 **Methods**

### 68 **Fungal isolates and commercial culture medium**

69 *Botrytis cinerea* wild type isolates used throughout this study were provided by the Plant  
70 Protection Research Center (PPRC) fungal collection at Hebron University. The first  
71 isolate, (PBC1) was isolated from infected Beans (*Phaseolous vulgaris*) growing under  
72 greenhouse in Hebron. The second isolate, (PBC3) was isolated from infected grape

73 berries (*Vitis vinefera L.*) growing in an open field in Hebron. Following isolation, the two  
74 isolates were grown on PDA medium and kept at  $20\pm 1$  °C under continuous light.

75 After 12 days, and when cultures sporulated, 5mm mycelium plug from each isolate culture  
76 was taken and placed in a fresh PDA culture plate; 24 hours later, one freely emerging  
77 conidium was transferred into another plate to get monosporic cultures. The monosporic  
78 cultures were grown on PDA medium amended with 10% homogenized bean leaves. Plates  
79 were then kept under continuous light in an incubator at  $20\pm 1$ °C for the coming  
80 experiments.

81 The third isolate used was B05.10 which is a universal known strain. It was derived from  
82 the wild-type isolate SAS56 by treatment with benomyl for haploidization (Quidde *et al.*,  
83 1999). This putative haploid wild type isolate B05.10 was provided by the lab. of Prof. P.  
84 Tudzynski (University of Munster, Germany).

### 85 **Conidial concentration**

86 The influence of conidial concentration on germination assays of *B. cinerea* isolates was  
87 assessed in a 24 well Sarstedt microtitre plate (Sarstedt, Newton. USA), according to  
88 (Doehlemann, 2006). Two plates of PDA medium amended with 10% homogenized bean  
89 leaves were inoculated with 100 µl of conidial suspension ( $1\times 10^6$  conidia/ml) from PBC3,  
90 PBC1 and B05.10 isolates. The inoculum was spread over the surface of the medium with  
91 the aid of a glass rod. After 11 days, conidia were harvested from each plate by 10 ml of  
92 SDW. Conidia were then filtered through a Nytex membrane to remove traces of mycelia  
93 and placed in a sterile plastic vial for each isolate.

94 Spore suspension was then washed three times with 10 ml of SDW and centrifuged (IEC  
95 Centra- CLD) for 3 minutes at 3000 rpm. The concentration of the conidial suspension was  
96 determined by a haemocytometer and diluted to the final concentrations of  $4\times 10^5$ ,  $2.5\times 10^4$ ,

97  $5 \times 10^3$  and  $2.5 \times 10^3$  conidia/ml. Spherical glass coverslips - 15mm (Roth, Karlsruhe,  
98 Germany) were placed in the bottom of each well of the 24-welled microtitre plate. A 25  $\mu$ l  
99 of each concentration were placed in the bottom of the well to which 475  $\mu$ l of 10mM D-  
100 Fructose solution were added to reach a final volume of 500  $\mu$ l and according to  
101 (Doehlemann, 2006). Plates were then incubated in the dark at  $20^\circ\text{C} \pm 1$  and conidial  
102 germination counted after 5 hours of incubation. Each treatment consisted of 4 replicates  
103 (wells) and 100 randomly selected conidia were counted in each of the 4 wells under an  
104 inverted microscope. A conidium was considered as germinated when the germ tube length  
105 was shorter, equal and/or exceeding the conidial diameter.

#### 106 **Age of conidia**

107 The influence of conidial age on germination of *B. cinerea*-isolate B05.10 conidia was  
108 assessed. The isolate B05.10 was grown on plates containing potato dextrose agar (PDA)  
109 amended with 10% homogenized bean leaves. Four plates of PDA medium were  
110 inoculated with 5 mm mycelium plug from a newly growing mycelium (two days old), and  
111 incubated at  $21^\circ\text{C}$  and continuous light. Conidia were then harvested after 7, 9, 10, 12, and  
112 14 days with 10 ml of SDW, and filtered through a Nynetex membrane to remove traces of  
113 mycelia.

114 Spore suspensions were then washed three times with 10 ml of SDW and centrifuged (IEC  
115 Centra- CLD) for 3 minutes at 3000 rpm; supernatant was discarded each time. Conidial  
116 concentrations were then determined with the aid of a haemocytometer [Tiefe Depth  
117 Protondeur 0.200 mm] and fixed at  $2.5 \times 10^4$  Conidia/ml. Spherical glass coverslips (15mm,  
118 Roth, Karlsruhe, Germany) were placed on each well of the 24-welled microtitre plate.  
119 Conidia (25  $\mu$ l of each age) were placed in the bottom of the well. Fructose was prepared  
120 and suspended in liquid Gamborg B5 basal salt mixture (GB5) (Duchefa Biochem. BV,

121 Haarlem, The Netherlands; Art: G0209.0050) to reach a final concentration of 10 mM.  
122 After that, 475  $\mu$ l of the 10mM fructose+GB5 solution were added to reach a final volume  
123 of 500  $\mu$ l. Sarstedt plates were then incubated in the dark at  $20\pm 1$  °C.

124 Using the same selected conidial ages, germination was monitored on a hydrophobic  
125 surface; polypropylene film was placed at the surface of a glass slide. Slides were then  
126 placed on a moist filter paper inside closed sterile petri dishes. Conidial suspension was  
127 prepared from the isolate B05.10 and fixed at a concentration of  $1\times 10^5$  Conidia/ml. The  
128 surfaces were then inoculated with 4 separate droplets of conidial suspension 25  $\mu$ l each and  
129 then placed in an incubator. A completely randomized design was used, each treatment  
130 consisted of 4 replicates (wells); germinated spores were counted out of 100 randomly  
131 selected spores under an inverted microscope.

### 132 **Microclimate pH**

133 The influence of microclimate pH on germination of *B. cinerea*, isolate B05.10 was  
134 determined in 1mM fructose solution. Fructose solutions were prepared and adjusted to pH  
135 ranges starting from 3, 4, 5, 6, 7, 8, 9 and up to 10 using 1M NaOH and 1M HCl. Conidia  
136 of *B. cinerea* (B05.10) were harvested from 10 days old sporulating cultures grown  
137 previously on (PDA+beans) medium with SDW and conidial concentration was fixed at  
138  $2.5\times 10^4$  conidia/ml. Spherical glass coverslips were placed in the bottom of each of the 24  
139 wells of the Sarstedt microtitre plate. After that, 25  $\mu$ l spore suspension was placed in the  
140 middle of each well and 475  $\mu$ l of Fructose solution were added to reach a final volume of  
141 0.5 ml. A completely randomized design was used with 3 replicates for each treatment.  
142 Numbers of germinated conidia were recorded after 5 hours.

143

144

## 145 **Sugars**

146 The role of carbon sources in conidial germination of *Botrytis cinerea* was investigated  
147 using three sugars: Fructose, Glucose and Sucrose in 5 molar concentrations 1 $\mu$ M, 10  $\mu$ M,  
148 100  $\mu$ M, 1mM and 10mM. Sugar solutions were prepared in DW and sterilized in the  
149 autoclave for 30 minutes at 127°C. *B. cinerea* was grown on (PDA+10% beans) and  
150 incubated at 21°C and continuous light for ten days. Spore suspensions from the isolates  
151 B05.10, PBC3 and PBC1 were prepared using SDW and adjusted to a final concentration of  
152 2.5 $\times$ 10<sup>4</sup> conidia/ml. Spherical glass coverslips were placed in the bottom of each of the 24  
153 wells of the Sarstedt microtitre plate. Spore suspension (25 $\mu$ l) was placed in the middle of  
154 each well and 475  $\mu$ l of each sugar treatment were added to reach a final volume of 0.5 ml.  
155 A completely randomized design was used with 4 replicates for each treatment. Numbers  
156 of germinated conidia were recorded after 5 and 25 hours.

## 157 **Salt cations**

158 The role of the cations, Ca<sup>2+</sup>, Mg<sup>2+</sup>, and Fe<sup>2+</sup> in conidial germination of *Botrytis cinerea* was  
159 investigated. Ca (CaCl<sub>2</sub>), Mg (MgCl<sub>2</sub>), and Fe (FeSO<sub>4</sub>.7H<sub>2</sub>O) were prepared into 6  
160 concentrations (0.001M, 0.01M, 0.1M, 100mM, and 1M). Solutions were prepared in  
161 distilled water and sterilized in the autoclave for 30 minutes at 127°C. *B. cinerea* was  
162 grown on (PDA+10% beans) and incubated at 21°C and continuous light for ten days.  
163 Conidial suspensions from the isolates B05.10 and PBC3 were harvested by SDW. Conidia  
164 were then filtered through Nytex membrane and washed three times to remove traces of  
165 mycelium. The concentration was adjusted to a final concentration of 1 $\times$ 10<sup>3</sup> conidia/ml.  
166 Spherical glass cover slips were placed in the bottom of each of the 24 wells of the Sarstedt  
167 microtitre plate. Spore suspension (25  $\mu$ l) was placed in the middle of each well and 475  $\mu$ l  
168 of each treatment were added to reach a final volume of 0.5 ml. A completely randomized

169 design was used with 4 replicates for each treatment. Numbers of germinated conidia were  
170 recorded after 40 hours of incubation at 21°C. At the same time, the average germ tube  
171 length of 10 random germinated conidia (replicates) was recorded.

### 172 **Inorganic nitrogen forms (NH<sub>4</sub> and NO<sub>3</sub>)**

173 The effect of the nitrogen forms, NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> on conidial germination of *Botrytis*  
174 *cinerea* was studied. The procedure is the same as that of the previous section. A  
175 completely randomized design was used with 4 replicates for each treatment. Numbers of  
176 germinated conidia were recorded after 25 hours of incubation at 21°C. At the same time,  
177 the average germ tube length of 10 random germinated conidia (replicates) was recorded.

### 178 **Statistical analysis**

179 The data of all experiments were analyzed statistically using analysis of variance (one way  
180 ANOVA) and fisher least significant difference (LSD) test with the aid of ([Sigma Stat 2.0](#)  
181 [for Windows® statistical package program, SPSS Inc., Chicago, IL, USA](#)).

### 182 **Results**

#### 183 **The effect of concentration of conidia on germination of *B. cinerea* conidia**

184 The influence of spore concentration of *B. cinerea*-isolates B05.10, PBC1 and PBC3 on  
185 conidial germination was determined in 10mM Fructose solution ([Fig. 1](#)). Results showed  
186 that conidial germination rates decreased with increasing spore concentrations. The highest  
187 germination rate was recorded at the spore concentration ( $2.5 \times 10^3$  conidia/ml) for all  
188 isolates. Generally, there were no significant differences in germination rates between the  
189 three *B. cinerea* isolates. It was evident that the three isolates responded similarly in which  
190 germination rates decreased with increasing spore concentrations ([Figure 1 and 2: C, D, and](#)  
191 [E](#)).

192



193 **The effect of age of conidia on germination of *B. cinerea* conidia**

194 Spore age could be another factor involved in early conidial germination in fungi. The  
195 influence of conidial age of *B. cinerea* (B05.10) on germination percentage was  
196 investigated. No significant differences in germination percentages were found between  
197 different conidial ages in sugar amended with Gamborg' B5- salt mixture (GB5). Conidial  
198 germination percentages, however, was significantly reduced in older conidia (67% after  
199 14days) compared to younger conidia (91% after 5 days) when germination was tested on a  
200 hydrophobic surface (Polypropylene). (see table 1 and 2).

201 **The effect of microclimate pH on germination of *B. cinerea* conidia**

202 The influence of microclimate pH on germination of *B. cinerea* conidia was assessed on  
203 Sarstedt plates. *B. cinerea* conidia were able to germinate well at pH values ranging from  
204 6-8; the highest germination rate was obtained at pH 7. However, B05.10 conidia  
205 germinated poorly at pH= 3 and 10. The experiment was repeated twice. Data on the  
206 average germination rates in different microclimate pH is presented in [Figure 3 and 4](#).

207 **The effect of sugars on germination of *B. cinerea* conidia**

208 The influence of the sugars (Fructose, Sucrose and Glucose) on conidial germination of *B.*  
209 *cinerea* was tested in various concentrations ([Fig.5 and 6](#)). Results showed that germination  
210 of conidia was stimulated in sugars in various proportions according to various  
211 concentrations compared to SDW. Sucrose was the best in inducing conidial germination  
212 even after 5 hpi only recording 87% compared to glucose 18% and fructose 59%. Almost  
213 all sugars have induced full germination (100%) after 24 hours of incubation at the highest  
214 concentration used (10mM) . The concentration (100  $\mu$ M) was the breaking point for all  
215 sugars to induce significant increase in conidial germination.

216

### 217 **The effect of cations on germination of *B. cinerea* conidia**

218 The cations  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{K}^+$ , and  $\text{Fe}^{2+}$  had no influence on conidial germination of *B. cinerea*  
219 isolates (B05.10 and PBC1) at the relatively low concentrations used (0.001-1mM). At  
220 10mM concentration, however, Fe reduced germination dramatically. At higher  
221 concentrations (>10mM), all cations showed toxicity and totally inhibited conidial  
222 germination (Fig. 7). Concerning germ tube elongation, only Fe was able to enhance  
223 germination at low concentrations, but as concentration increased germ tube elongation  
224 decreased until totally inhibited at concentrations (>10mM). All the other cations ( $\text{Ca}^{2+}$ ,  
225  $\text{Mg}^{2+}$  and  $\text{K}^+$ ), however, showed no influence on germ tube elongation at all concentrations  
226 tested (Fig. 7)

### 227 **The effect of inorganic nitrogen forms on germination of *B. cinerea* conidia**

228 The effect of  $\text{NH}_4$  and  $\text{NO}_3$  on germination of *B. cinerea* (B05.10 and PBC1) conidia and  
229 germ tube lengths was investigated (Fig. 8). Inorganic nitrogen forms had no influence on  
230 germination percentages of *B. cinerea* isolates at all concentrations tested. However, germ  
231 tube length growth was dramatically influenced by both nitrogen forms positively; germ  
232 tube length increased by almost 99% at the highest concentration of  $\text{NH}_4$  (1M) compared to  
233 the control (SDW).  $\text{NH}_4$  form of nitrogen enhanced germ tube growth to a larger extent  
234 than  $\text{NO}_3$  form of N for both *B. cinerea* isolates. Both *B. cinerea* isolates responded almost  
235 similarly in respect to percentage germination and germ tube growth.

236

### 237 **Discussion**

238 The ability of fungi to adhere to and germinate on leaves and other substrata is well  
239 documented and is thought to represent an important early event in plant-microbe  
240 interactions (Braun and Howard 1994; Jones 1994). Spore germination in *Botrytis cinerea*

241 follows a developmental sequence of spore swelling, localized outgrowth of the germ tube  
242 and subsequent polarized growth of the new hyphae. It was noted that, conidial germination  
243 rates of *B. cinerea*-isolates decreased with increasing spore concentrations without  
244 significant differences between isolates. At concentrations above  $4 \times 10^5$  conidia/ml, conidia  
245 were unable to germinate and appeared in clots. [Sharrock, et al. 2001](#) found that conidia of  
246 *B. cinerea* exhibit a self inhibition strategy during germination at high concentrations  
247 ( $1 \times 10^6$  conidia/ml) or more. It is assumed that at high concentrations, conidia tend to  
248 produce specific germination and/or growth inhibitors regardless of the richness of the  
249 substrate. Furthermore, several germination-self-inhibitors in other fungal species such as  
250 *Puccinia*, *Uromyces*, *Colletotrichum*, *Dictyostelium*, *Fusarium* and *Aspergillus* were  
251 investigated and reports showed that these inhibitors can be volatile or non-volatile  
252 ([Allen 1955](#); [Bacon et al. 1973](#); and [Barrios-Gonzales et al. 1989](#)). It was also concluded  
253 that self-inhibitors can affect other fungal processes, such as prevention of appressorium  
254 induction which make conidial germination unlikely to occur.

255 Spore age could be another factor involved in early conidial germination in fungi. It was  
256 found that conidial germination was significantly reduced in older conidia (67% after 14  
257 days) compared to younger conidia (91% after 5 days) when germination was tested on a  
258 hydrophobic surface (Polypropylene). However, no differences were noticed when spores  
259 germinated in Fructose and GB5. This suggests that nutritional factors may mask the effect  
260 of age and older conidia can germinate as well as younger conidia if the growth substrate  
261 was supplied with appropriate nutritional source. Using different germination conditions,  
262 [Shirashi et al., 1988](#) found that young *Botrytis* conidia, in general, germinated well at 20°C  
263 compared to old conidia.

264 As for pH, conidial germination was significantly impaired at high and/or low values  
265 (below 6 and above 8). Conidia germinated well at pH ranging from 6-8 with the highest  
266 germination rate at pH=7. In this direction, fungi very often can dynamically alter the local  
267 pH to fit its enzymatic arsenal, with the level of pathogenicity being related to the efficiency  
268 of the pH change. (Prusky et al. 2001). Generally, *Botrytis cinerea* is classified among  
269 acidic fungi (Prusky and Yakoby, 2003) and similar to other pathogenic fungi, such as  
270 *Penicillium expansum*, *P. digitatum*, *P. italicum*, and *Sclerotinia sclerotiorum* that use tissue  
271 acidification in their attack (Vautard and Fevre, 2003). This investigation, however, was  
272 restricted to the conidial germination in vitro. The ability of *B. cinerea* to germinate at  
273 various pH values emphasizes the previous findings stating that *Botrytis* spp. change the  
274 medium or site pH to facilitate the enzymatic activities.


275 Nutritional supplements, namely sugars are considered rich nutrients; germination of  
276 *Botrytis cinerea* conidia was stimulated in the three different sugars (fructose, sucrose and  
277 glucose) at various concentrations compared to the control (Sterile Distilled Water).  
278 Almost all sugars have induced full germination (100%) after 24 hours of incubation at the  
279 highest concentration used (10mM) knowing that the concentration (100  $\mu$ M) was the  
280 (breaking point) for all sugars to induce significant increase in conidial germination.  
281 Sugars at relatively low concentrations (i.e 10mM) induced early swelling of conidia and  
282 enhanced early germ tube branching. In this direction, it has been shown that Fructose  
283 induced germination of *B. cinerea* conidia more efficiently than any other monosaccharide  
284 (Blakeman, 1975). Germination induction by sugars was concentration dependent, and  
285 fructose was more effective than glucose. Similarly and among sugars, fructose has been  
286 pointed out as the best inducer of germination in *B. cinerea*, being more effective than  
287 glucose and other hexoses or disaccharides (Blakeman, 1975). One explanation for the

288 particular important activity of fructose in conidial germination could be that this sugar is  
289 preferentially taken up by a fructose-specific transport system. This is surprising since  
290 glucose is usually the most efficient hexose not only as a nutrient, but also as a signalling  
291 compound (Doehlemann et al. 2005). Using almost the same protocol for germination,  
292 Doehlemann, et al. 2006 found similar results after incubation for 24 hours. Induction of  
293 conidial germination by nutrients, in particular sugars, is well known in saprotrophic fungi  
294 (Osherov and May 2000). In rich media, most fungi germinate quickly, including  
295 phytopathogens such as *F. solani*, *C. graminicola* and *C. gloeosporioides* (Ruan et al.  
296 1995; Chaky et al. 2001; Barhoom and Sharon, 2004).

297 The mechanism of sugar sensing by *B. cinerea* conidia is unknown. As diverse sugars and  
298 acetate induce germination with similar efficiency, it appears unlikely that nutrient sensing  
299 occurs by plasma membrane proteins (Forsberg and Ljungdahl, 2001).

300 Regarding the addition of salt cations and from looking at the results, it was obvious that the  
301 tested cations ( $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{K}^+$ , and  $\text{Fe}^{2+}$ ) had no influence on conidial germination at a  
302 wide range of concentrations (0.001-1mM). However, at high concentrations (>10mM),  
303 germination declined sharply, especially with  $\text{Fe}^{2+}$  which suggests a level of toxicity  
304 induced at high concentrations. It is very likely that conidia before germination is not  
305 affected at low concentrations of cation availability in the growth substrate. However, after  
306 germination, germ tube growth becomes more sensitive to a wide range of cation  
307 concentrations in the growth media.  $\text{Fe}^{2+}$  seems to provide an important nutritional source  
308 for germ tube growth at low concentrations (0.001 M). Barakat and Almasri, 2009  
309 (unpublished data) found that at high concentrations (i.e. 1M) all these cations inhibited  
310 germination of *Botrytis* conidia and the level of toxicity varied between isolates. Shirani  
311 and Hatta (1987), found that at the concentration ( $5 \times 10^4$  conidia/ml) conidial germination of

312 *B. cinerea* was optimum (100%) in the presence of  $\text{Ca}^{2+}$  ( $\text{CaCl}_2$ ) and was relatively high  
 313 (66%) in  $\text{Mg}^{2+}$  ( $\text{MgSO}_4$ ) at the concentrations (0.1-0.7 g/liter). Conidial germination  
 314 responded almost similarly to nitrogen forms. While N-forms had no influence on  
 315 germination, germ tube growth and elongation responded positively with increasing  
 316 concentrations of both forms. This suggests that conidia may depend more on available  
 317 energy inside the spore to germinate but after germination, germ tube growth greatly depend  
 318 on nutritional elements available in the growth substrate.



338 **Figure 1.** Effect of spore concentration on conidial germination rates of *B. cinerea* isolates  
 339 grown on (PDA+10% bean leaves) medium and incubated in 10mM Fructose at  
 340  $20 \pm 1$  °C under continuous light after 20 hours of incubation. (vertical bars  
 341 represent  $\text{LSD} = 5.49$ ,  $n=3$ ).

343

344

345


A

B

C

D

E



346

347

348

349 **Figure 2.** *B. cinerea* (B05.10) conidial germination at different concentrations of conidia at  
 350 200X. Conidial concentrations: (A),  $5 \times 10^6$  conidia/ml; (B),  $1 \times 10^6$  conidia/ml;  
 351 (C),  $4 \times 10^5$  conidia/ml; (D),  $2.5 \times 10^4$  conidia/ml and (E),  $5 \times 10^3$  conidia/ml.  
 352

353

354 **Table 1.** Influence of conidial age on germination of *B. cinerea*-isolate B05.10 after 20  
 355 hours of incubation in 10 mM fructose solution+GB5

356

Age of B05.10 culture (days)	% Germination
5	97a
7	95a
10	96a
12	95a
14	93a

357

358 Means followed by the same letter in the same column are not significantly different  
 359 ( $P= 0.064$ ). GB5: Gamborgs B5-basic salt mixture.

360

361

362 **Table 2.** Influence of conidial age on germination of *B. cinerea* conidia isolate B05.10 after  
 363 20 hours of incubation on polypropylene surface.

364

Age of B05.10 culture (days)	% Germination
5	91 a
7	84 ab
10	92 a

<b>12</b>	78 bc
<b>14</b>	67 c


365 -Means followed by the same letter in the same column are not significantly different  
366 (LSD=11.309, n=4). GB5: Gamborgs B5-basic salt mixture.

367

368

369

370



371

372 **Figure 3.** Influence of microclimate pH on conidial germination of *B. cinerea*-isolate  
373 B05.10 in 1mM fructose solution after 5 hours of incubation. (LSD = 9.020, n=3).


374

375

376

377







378

379 **Figure 4.** Conidial germination of *B. cinerea* –isolate B05.10 under different pH values in  
 380 1mM fructose solution at 200 X .


381 **5 hpi** **24 hpi**  
 382 ● Fructose ■ Sucrose ▲ Glucose  
 383 *B. cinerea* (B05.10)




384




385 *B. cinerea* (PBC3)



386




387 *B. cinerea* (PBC1)



388

389 **Figure 5.** Influence of Fructose, Sucrose and Glucose solutions on germination of *B.*  
 390 *cinerea* conidia. (LSD=10.168, n=4, p<0.001). Experiment was done after 5 and  
 391 24 hours of incubation in various concentrations at 20±1 °C. SDW: Sterile  
 392 distilled water, GB5: Gamborg's B5 basic salt mixture; hpi: hours post  
 393 inoculation.


394  
 395  
 396




397

398 **Figure 6.** Conidial germination of *B. cinerea* (B05.10) in different concentrations of  
 399 sucrose. A): after 5 and B): after 24 hours at 200X.

400  
 401  
 402




403



404

405 **Figure 7.** Influence of  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{K}^+$ ,  $\text{Fe}^{2+}$  in various concentrations on conidial  
 406 germination and germ tube elongation of *B. cinerea* after 40 hours of incubation.  
 407 Conidial germination (LSD=13.527, n=4) ; Germ tube elongation (LSD=8.815,  
 408 n=10).  
 409  
 410  
 411  
 412



413

414

415

416

417 **Figure 8.** Influence of NH<sub>4</sub> and NO<sub>3</sub> in various concentrations on conidial germination and  
418 germ tube elongation of *B. cinerea* PBC1 (A, C), and B05.10 (B, D). Differences  
419 between means of germination percentages were not significant; bars in (B, D)  
420 represent the standard error of the mean with LSD=8.489.  
421

422

### 423 **Acknowledgment**

424 The authors acknowledge the financial support provided by the Deutsche  
425 Forschungsgemeinschaft (DFG) - grant number (Tu50/15).  
426

427

428

429

### 430 **References**

431

431 Allen, P.J. 1955. The role of a self-inhibitor in the germination of rust uredospores.  
432 *Phytopathology* 215, 259-266.

433

434 Bacon, C. W., Sussman, A.S. and A.G. Paul, 1973. Identification of a self-inhibitor from  
435 spores of *Dictyostelium discoideum*. *Journal of Bacteriology* 113, 1060-1063.

436

437 Barakat, R. and M.I. Al-Masri, Hebron. Personal communication (unpublished data) 2009.

438

439 Barhoom, S. and A. Sharon, 2004. cAMP regulation of 'pathogenic' and 'saprophytic'  
440 fungal conidia germination. *Fungal Genetics and Biology* 41,317-326.

441

442 Barrios-Gonzales, J., Martinez, C., Aguilera, A., and M. Raimbault, 1989. Germination of  
443 concentrated suspensions of spores from *Aspergillus niger*. *Biotechnology* 11, 551-  
444 554.

445

446 Blakeman, J.P. 1975. Germination of *B. cinerea* conidia *in vitro* in relation to nutrient  
447 conditions on leaf surfaces. *Transaction British Mycological Society* 65, 239-247.

448

449 Braun, E. J. and R. J. Howard, 1994. Adhesion of fungal spores and germlings to host plant  
450 surfaces. *Protoplasma* 181, 202-212.

451

452 Carlile, M.J. and S.C. Watkinson, 1994. *The Fungi*. Academic Press, London.UK

453

454 Chaky, J., Anderson, K., Moss, M., and L. Vaillancourt, 2001. Surface hydrophobicity and  
455 surface rigidity induce conidia germination in *Colletotrichum graminicola*.  
456 *Phytopathology* 91,558-564.

457

- 458 Di Lenna, P., Marciano, P. and P. Magro, 1981. Comparative investigation on  
459 morphological and physiological features of three isolates of *B. cinerea*.  
460 *Phytopathologische Zeitschrift* 100, 203-211.
- 461
- 462 Doehlemann, G., Berndt, P. and M. Hahn, 2006. Different signalling pathways involving  
463 a G $\alpha$ -protein, cAMP and a MAP kinase control germination of *B. cinerea* conidia.  
464 *Molecular Microbiology* 59, 821-835.
- 465
- 466 Doehlemann, G., Molitor, F. and M. Hahn, 2005. Molecular and functional characterization  
467 of a fructose specific transporter from the gray mold fungus *Botrytis cinerea*. *Fungal*  
468 *Genetics and Biology* 42, 601-610.
- 469
- 470 Filonow, A.B. 2002. Mycoactive acetate esters from apple fruit stimulate adhesion and  
471 germination of conidia of the gray mold fungus. *Journal of Agricultural Food*  
472 *Chemistry* 50, 3137-3142.
- 473
- 474 Forsberg, H., and P.O. Ljungdahl, 2001. Sensors of extracellular nutrients in  
475 *Saccharomyces cerevisiae*. *Current Genetics* 40, 91-109.
- 476
- 477 Grindle, M. 1979. Phenotypic differences between natural and induced variants of  
478 *B. cinerea*. *Journal of General Microbiology* 111, 109-120.
- 479
- 480 Jones, E.B. 1994. Fungal adhesion. *Mycological Research*, 98, 961-981.
- 481
- 482 Keressies, A., Zessen, A.I., Wagemakers, C.M. and J. Van Kan, 1997. Variation in  
483 pathogenicity and DNA polymorphism among *B. cinerea* isolates sampled inside and  
484 outside a glasshouse. *Plant Disease*, 81, 781-786.
- 485
- 486 Klimple, A. Gronover, C., Stewart, J., and B. Tudzynski, 2002. The adenylate cyclase  
487 (BAC) in *Botrytis cinerea* is required for full pathogenicity, *Molecular Plant*  
488 *Pathology* 6, 439-450.
- 489
- 490 Kosuge, T. and W.B. Hewitt, 1964. Exudates of grape berries and their effect on  
491 germination of conidia of *B. cinerea*. *Phytopathology* 54, 167-172.
- 492
- 493 Lorbeer, J.W. 1980. Variation in *Botrytis* and *Botryotinia*. In: Coley-Smith, J.R., Verhoeff,  
494 K. And Jarvis, W.R. (eds.). *The biology of Botrytis*. Academic Press Inc. (London)  
495 LTD. (pp 19-40).
- 496
- 497 Osherov, N. and G. May, 2000. Conidial germination in *Aspergillus nidulans* requires RAS  
498 signalling and protein synthesis. *Genetics* 155, 647-656.

- 499  
500 Polach, F.J. and G.C. Abawi, 1975. Occurrence and biology of *Botryotinia fuckeliana* on  
501 beans in New York. *Phytopathology* 65, 657-660.  
502
- 503 Prins, T.W., Tudzynski, P., von Tiedemann, A., Tudzynski, B., ten Have, A., Hansen, M.E.,  
504 Tenberge, K. and van J. Kan, 2000. Infection strategies of *B. cinerea* and related  
505 necrotrophic pathogens. In *Fungal Pathology* (Kronstaad, J.W., ed.). Dordrecht:  
506 Kluwer Academic Publishers (pp. 33-64).  
507
- 508 Prusky, D., McEvoy, J. L., Leverentz, B., and W. S. Conway, 2001. Local modulation of  
509 host pH by *Colletotrichum* species as a mechanism to increase virulence. *Molecular*  
510 *Plant-Microbe Interactions* 14, 1105-1113.  
511
- 512 Prusky, D., and N. Yakoby, 2003. Pathogenic fungi: leading or led by ambient pH?  
513 *Molecular Plant Pathology* 4, 509-516.  
514
- 515 Ruan, Y., Kotraiah, V., and D.C. Straney, 1995. Flavonoids stimulate spore germination in  
516 *Fusarium solani* pathogenic on legumes in a manner sensitive to inhibitors of cAMP-  
517 dependent kinase. *Molecular Plant-Microbe Interaction* 8, 929-938.  
518
- 519 Scharrock, K., Henzell, R. and F. Parry, 2001. Self-inhibition of germination of *Botrytis*  
520 *cinerea* conidia. *The Horticulture and Food Research Institute of New Zealand*.  
521 Hamilton, New Zealand. (pp25-26).  
522
- 523 Schumacher, J., Kokkelink, L., Huesmann, C., Jimenez-Teja, D., Collado, I., Barakat, R.,  
524 Tudzynski, P., and B. Tudzynski, 2008. The cAMP-dependent signaling pathway and  
525 its role in conidial germination, growth and virulence of the grey mould *B. cinerea* .  
526 *Molecular Plant-Microbe Interaction* 21, 1443-1459.  
527
- 528 Shirani, N. and T. Hatta, 1987. Mineral salt medium for the growth of *Botrytis cinerea* in  
529 *vitro*. *Annals of Phytopathological Society of Japan* 53, 191-197.  
530

531 Shirashi, M, Fucotomi M., and S. Akai 1988. Influence of temperature and conidium age  
532 on mycelium growth rates of *Botrytis cinerea* fr. Pers. *Annals of the*  
533 *phytopathological society of Japan* 40, 230-235.

534

535 Vautard-Mey G. and M. Fevre, 2003. Carbon and pH modulate the expression of the  
536 fungal glucose repressor encoding genes. *Current Microbiology* 46, 146-150.

537

538

539

540

UNDER PEER REVIEW