

1 **Genetic control for some traits using generation mean analysis in bread**
2 **wheat (*Triticum aestivum* L.).**
3
4
5

6 **ABSTRACT**
7

8 In order to study the inheritance and genetic analysis of drought tolerance indicators a
9 six generations of P₁, P₂, F₁, F₂, Bc₁ and Bc₂ of two wheat crosses i.e., Sakha 94 x
10 Tokwie (C₁) and Giza 168 x Tokwie (C₂) under normal irrigation (N) and drought stress
11 (D) were studied using generation mean analysis at Faculty of Agriculture, Sohag
12 University, Egypt. Genetic variation was found for No. of spikes/plant (NS), 100-seed
13 weight (SW), grain yield (GY), biological yield (BY), relative water content (RWC) and
14 chlorophyll content (CC) (N&D) in two crosses. High heterosis was observed for all
15 studied characters (N&D) except CC in two crosses. Genetic analysis showed
16 overdominance in the inheritance of all studied characters (N&D) in two crosses. High to
17 moderate heritability values in broad sense were detected for all characters in both
18 crosses. Narrow-sense heritability (C₁&C₂) ranged from 0.18 for CC (D) to 0.37 for
19 RWC (D) in C₁. The genetic advance (C₁&C₂) was high (more than 40%) for GY (N&D),
20 while NS, BY, RWC and CC (N&D) were moderate (14-40%), indicating the importance
21 of direct selection for these characters. The genetic models fitted for all studied characters
22 (N&D) in two crosses except RWC (D in C₁), indicated dominance and additive x
23 additive gene effects. Both additive x additive [i] and dominance x dominance [1] effects
24 were significant for all studied characters (N&D) in two crosses except RWC (D in C₁),
25 supporting the presence of duplicate type of epistasis. Since several important characters
26 are influenced by dominance and non-allelic gene interaction, it is advisable to delay
27 selection to later generation with increased homozygosity.

28
29 **Key words:** Wheat (*Triticum aestivum* L.), drought stress, generation mean analysis, gene action.
30
31
32
33
34
35
36
37

Introduction

In Egypt, wheat production is far below what is needed to meet the local consumption of the growing population resulting in increasing wheat imports. To formulate an efficient breeding program for developing drought-tolerance varieties, it is essential to understand the mode of inheritance, the magnitude of gene effects and their mode of action (Farshadfar et al., 2001, 2008b; Iqbal et al., 2007).

The plant breeder is interested in the estimation of gene effects in order to formulate the most advantageous breeding procedures for improvement of the attribute in question. Therefore, breeders need information about nature of gene action, heterosis, inbreeding depression, heritability and predicted genetic gain from selection for yield and yield components. Sprague (1963) listed three major factors that must be considered and which may limit progress in the analysis of quantitative genetic variation: the number of genes involved, the type of gene action, and the genotype- environment interaction.

The genetical studies based on the means and variances of basic generations, is a simple method for estimating the gene effects for a polygenic trait and has been reviewed in many crop species. The greatest merit of generation means analysis lies in its ability to estimate the epistatic effects (Mather and Jinks, 1982).

The possibility of epistasis accounting for a significant proportion of genetic variance of quantitative trait has been investigated extensively in previous studies in crop plants. Amount and type of epistasis can have a major consequence on both the reliability of predictions and the design of breeding program. Statistically, detection of epistasis using generation means analysis is more reliable and efficient than by the analysis of variance approach (Lamkey and Lee, 1993).

However, it has its own limitations and several assumptions. Triple test cross is a powerful method of genetic analysis, which provides unbiased estimates for epistasis. In addition, it also estimates the additive and dominance components of variation with high accuracy when epistasis is absent (Kearsey and Jinks, 1968).

The variance estimates attributed to environment, total genetic, additive and dominance deviation effects were obtained from the phenotypic variances for populations P1, P2, F1, F2, BC1 and BC2. These estimates allowed the determination of heritabilities in the broad and narrow sense, mean degree of dominance and minimum number of genes that

control each character, by using Burton's (1951) expression.

The objective of the present investigation was to investigate the genetic analysis of quantitative indicators of drought tolerance in wheat under drought condition using generation mean analysis.

Material and methods

The two Egyptian cultivars, Sakha 94 and Giza 168 were more adapted in Egypt and proved high yielding. However, the introduced line (Tokwie) is characterized as a drought tolerant. Therefore, the line introduced was crossed with the Egyptian cultivars in order to enlarge the variability for selection in the breeding program for these characters. The experiments reported herein were carried out during the three successive growing seasons of 2010/2011, 2011/2012 and 2012/2013. In 2010/2011, the parent genotypes of hexaploid wheat (*Triticum aestivum* L.) were sown to secure enough hybrid seed (Table 1). Two crosses namely Sakha 94 x Tokwie (Cross 1) and Giza 168 x Tokwie (Cross 2) were developed at Faculty of Agriculture, Sohag University, Egypt.

In 2011/2012 season, F_1 plants were selfed to produce F_2 seeds and backcrossed to the parents to produce BC_1 and BC_2 seeds. In 2012/2013 season, The parents (P_1 and P_2), the first (F_1) and second (F_2) generation hybrids and the first ($P_1 \times F_1 = BC_1$) and second ($P_2 \times F_1 = BC_2$) backcrosses were grown in two experiments in a randomized complete blocks design with two replicates for each one. Each replicate consisted of 20 grains in one row for each of the parents and F_1 , 40 grains in two rows of each of back cross and 80 grains in four rows for the F_2 population. Rows were 2.0 m long and 30 cm apart and 10 cm between plants. The first experiment was under normal irrigation (N) (gave irrigation when ever required), the second experiment was under drought stress (D) (after the emergence of 50% of the spikes, the water stress treatment received no more water until harvesting). The soil was fertilized at the rate of 20 kg/fed (15% P_2O_5) and 80 kg/fed (33.5% ammonium nitrate) and weeds were controlled by hand.

Data were recorded on 5 competitive individual plants for non-segregate basis as (P_1, P_2 and F_1) and 10 plants for BC_1 and BC_2 and 60 plants for F_2 population for each replicate follows:

1-No. of spikes/plant (NS).

101 2-100-seed weight (SW) in grams.
 102 3-Grain yield/plant (GY) in grams.
 103 4-Biological yield/plant (BY) in grams.
 104 5-Relative water content (RWC): A 4 cm segment of the youngest leaf was taken and cut
 105 into 2 cm segments and weighed (fresh weight = FW). Then the segments were placed in
 106 distilled water for 4 hours and reweighed to obtain turgor weight (TW). Thereafter the
 107 leaf segments were oven dried and weighed (dried weight = DW). RWC was calculated
 108 using the formula of Ritchie et al. (1990), $RWC \% = [(FW - DW) / (TW - DW)] \times 100$.
 109 6-Chlorophyll content (CC). Chlorophyll content was measured using a SPAD-502
 110 chlorophyll meter (Minolta, Japan). For this measurement the average of three leaves per
 111 plant per replication per treatment was taken.

112 **Statistical analysis:**

113 Analysis of variance and mean comparison of the characters was done using SAS
 114 Software. Generation mean analysis was performed using Mather and Jinks method
 115 (1982). In this method the mean of each character is indicated as follows:

$$116 \quad Y = m + \alpha [d] + \beta [h] + \alpha^2 [i] + 2\alpha \beta [j] + \beta^2 [l]$$

118
 119 Where:

120 Y = The mean of one generation
 121 m = The mean of all generation
 122 d = The sum of additive effects
 123 h = The sum of dominance effects
 124 i = The sum of additive x additive interaction (complementary)
 125 l = The sum of dominance x dominance interaction (duplicate)
 126 j = Sum of additive x dominance and α , $2\alpha \beta$ and β^2 are the coefficients of genetic
 127 parameters.

128 The genetic parameters (m, [d], [h], [I], [j], [l]) were tested for significance using a t-test.
 129 To estimate the parameters and to select the most suitable model the least squares method
 130 and the joint scaling test of Mather and Jinks (1982) were employed.

131 Potence ratio, was estimated by using the formula of Smith (1952).

132 Stress Tolerance index (STI) for grain yield were computed as formula using by
 133 Farshadfar, et al. (2001), $STI = (GY_N)(GY_D)/(GY_N)^2$

134 where GY_N is grain yield under normal irrigation and GY_D is grain yield under drought.

135 Broad-sense (H_b^2) and narrow-sense (H_n^2) heritability were estimated by Warner (1952)
 136 formulas:

137 $H_b^2 = [V_{F2} - (V_{P1} + V_{P2} + V_{F1})/3] / V_{F2}$

138 $H_n^2 = [2V_{F2} - (V_{BC1} + V_{BC2})] / V_{F2}$

139 Genetic advance was calculated (Johanson, 1955) with a selection intensity of $i=5\%$ for
140 all the characters as: $G_A = i.H_b.\sqrt{V_{F2}}$

141 The components of variation for six generations were calculated by the formulae of F2
142 variance were obtained by the following formula of Mather and Jinks (1982) as:

143 $E = 1/3 (V_{P1} + V_{P2} + V_{F1})$

144 $D = 4V_{F2} - 2 (V_{BC1} + V_{BC2})$

145 $H = 4(V_{F2} - 1/2V_D - V_E)$

146 $F = V_{BC1} - V_{BC2}$

147 Where:

148 D - Additive genetic variance

149 H - Dominance variance

150 E - Environmental component of variance

151 F - Correlation between D and H over all loci

152 **Results and discussion**

153 The analysis of variance (Table 2) revealed significant differences for two environments
154 and generations for No. of spikes/plant (NS), 100-seed weight (SW), grain yield (GY),
155 biological yield (BY), relative water content (RWC) and chlorophyll content (CC) in two
156 crosses, indicating the existence of genetic variation and possibility of selection for
157 drought tolerance. The genotypes x environments interaction was also significant for all
158 studied characters in C_2 , except for GY, displaying their similar response and different
159 responses of other traits. While, the genotypes x environments interaction was non-
160 significant for all studied characters in C_1 . Genetic variation was found in wheat for NS,
161 SW, BY and GY by Tammam, 2005; Farshadfar et al., 2008a; Amin, 2013 and for RWC
162 by Manette et al., 1988; Farshadfar et al., 2001.

163 The data six generations means (Table 3) showed that F1 hybrids were higher than mid-
164 parent and or best parent for all studied characters under both conditions in two crosses
165 except CC. These results showed the presence of heterotic effects for these characters.

166 In fact the development of any plant breeding program is dependent upon the existence of
167 genetic variability. The efficiency of selection and expression of heterosis also largely

upon the magnitude of genetic variability present in the plant population (Singh and Narayanan, 1993; Singh and Chaudhary, 1999; Farshadfar et al., 2001, 2008b; Amin, 2013). The potence ratio presented in table (3), its values ranged from less than one (0.11) for CC (D in C₂) to more than one (36.91) for RWC (D in C₂), indicating the presence of over dominance for all studied characters in two Crosses under normal (N) and drought stress (D) except CC (D in C₁) was partial dominance. These results are in line with those obtained by Ketata et al., 1976, Moshref, 1996, Tammam, 2005 and Amin, 2013.

The highest stress tolerance index (Table 4) was revealed by the F₁ hybrid (STI=0.85 in C₁ and 0.83 in C₂), displaying the presence of heterobeltiosis for drought resistance in the F₁ hybrid, followed by P₂ (0.81) in C₁ and P₂ (0.81) and P₁ (0.80) in C₂.

The degree of dominance (h/d), broad-sense (Hb) and narrow-sense (Hn) heritabilities, genetic advance (GA) and genetic components of variation are presented in Tables (5&6), which shows that the degree of dominance (h/d) for all studied characters was greater than one in two crosses (N&D) except NS (N in C₂), indicating the presence of the overdominance type of gene action in the inheritance of these traits. Selection of these characters must therefore be delayed until the F₃ or F₄ generation. This delay permits a loss of non-additive genetics variance through inbreeding, so that the additive genetics variance can be more clearly evaluated. these results are in harmony with those obtained by Zaazaa et al., (2012), whereas they revealed that, the complex genetic behavior particularly additive and dominance components could be successfully exploited in later generation.

NS (N in C₂) was controlled by the additive type of gene action; the pedigree method of selection can be used for improved of this trait, While for characters under control of the non-additive type of gene action, biparental mating offers good prospects for increasing the frequency of genetic recombination, hastening the rate of genetic improvement, through it may be necessary to resort to heterosis breeding (Gill et al., 1972; Sharma and Singh, 1976; Srivastava et al., 1992; Farshadfar et al., 2001; Tammam, 2005; Kheiralla et al., 1993; Amin, 2013).

Heritability estimate indicates the progress from selection for plant characters is relatively easy or difficult to make in breeding program. Plant breeders, through

experience, can perhaps rate a series of their response to selection. Heritability gave a numerical description of this concept. Assessment of heritability of various traits is of considerable important in crop improvement program, for example, to predict response to selection, Nyguist, 1991. High to moderate broad-sense heritability estimates for all studied characters in two Crosses (N&D) (Tables 5&6) showed that effective progress can be mad through selection. Moderate narrow-sense heritability (0.2-0.5) was show for all studied characters in two crosses (N&D) except CC (D) in Cross 1 and RWC (D) in Cross 2 indicated low heritability estimate (less than 0.2) (Tefra and Peat, 1997).The difference between H_n and H_b exhibits the involvement of the dominance effect in the genetic constitution of these characters.

The variation observed between the genotypes for the characters investigated exhibited that selection maybe effective for the improvement of drought tolerance (Umarahan et al., 1997; Farshadfar et al., 2001; Farshadfar et al., 2008b; Farshadfar 2012), however, the selection efficiency is related to the magnitude of heritability and genetic advance (Johnson et al., 1955; Singh and Narayanan, 1993). Heritability estimates along with genetic advance are important selection parameters and normally more helpful in predicting the gain under selection than heritability estimates alone. However, heritability estimates are influenced by the type of genetic material, sample size, method of sampling, conduct of experiment, method of calculation and effect of linkage. Genetic advance which refers to the improvement in the mean genotypic value of selected individuals over the parental population is influenced by the genetic variability, heritability and selection intensity (Alza and Martinez, 1997; Sharma, 2003).

The rate of genetic advance is connected with heritability (Mather and links, 1982). The genetic advance (C_1 & C_2) was high (more than 40%) for GY (N&D), while NS, BY, RWC and CC (N&D) were moderate (14-40%), indicating the importance of direct selection for these characters and the significance of indirect selection for SW (N&D) in two crosses with low genetic advance (less than 14%) through correlated response with characters having high heritability and genetic advance (Sharma et al., 1991; Farshadfar et al., 2001 and 2008a; Sood et al., 2006; Golparvar 2012).

Degree of dominance and variance components are presented in Tables (5&6), Ew, D and H are environmental, additive and dominance components, respectively. F is an indicator

of correlation between D and H over all loci. If F is zero it means that dominant genes are in the parent with high performance, while negative F exhibits that dominant genes are in the low performance parent. If the ratio of $F/\sqrt{D \times H}$ is equal to or near one confirms that the magnitude and sign of dominance for all the genes monitoring the character is equal, therefore, the ratio $\sqrt{H/D}$ is a good estimator of dominance. If $F/\sqrt{D \times H}$ is equal to zero or close to zero, the magnitude and sign of the genes controlling the character is not equal and hence $\sqrt{H/D}$ explains average dominance. The h/d ratio estimates the degree of dominance (Singh and Chaudhary, 1999; Farshadfar et al., 2001, 2008a; Naroui Rad et al., 2013). The ratio of $\sqrt{H/D}$ for all studied characters (N&D) in two crosses showed average dominance except NS (D), GY (D) and CC (N&D) in C₁ and GY (N), RWC (N) and CC (N&D) in C₂ showed over dominance.

The estimates of heterosis and inbreeding depression together provide information about type of gene action involved in the expression of various quantitative traits. The percentage of heterosis with regard to High Parent (HP) and Mid-Parent (MP) and Inbreeding Depression (ID) (Fig. 1,2,3 and4) exhibited that mid-parent and high parent heterosis were positive for NS, SW, BY, GY, RWC and CC in two crosses under both conditions except CC was negative (D) (C₁&C₂) compared with high parent. Inbreeding depression was positive for all studied characters.

The joint scaling test (Mather and Jinks, 1982) was employed to estimate the mean (m), additive effect (d), dominance effect (h), additive x additive (i), additive x dominance (j) and dominance x dominance (l) values (Tables 7&8). The results of A, B, C and D scaling test for the two wheat crosses under both environments, revealed that significant of any of these tests indicates the presence of non-allelic gene interactions or epistasis on the scale of measurement used. Results of scaling test, showed that additive-dominance model is inadequate for explaining the inheritance of all studied characters, indicating the present of non-allelic gene interaction in two crosses under two environments. Lal et al., (2013) studied the generation mean analysis in heat tolerance in wheat; they showed the adequacy of additive-dominance model for grain yield and its components.

The mean parameters (m) for all studied attributes of two crosses and environments (Tables 7&8) which reflect the contribution due to the over all mean plus the locus effects and interaction of the fixed loci were significant. The estimated of dominance

gene action (h) was significant for the all studied characters (N&D) in two crosses, indicating the importance gene effects in inheritance of these characters. The significant [d] and [h] in the inheritance of RWC (D in C₂) revealed that both types of additive and dominance effects are involved in the genetics of RWC (Farshadfar et al., 2001; 2003; 2008b; Tammam, 2005; Amin, 2013).

The genetic models fitted (Tables 7&8) for all studied characters (N&D) in two crosses except RWC (D in C₁), indicated dominance and additive x additive gene effects. indicated dominance and additive x additive gene effects. It is there fore suggested that selection should be carried out in late generation and the interaction should be fixed by selection under selfing conditions. The epistatic effect (dominance x dominance [1]) was significant for all studied characters (N&D) in two crosses, which confirm the important role of dominance x dominance gene interaction in the genetic system controlling, these result were reported by Srivastava et al., 1992; Kearsey and Pooni, 2004; Tammam, 2005; Amin, 2013. Both additive x additive [i] and dominance x dominance [1] effects were significant for all studied characters (N&D) in two crosses except RWC (D in C₁), supporting the presence of duplicate type of epistasis. This complementary interaction increases the variation between the generation and in the segregating population. The cross, which showed most promising in terms of narrow sense heritability and genetic gain, also showed highest means under both conditions, chance to find stress tolerant breeding material in segregating populations of this cross are promising. these finding are in line with Dashti et al., (2012), they studied genetic analysis of salt tolerance, and refer to High narrow sense heritability may be used as a useful indicator index for the selection of salt tolerant genotypes at the vegetative growth stage in wheat.

Reference

- Alza JO, Martinez JMF.** 1997. Genetic analysis of yield and related traits in sunflower (*Helianthus annuus* L.) in dryland and rigated environments. *Euphytica*, 95: 243-251.
- Amin IA.** 2013. genetics behaviour of some agronomic traits in two durum wheat crosses under heat stress. *Alex.J. Agric. Res.*, 58(1): 53-66.
- Burton GW.** 1951. Quantitative inheritance in pearl millet (*Pennisetum glaucum*).

292 Agronomy Journal, v.43:409-417.

293 **Dashti H, Bihamta MR, Shirani H, Majidi MM.** 2012. Genetic analysis of salt
 294 tolerance in vegetative stage in wheat (*Triticum aestivum* L) . POJ 5(1):19-23.

295 **Farshadfar E.** 2012. Application of integrated selection index and rank sum for
 296 screening drought tolerant genotypes in bread wheat. Intl J Agri Crop Sci. Vol., 4 (6),
 297 325-332.

298 **Farshadfar E, Ghandha M, Zahravi M, Sutka J.** 2001. Generation mean analysis of
 299 drought tolerance in wheat (*Triticum aestivum* L.). Acta Agron. Hung,49: 59-66.

300 **Farshadfar E, Mohammadi R, Aghaee M, Sutka J.** 2003. Identification of QTLs
 301 involved in physiological and agronomic indicators of drought tolerance in Rye
 302 using multiple selection index. Acta Agron. Hung., 51: 419-428.

303 **Farshadfar E, Aghaie M, Sharifi M, Yaghotipoor A.** 2008a. Assessment of salt
 304 tolerance inheritance in barley via generation mean analysis. J. Biol. Sci., 8: 461-
 305 465.

306 **Farshadfar E, Mahjouri S, Aghaee M.,** 2008b. Detection of epistasis and estimation of
 307 additive and dominant components of genetic variation for drought tolerance in
 308 Durum wheat. J. Biol. Sci., 8: 598-603.

309 **Gill KS, Dhillon SS, Bains KS.** 1972. Combinig ability and inheritance of yield
 310 components in crosses involving Indian and exotic wheat germplasm. Indian J.
 311 Genet. Pl. Breeding, 32: 421-430.

312 **Golparvar AR.** (2012): Heritability and mode of gene action determination for grain
 313 filling rate and relative water content in hexaploid wheat. - Genetika, Vol 44, No.
 314 1, 25 -32.

315 **Iqbal M, Navabi A, Salmon DF, Young RC, Mardoch BM, Moore SS, Spaner D.**
 316 2007. Genetic analysis of flowering and maturity time in high latitude spring
 317 wheat. Euphytica, 154: 207-218.

318 **Johnson HW, Robinson HF, Comstock RE.** 1955. Estimates of genetic and
 319 environmental variability in soybean. Agron. J., 47: 314-318.

320 **Kearsey MJ, Jinks JL.** 1968. A general method of detecting, additive, dominance and
 321 epistatic variation for metric traits.I. Theor. Heredity, 23: 403-409.

322 **Kearsey M, Pooni HS.** 2004. The Genetical Analysis of Quantitative Traits. 2nd Edn.,

- 323 Chapman and Hall, U.K., ISBN: 0-7487-4082-1.
- 324 **Ketata H, Smith EL, Edwards LH, McNew RW.** 1976. Detection of epistatic, additive
325 and dominance variation in winter wheat (*Triticum aestivum* L.). Crop Sci., 16: 1-
326 4.
- 327 **Kheiralla KA, El-Defrawy MM, Sherif THI.** 1993. Genetic analysis of grain yield,
328 biomass and harvest index in wheat under drought stress and normal moisture
329 conditions. Assiut J. Agric. Sci., 24: 163-183.
- 330 **Lal C, Maloo SR, Kumar V.** 2013. Generation mean analysis for some heat tolerance
331 and quantitative characters in bread wheat (*Triticum aestivum* L.). J. Wheat Res. 5
332 (2): 22-26.
- 333 **Lamkey KR, Lee M.** 1993. Quantitative genetics, molecular markers and plant
334 improvement. In: Focussed plant improvement- towards responsible and
335 sustainable agriculture. ed. imrie, b.c. and hacker, j.b. PP. 104-115.
- 336 **Manette A, Schonfeld C, Richard J, Carre B, Morhinweg W.** 1988. Water relations in
337 winter wheat as drought resistance indicators. Crop Sci., 28: 526-531.
- 338 **Mather K, Jinks JL.** 1982. Biometrical Genetics. 3rd Edn. Chapman and Hall Ltd.,
339 ISBN-10: 0412228904.
- 340 **Moshref MK.** 1996. Genetical and statistical studies in wheat. Ph.D. Thesis, Al-Azhar
341 Univ., Egypt.
- 342
- 343 **Naroui Rad MR, Abdul Kadir M, Rafii MY, Jaafar ZEH , Naghavi5 MR, Ahmadi**
344 **F.** 2013. Genotype \times environment interaction by AMMI and GGE biplot analysis
345 in three consecutive generations of wheat (*Triticum aestivum*) under normal and
346 drought stress conditions. AJCS 7(7):956-961.
- 347 **Nyguist WE.** 1991. Estimation of heritability and prediction of response in plant
348 populations. CRC Critical Reviews in Plant Science, 10(3): 235-322.
- 349 **Ritchie SW, Nguyen HT, Holiday AS.** 1990. Water relations in winter wheat as drought
350 resistance indicators. Crop Sci., 28: 526-531.
- 351 **Sharma G, Singh RB.** 1976. Inheritance of plant height and spike length in spring
352 wheat. Indian J. Genet. Pl. Breeding, 36: 173-183.
- 353 **Sharma BD, Sood BC, Halotra VV.** 1991. Studies on variability, heritability and

- genetic advance in chickpea research. Indian J. Pulses Res., 3: 1-6.
- Sharma SN.** 2003. Genetics of spike length in durum wheat. Euphytica, 130: 155-161.
- Singh P, Narayanan SS.** 1993. Biometrical Techniques in Plant Breeding. 1st Edn., Kalayani publishers, New Dehli, India.
- Singh RK, Chaudhary BD.** 1999. Biometrical Methods in Quantitative Genetic Analysis. 1st Edn., Kalyani Publisher, New Dehli, India, ISBN: 81-7663-307-0.
- Smith, H.H.,** 1952. Fixing Transgressive Vigour in *Nicotiana Rustica*. In: Heterosis. Iowa State College Press, Ames, IA, USA.
- Sood S, Kalia NR, Bhateria S, Kumar S.** 2006. Detection of genetic components of variation for some biometrical traits in *Linum usitatissimum* L. in sub-mountain Himalayan region. Euphytica, 155: 107-115.
- Sprague GF.** 1963. Orientation and objectives In “Statistical genetics and plant breeding” Nat. Acad. Sci. N.R.C. Pub. 982. IX-XV.
- Srivastava RB, Sharma SC, Yunus M.** 1992. Additive and non-additive gene effects for yield and yield components in two crosses of wheat (*T. aestivum* L.). India J. Genet., 52: 297-301.
- Tammam AM.** 2005. Generation mean analysis in bread wheat under different environmental conditions. Minufiy J. Agri. Res., 30(3): 937-956.
- Tefra H, Peat HE.** 1997. Genetics of grain yield and other agronomic characters in (*Eragrostis tef* Zuccotrotter) II the triple test cross. Euphytica, 96: 193-202.
- Umarahan P, Ariyanayagam RP, Haque SQ.** 1997. Genetic analysis of yield and its components in vegetable cowpea (*Vigna unguiculata* L. walp). Euphytica 96: 207-213.
- Warner JN.** 1952. A method of estimating heritability. Agron. J., 44: 427-430.
- Zaazaa EI, Hager MA, El-Hashash EF.** 2012. Genetical Analysis of Some Quantitative Traits in Wheat Using Six Parameters Genetic Model. American-Eurasian J. Agric. & Environ. Sci., 12 (4): 456-462.

Table 1: Pedigree and origin of the genotypes used in the two bread wheat crosses.

Cross	Parental name	Pedigree	Origin
Cross 1	Sakha 94 (P1)	Opata/Rayon//Kauz	Egypt
	Tokwie (P2)	-----	South Africa
Cross 2	Giza 168 (P1)	Mill/Buc//Seri	Egypt
	Tokwie (P2)	-----	South Africa

Table 2: Analysis of variance for various characters investigated.

Table 2. Analysis of variance for various characters investigated.							
SOV	df	Mean square					
		NS	SW	BY	GY	RWC	CC
Cross 1							
Environments (A)	1	9.61**	5.58**	10859.21**	715.70**	1234.65**	659.63**
Error	2	0.05	0.08	7.02	45.63	3.33	2.41
Generations (B)	5	8.94**	0.82**	283.17*	191.21**	120.94**	227.39**
A x B	5	0.35 ^{ns}	0.13 ^{ns}	25.55 ^{ns}	7.54 ^{ns}	12.54 ^{ns}	11.98 ^{ns}
Error	20	0.25	0.07	84.49	14.90	15.23	1.05
Cross 2							
Environments (A)	1	14.06**	9.06**	11600**	620.63**	1441.29**	1416.27**
Error	2	0.05	0.001	149.70	0.08	11.95	2.94
Generations (B)	5	18.49**	0.53	269.52**	207.59**	532.92**	179.34**
A x B	5	1.03**	0.10**	20.87**	5.71 ^{ns}	53.83**	58.96**
Error	20	0.17	0.02	48.62	4.13	10.39	4.26

* and ** significant at 5% and 1% levels of probability, respectively.

Table 3: Mean comparison of the characters studied.

Generations	Characters											
	NS		SW		BY		GY		RWC		CC	
	N	D	N	D	N	D	N	D	N	D	N	D
Cross 1												
Gemmeiza 9 (P ₁)	14.87	13.17	6.10	4.84	100.97	64.60	44.49	33.76	67.97	54.67	46.47	40.17
Inbred line 1 (P ₂)	10.67	10.17	5.63	4.80	96.98	60.91	37.70	30.58	71.21	61.73	59.70	49.37
F ₁ (P ₁ x P ₂)	16.07	14.67	6.30	5.56	109.32	68.29	48.41	41.03	74.02	66.91	52.57	45.40
F ₂	13.17	12.17	5.11	4.63	82.69	52.95	32.52	23.16	69.85	55.21	39.37	33.93
P ₁ x F ₁ (BC ₁)	14.67	13.67	5.74	4.91	96.16	65.05	40.66	30.49	67.14	53.35	46.20	37.43
P ₂ x F ₁ (BC ₂)	12.67	12.07	5.36	4.78	95.70	61.61	36.93	28.18	68.50	56.55	54.30	40.93
LSD _{0.05}	1.88	1.01	0.45	0.42	4.25	5.19	4.31	3.85	2.01	1.36	3.08	3.94
Potence ratio	-1.57	-2.00	1.83	4.27	5.19	3.00	1.89	4.32	-2.73	14.97	-1.10	1.41
Cross 2												
Sids 1 (P ₁)	14.67	12.67	5.75	4.62	101.76	67.85	42.40	34.07	66.97	52.73	53.70	39.33
Inbred line 2 (P ₂)	10.67	10.17	5.63	4.80	96.98	60.91	37.70	30.58	71.21	61.73	59.70	49.37
F ₁ (P ₁ x P ₂)	16.67	14.17	6.30	4.95	111.44	79.03	47.11	38.91	86.14	78.36	67.73	44.40
F ₂	14.17	13.67	5.13	4.52	96.58	59.19	30.39	22.14	56.74	51.66	43.87	39.33
P ₁ x F ₁ (BC ₁)	14.17	13.17	6.09	5.00	100.03	62.42	35.41	26.40	74.97	55.87	52.53	40.43
P ₂ x F ₁ (BC ₂)	12.67	11.67	5.75	4.73	98.63	60.62	39.72	28.80	81.87	61.60	55.37	44.77
LSD _{0.05}	1.90	1.20	0.51	0.46	3.52	2.59	2.52	2.32	1.49	2.41	2.93	2.89
Potence ratio	-2.00	-2.20	-10.22	2.70	5.05	4.22	-2.40	-3.78	8.04	36.91	19.46	0.11

Table 4: Grain yield/plant under normal (GY_N) and drought stress (GY_D), and stress tolerance index (STI) for each generation.

Generations	GY _N	GY _D	STI	Generations	GY _N	GY _D	STI
Cross 1				Cross 2			
Gemmeiza 9 (P ₁)	44.49	33.76	0.76	Sids 1 (P ₁)	42.40	34.07	0.80
Inbred line 1 (P ₂)	37.70	30.58	0.81	Inbred line 2 (P ₂)	37.70	30.58	0.81

F ₁ (P ₁ x P ₂)	48.41	41.03	0.85	F ₁ (P ₁ x P ₂)	47.11	38.91	0.83
F ₂	32.52	23.16	0.71	F ₂	30.39	22.14	0.73
P ₁ x F ₁ (BC ₁)	40.66	30.49	0.75	P ₁ x F ₁ (BC ₁)	35.41	26.40	0.75
P ₂ x F ₁ (BC ₂)	36.93	28.18	0.76	P ₂ x F ₁ (BC ₂)	39.72	28.80	0.73

Table 5: Genetic parameters and components of variation for all studied characters in the cross 1 under normal (N) and drought stress (D) conditions.

Characters		h/d	H _b	H _n	G _A	D	H	F	E _w	√H/D	F/√HxD
NS	N	+2.65	0.69	0.28	25.06	19.80	9.07	+1.40	5.42	0.68	0.11
	D	+3.63	0.67	0.20	22.83	13.40	17.53	-0.80	5.50	1.14	-0.05
SW	N	+5.79	0.78	0.33	8.87	7.21	2.80	-0.11	1.19	0.62	-0.03
	D	+12.31	0.83	0.32	12.55	9.30	5.76	+0.60	1.27	0.79	-0.08
BY	N	+134.72	0.79	0.36	20.29	17.82	3.75	-1.60	2.56	0.46	-0.20
	D	+13.69	0.74	0.33	25.05	21.56	5.51	-1.55	4.34	0.51	0.14
GY	N	+8.94	0.79	0.29	56.63	40.35	29.26	-2.36	7.49	0.85	-0.07
	D	+13.67	0.74	0.24	46.60	28.68	33.12	+0.99	7.90	1.08	0.03
RWC	N	+2.70	0.74	0.32	21.36	18.03	5.42	-1.88	3.73	0.55	-0.19
	D	-2.40	0.76	0.37	23.76	22.37	1.39	-1.16	3.74	0.25	-0.21
CC	N	-5.31	0.73	0.21	23.43	13.25	19.00	-1.20	4.13	1.20	-0.08
	D	-6.18	0.74	0.18	19.92	9.20	20.27	2.24	3.36	1.48	0.16

Table 6: Genetic parameters and components of variation for all studied characters in the cross 2 under normal (N) and drought stress (D) conditions.

Characters		h/d	H _b	H _n	G _A	D	H	F	E _w	√H/D	F/√HxD
NS	N	+0.67	0.73	0.30	27.98	22.50	9.33	-0.75	5.00	0.64	-0.05
	D	-1.50	0.70	0.29	25.06	20.70	7.27	-0.35	5.17	0.59	-0.03
SW	N	+11.06	0.81	0.32	10.66	8.09	4.51	+0.03	1.21	0.75	0.005
	D	+6.07	0.75	0.29	6.71	5.01	3.01	+0.50	1.08	0.78	0.13
BY	N	+16.48	0.66	0.31	17.92	16.21	2.37	-0.32	2.91	0.63	-0.02
	D	+13.32	0.77	0.26	22.80	14.91	13.45	-1.67	4.50	0.38	-0.27
GY	N	-8.51	0.71	0.24	47.64	30.87	30.76	-2.37	9.52	1.00	-0.08
	D	-11.79	0.75	0.30	53.60	41.71	20.66	+0.75	8.47	0.70	-0.03

RWC	N	-15.04	0.79	0.25	23.83	14.53	17.22	+2.08	3.16	1.09	0.11
	D	-8.63	0.77	0.19	19.62	20.62	8.74	+1.30	2.81	0.65	0.10
CC	N	-18.15	0.69	0.20	19.73	11.20	15.91	+0.88	4.25	1.19	0.07
	D	-3.03	0.80	0.26	20.01	12.34	14.16	+0.03	2.41	1.07	0.002

Table 7: Estimates of scaling test and types of gene action using generation means for all studied characters cross 1 under normal (N) and drought stress (D) conditions.

Characters		Scaling test				Genetic parameters					
		A	B	C	D	m	[d]	[h]	[i]	[j]	[l]
NS	N	-1.60**	-1.40**	-5.00**	-1.00**	13.17**	2.00	5.30**	2.00**	-0.10	1.00**
	D	-0.50**	-0.70**	-4.00**	-1.40**	12.17**	1.60	5.80**	2.80**	0.10	-1.60**
SW	N	-0.91**	1.20**	-3.88**	-0.88**	5.11**	0.38	2.20**	1.77**	0.14	0.45**
	D	-0.57**	-0.80**	-2.23**	-0.43**	4.63**	-0.13	1.60**	0.86*	0.12	0.52**
BY	N	-17.97**	-14.91**	-85.85**	-26.49**	82.69**	0.47	63.32**	52.97**	-1.53	-20.09**
	D	-2.79**	-5.97**	-50.30**	-20.77**	52.95**	3.44	47.08**	41.54**	1.60	-32.78**
GY	N	-11.58**	-10.26**	-46.95**	-12.56**	32.52**	3.74	33.43**	25.11**	-0.66	-3.28**
	D	-11.81**	-13.25**	-49.78**	-12.36**	23.16**	2.31	31.58**	24.72**	0.72	-0.34**
RWC	N	-7.71**	-8.23**	-7.84**	4.05**	69.85**	-1.36	-3.67*	-8.10**	0.26	24.04**
	D	-14.89**	-15.54**	-29.40**	0.51**	55.21**	-3.21	7.69**	-1.03	0.32	31.46**
CC	N	-6.63**	-3.67**	-53.83**	-21.77**	39.37**	-8.10	43.02**	43.53**	-1.48	-33.23**
	D	-10.70**	-12.90**	-44.60**	-10.50**	33.93**	-3.50	21.63**	21.00**	1.10	2.60**

* and ** significant at 5% and 1% levels of probability, respectively.

Table 8: Estimates of scaling test and types of gene action using generation means for all studied characters in the cross 2 under normal (N) and drought stress (D) conditions

Characters		Scaling test				Genetic parameters					
		A	B	C	D	m	[d]	[h]	[i]	[j]	[l]
NS	N	-3.00**	-2.00**	-2.00**	1.50**	14.17**	1.50	1.00*	-3.00**	-0.50	8.00**
	D	-0.50**	-1.00**	3.50**	2.50**	13.67**	1.50	-2.25**	-5.00**	-0.25	6.50**
SW	N	0.13**	-0.43**	-3.88**	-1.57**	5.13**	0.34	3.76**	3.15**	0.28	-2.85**
	D	0.43**	-0.29**	-2.23**	-0.70**	4.52**	0.27	1.64**	1.40**	0.36	-1.54**
BY	N	-13.14**	-11.16**	-35.30**	-5.50**	96.58**	1.40	23.07**	11.00**	-0.99	13.30**
	D	-22.04**	-18.70**	-50.05**	-4.66**	59.19**	1.80	23.98**	9.32**	-1.67	31.41**

GY	N	-18.70**	-3.37**	-50.77**	-14.36**	30.39**	-4.32	36.77**	28.71**	-7.67	-6.65**
	D	-20.19**	-11.89**	-53.91**	-10.92**	22.14**	-2.41	28.42**	21.83**	-4.15	10.25**
RWC	N	-3.18**	6.39**	-83.50**	-43.36**	56.74**	-6.90	103.7**	86.71**	-4.78	-89.92**
	D	-19.35**	-16.88**	-64.55**	-14.16**	51.66**	-5.73**	49.44**	28.31**	-1.23	-7.91**
CC	N	-16.37**	-16.70**	-73.40**	-20.17**	43.87**	-2.83	51.37**	40.33**	0.16	-7.27**
	D	-2.87**	-4.23**	-20.17**	-6.53**	39.33**	-4.33	13.12**	13.07**	0.68	-5.97**

* and ** significant at 5% and 1% levels of probability, respectively




Fig 1&2: Percentage of heterosis and inbreeding depression under two environments in Cross 1 for characters investigated.




Fig 3&4: Percentage of heterosis and inbreeding depression under two environments in Cross 2 for characters investigated.