

Effect of soil zinc and boron on the yield and uptake of wheat in an acid soil of West Bengal, India

Abstract: The production of wheat (*Triticum aestivum* L.), an important staple food in the world, is often restricted due to micronutrients status in soil. Micronutrient deficiency in soil including boron (B) and zinc (Zn) is quite widespread in Asian countries including India due to prevalent soil and environmental conditions. A field experiment was conducted following randomized complete block design over a two-year period in an acid soil of *Terai* region of West Bengal to study the effect of zinc and boron on the yield and uptake of nutrients by wheat. The highest grain yield (4.4 t ha^{-1}) was obtained after the combined application of Zn and B over that of other treatment combinations (variable rates of B and Zn application with nutrients) or control (no nutrients, B and Zn). Application of one micronutrient might have accelerated the uptake of other micro- and macro-nutrients (such as B, Zn, nitrogen, phosphorus and potassium) resulting in higher yield. A positive correlation was observed between the grain yield and the uptake of different nutrients with the weakest with Zn. An enhancement of the nutrients in soils was also observed at the harvest. High response from a combined application of B and Zn clearly demonstrated the necessity of micronutrients for improving production in the studied regions with acid soils. Therefore, an application of a mixture of micronutrients is recommended over a single micronutrient for the acid soil regions of West Bengal in order to get a better response from the applied nutrient sources and thus the production.

Key words: micronutrients, synergistic effect, *Terai* region, deficiency, growth stages

Introduction

Wheat (*Triticum aestivum* L.) is the most important staple food for humans and is grown on more land than any other commercial crops in the world. It was grown on 216.6 million ha land in 2012 [1] producing 674.9 million tonnes of wheat globally, the third most produced cereal (perhaps any food crop) after maize (875.1 million tonnes) and rice (718.3 million tonnes) [1]. In 2012, India produced 94.9 million tonnes of wheat from a cultivated area of 29.9 million ha, the largest area devoted to wheat production by any country in the world [2].

Comment [MA1]: Check spacing

Comment [MA2]: spacing

Comment [MA3]: spacing; an before "e:"

Comment [MA4]: spacing

Comment [MA5]: spacing

31 With the demand of ever-increasing population, the present day agriculture became more
32 intensive and mined available nutrients from soil over years. However, one of the major
33 triggering factors behind the dramatic improvement in the production and yield of wheat was the
34 supply of artificial nutrient source for plant growth and development especially the use of
35 synthetic nitrogen fertilizer. Potash and phosphorus fertilizer in addition to the nitrogen fertilizer
36 supplied the major nutrients for the growth, development and production of wheat. In addition to
37 these major (macro) nutrients, there are some nutrients, which are essential for wheat growth but
38 needed only in very small (micro) quantities. Among these, boron (B), zinc (Zn), iron, (Fe),
39 copper (Cu), manganese (Mn), and chlorine (Cl) are known to have effect on the grain- as well
40 as straw-yield of wheat. These micronutrients play a pivotal role in the yield improvement of
41 wheat crop [3]. They are needed in trace amount while the adequate supply improves nutrient
42 availability and positively affects the cell physiology that is reflected in yield as well [4, 5]. A
43 number of micronutrients are part of the photosynthesis and respiration processes, chlorophyll
44 formation, nucleic acid and protein synthesis, nitrogen-fixation and other biochemical pathways
45 [6-8]. However, the deficiencies of micronutrients are wide spread in many Asian countries
46 including India due to calcareous nature of soils, high pH, low organic matter, salt stress,
47 prolonged draught, high bicarbonate content in irrigation water and imbalanced application of
48 NPK fertilizers [9, 10]. The deficiency of micronutrients can induce the stress in plants including
49 low crop yield and quality, imperfect plant morphological structure (such as fewer xylem vessels
50 of small size), widespread infestation of various diseases and pests and low fertilizer use
51 efficiency.

Comment [MA6]: spacing

52 Zinc is one of the important micronutrients, which is important in the production of various
53 crops including wheat [11, 12]. It improves the number of grains per spike [13]. In addition to
54 the yield [14, 15], adequate supply of Zn can improve the water use efficiency of wheat plants
55 [16]. It also provides thermo-tolerance to the photosynthetic apparatus [17]. It is important in
56 plant metabolism and thus the growth and production of wheat [18]. The Zn is the third most
57 common deficient nutrient after N and P [19, 20]. Zinc deficiency in plants not only reduces the
58 grain yield, but also the nutritional quality of crops [21].

Comment [MA7]: spacing

59 Boron is another important micronutrient that is essential for plant growth and improves the
60 production efficiency of wheat. However, the deficiency of B is the most frequently encountered
61 in field [22]. Boron is essential for cell division and elongation of meristematic tissues, floral

62 organs and the flower male fertility, pollen tube germination and its elongation and the seed and
63 fruit formation. Lack of B can cause the ‘wheat sterility’ resulting in increased number of open
64 spikelets and decreased number of grains per spike [23]. The B deficiency in soil can affect
65 seedling emergence and cause an abnormal cellular development in young wheat plant [24]. It
66 also inhibits root elongation by limiting cell division in the growing zone of root tips [25].
67 Deficiency of B is known to inhibit the leaf expansion and reduction in photosynthesis. In the
68 field, sexual reproduction is often affected by low B reducing the grain yield significantly
69 without any visual symptoms expressed during vegetative growth.

70 The *Terai* region is located at the south of the outer foothill of the Himalaya and Siwalik hills
71 and the north of the Indo-Gangetic plain. It spreads over a number of states in India including
72 Himachal Pradesh, Haryana, Uttaranchal, Uttar Pradesh, West Bengal, Sikkim and Assam. It also
73 covers a major part in Nepal, Bhutan and Bangladesh. The *Terai* region is the habitat of millions
74 of people. It is a very productive region and agriculture is the base of the economy of the
75 habitants. Rice and wheat are important crops of this region. The rice-wheat system is the most
76 important cropping pattern in this region and considered to be the major determinant factor of the
77 agriculture-based economy. However, the intensive cultivation practices overexploited the
78 natural soil resource base, which was further enhanced by the imbalanced use of inputs [22, 26].

79 The deficiency of B and Zn in soils of different agro-climatic zones is not rare and *Terai*
80 region is not an exception in this regard. Deficiency of different micronutrients has been reported
81 from this region. Among the most prevalent ones, the deficiency of Zn is estimated to be the
82 highest [27, 28]. Incidence of B deficiency from the areas of West Bengal and Bihar has also
83 been reported [26]. Comprehensive study on the effect as well as the interaction of these
84 nutrients on the production of wheat at this part of the world would help understanding
85 constraints of cultivation and decreasing the yield gap to secure food for the future.

86 Based on the above perspectives the present study was undertaken in the *Terai* region of
87 West Bengal 1) to assess the effect of Zn and B on the yield of wheat, 2) to examine the interaction
88 effect of Zn and B on the yield of wheat and 3) to evaluate the residual status of Zn and B in soil
89 at different stages of wheat crop.

90 **Materials and Methods**

91 **Experimental site:**

Comment [MA8]: spacing

Comment [MA9]: spacing

A field experiment was carried out at the agricultural farm of Uttar BangaKrishiViswavidyalaya, Pundibari, Cooch Behar, West Bengal, India. The farm is located within the Terai Agro-climatic zone and its geographic location is 26°19'86" N latitude and 89°23'53" E longitude. The elevation of the farm is 43 meters above the mean sea level. The field experiment was carried out in the same field during the winter season (*Rabi season*) of 2010-11 and 2011- 12.

Comment [MA10]: spacing

Comment [MA11]: spacing

Comment [MA12]: spacing

Experimental plots:

The local topography of the study area is almost flat with good drainage facilities. The soil of the experimental site is sandy loam in texture (sand- 60%, Silt- 21% and Clay- 19%). Before laying out the experimental plots, a set of surface soil samples was collected over the whole experimental area, composite together and tested in the laboratory following the methods described in the following sub-section. The measured physical, chemical and physico-chemical properties (Table 1) were used as the baseline measurement for the experimental plots.

Table 1: Initial characteristics of experimental soil for two years

Characteristics	Measurements	
	2010-11	2011-12
pH	5.00	5.00
EC (dSm ⁻¹)	0.05	0.05
Organic Carbon (%)	1.04	1.02
Nitrogen (kg ha ⁻¹)	206.98	188.16
Phosphorus (kg ha ⁻¹)	0.76	0.89
Potassium (kg ha ⁻¹)	89.60	88.48
Boron (kg ha ⁻¹)	0.68	0.62
Zinc (kg ha ⁻¹)	0.73	0.84

A set of 30 experimental plots (5 m × 4 m) were laid out following randomized complete block design (RCBD) for this experiment. Ten treatment combinations (Table 2) were developed following three doses of B (0 kg ha⁻¹, 5 kg ha⁻¹ and 10 kg ha⁻¹), three doses of Zn (0 kg ha⁻¹, 12.5 kg Zn Sulphate ha⁻¹ and 25 kg Zn Sulphate ha⁻¹) and a treatment without application of any nutrients (T₁₀). Though the treatment T₁ (B₀Zn₀) received recommended doses of N, P, and K, treatment T₁₀ (control) did not receive any nutrient or micronutrients (B and Zn). The treatments were replicated three times in this field experiment.

115 **Table 2: Details on the experimental plots and treatment combinations**

Experimental details	
Crop	: Wheat (<i>Triticum aestivum</i> L.)
Variety	: NW 1014
Experimental design	: Randomized Complete Block Design
Total Area	: 801m ²
Plot size	: 5 m × 4 m
Number of replication	: 3
Spacing	: 23 cm (Row to Row)
Treatments	: T ₁ -B ₀ Zn ₀ , T ₂ -B ₀ Zn ₁ , T ₃ -B ₀ Zn ₂ , T ₄ -B ₁ Zn ₀ , T ₅ -B ₁ Zn ₁ , T ₆ -B ₁ Zn ₂ , T ₇ -B ₂ Zn ₀ , T ₈ -B ₂ Zn ₁ , T ₉ -B ₂ Zn ₂ , T ₁₀ - Control. B ₀ = without boron Zn ₀ = without zinc sulphate B ₁ = 5 kg ha ⁻¹ of boron Zn ₁ = 12.5 kg ha ⁻¹ of zinc sulphate B ₂ = 10 kg ha ⁻¹ of boron Zn ₂ = 25 kg ha ⁻¹ of zinc sulphate

Comment [MA13]: spacing

116

117 **Field operations:**

118 The land preparation for this experiment was started with a deep ploughing (21 and 22,
119 December 2010 and 12 and 13 December 2012) using a tractor. A laddering (similar to levelling
120 of soil surface) was performed after a day of soil drying following two secondary tillage using a
121 power tiller in order to prepare a good soil tilth. The weeds and stubbles were removed by hand
122 picking and the final laddering was performed to prepare the seed bed. Bunds and channels were
123 prepared manually to prepare the experimental plots following the specifications mentioned in
124 Table 1. Nitrogen (N, 100 kg ha⁻¹), phosphorus (P, 60 kg ha⁻¹) and potassium (K, 30 kg ha⁻¹), in the
125 form of urea, single super phosphate, muriate of potash; B as Borax (10 kg/ha, sodium borate),
126 and Zn as zinc sulphate (25 kg ha⁻¹) were applied to the soil as per the treatments. Full dose of P,
127 K, B and Zn and half of the recommended dose of N were surface applied as basal dose and
128 incorporated in the soil. The remaining half of the recommended dose of N was applied as top
129 dressing at 21 days after sowing (DAS), after completion of the first weeding.

130 The wheat variety of NW-104 was used for this experiment at the rate of 100 kg ha⁻¹. Sowing
131 was completed in rows (spacing 23 cm) in North-South using a duck-foot tyne at a depth of 2.5
132 to 3 cm. Two weeding operations were performed manually on 21 DAS and 45 DAS. Two
133 irrigations were applied on 21 DAS (after weeding and fertilizer application) and 65 DAS. The
134 excess water was drained out using drainage channels.

135 The soil and plant samples were collected for laboratory analysis on 21 (CRI- crown root
 136 initiation stage), 55 (tillering stage), 70 (booting stage) and 110 (maturity) DAS. Leaving the
 137 border rows, half of the area in each plot was marked for recording biometrical observation
 138 including destructive plant sampling and other half for recording yield components and yield of
 139 wheat. The height (from ground level) of five randomly selected plants were recorded and
 140 averaged from each plot. The measured plants were tagged after first measurement for
 141 subsequent measurements. Dry weight of both roots and shoots were also recorded. The number
 142 of tillers per m² was recorded from 10 randomly selected plants. The crop was harvested from
 143 net plot area discarding the border row. The number of spikes per plant was recorded from 10
 144 randomly selected plants and converted to number of spikes per m². Length of spikes was
 145 measured prior to harvest and average length was calculated. Number of grains per spikes as well
 146 as 1000 grain dry weight were also recorded for each treatment. The final yield of wheat and
 147 straw was recorded after sun drying and thrashing. The yields were recorded and calculated as
 148 tonne per ha following,

149
$$\text{Grain yield (t ha}^{-1}\text{)} = (\text{Plot yield (kg)} \times 10000 / \text{Plot size (m}^2\text{)} \times 1000)$$

150 **Analytical methods:**

151 Collected soil and plant samples were tested for a series of parameter in laboratory. pH and
 152 electrical conductivity (EC) of soil samples was determined in suspensions (soil:water 1:2.5)
 153 using a Systronics glass electrode-pH meter and a Systronics digital conductivity meter (Model
 154 no. 304), respectively [29]. Organic carbon (OC) content of soil samples was estimated by
 155 Walkley and Black's titration method [30]. Mechanical analysis of soil samples was carried out
 156 following the hydrometer method [31]. The textural class of the soils was ascertained from the
 157 particle-size distribution of sand, silt and clay particles. Available nitrogen (N) in soil and plant
 158 samples was determined by alkaline KMnO₄ method following Subbiah and Asija [32]. Available
 159 P in soil and plant was determined by extracting the samples with a mixture of 0.03 M NH₄F and
 160 0.025 M HCl [33] followed by colorimetric measurement at 880 nm using spectrometer
 161 (Systronics Model No. 167) [34]. Available K in soil and plant was measured using a flame
 162 photometer (Systronics Model No. 128) [34]. The extraction was carried out with neutral normal
 163 ammonium acetate. DTPA-(Diethylenetriaminepentaacetic acid) extractable Zn⁺² of soil and
 164 plant samples were determined by extraction with the extractant containing 0.005M DTPA,
 165 0.01M CaCl₂ and 0.1M Triethanol amine buffered at pH 7.3 [35] followed by the measurement

166 using Atomic Absorption Spectrophotometer (AAS). Available Boron in soil and plant was
167 extracted by boiling a known amount of samples with double distilled water (in 1:2.5 ratio)
168 prepared by quartz glass distillation apparatus, for five minutes under a reflux condenser,
169 followed by cooling and filtration [36]. The concentration was measured using AAS.

170 **Statistical analyses**

171 Statistical analysis for the collected data was performed in SigmaPlot (Systat Software Inc.).
172 The significant difference between the treatments was tested using ANOVA and LSD. The
173 interaction between the effect of B and Zn was tested using two-way ANOVA. The correlation
174 between the yield components and nutrient uptakes were also calculated. The figures were
175 prepared using the SigmaPlot.


176 **Results and Discussion**

177 The yield components and grain yield of wheat are shown in Table 3. A significant difference
178 was observed among the treatment combinations on yield components and grain yield of wheat.
179 The maximum mean grain yield (4.4tha^{-1}) was observed in the treatment T_6 (B_1Zn_2), while
180 minimum was observed in the control (1.7tha^{-1}). Relatively higher yield was obtained from the
181 treatments T_1 (B_0Zn_0) to T_9 (B_2Zn_2) over that of the control (T_{10}). The lowest harvest index was
182 observed in T_4 (B_1Zn_0) and the highest in controls (T_{10}). The application of B and Z in
183 combination significantly ($p < 0.05$) increased the grain yield of wheat. The grain yield increase
184 with B and Zn addition was reported by Chaudry et al. [37]. Boron concentration has been reported
185 to increase grain yield of durum wheat by 16% [38]. This may be due to the requirement of B in
186 wheat during the vegetative stage leading to high response to the grain yield [39, 40]. Therefore,
187 even a small amount of Zn and B directly affected the grain yield. Mandal [41] reported a direct
188 relationship between the number of grains and tillers and the wheat yield under B deficient soils
189 of Terai region of West Bengal.

190 The effect of B on the grain and straw yield was significant at $\alpha = 0.05$ (95% significant
191 level). However, the scenario was little different for Zn. For example, without any addition of B,
192 Zn had no effect on crop yield, while with regular dose of B (5 kg Bha^{-1}) application, the yield
193 increased linearly. This indicated that with a regular dose of B, the efficiency of Zn increased (at
194 least for the application rate considered in this study). However, with excess (more than regular)
195 application of B, the effect of Zn decreased indicating antagonistic effect between the
196 micronutrient at high dose, specifically B. The two-way ANOVA following a general linear

model with $\alpha = 0.05$ showed a significant interaction between the effect of B and Zn on the grain and straw yield of wheat. This means that the difference in the mean values among the different levels of B and Zn is great enough to exclude the possibility that the difference is just not due to random sampling variability after allowing for the effects of differences in Zn and B, respectively. Therefore, care should be taken in deciding the amount of micronutrient application, which may have different effect. From this study, it could be suggested to choose a regular dose of B for better efficiency of Zn. Sometime a high dose of Zn could be even beneficial with a controlled application of B.

In spite of the highest dry biomass production until the booting stage in T_6 (B_1Zn_2), the T_8 (B_2Zn_1) produced the highest dry straw at maturity (Fig. 1). Combination of B and Zn might have boosted the vegetative growth during the early stage, while the high amount of Zn along with a regular dose of B improved the yield and yield components of wheat at maturity [14, 15]. A combination of regular dose of Zn and B (T_5) could not produce high amount of straw compared to other treatments with single or double dose of either Zn or B or in combination (Fig. 1). For example high straw yield with very little difference was observed among treatments T_4 (B_1Zn_0), T_6 (B_1Zn_2), T_7 (B_2Zn_0) and T_8 (B_2Zn_1). The lowest biomass production was recorded in control (T_{10}) at all stages of crop growth.



216 **Fig. 1:**Effect of treatments on average straw yield (kg ha^{-1}) over two years at different stages of
217 wheat growth. The standard deviation of measurement is shown as error bars. The CRI stage
218 indicates crown root initiation.

219
220 A significant difference in the nutrient uptake was recorded in different treatments and at
221 different growth stages (Fig. 2). The highest uptake (kg ha^{-1}) of N over the entire growth period
222 was recorded in treatment T_6 (B_1Zn_2) and minimum in T_{10} (control). The maximum amount of N
223 uptake at different growth stages was not consistent. For example, T_2 (B_0Zn_1) was recorded with
224 the highest amount of N uptake during booting stage. While the highest amount of P uptake was
225 recorded in T_7 (B_2Zn_0), the highest amount of K uptake was recorded in T_8 (B_2Zn_1). Similar to N
226 uptake, a variable amount of P and K uptake was also recorded at different growth stages in
227 different treatments. The highest amount of B and Zn uptake was recorded in treatment T_9
228 (B_2Zn_2). High amount of B and Zn application might show some synergistic effect to provide
229 higher amount of uptake. The treatment T_{10} (control) always recorded with the least amount of
230 nutrient uptake.

Table 3: Effect of treatments on the yield components and grain yield (tha⁻¹) of wheat. The data from 2010-11 and 2011-12 are shown along with the mean (average) over two years. S.D. stands for standard deviation.

Treatments	TillerSq. m. ⁻¹			GrainsSpike ⁻¹			1000 Grain weight			Grain Yield (tha ⁻¹)			Harvest Index (%)		
	2010-11	2011-12	Mean (S.D.)	2010-11	2011-12	Mean (S.D.)	2010-11	2011-12	Mean (S.D.)	2010-11	2011-12	Mean (S.D.)	2010-11	2011-12	Mean (S.D.)
T1	171	168	169.5 (2.12)	37	36	36.5 (0.70)	45.72	45.75	45.74 (0.02)	2.89	2.77	2.83 (0.08)	27.7	27.6	27.6 (0.07)
T2	156	156	156.0 (0)	41	42	41.5 (0.70)	42.74	42.80	42.77 (0.04)	2.73	2.80	2.77 (0.05)	29.5	29.8	29.7 (0.21)
T3	165	162	163.5 (2.12)	39	39	39.0 (0)	45.92	45.72	45.82 (0.14)	2.95	2.89	2.92 (0.04)	30.6	30.1	30.4 (0.35)
T4	143	145	143.8 (1.41)	43	43	43.0 (0)	44.54	44.00	44.27 (0.38)	2.73	2.74	2.74 (0.01)	25.2	24.9	25.1 (0.21)
T5	158	150	153.8 (5.66)	42	42	42.0 (0)	51.50	51.51	51.51 (0.01)	3.41	3.25	3.33 (0.11)	44.1	43.5	43.8 (0.42)
T6	188	185	186.3 (2.12)	48	51	49.5 (2.12)	47.64	47.34	47.49 (0.21)	4.29	4.47	4.38 (0.13)	34.6	35.2	34.9 (0.42)
T7	176	174	174.8 (1.41)	50	53	51.5 (2.12)	43.80	43.83	43.82 (0.02)	3.84	4.04	3.94 (0.14)	32.3	32.8	32.6 (0.35)
T8	134	137	135.3 (2.12)	45	46	45.5 (0.70)	44.74	44.20	44.47 (0.38)	2.69	2.79	2.74 (0.07)	24.4	24.8	24.6 (0.28)
T9	143	142	142.3 (0.71)	52	50	51.0 (1.41)	41.38	39.38	40.38 (1.41)	3.07	2.80	2.93 (0.19)	29.7	27.6	28.7 (1.48)
T10	122	124	122.8 (1.41)	36	39	37.5 (2.12)	37.00	36.00	36.50 (0.71)	1.62	1.74	1.68 (0.08)	44.1	46.4	45.2 (1.63)

239 **Table 4:**Effect of treatments on the uptake of nutrients (kg ha⁻¹) by seed.The data from 2010-11 and 2011-12 are shown along with the
240 average over two years. S.D. stands for standard deviation.


Treatments	Nitrogen			Phosphorus			Potassium			Boron			Zinc		
	2010-11	2011-12	Mean (S.D.)	2010-11	2011-12	Mean (S.D.)	2010-11	2011-12	Mean (S.D.)	2010-11	2011-12	Mean (S.D.)	2010-11	2011-12	Mean (S.D.)
T1	70.47	59.66	65.06 (7.64)	1.40	1.74	1.57 (0.24)	23.14	24.90	24.02 (1.24)	0.280	0.255	0.267 (0.02)	0.297	0.228	0.262 (0.05)
T2	58.17	58.89	58.53 (0.51)	1.58	1.50	1.54 (0.06)	19.14	22.43	20.78 (2.33)	0.249	0.253	0.251 (0.01)	0.314	0.168	0.241 (0.10)
T3	64.54	59.85	62.19 (3.32)	1.40	1.24	1.32 (0.11)	19.21	21.66	20.44 (1.73)	0.292	0.271	0.282 (0.02)	0.406	0.361	0.384 (0.03)
T4	69.54	66.06	67.80 (2.46)	2.29	1.90	2.09 (0.28)	25.93	27.43	26.68 (1.06)	0.211	0.239	0.225 (0.02)	0.300	0.645	0.472 (0.24)
T5	83.94	72.69	78.32 (7.95)	2.25	1.88	2.07 (0.26)	18.74	21.09	19.92 (1.66)	0.237	0.289	0.263 (0.04)	0.554	0.292	0.423 (0.18)
T6	121.25	116.31	118.78 (3.49)	2.61	2.46	2.54 (0.08)	38.59	44.67	41.63 (4.30)	0.231	0.385	0.308 (0.11)	0.482	0.402	0.442 (0.06)
T7	104.39	101.86	103.12 (1.79)	2.53	2.41	2.47(0.11)	28.83	36.38	32.60 (5.34)	0.215	0.246	0.230 (0.02)	0.413	0.333	0.373 (0.06)
T8	63.97	70.97	67.47 (4.50)	1.20	1.11	1.16 (0.06)	18.81	30.64	24.73 (8/36)	0.190	0.232	0.211 (0.03)	0.255	0.195	0.225 (0.04)
T9	77.27	72.02	74.65 (3.71)	1.77	1.47	1.62 (0.21)	18.40	26.56	22.48 (5.77)	0.171	0.197	0.184 (0.02)	0.284	0.175	0.229 (0.07)
T10	13.14	9.75	11.45 (2.40)	0.76	0.79	0.77 (0.02)	7.28	9.58	8.43 (1.62)	0.061	0.060	0.060 (0.00)	0.020	0.065	0.043 (0.03)

241

242 A significant difference was observed in the uptake of different nutrients by seed (Table 4).
 243 The highest uptake of almost all nutrients (N, P, K, and B) was recorded in treatment T6 (B₁Zn₂)
 244 except for Zn, the highest uptake of which was recorded in treatment T₄ (B₁Zn₀). The highest
 245 production as well as the interaction between the micro-nutrients (B and Zn) in treatment T₆
 246 facilitated higher amount of nutrients uptake in seed[42]. The lowest uptake of all nutrients was
 247 recorded in treatment T₁₀ (control). A similar trend was observed for the uptake of nutrients by
 248 straw (Table 5).

249 **Table 5:**Effect of treatments on the uptake of nutrients (kg ha⁻¹) by straw along with the
 250 average grain yield and straw yield over two years. S.D. stands for standard deviation and
 251 presented in bracket.
 252

Treatments	Grain yield (S.D.)	Straw yield (S.D.)	Nitrogen (S.D.)	Phosphorus (S.D.)	Potassium (S.D.)	Boron (S.D.)	Zinc (S.D.)
T1	2829.83 (89.10)	7410.68 (220.6)	60.26 (7.66)	0.34(0.01)	109.61 (2.83)	0.118 (0.02)	1.130 (0.01)
T2	2768.95 (49.49)	6558.93 (45.6)	19.26 (6.36)	0.38(0.00)	107.67 (2.34)	0.252 (0.03)	2.134 (1.60)
T3	2921.77 (46.67)	6702.67 (13.2)	51.61 (1.43)	0.21(0.03)	111.44 (12.46)	0.172 (0.02)	0.821 (0.21)
T4	2736.29 (9.89)	8177.83 (107.7)	27.29 (28.78)	0.36(0.05)	131.95 (14.75)	0.296 (0.26)	1.723 (0.71)
T5	3325.93 (114.55)	4268.08 (68.7)	25.67 (2.12)	0.21(0.00)	73.80 (1.44)	0.230 (0.01)	0.581 (0.06)
T6	4377.06 (126.57)	8171.53 (84.7)	84.69 (7.35)	0.23(0.01)	141.31 (3.87)	0.477 (0.03)	1.756 (2.11)
T7	3942.73 (140.71)	8160.33 (146.6)	20.36 (22.25)	0.40(0.07)	149.46 (13.66)	0.413 (0.00)	1.947 (2.34)
T8	2736.62 (68.59)	8376.87 (96.9)	34.00 (1.26)	0.33(0.00)	155.96 (8.56)	0.227 (0.03)	5.001 (6.15)
T9	2931.12 (190.92)	7288.23 (44.6)	25.52 (4.48)	0.17(0.00)	120.57 (5.46)	0.567 (0.01)	3.448 (4.61)
T10	1679.67 (86.97)	2033.42 (28.2)	2.28 (0.03)	0.06(0.04)	30.75 (0.89)	0.049 (0.00)	0.246 (0.31)



253
 254 **Fig. 2:** Effect of treatments on the uptake of nutrients (N, P, K, B, and Zn) at different growth
 255 stages (CRI, Tillering, Booting and Maturity) of wheat. The standard deviation of measurement is
 256 shown as error bars. The CRI stage indicates crown root initiation.




Fig. 3: Effect of treatments on the residual nutrient (N, P, K, B, and Zn) status at different growth stages (CRI, Tillering, Booting and Maturity) of wheat. The standard deviation of measurement is shown as error bars. The CRI stage indicates crown root initiation.

The B and Zn concentration in seeds (Table 4) and straw (Table 5) were calculated after dividing the total uptake of nutrients by the total grain and straw production. It clearly showed that with the increasing production, the concentration of nutrients, both B and Zn in seed and straw decreased.

265 There was significant difference in the residual N status of soil at different treatment plots
266 and at different growth stages. Initial application of N resulted a high amount of residual N at the
267 CRI stage and gradually decreased towards maturity, which had the least amount of residual N
268 (Fig. 3). Minimum demand of the applied N at the beginning of the growth stages resulted in a
269 high amount of residual N at the CRI stage, while the high demand towards maturity left the least
270 amount of residual N. High demand during the peak growth stages such as tillering and booting
271 resulted in a very similar amount of residual N, which was lower than that at CRI stage. The
272 highest amount of residual N was recorded in treatment T_3 (B_0Zn_2) and the lowest amount was
273 recorded in treatment T_{10} (control). The residual K status in soil at different growth stages of
274 wheat showed a very similar trend as that of N. The CRI stage was recorded with the highest
275 amount of residual K, which in general decreased towards maturity. There was a significant
276 difference between the treatments at different growth stages. Treatment T_7 (B_2Zn_0) was recorded
277 with the highest of amount of residual K at the CRI stage, while Treatment T_1 (B_0Zn_0) was
278 recorded with the highest amount of K at other growth stages (Fig. 3). The absence of
279 micronutrients in treatment T_1 might have inhibited the uptake resulting in a high amount of
280 residual K.

281 The residual P in soil showed a little different trend than N and K (Fig. 3). There was no
282 specific trend of residual P at different growth stages. In general, a higher amount of residual P
283 was recorded at the CRI stage compared to tillering and booting stage. This might be due to the
284 presence of unavailable form of P at the beginning of the growth stage. While the difference
285 between the growth stages of wheat was not significant, the difference between the treatments
286 was significant. The highest amount of P was recorded in treatment T_2 (B_0Zn_1) for the CRI stage
287 while treatment T_4 (B_1Zn_0) at the maturity. The lowest amount of residual P was recorded in
288 treatment T_{10} (control).

289 There was a significant difference in the residual B status in soil at different nutrient
290 treatment combinations. However, the difference was not significant at different growth stages.
291 There was no specific trend on the residual amount of B among the growth stages (Fig. 3). For
292 example, while the treatment T_8 (B_2Zn_1) was recorded with the highest amount of residual B at
293 the CRI stage, treatment T_1 (B_0Zn_0) was recorded with the highest amount of B at the tillering
294 stage (Fig. 3). A similar trend in the residual Zn content was observed at different growth stages
295 and at different treatments. For example, the highest amount of residual Zn was observed in

296 treatment T₄ (B₁Zn₀) at the CRI stage, while the highest amount of residual Zn was observed in
 297 treatment T₈ (B₂Zn₁) at the booting stage. A growth stage dependent Zn demand and the residual
 298 Zn were also reported by Ozturk et al. [43]. The variation in the residual Zn might also be due to
 299 the combined effect of pH, EC, organic carbon and P, which ultimately controls the Zn
 300 availability [44]. The lowest amount of residual B and Zn was observed in treatment T₁₀
 301 (control). In general a lower amount of Zn was recorded at maturity, which indicates a demand
 302 of Zn in the production of crop.

303 **Table 6:** Correlation (*r*) between soil available nutrient status at harvest and the nutrient content
 304 in straw averaged over two years

	Plant N	Plant P	Plant K	Plant B	Plant Zn
Soil N	0.41**	0.59**	0.66**	0.39**	0.33*
Soil P	0.36**	0.63**	0.67**	0.60**	0.11
Soil K	0.28*	0.39**	0.38**	0.31*	0.64**
Soil B	0.68**	0.57**	0.63**	0.26*	0.25*
Soil Zn	0.10	0.50**	0.48**	0.35	0.64**

305 * p < 0.05, ** p < 0.01

306
 307 **Table 7:** Correlation (*r*) between soil available nutrient status at harvest and the nutrient content
 308 in seed averaged over two years

	Soil N	Soil P	Soil K	Soil B	Soil Zn
Seed N	0.54**	0.76**	0.23*	0.56**	0.37**
Seed P	0.40**	0.83**	-0.03	0.47**	0.07
Seed K	0.48**	0.71**	0.19	0.63**	0.25*
Seed B	0.80**	0.74**	0.32*	0.69**	0.41**
Seed Zn	0.70**	0.80**	-0.06	0.64**	0.15

309 * p < 0.05, ** p < 0.01

310 A positive correlation was observed between the uptake of different nutrients and the grain
 311 and straw yield irrespective of different treatments (Fig. 4). The correlation coefficient (*r*) was as
 312 high as 0.97 between K uptake (kg ha⁻¹) and the straw yield (tha⁻¹). There was a very weak
 313 correlation between the Zn uptake and the grain yield (Fig. 4). The uptake of nutrients was
 314 governed by the soil, environmental and management practices. For example, the availability of
 315 B was determined by the availability of Zn in soil [28]. Santra et al. [45] also reported an
 316 increased amount of DTPA extractable Zn with the application of B. The relationship between B
 317 and Zn was found to be synergistic making high amount of Zn available in soil. A high
 318 correlation was also observed between the residual nutrient status in soil and the nutrient status is
 319 straw (Table 6) or between residual status in soil and the nutrient status is seed (Table 7).




Fig. 4: Correlation coefficient (r) between the nutrient (N, P, K, B, and Zn) uptake (kg ha⁻¹) by plants and the grain and straw yield (t ha⁻¹) of wheat.

Conclusions

This study examined the effect of boron and zinc on the yield and uptake of different nutrients by wheat in the acid soil region of West Bengal, India. The yield components and grain yield of wheat showed a significant difference among the treatment combinations. The maximum average grain yield (4.4tha^{-1}) over two years was observed in the treatment T_6 with higher amount of Zn application along with recommended dose of Boron application. The minimum grain yield was observed in treatment T_{10} (1.7tha^{-1}). A relatively higher yield was obtained from the treatments with any nutrient combination over that of the control (T_{10}). Along with the difference in grain yield, a significant difference in straw yield was also observed among the treatments. The application of boron and zinc might show some synergistic effects leading to high grain and straw yield in the acid soil region. High response from a combined application of B and Zn clearly demonstrated the necessity of micronutrients for improving production in the studied regions with acid soils. The presence of micro-nutrients and their combination also affected the uptake of nutrients in different growth stages of wheat. The interaction effect was also visible in the uptake nutrients by seeds. A positive correlation was observed between the uptake of nutrients and the yield of grain and straw in this study region with acid soils. The residual nutrient status showed a build-up of nutrients in soils. Therefore, an application of a mixture of micronutrients is recommended over a single micronutrient for the acid soil regions of West Bengal in order to get a better response from the applied nutrient sources and thus the production. This result may also be applied for the other grain crops in this region. However, the response of multiple nutrient combinations on the crop growth and production are required to study in future for better understanding the nutrient dynamics in the acid soil regions of West Bengal.

References

1. FAO (Food and Agriculture Organization of the United Nations) (2012) Food and Agriculture Organization of the United Nations Crop Production and Trade Statistics. (available from: <http://faostat.fao.org/site/339/default.aspx>)
2. Index Mundi (2011) Afghanistan Wheat Area Harvested by Year, last viewed 10 Aug. 2011. <http://www.indexmundi.com/agriculture/?country=af&commodity=wheat&graph=area-harvested>.
3. Rehm, G., and Sims, A. (2006) Micronutrients and production of hard red spring wheat. Minnesota Crop News, University of Minnesota.

- 356 4. Taiwo, L.B., Adediran, J.A., Akande, M.O., Banjoko, V.A., and Oluwatosin, G.A. (2001)
 357 Influence of legume fallow on soil properties and yield of maize in south-western Nigeria.
 358 *Journal of Agriculture in Tropics and Subtropics* **102(2)**: 109-117.
- 359 5. Adediran, J.A., Taiwo, L.B., Akande, M.O., Idowu, O.J., and Sobulo, R.A. (2004)
 360 Application of organic and inorganic fertilizer for sustainable yield of maize and cowpea in
 361 Nigeria. *Journal of Plant Nutrition* **27(7)**: 1163-1181.
- 362 6. Goldbach, H.E., Blaser-Grill, J., Lindemann, N., Porzelt, M., Horrmann, C., Lupp, B., and
 363 Gessner, B. (1991) Influence of boron on net proton release and its relation to other
 364 metabolic processes. *Current Topics in Plant Biochemistry and Physiology* **42**: 925-929.
- 365 7. Reddy, S.R. (2004) Principles of Crop Production – Growth Regulators and Growth
 366 Analysis, 2nd Ed. Kalyani Publishers, Ludhiana, India.
- 367 8. Ali, S., Shah, A., Arif, M., Miraj, G., Ali, I., Sajjad, M., Farhatullah, M., Khan, Y., and
 368 Khan, N.M. (2009) Enhancement of wheat grain yield and yield components through foliar
 369 application of zinc and boron. *Sarhad Journal of Agriculture* **25(1)**: 15-19.
- 370 9. Ahmadikhah, A., Narimani, H., Rahimi, M.M., and Vaezi, B. (2010) Study on the effects of
 371 foliar spray of micronutrient on yield and yield components of durum wheat. *Archives of*
 372 *Applied Science Research* **2(6)**: 168-176.
- 373 10. Nadim, M.A., Awan, I.U., Baloch, M.S., Khan, E.A., Naveed, K., and Khan, M.A. (2012)
 374 Response of Wheat (*Triticumaestivum*L.) to different Micronutrients and their Application
 375 Methods. *The Journal of Animal & Plant Sciences* **22(1)**: 113-119.
- 376 11. Kanwar, J.S., and Randhawa, N.S. (1974) Micronutrient research in soils and plants in India.
 377 Indian Council of Agricultural Research, New Delhi. p. 18-32.
- 378 12. Takkar, P.N., Mann, M.S., and Randhawa, N.S. (1971) How zinc deficiency affects wheat
 379 yields. *Indian Farming* **21(9)**: 31-32.
- 380 13. Asad, A., and Rafique, R. (2002) Identification of micronutrient deficiency of wheat in the
 381 Peshawar valley. *Communication in Soil Science and Plant Analysis* **33**: 349-364.
- 382 14. Kaya, Y., Arisoy, R.Z., and Gocmen, A. (2002) Variation in grain yield and quality traits of
 383 bread wheat genotypes by zinc fertilization. *Pakistan Journal of Agronomy* **1**: 142-144.
- 384 15. Singh, Y.P. (2004) Effect of nitrogen and zinc on wheat irrigated with alkali water. *Annals*
 385 *of Agricultural Research* **25**: 233-236.
- 386 16. Bagci, S.A., Ekiz, H., Yilmaz, A., and Cakmak, I. (2007) Effect of zinc deficiency and
 387 drought on grain yield of field-grown wheat cultivars in Central Anatolia. *Journal of*
 388 *Agronomy and Crop Science* **193**: 198-206.
- 389 17. Graham, W.A., and McDonald, G.K. (2001) Effects of zinc on photosynthesis and yield of
 390 wheat under heat stress. In: *Proceedings of the 10th Australian Agronomy Conference*,
 391 Hobart.
- 392 18. Genc, Y., McDonald, G.K., and Graham, R.D. (2006) Contribution of different mechanisms
 393 to zinc efficiency in bread wheat during early vegetative stage. *Plant and Soil* **281**: 353-
 394 367.

- 395 19. NFDC(1998) Micronutriments in Agriculture: Pakistan Perspective. NFDC Publication No.
396 4/98, Islamabad.
- 397 20. Rafique, E., Rashid, A., Ryan, J., and Bhatti, A.U. (2006): Zinc deficiency in rain-fed wheat
398 in Pakistan: Magnitude, spatial variability, management, and plant analysis diagnostic
399 norms. *Communications in Soil Science and Plant Analysis* **37**: 181–197.
- 400 21. Erenoglu, B., Cakmak, I., Romheld, V., Derici, R., and Rengel, Z. (1999) Uptake of zinc by
401 rye, bread wheat and durumwheat cultivars differing in zinc efficiency. *Plant and Soil* **209**:
402 245–252.
- 403 22. Gupta, V.K., and Mehla, D.S. (1993) Depletion of micronutrients from soil and their uptake
404 in rice-wheat rotation. *Journal of the Indian Society of Soil Science* **4**: 704–706.
- 405 23. Rerkasem, B., and Jamjod, S. (2004) Boron deficiency in wheat: a review. *Field Crops*
406 *Research* **89**: 173–186.
- 407 24. Snowball, K., and Robson, A.D. (1983) Symptoms of nutrient deficiencies:
408 Subterranean clover and wheat. Institute of Agriculture, University of Western Australia:
409 Nedlands, Western Australia.
- 410 25. Dell, B., and Huang, L.B. (1997) Physiological response of plants to low boron. *Plant and*
411 *Soil* **193**: 103–120.
- 412 26. Nayyar, V.K., Arora, C.L., and Kataki, P.K. (2001) Management of Soil Micronutrient
413 Deficiencies in the Rice-Wheat Cropping System. p. 87–131. In: Kataki, P.K. (Eds.). The
414 Rice-Wheat Cropping Systems of South Asia: Efficient Production Management, Food
415 Products Press, New York, USA.
- 416 27. Takkar, P.N. (1982) Micronutrients: Forms, Contents Distribution in Profile, Indices of
417 Availability and Soil Test Methods. p. 361–391. In: Reviews of Soil Research in India.
418 Part I. 12th International Congress of Soil Science, New Delhi, India.
- 419 28. Singh, M.V. (1999) Micronutrient deficiency delineation and soil fertility mapping.
420 In: National symposium on zinc fertilizer industry - Whither To (Ramendra Singh
421 and Abhay Kumar, eds). Session II.
- 422 29. Jackson, M.L. (1967) *Soil Chemical Analysis*. Prentice-Hall of India, New Delhi.
- 423 30. Nelson, D.W., and Sommers, L.E. (1982) Total carbon, organic carbon and organic matter.
424 In: *Methods of Soil Analysis* (Page et al. Eds.), part 2, second edition, Agronomy Monograph
425 9, ASA and SSSA, Madison, Wisconsin. p. 539–594.
- 426 31. Dewis, J., and Freitas, F. (1984) *Physical and Chemical Methods of Soil and*
427 *Water Analysis*. Oxford and IBH Pub. Co., New Delhi. p. 51–106.
- 428 32. Subbiah, B.V., and Asija, G.L. (1956) A rapid procedure for the determination of available
429 nitrogen in soils. *Current Science* **25**: 259–260.
- 430 33. Bray, R.H., and Kurtz, L.P. (1945): Determination of total, organic and available forms of
431 phosphorus in soils. *Soil Science* **59**: 39–45.
- 432 34. Black, C.A. (1965) *Methods of soil analysis*. Part II, Agronomy Ser. No. 9. American
433 Society of Agronomy Inc. Madison, Wisconsin, U.S.A.

- 434 35. Lindsay, W.L., and Norvell, W.A. (1978) Development of DTPA soil test for zinc, iron,
435 manganese and copper. *Soil Science Society of America Journal* **42**: 421-428.
- 436 36. Saha, A.K. (2008) Methods of Physical and Chemical Analysis of Soil. Kalyani Publishers.
437 P. 1-469.
- 438 37. Chaudry, E.H., Timmer, V., Javed, A.S., and Siddique, M.T. (2007) Wheat response to
439 micronutrients in rain-fed areas of Punjab. *Soil and Environment* **26**: 97-101.
- 440 38. Topal, A., Gezgin, S., Akgun, N., Dursun, N., and Babaoglu, M. (2002) Yield and yield
441 attributes of durum wheat (*Triticum durum* Desf.) as affected by boron application. In:
442 Goldbach, H.E., Rerkasem, B., Wimmer, M.A., Brown, P.H., Thellier, M., Bell, R.W.
443 (Eds.), All Aspects of Plant and Animal Boron Nutrition.
- 444 39. Marten, J.M., and Westermann, D.T. (1991) Fertilizer Applications for Correcting
445 Micronutrient Deficiencies. In Micronutrients in Agriculture, Mordtvedt; Mordtvedt, J.J.,
446 Giordano, P.M., Lindsay, W.L. (Eds.) Soil Science Society of America Incorporation:
447 Madison, WI. 549-592.
- 448 40. Kutman, U. B., Yildiz, B., Ozturk, L., Cakmak, I. (2010) Bio-fortification of durum wheat
449 with zinc through soil and foliar applications of nitrogen. *Cereal Chemistry* **87**: 1-9.
- 450 41. Mandal, A.B. (1989) *Bangladesh Journal of Botany* **18**: 247 - 252.
- 451 42. Hosseini, S.M., Maftoun, M., Karimian, N., Rounaghi, A., Emam, Y. (2007) Effect of zinc
452 \times boron interaction on plant growth and tissue nutrient concentration of corn. *Journal of*
453 *Plant Nutrition* **30**: 773-781.
- 454 43. Ozturk, L., Yazici, M.A., Yucel, C., Torun, A., Cekic, C., Bagci, A., Ozkan, H., Braun, H.J.,
455 Sayers, Z., and Cakmak, I. (2006) Concentration and localization of zinc during seed
456 development and germination in wheat. *Physiology of Plant* **128**: 144-152.
- 457 44. Sakal, R., Singh, A.P., and Singh, S.P. (1988) Distribution of available Zinc, Copper, Iron
458 and Manganese in Old Alluvial soils as related to certain soil characteristics. *Journal of the*
459 *Indian Society of Soil Science* **36**: 59-63.
- 460 45. Santra, G.H., Das, D.K. and Mandal, B.K. (1989) Relationship of boron with iron,
461 manganese, copper and zinc with respect to their availability in rice soil. *Environmental*
462 *Ecology* **7**: 874 - 877.