

ABSTRACT

A field experiment was carried out during the period from November 2012 to May 2013 in Agroecological Zone 20 (Eastern Surma-Kushiyara Floodplain) to observe the varietal performances of high yielding and local varieties of Boro rice. Four varieties viz. BRRI dhan28, BRRI dhan29, Khoiaboro and Begunbichi, and transplanting of three seedling ages viz. 15, 20 and 25 days old were included as treatments in the experiment. The experiment was laid out in a factorial RCBD (Randomized Complete Block Design) with three replications. The results revealed that BRRI dhan29 produced significantly highest grain yield (6.25 t ha^{-1}) attributed by the higher number of effective tillers hill⁻¹, grains panicle⁻¹ and 1000-grain weight. BRRI dhan28 produced the second highest grain yield (5.37 t ha^{-1}) while the local variety Begunbichi produced the lowest grain yield (2.26 t ha^{-1}) in spite of its highest number of grains panicle⁻¹, because of its small sized grain. Total number of spikelets panicle⁻¹, number of grains panicle⁻¹, unfilled spikelets panicle⁻¹, grain and straw yield varied significantly but other characters did not among different ages of seedlings. The highest grain yield of 4.49 t ha^{-1} was obtained from planting 25 days old seedlings ascribed to higher number grains panicle⁻¹ mainly. Grain yield of 4.23 t ha^{-1} was obtained from planting 20 days old seedlings which was statistically similar to that of planting 25 and 15 days old seedlings. Interaction of variety and seedling age produced significant effect on most of the characters except plant height, number of non-effective tillers hill⁻¹, length of panicle and grain yield. This indicated that all varieties require planting 25 days old seedlings for obtain higher grain yield. Cost and return analysis showed that BRRI dhan29 gave maximum gross return, net return, and Benefit Cost Ratio of US\$ 1665.00 ha⁻¹, US\$ 699.28 and 1.72, respectively with planting 25 days old seedlings. Local variety Begunbichi showed more profitability than BRRI dhan28 and local Khoiaboro varieties.

Key words: Boro rice, Variety, Seedling age, Yield

1. INTRODUCTION

Rice is the staple food of about 149.69 million people of Bangladesh and it is being grown in about 75% of the total cropped area and more than 80% of the total irrigated area [16]. Almost all the farm families of 13 million grow rice in the country. It provides nearly 40% of national employment (48% of rural

employment), about 70-76% of total calorie supply and 66% of protein intakes of an average person in the country [16, 10]. Rice sector contributes one-half of the agricultural GDP and one-sixth of the national income in Bangladesh [17]. Thus, rice plays a vital role in the livelihood of the Bangladeshi people. Rice production needs to be increased more as the population of Bangladesh is still growing by two millions in every year and may increase by another 30 millions over the next 20 years. There are less possibilities of bringing more lands under cultivation of rice; much of the additional rice requirement will have to be met by increasing the average yield from the existing land. Although rice is grown on large area in Bangladesh, its average yield is still far below the levels attained in other rice producing countries in spite of having many high yielding varieties. The average yield in Bangladesh is about 2.74 t ha⁻¹ as compared to Japan (5.93 t ha⁻¹) and Korea (6.12 t ha⁻¹) [19]. To combat the situation it requires adoption of modern technologies such as better management package, high yielding cultivars of both inbred and hybrid, and higher input use [42]. There are several reasons behind this but one of the most important reasons is that the seedling age is not managed properly to get vigorous seedlings for uniform stand and better bush establishment. Among the various factors that influence rice productivity, seedling age has tremendous effect on plant height, tiller production, panicle length, grain formation and other yield attributing characters [1]. Younger seedlings may not be able to withstand transplanting shock whereas too old seedlings may not be able to produce its yield potential to the peak. Transplanting seedlings in proper age can provide appropriate ground for achieving potential production by reducing the death of tillers. Chopra *et al.* (2002) evaluated the yield and quality of seeds of the rice cv. Pusa 44 by transplanting seedlings at 25, 35, 45, 55, and 65 days [7]. They found that transplanting seedlings at 35 days resulted greater number of panicles hill⁻¹, panicle length, 1000 seed weight and seed yield than 55 to 65-day old seedlings. Farmers transplant seedlings at different ages but more often with those of at 25 to 50 days older in lowland rice [9, 41, 35]. Many researchers reported that grain yield increased by transplanting younger seedlings of 25 days [36, 2, 26, 39]. On the other hand some studies exposed that grain yield was not affected by transplanting even 30-60 days old seedlings [6]. Recent studies on the System of Rice Intensification (SRI) also showed that yield and yield components of rice might be increased by transplanting seedlings as younger as 14 days as compared to older seedlings of 21-23 days [22]. McHugh (2002) also observed in Madagascar that 8 to 15 days old seedlings transplanted at 25 hills m² produced the highest yields [24]. Bangladesh Rice Research Institute (BRRI) has recommended to decide seedling age of rice for transplanting according to growing season. BRRI (1991; 1992)

recommended for transplanting 20-30 days old seedlings in Aus season, 20-35 days old seedlings in T. Aman season and 40-45 days old seedlings in Boro season [4, 5]. It is generally seen that Researchers' recommendations are not following by farmers [17] and it has been reported that farmers even use 80 days old seedlings of Boro rice for transplanting [18]. In Bangladesh, younger seedlings transplantation in Boro season is very difficult and it is labour intensive because of stunted growth of seedlings due to cold weather. To avoid the situation older seedlings with optimum growth need to be transplanted. It was reported that paddy yield was decreased significantly after transplanting of younger seedlings due to its higher mortality rate in the field while transplanting of older seedlings resulted in better performance [21]. In most of the above citations transplanting rice at different ages of HYVs/modern varieties have shown variation in their performances in respect of yield but local varieties of Boro season have not been tested. The major objectives of the study were to know the effect of seedling age at transplanting on the growth and yield performances of high yielding and local varieties of Boro season in Sylhet region, Bangladesh.

2. MATERIALS AND METHODS

The experiment was conducted during the period from November 2012 to May 2013 at Patnipara, Chicknagul union under Jointapur upazila of Sylhet district, 18 km far North-East from Sylhet Agricultural University, Sylhet. Geographically the location is situated at 23° to 25°1' North and 90°57' to 92°28' East longitude and latitude, respectively with an elevation of 34 m above the mean sea level. The experimental field had fairly leveled topography: medium low land and good drainage system. The experimental plot was under the Agro-ecological zone 20 and the soil type was silty clay loam in texture and pH of the soil was about 5.5-6.5. Organic matter content of the soil was moderate. Levels of cation exchange capacity (CEC) and Zn was medium while the status of P, K and B was low. Sylhet has a tropical climate and as the monsoon clouds blow in the area throughout the year, there is considerable rainfall in most of the months of the year while June and July receive the highest amount. This area is much cooler and hotter than the other parts of Bangladesh. Monthly maximum and minimum temperature, rainfall and relative humidity during the crop growing period have been presented in Table 1.

Table 1. Monthly average rainfall, minimum and maximum temperatures and relative humidity during the study period from November 2012-June 2013

Month	Year	Rainfall (mm)	Air Temperature (°C)			Relative Humidity (%)
			Maximum	Minimum	Average	
November	2012	11.9	29.2	18.9	24.1	68
December	2012	Nil	25.0	14.5	19.8	75
January	2013	Nil	25.6	11.8	18.7	63
February	2013	2.3	31.2	15.9	23.6	49
March	2013	1.9	36.2	19.7	27.8	47
April	2013	13.9	33.0	21.9	27.5	59
May	2013	34.2	30.4	22.7	26.6	78
June	2013	26.9	33.9	25.8	29.9	75

Source: Department of Meteorology, Sylhet

The treatments included in the experiment were as follows.

Factor A. Variety: 4

i. BRRI dhan28 (V_1)

ii. BRRI dhan29 (V_2)

iii. Khoiaboro (V_3)

iv. Begunbichi (V_4)

Factor B. Seedling age at transplanting: 3

i. 15 days old seedlings (15DOS)

ii. 20 days old seedlings (20DOS)

iii. 25 days old seedlings (25DOS)

Among the varieties BRRI dahn28 and BRRI dahn29 were the High Yielding Varieties (HYV) and Khoiaboro and Begunbichi (aromatic) were the local or indigenous varieties of rice. The experiment was laid out according to Randomized Complete Block Design (Factorial). The unit plot size was 3 m x 2 m. Seed was used at the rate of 10 kg ha⁻¹ having germination percentage of 93%, 95%, 92% and 95% for BRRI dhan28, BRRI dhan29, Khoiaboro and Begunbichi, respectively. Pregerminated seeds of all varieties were sown in nursery beds on 23 November 2012 (for 25 days old seedlings), 28 November 2012 (for 20 days old seedling) and 03 December 2012 (for 15 days old seedlings). Frequent irrigation was done to maintain enough moisture content in the seed bed. Field was prepared by power tiller on 1 December 2012 with a power tiller i.e. 15 days before transplanting. Organic manures and inorganic

fertilizers were applied in the field. The source of organic manures was cowdung and applied at the rate of 10 t ha⁻¹ as basal application 10 days before final land preparation. Fertilizers were applied as suggested by BRRI (2011) [3] at the rate of 138-20-60-20-4 kg ha⁻¹ NPKS & Zn for the variety of BRRI dhan29, 121-20-60-20-4 kg ha⁻¹ NPKS & Zn for BRRI dhan28, 52-15-15 kg ha⁻¹ of NPK, respectively and 5 t ha⁻¹ of Cowdung as basal for both the local varieties of Khoiaboro and Begunbichi. N, P, K, S and Zn were applied through Urea, TSP, MoP, Gypsum and ZnSO₄, respectively. Urea was applied into three installments- 1/3rd at final land preparation, 1/3rd at 21 days after transplanting (DAT) (tillering stage) and 1/3rd at 36 DAT (active tillering stage) in each plot. The nursery beds were made wet by application of water both in morning and evening on the previous day of uprooting the seedlings. Seedlings were uprooted carefully so that minimum damage was done to the root system and uprooted seedlings kept in shade before transplanting. The methods were followed for each case of uprooting and transplanting for different ages of seedling. The seedlings uprooted from the nursery bed were transplanted on the same day. Single seedling of each 15, 20 and 25 days old was transplanted in a square pattern maintaining 25 cm × 25 cm spacing on the well puddled plots on 16 December 2012. During transplanting of seedling the plot was saturated with sufficient 2-3 cm depth of water. To maintain the desired plant population in each plot, gap filling was done within 10 days of transplanting as some hills died off using seedling of the same source of the respective age. The first manual weeding was done at 21 DAT after which first top dress of urea was done. Second top dress of urea was done after second weeding at 36 DAT for each crop. At both weeding and top dressing sufficient moisture was ensured in each plot. Water management was done properly following flood irrigation method with the help of shallow tubewell from surface water. After transplanting 3-4 cm water depth was maintained throughout the life cycle of the crop but removed 10 days before maturity. For controlling insect-pest granular insecticide Carbofuran 5G (Furadan) was applied at the rate of 16 kg ha⁻¹ by maintaining 4-5 cm water depth in the crop field at maximum tillering stage. Crop maturity was determined when 80% grain of all panicles in a plot turned into golden yellow in colour except Khoiaboro. Grain colour of Khoiaboro rice was blackish yellow. At this stage culm and leaves were also turn into yellow colour. At maturity ten random hills were sampled for collection of data on yield and yield attributes. BRRI dhan28 was harvested on 12, 19, 25 April 2013 respectively for the seedling age of 25 days old, 20 days old and 15 days old, respectively. BRRI dhan29 was harvested on 2, 9 and 15 May 2013 April, respectively of the seedling age of 25 days old, 20 days old and 15 days old. Khoiaboro was harvested on 12, 16 and 18 April 2013 of the seedling age of 25 days old, 20 days old and 15 days old, respectively and Begunbichi

was harvested on 12, 20 and 24 April 2013 of the seedling age of 25 days old, 20 days old and 15 days old, respectively. First of all, border row from each side were harvested and these were excluded from final threshing. Remaining net plot area was harvested manually at ground level using sickle and kept separately for recording crop yield plot wise. Then grains were separated from each bundle by beating with bamboo sticks and grains were dried in the sun. Then moisture was recorded with moisture meter (GMK-303RS) and grain weight of individual plot was adjusted at 12% moisture content. After thorough sun drying straw weight was recorded separately. Finally, grain and straw weights in kg plot^{-1} of the individual plot were converted into t ha^{-1} . Data were collected on the growth, yield and yield attributes as follows.

- i. Number of tiller plant^{-1} at every 10-day intervals
- ii. Days to 50% flowering (when at least 50% tillers had panicle in each plot).
- iii. Days to maturity
- iv. Plant height at harvest
- v. Total number of tillers hill^{-1}
- vi. Number of effective tillers hill^{-1}
- vii. Number of non-effective tillers hill^{-1}
- viii. Length of panicle
- ix. Total number of spikelets panicle^{-1}
- x. Number of grains panicle^{-1}
- xi. Number of unfilled grains panicle^{-1}
- xii. 1000 grain weight
- xiii. Grain weight plot^{-1}
- xiv. Straw weight plot^{-1}
- xv. Biological yield
- xvi. Harvest index

Number of total tillers was counted from the selected five hills at every 10 day intervals. Tillers were counted by spreading the base of each standing hill so that small tiller may not be left out. Number of tillers counted in each date from five hills was averaged for individual plot and this data were statistically analyzed.

Harvest index (HI) was calculated on the basis of grain and straw yields using the following formula and expressed in percentage [13].

$$\text{Harvest index} = \frac{\text{Grain yield}}{\text{Biological yield}} \times 100$$

Where, Biological Yield= Grain yield + Straw yield


Statistical analysis

The collected data were tabulated and these were analyzed using computer software MSTATC. Mean separations were done at 5% level of significance by Least Significant Difference (LSD) Test wherever F values were significant at either 0.01% or 0.05% level of probability.

3. RESULTS AND DISCUSSION

Number of tillers hill⁻¹ at 10-day intervals

Individual effect of variety and seedling age has not discussed here as interaction effect of variety and seedling age was found significant for number of tillers hill⁻¹ at 10-day intervals at most of the cases. Interaction effect of variety and seedling age was found non-significant for number of tillers hill⁻¹ at 40 DAT but at 50 DAT (Figure 1). The highest number of tillers (14.00 hill⁻¹) was recorded from the combination of V₂×25DOS and the lowest (1.89 hill⁻¹) was obtained from the combination of V₃×15DOS at 50 DAT. The result indicated that at 60 DAT, the combination of V₂×25DOS produced maximum number of tillers (25.66 hill⁻¹) which was statistically similar to that of V₂×20DOS (23.67 hill⁻¹) and minimum (6.88 hill⁻¹) was found in the combination of V₃×15DOS. Number of tillers hill⁻¹ was significantly affected by the interaction of variety and seedling age at 70 DAT. The maximum number of tillers (33.67 hill⁻¹) was recorded from the combination of V₁×25DOS which was statistically similar to that of V₂×20DOS (29.78 hill⁻¹) and V₂×25DOS (32.22 hill⁻¹). The minimum number of tillers (14.56 hill⁻¹) was obtained from the combination of V₄×15DOS similar to that of V₂×15DOS (16.65 hill⁻¹) and V₃×15DOS (17.33 hill⁻¹).



The interaction effect of variety and seedling age was non-significant for number of tillers hill⁻¹ at 80 DAT while the same was significant at 90 DAT (Figure 1). The highest number of tillers (33.89 hill⁻¹) was recorded in the combination of V₂×25DOS which was statistically similar to that of V₂×15DOS (33.22 hill⁻¹), V₁×25DOS (31.33 hill⁻¹) and V₃×25DOS (31.44 hill⁻¹) combinations. On the other hand, the lowest number of tillers (22.11 hill⁻¹) was found in the combination of V₄×15DOS which was statistically at par with that of V₄×20DOS (22.44 hill⁻¹). The result revealed that older seedling produced more number of tillers hill⁻¹. Interaction of variety and seedling age produced significant effect on the number of tillers hill⁻¹ at 100 DAT and the results showed that the highest number of tillers (37.22 hill⁻¹) was produced by the combination of V₃×20DOS while the lowest (20.33 hill⁻¹) was produced by the combination of V₄×15DOS. The highest number of tiller (40.67 hill⁻¹) was obtained due to the treatment combination of V₃×20DOS which was significantly different from the others while the lowest (17.00 hill⁻¹) was obtained in the combination of V₄×15DOS at 110 DAT.

Phenology, yield components and yield

Varietal performances

Maximum days to 50% flowering (115.0) was recorded from the variety Begunbichi and minimum days to flowering (97.3) was recorded from the variety Khoiaboro (Table 2). Both the varieties BRRI dhan29 and Begunbichi took maximum days for their maturity (141.8) while Khoiaboro took the minimum days for maturity (122.0) (Table 2). Plant height was significantly varied among the varieties irrespective of seedling age. Result revealed that the local variety Begunbichi produced the tallest plant (154.49 cm) which was significantly different from the others. Moderate plant height was found in Khoiaboro whilst the shortest plant (96.82 cm) was found in BRRI dhan29 which was statistically similar to that of BRRI dhan28 (100.33 cm). Both the local varieties produced the taller plant and HYV's produced the shorter might be due to genetic variations of the varieties (Table 2). Total number of tillers hill⁻¹ included effective and non-effective tillers was significantly differed among the varieties. It is evident that maximum number of tillers (27.20 hill⁻¹) was obtained from the local variety Khoiaboro which was significantly different from the others. Minimum total number of tillers (20.70 hill⁻¹) was found in the local aromatic variety Begunbichi. BRRI dhan28 and BRRI dhan29 produced statistically similar number of total tillers (22.30 and 22.70 hill⁻¹, respectively) to the local variety Begunbichi (Table 2). The highest number of effective tillers (22.51 hill⁻¹) was found in the variety Khoiaboro which was significantly different from the others (Table 2). The Begunbichi produced the lowest number of effective tillers (16.41 hill⁻¹) while both the varieties BRRI dhan28 and BRRI dhan29 produced moderate number of effective tillers hill⁻¹. Venugopal and Singh (1985) obtained the highest number of effective tillers in short duration rice variety [40]. There was significant variation among the varieties in respect of number of non-effective tillers hill⁻¹. Both the varieties Khoiaboro and Begunbichi produced statistically similar number of non-effective tillers (4.7 and 4.3 hill⁻¹, respectively) having the highest in the variety Khoiaboro. The lowest number of non-effective tillers (3.0 hill⁻¹) was found in the variety BRRI dhan29 which was statistically similar to the variety BRRI dhan28. The variety BRRI dhan28 also produced similar number of non-effective tillers hill⁻¹ to the varieties Khoiaboro and Begunbichi. The varieties differed significantly in terms of length of panicle. BRRI dhan29 and Begunbichi had statistically similar panicle length having the highest value (25.0 cm) in the variety Begunbichi. The variety Khoiaboro had the lowest panicle length (20.0 cm) which was similar to that of BRRI dhan28 (22.2 cm) (Table 3). There was also significant variation in terms of total number of filled and unfilled spikelets panicle⁻¹. The variety Begunbichi produced maximum total number of spikelets (200.89

panicle⁻¹) while the variety Khoiaboro produced minimum (87.57 panicle⁻¹). BRRI dhan29 produced the second highest total number of **spikelets** (177.92 panicle⁻¹) which was significantly different from that of BRRI dhan28 (141.29) (Table 3). Variation was found significant among all varieties in respect of number of **grains** panicle⁻¹. Significantly highest number of **grains** (163.92 panicle⁻¹) was found in the variety Begunbichi followed by BRRI dhan29 while the lowest number (69.18 panicle⁻¹) was found in the variety Khoiaboro (Table 3).

Table 2. Phenology and yield components of rice varieties during Boro season 2012-2013

Varieties	Days to 50% flowering	Days to maturity	Plant height (cm) at harvest	Total number of tillers hill ⁻¹	Number of effective tillers hill ⁻¹	Number of non-effective tillers hill ⁻¹
V ₁	101.7 ^b	126.8 ^b	100.33 ^c	22.14 ^b	18.11 ^b	4.0 ^{ab}
V ₂	114.3 ^a	141.8 ^a	96.82 ^c	21.69 ^b	18.72 ^b	3.0 ^b
V ₃	97.3 ^c	122.0 ^c	146.02 ^b	27.20 ^a	22.51 ^a	4.7 ^a
V ₄	115.0 ^a	141.8 ^a	154.49 ^a	20.72 ^b	16.41 ^c	4.3 ^a
CV(%)	2.11	1.55	3.73	9.70	8.85	27.54
LSD _{0.05}	2.206	2.015	4.542	2.121	1.638	1.077

Note: V₁= BRRI dhan28, V₂=BRRI dhan29, V₃= Khoiaboro, V₄= Begunbichi; Figures within the same column having same or no letter(s) do not differ significantly at 5% level of probability.

Maximum number of unfilled spikelets (56.8 panicle⁻¹) was found in BRRI dhan29 and the local variety Khoiaboro produced the minimum (18.4 panicle⁻¹) (Table 3). The second highest number of unfilled **spikelets** of 37.3 panicle⁻¹ was found in the variety Begunbichi and it was 32.7 panicle⁻¹ in BRRI dhan28.

Table 3. Yield components of rice varieties during Boro season 2012-2013

Variety	Length of panicle (cm)	Total number of spikelets panicle ⁻¹	Number of grains panicle ⁻¹	Number of unfilled spikelets panicle ⁻¹	1000 grain weight (g)
V ₁	22.2 ^b	141.29 ^c	108.63 ^c	32.69 ^c	22.01 ^a
V ₂	24.7 ^a	177.92 ^b	121.09 ^b	56.82 ^a	22.38 ^a
V ₃	22.0 ^b	87.57 ^d	69.18 ^d	18.40 ^d	20.81 ^b
V ₄	25.0 ^a	200.89 ^a	163.92 ^a	37.30 ^b	12.10 ^c
CV(%)	5.21	3.88	6.53	9.26	4.05
LSD _{0.05}	1.197	5.769	7.382	3.287	0.765

Note: V₁= BRRI dhan28, V₂=BRRI dhan29, V₃= Khoiaboro, V₄= Begunbichi; Figures within the same column having same or no letter(s) do not differ significantly at 5% level of probability.

Statistically similar 1000 grain weight was found in both varieties BRRI dhan28 and BRRI dhan29 and having maximum (22.38 g) in BRRI dhan29 (Table 3). The variety Begunbichi gave the minimum 1000 grain weight (12.10 g) which was significantly different from others. Significant variation was also observed among varieties in terms of grain yield. The result showed that the highest grain yield (6.25 t ha⁻¹) was produced in BRRI dhan29 followed by BRRI dhan28 (5.37 t ha⁻¹) whilst the lowest (2.26 t ha⁻¹) was obtained in the local variety Begunbichi (Table 4). The highest grain yield in BRRI dhan29 was possibly attributed by the higher number of effective tillers hill⁻¹ and grains panicle⁻¹. In spite of lower number of effective tillers hill⁻¹ in BRRI dhan28 than Khoiaboro grain yield was compensated in BRRI dhan28 probably due to its higher number of grains as well as larger grain size. On the contrary, the local variety Begunbichi had the highest number of grains panicle⁻¹ but due to its lower number of tillers hill⁻¹ and smallest grain size the variety produced the lowest grain yield (Table 4). The variety Khoiaboro produced maximum straw yield (6.86 t ha⁻¹) among the variety irrespective of seedling age which was statistically similar to that of BRRI dhan29 (6.82 t ha⁻¹) (Table 4). Moderate straw yield (5.71 t ha⁻¹) was found in BRRI dhan28 and the minimum straw yield (4.88 t ha⁻¹) was found in the local variety Begunbichi. Local variety Khoiaboro produced maximum straw yield might be due to its taller plant stature but in spite of taller plant in Begunbichi lowest straw yield was produced might be due to its thin plant stature.

Table 4. Yield and harvest index of rice varieties during Boro season 2012-2013

Varieties	Grain yield (t ha ⁻¹)	Straw yield (t ha ⁻¹)	Biological yield (t ha ⁻¹)	Harvest index (%)
V ₁	5.37 ^b	5.71 ^b	11.08 ^b	48.60 ^a
V ₂	6.25 ^a	6.82 ^a	13.07 ^a	47.95 ^a
V ₃	2.90 ^c	6.86 ^a	9.77 ^c	30.09 ^b
V ₄	2.26 ^d	4.88 ^c	7.14 ^d	31.78 ^b
CV(%)	10.54	8.42	7.06	7.67
LSD _{0.05}	0.432	0.499	0.708	2.968

Note: V₁= BRRI dhan28, V₂=BRRI dhan29, V₃= Khoiaboro, V₄= Begunbichi; Figures within the same column having same or no letter(s) do not differ significantly at 5% level of probability.

Variation on biological yield was also found significant among the varieties. BRRI dhan29 gave the maximum biological yield (13.07 t ha⁻¹) while the minimum biological yield (7.14 t ha⁻¹) was found in the variety Begunbichi (Table 4). BRRI dhan28 had biological yield of 11.08 t ha⁻¹ followed by that of

Khoiaboro (9.77 t ha⁻¹) which was significantly different from each others. The result revealed that variety BRRI dhan28 gave the highest HI (48.67%) and it was identical to that of BRRI dhan29 (47.95%). There were statistically identical harvest indices of 30.09% and 31.78% of the local varieties Khoiaboro and Begunbichi (Table 4). This results indicated that assimilate partitioning is more in the grains of HYV's than the local which in turn resulted larger size of seed as well as higher grain yield in HYV's.

Effect of seedling age

Planting 20 days old seedlings took maximum duration for 50% flowering (110.9 days) and 15 days old seedlings took minimum (103.5 days) (Table 5). The results confirmed with the findings of Raju *et al.* (1989) who stated that days to flowering delayed in case of planting older seedlings [29]. But Padalia (1981) observed that days from sowing to flowering decreased with the increase of seedling age at planting [27]. Planting 25 days old seedlings took maximum duration for maturity (136.4 days) while planting 15 days old seedlings took minimum (130.3 days) (Table 5). Plant height did not vary significantly by the seedling age. However, plant height ranged from 123.06 cm in planting 15 days old seedlings to 125.97 cm in planting 20 days old seedlings (Table 5). Planting 25 days old seedlings produced shorter plant of 124.23 cm than that of planting 20 days old seedlings. Similar result has been reported by Murthy *et al.* (1993) [25]. Gani *et al.* (2002) reported that younger seedlings produced taller plant than older [12]. Total number of tillers hill⁻¹ as well as number of effective tillers hill⁻¹ did not vary significantly due to variation of seedling age. The results revealed that number of effective tillers ranged from 18.14 hill⁻¹ in planting 20 days old seedlings to 19.55 hill⁻¹ in planting 25 days old seedlings (Table 5). The results are in partial conformity with that of Mannan and Siddique (1991) [23]. On the contrary, Das *et al.* (1988) obtained higher tillers hill⁻¹ in younger seedling [8].

Table 5. Phenology and yield attributes of rice as influenced by seedling age during Boro season 2012-2013

Age of seedlings	Days to 50% flowering	Days to maturity	Plant height (cm) at harvest	Total number of tillers hill ⁻¹	Number of effective tillers hill ⁻¹	Number of non-effective tillers hill ⁻¹
15DOS	103.5 ^c	130.3 ^c	123.06	23.09	19.13	3.9
20DOS	106.8 ^b	132.6 ^b	125.97	22.22	18.14	4.0
25DOS	110.9 ^a	136.4 ^a	124.23	23.50	19.55	3.9
CV(%)	2.11	1.55	3.73	9.70	8.85	27.54

LSD _{0.05}	1.910	1.745	NS	NS	NS	NS
---------------------	-------	-------	----	----	----	----

Note: 15DOS= 15 days old seedlings, 20DOS= 20 days old seedlings, 25DOS= 25 days old seedlings; Figures within the same column having same or no letter(s) do not differ significantly at 5% level of probability; NS = Not significant.

Number of non-effective tiller hill⁻¹ variation was not significant and it was found that number of non-effective tiller was about 4.0 hill⁻¹ for different ages of seedling (Table 5).

Table 6. Yield attributes of rice as influenced by seedling age during Boro season 2012-2013

Age of seedlings	Length of panicle (cm)	Total number of spikelets panicle ⁻¹	Number of grains panicle ⁻¹	Number of unfilled spikelets panicle ⁻¹	1000 grain weight (g)
15DOS	23.6	143.74 ^c	111.8 ^b	32.0 ^c	19.52
20DOS	23.3	151.57 ^b	114.8 ^{ab}	36.8 ^b	19.07
25DOS	23.7	160.44 ^a	120.6 ^a	40.1 ^a	19.39
CV(%)	5.21	3.88	6.53	9.26	4.05
LSD _{0.05}	NS	4.996	6.393	2.847	NS

Note: 15DOS= 15 days old seedlings, 20DOS= 20 days old seedlings, 25DOS= 25 days old seedlings; Figures within the same column having same or no letter(s) do not differ significantly at 5% level of probability; NS = Not significant.

Length of panicle did not vary significantly due to variation in the age of seedling in this experiment. Rao and Raju (1987) also recorded similar findings and they stated that seedling age produced no significant effect on panicle length of rice [30]. But Singh *et al.* (2004) concluded that planting 21 days old seedlings produced higher panicle length than that of planting 31, 41 and 51 days old seedlings [34]. A significant variation was found in terms of total number of spikelets panicle⁻¹. The highest number of spikelets (160.4 panicle⁻¹) was obtained from planting 25 days old seedlings. Planting 15 days old seedlings produced the lowest number of spikelets (143.7 panicle⁻¹) (Table 6). The results exhibited that there was significant variation in terms of number of grains panicle⁻¹. The highest number of grains (120.4 panicle⁻¹) was found in the planting 25 days old seedlings and the lowest number of grains (111.7 panicle⁻¹) was found in planting 15 days old seedlings (Table 6). Number of grains of 114.7 panicle⁻¹ was produced in planting 20 days old seedlings. The result did not agree with many other scientists [15, 29, 33]. Planting 25 days old seedlings had significantly highest number of unfilled spikelets (40.1 panicle⁻¹) while planting 15 days old seedlings produced the lowest (32.0 panicle⁻¹) (Table 6). Reddy and Narayana (1981) observed that spikelet sterility decreased with the increased seedling age [32]. But Gill and Shahi (1987) opined that spikelet sterility increased in the older seedlings [14]. Seedling age also failed to produce significant variation in respect of 1000 grain weight. It was found that planting 15, 20 and 25 days old seedlings gave 19.52, 19.07 and 19.39 g 1000 grain weight, respectively (Table 6).

The result did not agree with the findings of Sunder Singh *et al.* (1983) who opined that 1000 grain weight increased significantly with the increase of seedling age [37]. On the contrary, Kamdi *et al.* (1991) reported that 1000 grain weight reduced with transplanting older seedlings [20]. Seedling age showed a significant influence on grain yield. The result presented in Table 7 showed that grain yield increased with the increase of seedling age. Planting 25 days old seedlings gave the highest grain yield (4.49 t ha⁻¹) and it was significantly different from other treatments. Planting 20 days old seedlings produced grain yield of 4.23 t ha⁻¹ which was statistically similar to that of both planting 25 and 15 days old seedlings. The lowest grain yield of 3.86 t ha⁻¹ was obtained from planting 15 days old seedlings (Table 7). Higher grain yield in planting 25 days old seedlings was ascribed to mainly by the higher number of grains panicle⁻¹. Initial higher leaf area and photosynthesis, and less respiration loss for tiller production than 15 and 20 days old seedlings helped to produce more early dry matter accumulation which in turn might augment formation of more number of grain in planting 25 days old seedlings. The results are in close conformity with that of Teetharappan and Palaniappan (1984) who stated that planting 25 days old seedlings gave the highest grain yield of rice [38]. Prasad *et al.* (1992) reported that grain yield increased with the seedling age at transplanting up to 35 days old [28]. Rashid *et al.* (1990) opined that planting 40 days old seedlings gave higher grain yield than that of planting 20 or 60 days old seedlings [31].

Table 7. Yield and harvest index of rice as influenced by seedling age during Boro season 2012-2013

Age of seedlings	Grain yield (t ha ⁻¹)	Straw yield (t ha ⁻¹)	Biological yield (t ha ⁻¹)	Harvest index (%)
15DOS	3.86 ^b	5.25 ^c	9.11 ^c	40.79
20DOS	4.23 ^{ab}	6.19 ^b	10.40 ^b	39.33
25DOS	4.49 ^a	6.78 ^a	11.27 ^a	38.70
CV(%)	10.54	8.42	7.06	7.67
LSD _{0.05}	0.374	0.435	0.613	NS

Note: 15DOS= 15 days old seedlings, 20DOS= 20 days old seedlings, 25DOS= 25 days old seedlings; Figures within the same column having same or no letter(s) do not differ significantly at 5% level of probability; NS = Not significant.

The highest straw yield of 6.78 t ha⁻¹ was obtained from planting 25 days old seedlings while the lowest of 5.25 t ha⁻¹ was obtained from planting 15 days old seedlings. Planting 20 days old seedlings produced 6.19 t ha⁻¹ straw yield which was significantly different from all other seedling ages (Table 7). The result

indicated that planting 20 days old seedlings had a little bit higher plant height as well as total number of tillers hill⁻¹ which might be attributed to produce more straw yield. The lowest straw yield was obtained from planting 15 days old seedlings because of little bit lower plant height and tillering capacity than others. Planting 40 days old seedlings produced higher straw yield than that of planting 20 or 60 days old seedlings [31]. Furuk *et al.* (2009) also stated that planting 2 weeks old seedlings gave the lowest straw yield than planting 4 weeks old seedlings of rice [11]. Biological yield was significantly influenced by seedling age. The highest biological yield (11.27 t ha⁻¹) was obtained from planting 25 days old seedlings whilst the lowest biological yield (9.11 t ha⁻¹) was recorded from planting 15 days old seedlings. The result clearly indicated that biological yield was increased with increase of seedling age from planting 15 to 25 days old (Table 7). Harvest index (HI) was not influence significantly due to seedling age (Table 7). The highest HI (40.79%) was obtained from planting 15 days old seedlings. Planting of both 20 and 25 days old seedlings gave harvest indices of 39.33% and 38.70%, respectively.

Interaction effect of variety and seedling age

The result exhibited that BRRI dhan29 took maximum days for 50% flowering (121.3 days) with planting 25 days old seedlings closely followed by Begunbichi (119.0 days) (Table 8). Khoiaboro took the minimum duration for 50% flowering (91.67 days) at planting 15 days old seedlings. It was found that the variety BRRI dhan29 took the maximum days for maturity (146.0) closely followed by the variety Begunbichi (145.3) at planting 25 days old seedlings. The variety Khoiaboro took minimum days (116.7) for its maturity (Table 8). Variations of plant height at harvest, total number of tillers hill⁻¹ and number of non-effective tillers hill⁻¹ due to the interaction of variety and seedling age were not significant. Interaction of varieties and ages of seedling exerted significant influence on number of effective tillers hill⁻¹. The results revealed that the combination of V₃×25DOS gave the highest number of effective tillers (25.47 hill⁻¹) while the combination V₄×20DOS gave the lowest (15.67 hill⁻¹) (Table 8). The combinations of V₁×25DOS, V₄×15DOS and V₄×25DOS also produced statistically similar number of effective tillers hill⁻¹ to that of V₄×20DOS. Actually there was no consistent trend in respect of the number of effective tillers hill⁻¹ with different seedling ages for different varieties.

Table 8. Phenology and yield attributes of rice as influenced by the interaction of variety and seedling age during Boro season 2012-2013

Interaction (Variety × Seedling age)	Days to 50% flowering	Days to maturity	Plant height (cm) at harvest	Total number of tillers hill ⁻¹	Number of effective tillers hill ⁻¹	Number of non-effective tillers hill ⁻¹
---	--------------------------	---------------------	------------------------------------	--	--	--

V ₁ ×15DOS	101.0 ^d	126.3 ^{de}	101.43	23.63	19.37 ^{bcd}	4.27
V ₁ ×20DOS	101.3 ^d	125.7 ^{de}	100.60	22.76	18.77 ^{bcde}	4.00
V ₁ ×25DOS	102.7 ^d	128.3 ^d	98.97	20.03	16.20 ^{ef}	3.83
V ₂ ×15DOS	111.7 ^c	140.7 ^{bc}	93.63	22.00	19.27 ^{bcd}	2.73
V ₂ ×20DOS	110.0 ^c	138.7 ^c	98.10	20.03	17.07 ^{cdef}	2.97
V ₂ ×25DOS	121.3 ^a	146.0 ^a	98.73	23.03	19.83 ^{bc}	3.20
V ₃ ×15DOS	91.67 ^e	116.7 ^f	142.53	25.50	21.00 ^b	4.50
V ₃ ×20DOS	99.67 ^d	123.3 ^e	151.30	26.23	21.07 ^b	5.17
V ₃ ×25DOS	100.7 ^d	126.0 ^{de}	144.23	29.86	25.47 ^a	4.40
V ₄ ×15DOS	109.7 ^c	137.3 ^c	154.63	21.23	16.87 ^{def}	4.37
V ₄ ×20DOS	116.3 ^b	142.7 ^{ab}	153.87	19.86	15.67 ^f	4.20
V ₄ ×25DOS	119.0 ^{ab}	145.3 ^a	154.97	21.06	16.70 ^{def}	4.37
CV(%)	2.11	1.55	3.73	9.70	8.85	27.54
LSD _{0.05}	3.821	3.491	NS	NS	2.837	NS

Note: V₁= BRRI dhan28, V₂=BRRI dhan29, V₃= Khoiaboro, V₄= Begunbichi, 15DOS= 15 days old seedlings, 20DOS= 20 days old seedlings, 25DOS= 25 days old seedlings; Figures within the same column having same or no letter(s) do not differ significantly at 5% level of probability; NS = Not significant.

Interaction of variety and seedling age was found non-significant in respect of length of panicle and total number of **spikelets** panicle⁻¹. Number of grain panicle⁻¹ also did not vary significantly due to interaction of variety and seedling age (Table 9). Number of unfilled **spikelets** panicle⁻¹ varied significantly due to the interaction of variety and seedling age. The results exhibited that the variety V₂ (BRRI dhan29) had significantly highest number of unfilled **spikelets** panicle⁻¹ (62.5) along with planting 25 days old seedlings (Table 9). It is evident that variety V₃ (Khoiaboro) produced the lowest number of unfilled **spikelets** panicle⁻¹ (11.0) with planting 15 days old seedlings which was statistically identical to that of planting 20 days old seedlings (14.6 panicle⁻¹) of the same variety. A moderate number of unfilled **spikelets** panicle⁻¹ was observed in both the varieties V₁ (BRRI dhan28) and V₄ (Begunbichi) with all seedling ages.

Table 9. Yield attributes of rice as influenced by the interaction of variety and seedling age during Boro season 2012-2013

Interaction (Variety × Seedling age)	Length of panicle (cm)	Total number of spikelets panicle ⁻¹	Number of grains panicle ⁻¹	Number of unfilled spikelets panicle ⁻¹	1000 grain weight (g)
V ₁ ×15DOS	22.10	136.30 ^f	105.5 ^e	30.9 ^d	22.27
V ₁ ×20DOS	21.83	148.63 ^e	115.7 ^{de}	32.87 ^d	22.10
V ₁ ×25DOS	22.63	138.93 ^{ef}	104.7 ^e	34.2 ^d	21.67
V ₂ ×15DOS	25.73	166.70 ^d	113.5 ^{de}	53.2 ^b	21.97
V ₂ ×20DOS	23.93	174.93 ^d	120.2 ^{cd}	54.7 ^b	23.03
V ₂ ×25DOS	24.57	192.13 ^{bc}	129.6 ^c	62.5 ^a	22.13
V ₃ ×15DOS	21.83	86.70 ^h	75.7 ^f	11.0 ^e	21.17
V ₃ ×20DOS	22.33	66.03 ⁱ	51.5 ^g	14.6 ^e	20.17
V ₃ ×25DOS	21.87	109.97 ^g	80.4 ^f	29.6 ^d	21.10
V ₄ ×15DOS	24.57	185.27 ^c	152.4 ^b	32.9 ^d	12.67
V ₄ ×20DOS	24.97	216.67 ^a	171.7 ^a	45.0 ^c	10.97
V ₄ ×25DOS	25.53	200.73 ^b	167.7 ^a	34.0 ^d	12.67
CV(%)	5.21	3.88	6.53	9.26	4.05
LSD _{0.05}	NS	9.993	12.79	5.693	NS

Note: V₁= BRRI dhan28, V₂=BRRI dhan29, V₃= Khoiaboro, V₄= Begunbichi, 15DOS= 15 days old seedlings, 20DOS= 20 days old seedlings, 25DOS= 25 days old seedlings. Figures within the same column having same or no letter(s) do not differ significantly at 5% level of probability. NS = Not significant.

The results revealed that interaction of variety and seedling age failed to produce significant effect on 1000 grain weight. The values of 1000 grain weight presented in Table 9 indicated that the varieties V₁, V₂ and V₃ had comparatively larger sized grain (ranged from 20.17 g to 23.03 g) while V₄ had small sized grain (ranged from 10.97 g to 12.67 g). Grain yield was not significantly varied due to interaction of variety and seedling age (Table 10). The results indicated that all varieties included in the experiment required a particular seedling age for producing maximum grain yield.

Table 10. Yield and harvest index of rice as influenced by the interaction of variety and seedling age during Boro season 2012-2013

Interaction (Variety × Seedling age)	Grain yield (t ha ⁻¹)	Straw yield (t ha ⁻¹)	Biological yield (t ha ⁻¹)	Harvest index (%)
V ₁ ×15DOS	5.24	5.50 ^{cd}	10.74 ^b	49.19
V ₁ ×20DOS	5.42	5.59 ^{cd}	11.01 ^b	49.28
V ₁ ×25DOS	5.44	6.04 ^{bc}	11.48 ^b	47.33

V ₂ ×15DOS	5.58	5.63 ^{cd}	11.20 ^b	49.83
V ₂ ×20DOS	6.23	7.96 ^a	14.19 ^a	43.78
V ₂ ×25DOS	6.94	6.88 ^b	13.81 ^a	50.23
V ₃ ×15DOS	2.66	5.84 ^{cd}	8.50 ^{cd}	31.32
V ₃ ×20DOS	2.98	6.13 ^{bc}	9.10 ^c	32.70
V ₃ ×25DOS	3.07	8.62 ^a	11.69 ^b	26.24
V ₄ ×15DOS	1.97	4.04 ^e	6.01 ^e	32.81
V ₄ ×20DOS	2.30	5.00 ^d	7.30 ^d	31.54
V ₄ ×25DOS	2.51	5.59 ^{cd}	8.10 ^{cd}	31.00
CV(%)	10.54	8.42	7.67	7.67
LSD _{0.05}	NS	0.865	1.227	NS

Note: V₁= BRRI dhan28, V₂=BRRI dhan29, V₃= Khoiaboro, V₄= Begunbichi, 15DOS= 15 days old seedlings, 20DOS= 20 days old seedlings, 25DOS= 25 days old seedlings; Figures within the same column having same or no letter(s) do not differ significantly at 5% level of probability. NS = Not significant.

Effect of interaction between variety and seedling age on straw yield was found significant. The highest straw yield (8.62 t ha⁻¹) was obtained from the combination of V₃×25DOS. The lowest straw yield (4.04 t ha⁻¹) was recorded from the combination of V₄×15DOS (Table 10). The combinations of V₁×15DOS, V₁×20DOS, V₁×25DOS, V₂×15DOS, V₃×15DOS and V₃×20DOS produced statistically similar straw yield. Significant variation was found in respect of biological yield due to interaction effect of variety and seedling age. The highest (14.19 t ha⁻¹) biological yield was obtained from the combination of V₂×20DOS which was statistically identical to that of V₂×25DOS (13.81 t ha⁻¹). The lowest biological yield (6.01 t ha⁻¹) was recorded from the combination of V₄× 15DOS (Table 10). The results indicated that the combinations of V₁×20DOS, V₁×25DOS and V₃×25DOS produced statistically similar biological yields of 11.01, 11.48 and 11.69 t ha⁻¹, respectively. The combinations of V₁×15DOS and V₂×15DOS also produced similar biological yields of 10.74 and 11.20 t ha⁻¹, respectively. Biological yields of 7.30 and 8.10 t ha⁻¹ of the combinations of V₄×20DOS and V₄×25DOS were statistically similar. Interaction effect of variety and seedling age also produced significant influence on harvest index (HI). The highest HI (50.23%) was obtained from the combination of V₂×25DOS which was similar to the combinations of V₁×15DOS, V₁×20DOS and V₂×15DOS (Table 10). The lowest HI (26.24%) was obtained from the combination of V₃×25DOS which was significantly different from other combinations. The combinations of V₃×15DOS, V₃×20DOS, V₄×15DOS, V₄×20DOS and V₄×25DOS produced statistically similar HI's of 31.32%, 32.71%, 32.81%, 31.54% and 31.00%, respectively.

Economic performance

Total cost of cultivation was calculated maximum (US\$ 965.73 ha⁻¹) in the variety BRRI dhan29 followed by the variety BRRI dhan28 (US\$ 957.29 ha⁻¹). The maximum production cost incurred in the HYV's due to the requirement of more inputs for their production (Table 11). Maximum gross return (US\$ 1665.00 ha⁻¹), net return (US\$ 699.28 ha⁻¹) and BCR (1.72) were also obtained from the same variety BRRI dhan29 with planting 25 days old seedlings. The higher profitability obtained in BRRI dhan29 was due to its higher yield. It was found that cultivation of local variety 'Begunbichi' was more profitable than BRRI dhan28 with planting 25 days old seedlings and than Khoiaboro at all seedling ages. This was due to more market price of the scented grain of Begunbichi (US\$ 0.43 kg⁻¹) compared to BRRI dhan28 (US\$ 0.23 kg⁻¹). Cultivation of Khoiaboro was found less profitable due to its lower productivity as well as low market price because of its coarse size grains.

Table 11. Cost and return analysis of production of different rice varieties during Boro season 2012-2013

Interaction		Yield (t ha ⁻¹)		Total cost of cultivation (US\$ ha ⁻¹)	Gross return (US\$ ha ⁻¹)			Net return (US\$ ha ⁻¹)	BCR
Variety	Seedling age	Grain	Straw		Grain	Straw	Total		
V ₁	15DOS	5.24	5.50	957.29	1205.20	55.00	1260.20	302.91	1.32
	20DOS	5.42	5.59	957.29	1246.60	55.90	1302.50	345.21	1.36
	25DOS	5.44	6.04	957.29	1251.20	60.40	1311.60	354.31	1.37
V ₂	15DOS	5.58	5.63	965.73	1283.40	56.30	1339.70	373.98	1.39
	20DOS	6.23	7.96	965.73	1432.90	79.60	1512.50	546.78	1.57
	25DOS	6.94	6.88	965.73	1596.20	68.80	1665.00	699.28	1.72
V ₃	15DOS	2.66	5.87	611.66	558.60	58.70	617.30	5.64	1.01
	20DOS	2.98	6.13	611.66	625.80	61.30	687.10	75.44	1.12
	25DOS	3.07	8.62	611.66	644.70	86.20	730.90	119.24	1.19
V ₄	15DOS	1.97	4.04	723.94	847.10	40.40	887.50	163.56	1.23
	20DOS	2.30	5.00	723.94	989.00	50.00	1039.00	315.06	1.44
	25DOS	2.51	5.53	723.94	1079.30	55.30	1134.60	410.66	1.57

Note: V₁= BRRI dhan28, V₂=BRRI dhan29, V₃= Khoiaboro, V₄= Begunbichi, 15DOS= 15 days old seedlings, 20DOS= 20 days old seedlings, 25DOS= 25 days old seedlings.

Selling price: Rice grain – US\$ 0.23 kg⁻¹ for both BRRI dhan28 and BRRI dhan29; US\$ 0.21 kg⁻¹ for Khoiaboro; US\$ 0.43 kg⁻¹ for Begunbichi; Straw- US\$ 0.01 kg⁻¹; 1 US\$= BDT 80; BCR = Benefit-Cost Ratio.

CONCLUSIONS

On the basis of the results obtained from the experiment lead to conclude that 25 days old seedlings was found to produce the highest grain yield and therefore, all High Yielding and local varieties are suggested to be grown with 25 days old seedlings. BRRI dhan29 gave the maximum economic benefit followed by Begunbichi a local aromatic Boro rice variety. Considering the profitability the local variety Begunbichi may also be suggested for cultivation in other parts of the country as there is only a high yielding fine rice variety BRRI dhan50.

REFERENCES

1. Ali MY, Rahman MM, Haq MF. Effect of time of transplanting and age of seedling on the performance of late planted aman rice. Bangladesh J. Sci. Ind. Res. 1995;30(1):45-58.
2. Ashraf M, Khalid A, Ali K. Effect of seedling age and density on growth and yield of rice in saline soil. Pakistan J. Biol. Sci. 1999;2:860–862.
3. BRRI. ‘Adhunik Dhaner Chash’ (Modern rice cultivation). Bangladesh Rice Research Institute, Joydebpur, Gazipur. 9th Edt. 2011: p.7.
4. BRRI. ‘Adhunik Dhaner Chash’(Modern Rice Cultivation). In Bengali. Pub. No.5. Bangladesh Rice Res. Inst. Joydebpur, Gazipur, Bangladesh; 1991: pp.45.
5. BRRI. Annual Report for 1990. Bangladesh Rice Res. Inst. Joydebpur, Gazipur, Bangladesh; 1992: pp.329.
6. Chandra D, Manna GB. Effect of planting date seedling age and planting density on late planted wet season Rice. Intl. Rice Res. Newsl. 1988;13:30-31.
7. Chopra MK, Chapra N, Sinha SN. Influence of dates of transplanting on production and quality of scented rice (*Oryza sativa*) seed. Indian J. Agric. Sci. 2002;73(1):12-13.
8. Das K, Biswal D, Pradhan T. Effect of plant density and age of seedlings on the growth and yield of rice variety, Parjit. Oryza. 1988;25:191-194.
9. De Datta SK. Principles and Practices of Rice Production. New York, USA. International Rice Research Institute, John Willy & Sons, Inc.;1981.

- 542 10. Dey MM, Miah MN, Mustafi BAA, Hossain M. Rice production constraints in Bangladesh:
543 Implication for further research priorities. In: Rice Research in Asia: Progress and Priorities. CAB
544 International and IRRI Philippines; 1996.
- 545 11. Furuk MO, Rahman MA, Hassan MA. Effect of seedling age and number of seedling per hill on the
546 yield and yield contributing characters of BRRI dhan33. Intl. J. Sustain. Crop Prod. 2009;4:58-61.
- 547 12. Gani A, Rahman A, Dahono, Rustam, Hengsdijk H. Synopsis of water management experiments in
548 Indonesia. In: Water-wise Rice Production, IRRI; 2002:pp. 29-38.
- 549 13. Gardner FP, Pearce RB, Mitchell RL. Physiology of Crop Plants. 1st Edn., The Iowa State
550 University Press, ISBN-10: 081381376X, USA.; 1985: pp. 327.
- 551 14. Gill PS, Shahi HN. Effect of nitrogen levels on relation to age of seedling and time of transplanting
552 on the growth, yield and milling characteristics of rice. Indian J. Agric. Sci. 1987;57(9):630-634.
- 553 15. Hariom R, Joon K, Singh OP. Effect of age of seedlings and different date of transplanting on the
554 growth and yield of rice. Indian J. Agron. 1989;34(3):325-327.
- 555 16. Hossain M, Deb UK. Liberalization of Rice Sector: Can Bangladesh with stand Regional
556 Competition? Poster paper presented at PETRRA Communication Fair 2003 held at Hotel Sheraton,
557 Dhaka on Aug. 10-11; 2003.
- 558 17. Hossain SMA, Salam MU, Sattar M, Ahmed JU. Hill population in rice researchers' findings and
559 farmers' technology. Bangladesh J. Agril. Sci. 1989;16:213-217.
- 560 18. Hossain SMA, Sattar M, Ahmed JU. Spacing and seedling age in rice. Component Technology
561 Research. CSRDP, BAU, Mymensingh, Bangladesh. 1983: pp.12-13.
- 562 19. IRRI. Annual Report. Manila (Philippines): International Rice Research Institute;1997.
- 563 20. Kamdi JT, Hatwar KG, Bodhe GN, Patil SM. Effect of age of seedling at transplanting on yield of
564 rice varieties. J. Soils and Crops. 1991;1(2):154-156.
- 565 21. Kewat ML, Agrawal SB, Agrawal KK, Sharma RS. Effect of divergent plant spacings and age of
566 seedlings on yield and economics of hybrid rice (*Oryza sativa*). Indian J. Agron. 2002;47:367-371.
- 567 22. Makarim AK, Balasubramanian V, Zaini Z, Syamsiah I, Diratmadja IGPA, Arafah H, Wardana IP, Gani
568 A. System of Rice Intensification (SRI): Evaluation of seedling age and selected components in Indonesia.
569 In: Bouman BAM, A Hengsdijt A, B Hardy, PS Bindraban, TP Tuong and JK Ladha (Eds.), Water-wise
570 Rice Production, IRRI; 2002:pp. 129-139.
- 571 23. Mannan MA, Siddique JB. Effect of seedling age and date of planting on the growth and yield of
572 photo-period sensitive rice. Bangladesh Rice J. 1991;2(1-2):104-106.

24. McHugh O. Farmer alternative wet/dry, non flooded and continuously flooded irrigation practices in traditional and intensive systems of rice cultivation in Madagascar. MS Thesis. Cornell University. Ithaca, NY, USA; 2002.
25. Murthy KK, Pillai KG, Srinivasn TE and Ramprasad AS. Rice varieties and technological improvement for late planting during the main kharif season. *Seeds and Farms*.1993;49: 9-13.
26. Nandini DK, Singh AI. Influence of seedling age and plant density on the performance of rice. Central Agricultural University, College of Agriculture, Imphal, Manipur, India; 2000.
27. Padalia CR.. Effect of age of seedling on the growth and yield of transplant rice. *Oryza*. 1981;18(3):165-167.
28. Prasad K, Singh R, Singh S. Effect of seedling age and number of seedling per hill on the yield of rice in sodic soil. *Curr. Agric*. 1992;16(1-2):67-70.
29. Raju RA, Reddy GV, Reedy MN. Response of long duration rice to spacing and age of seedlings. *Indian J. Agron*. 1989;31(4):506-507.
30. Rao CP, Raju MS. Effect of age of seedling, nitrogen and spacing on rice. *Indian J. Agron*. 1987;32:100–102.
31. Rashid MA, Aragon ML, Denning GL. Influence of variety, seedling age and nitrogen on growth and yield of rice grown on saline soil. *Bangladesh Rice J*. 1990;1(1): 37-47.
32. Reddy SN, Narayana P. Effect of age of seedlings and different dates of transplanting on yield and yield components of rice (*Oryza sativa* L.). *Res. Bull. Mahathwada Agril. Univ*. 1981;5(5/12):18-21. [Rice Abst. 7(12): 152. 1984].
33. Roy BC, Sattar SH, Gaffer MA, Islam MA. Seedlings age effect on yield of irrigated rice. *Bangladesh Rice J*. 1992: 3(1 & 2):83-88.
34. Singh KK, Yadav SK, Tomar BS, Singh JN, Singh PK. Effect of seedling age on seed yield and seed quality attributes in rice (*Oryza sativa*) cv. Pusa Basmati-1. *Seed Res*. 2004;32(1):5-8.
35. Singh RS, Singh SB. Effect of age of seedlings, N levels and time of application on growth and yield of rice under irrigated condition. *Oryza*. 1999;36(4):351-354.
36. Singh RS, Singh SB. Response of rice (*Oryza sativa* L.) to age of seedlings and level and time of application of nitrogen under irrigated condition. *Indian J. Agron*. 1998;43(4):632-635.
37. Sundar Singh SD, Gurujan B, Sabaiah V, Subramanian V. Note on the performance of rice Co-40 at different age of planting in single crop periar canal command area. *Madrass Agril. J*. 1983;70:821-822.

- 604 38. Teetharappan TS, Palaniappan SP. 1984. Optimum seedling age transplanting short duration rice.
605 Int. Rice Res. Newsl. 1984:9:2-29.
- 606 39. Thanunathan K, Sivasubramanian V. Age of seedling and crop management practices for high
607 density (HD) grain in rice. Crop Res. 2002:24:421–424.
- 608 40. Venugopal K, Singh RD. Effect of plant density and age of seedling on the yield of DR 92 rice
609 variety in Sikkim. *Oryza*. 1985:22:162-165.
- 610 41. Wagh RG, Khanvilkar SA, Thorat ST. Effect of age of seedlings at transplanting, plant densities
611 and nitrogen fertilization on the yield of rice variety R711. *Oryza*. 1988:25:188-190.
- 612 42. Wang S, Cao W, Jiang D, Dai T, Zhu Y. Physiological characteristics and high-yield techniques with
613 SRI rice. *In: Assessments of the System of Rice Intensification. Proc. Intl. Conf., Sanya, China.*
614 Apr. 1-4; 2002: pp.116-124. [www. bank-brri.org](http://www.bank-brri.org).