

**INFLUENCE OF POULTRY DROPPINGS ON SOIL CHEMICAL PROPERTIES
AND PERFORMANCE OF RICE (*Oriza Sativa* L.) IN SOKOTO, SUDAN
SAVANNA ZONE OF NIGERIA**

M. Audu, M. Haliru and A.M. Isah

**Department of Soil Science and Agricultural Engineering, Usmanu Danfodiyo University
Sokoto, Nigeria.**

Abstract

Proper management of poultry manure could serve as a sustainable source of fertilizer for increased rice production in Sokoto State of Nigeria. As a result, an experiment was conducted in a screen house at the Botanical Garden, Biological Science Department of the Sokoto State Polytechnic, to determine the influence of poultry dropping on some chemical properties of soil and performance of rice (*Oriza Sativa* L.). The treatments consisted of three levels of poultry dropping: 2, 5 and 10 tha^{-1} and a control (without fertilizer). The experiment was laid in a completely randomized design (CRD) replicated three times. Plant parameters like plant height, number of tillers and number of leaves were taken fortnightly until harvest at 16th week after planting (WAP). Grain and stalk yield were taken at harvest. The nutrient composition of the soil samples and poultry droppings used was determined using standard laboratory procedures. The result revealed that application of poultry manure had significant ($P > 0.05$) effect on soil organic carbon, available phosphorus, exchangeable bases (Ca, Mg, K, Na), CEC and rice performance in which application of 10 tha^{-1} recorded the highest rice stalk and grain yield. This research therefore, concluded that application of poultry dropping is an important means of improving soil fertility and that, application of 10 tha^{-1} can bring about high growth performance and yield of rice in the study area.

Key words: Poultry Droppings, Levels, Soil, Rice and Sudan savanna

INTRODUCTION

Farming is the main occupation of people of Sokoto State of Nigeria and mostly, they grow cereal and vegetable crops. One of the major cereal crops grown in this area is rice. Rice belongs to the grass family poaceae from the genus *Oryza* of which two species (*Oryza sativa* and *Oryzaglaberrima*) are cultivated. It is normally grown as annual plant in the tropical areas; it can survive as a perennial crop (IRRI, 2008). Rice is the staple food of over half of the world population and provides 20% of the world dietary energy supply, compare to wheat 19 and maize 5% (FAO, 2004). In Nigeria, rice is a major cereal crop consumed by over 120 million population of the country (EIARD, 2013), placing a high demand for this crop. However, production of rice requires fertile soil that are rich in nutrients, and soils of the Sudan savanna are constrained by erosion, degradation of physical condition, deterioration of nutrient status and changes in the composition and number of soil organisms (Ogunwole *et al.*, 2005; Adeniyi, 2008) which limit their productivity. Usman *et al.* (2007) stated that soils of the savanna region of Nigeria are relatively low in nutrients and organic matter content. Thus, application of fertilizers either in organic or inorganic form could be a sure way of meeting nutrient requirements for high productivity of rice in Sokoto state of Nigeria. However, short supply and high cost of fertilizers during the growing season are the major limitations to small holder farmers in this region (Sobulo and Osiname, 1985, Maobe *et al.*, 2000). Therefore, the need to explore alternative sources of nutrient supply remains imperative.

It has been established that organic manure can serve as alternative practice to mineral fertilizers (Naeem *et al.*, 2006) for improving soil structure (Dauda *et al.*, 2008) and microbial biomass (Suresh *et al.*, 2004). Organic manure application on the farm has yielded good response of crops and residual effect on soils. Agboola and Obatolu (1989), Lombin *et al.* (1991), Ojeniyi and Adeniyi (1999), and Kwari (2003) have all demonstrated the use of organic manure as a sound strategy for maintaining soil fertility and crop productivity. Akanni (2005) mentioned that manure application improved organic matter (OM), N, P and exchangeable cation concentration of soil that could benefit growing crops. Poultry manure had been reported to improve growth and yield of maize (Ezeibekwe *et al.*, 2009). It also improves the chemical and biological qualities of the soil which increases crop productivity than chemical fertilizers (Obi and Ebo, 1995). In a similar way, Agbede *et al.* (2008) reported that application of poultry

manure(7.5t/ha)increased growth parameter (plant height, stem girth, leaf area) of sorghum in south-west Nigeria. Further, Boateng *et al.* (2006) also reported increased in maize plant height due to organic fertilizer (poultry manure) in Ghana.

Livestock production is one of the major occupation of the people of Sokoto State of Nigeria and therefore relative availability of animal and poultry wastes in the area (Okafor, 1999), but are poorly managed for effective utilization due to lack of scientific basis for advising farmers on aspects such as appropriate application rates, storage techniques and application methods (Maerere, 2001). Adeboye *et al.* (2006) suggested that researches on soil fertility should be focused on locally available and affordable internally sourced materials to improve the production of cereals which are important in the diets of millions of people in the Nigeria.

In order to supply food for steadily increasing human population from fixed or limited land resources, improvement in existing management practices is essential through the use of locally available and affordable resources. This can be achieved through proper rate of nutrient selection which depends on the knowledge of nutrient supplying power of the soil and on which the crop is to be grown (Tisdale *et al.*, 2003). This study was therefore conceived to determine the influence of poultry dropping on soil chemical properties and performance of rice in Sokoto, Sudan Savanna agro-ecological zone of Nigeria.

MATERIALS AND METHODS

Site Description

The experiment was conducted during the 2012/2013 dry season in a screen house at the Botanical Garden of the Biological Science Department, Sokoto State Polytechnic, Sokoto. Sokoto State is located between Latitudes 11° 30¹N and 13° 50¹N and Longitudes 4° 0¹E and 6° 0¹E, 315m above sea level. Sokoto falls in the Sudan savanna agro-ecological zone of Nigeria (Ojanuga, 2005) that is characterized by erratic and scanty rainfall that last for about four months (Mid June-September) and dry period (October- May). The annual rainfall of the area is highly

variable over the years and averaged around 700mm (Singh, 1995) with minimum and maximum temperatures of the year fluctuating between 15 and 40°C, respectively (Arnborg, 1988).

Screen House Procedure and Soil Analysis

The soil sample used for this experiment was collected from a fallow land within the Faculty of Agriculture Teaching and Research Lowland Farm, Usmanu Danfodiyo University, Sokoto, at 0-15cm depth. The experiment was conducted at the Botanical garden of the Biological Science department, Sokoto State Polytechnic in 2013/2014 dry season (December – March). The soil was air dried, crushed and passed through a 2mm sieve. A sub-sample was analyzed for some physical and chemical properties using the methods described by IITA (1989). Particle size distribution was determined using the Buoyoucos hydrometer method. Soil pH was determined using glass electrode pH meter. Cation exchange capacity was determined by the neutral ammonium acetate saturation (NH₄OAC) method. Organic carbon was determined using Walkley and Black method. Exchangeable calcium and magnesium were determined using EDTA titration method, while exchangeable sodium and potassium was determined using flame photometer. Total N was determined using micro Kjeldahl digestion method, and available P was determined by Bray No.1 method. Seven and half (7.5 kg) kilograms each of the soil was placed in plastic container (10 L) according to the number of the treatments. Soil samples were also collected from each pot at harvest (16 WAP) and analyzed for pH, organic C, total nitrogen, available phosphorus, CEC and exchangeable bases contents. The poultry dropping used for the experiment was also analyzed for nitrogen, phosphorus and potassium contents.

Experimental Set-up, Data Collection and Analysis

Treatment consisted of three levels of poultry manure: 2, 5 and 10t ha⁻¹ corresponding to 7.5g, 18.75g and 37.5 g in 7.5kg of soil per pot respectively and a control (without manure). An improved rice variety (Faro₄₄) was planted as a test crop. The experiment was laid in completely randomized design (CRD) replicated three times. The poultry dropping was mixed evenly with soil and watered to field capacity and allow for a period of one week before planting. Five seeds per pot were sown and latter thinned to three stands per pot at two weeks after planting. Weeds

were controlled manually by hand picking and the plants were irrigated when necessary. Growth parameters, such as plant height, number of leaves and tillers per plant were recorded at 2 weeks interval. The crop was harvested 16 week after planting and some yield parameters were taken (stalk and grain yield). The data obtained were subjected to analysis of variance (ANOVA) using SAS (2003) procedure for CRD. Significant difference in the treatments means was further separated using least significant difference (LSD) (Gomez and Gomez, 1984).

RESULT AND DISCUSSION

Table 1: Initial Soil and Poultry Dropping Analysis

SoilParameters	Value
pH (H ₂ O) 1:1	7.01
Organic carbon (%)	2.87
Total nitrogen (%)	0.09
Available phosphorous (mgkg ⁻¹)	0.99
Cation exchange capacity (CEC) (Cmol kg ⁻¹)	9.42
Exchangeable bases (Cmol kg ⁻¹)	
Calcium (Ca ²⁺)	1.35
Magnesium (Mg ²⁺)	0.55
Potassium (K ⁺)	0.31
Sodium (Na ⁺)	0.96
Sand (%)	9.0
Silt (%)	5.0
Clay (%)	86.0
Texture	Clay
Poultry Dropping	
Total nitrogen (%)	0.64
Available phosphorous (mgkg ⁻¹)	1.68
Potassium (Cmolkg ⁻¹)	0.54

Result of the initial physical and chemical properties of the soil is presented in Table 1. The result indicated that organic carbon content of the soil was high while, total nitrogen, and available phosphorus was very low (Table 1). Exchangeable potassium and sodium were high while magnesium was moderate. Cation exchange capacity was moderate and exchangeable calcium was low based on the standard ratings of Esu (1991).

Table 2: Influence of Poultry Manure on Chemical Properties of Soil

Exchangeable bases

Treatment t ha ⁻¹	pH	OC %	Total N	Avail. P mgkg ⁻¹	Ca cmol(+)kg ⁻¹	Mg	K	Na	CEC
0	7.08	1.84c	0.08	0.41d	1.23c	0.56b	0.35b	0.55b	4.25b
2	7.15	2.49b	0.09	0.46c	1.44b	0.76a	0.56a	0.62b	4.70b
5	7.25	2.71ab	0.10	0.51b	1.65a	0.75a	0.55a	0.58b	5.45a
10	7.26	2.82a	0.12	0.57a	1.77a	0.84a	0.57a	0.75a	5.27a
SE	0.21	0.09	0.04	0.01	0.04	0.03	0.02	0.02	0.13
SIG.	NS	*	NS	*	*	*	*	*	*

Means followed by same letter (s) within the same row are not significantly different at 5% level of probability.

NS=not significant.

*= significant at 5% level.

Influence of poultry dropping on chemical properties of soil is presented in Table 2. The result indicated that treatments had significant ($P < 0.05$) effect on all the considered chemical parameters of the soil except pH and total nitrogen. However, increased in pH was recorded due to treatments application as compared to the initial values with the highest increase due to application of 10t ha⁻¹. Application of 10t ha⁻¹ also gave the highest value of organic carbon, total N, available phosphorous, exchangeable bases and cation exchange capacity of the soil (2.82%, 0.12%, 0.57mg kg⁻¹, 1.77, 0.84, 0.57, 0.75 and 5.27cmol(+)kg⁻¹) respectively. This could be attributed to the influence of poultry dropping on soil fertility as it mineralized, resulting to an improvement in the soil condition and microbial activities. However, the least values were obtained in pots where no poultry dropping was applied (control). The result of this finding was similarly reported by Balasubramanian and Singh (1978) and Wild (1988) that, application of farm yard manure increased the availability of phosphorous in soil solution and reduced phosphorous adsorption in an experiment conducted on an Ultisol in Nigeria. Pierre and Morrean (1997) observed that the addition of farmyard manure combined with mulch had enhanced the physical properties of soil. The authors reported that moisture retention, water infiltration and cation exchange capacity in the soil improved following the application of organic matter (3 and 4 tons

ha⁻¹) on a farmland. Poultry manure application is known to improve SOM, micro-nutrient status and micro-nutrient qualities of the soil (Maerereet *al.*, 2001;Adeniyi and Ojeniyi, 2003). Adesodunet *al.*(2005) found that application of poultry manure to soil increased soil organic matter, N and P contents and aggregate stability.

Table 3: Influence of Poultry Manure on Growth Performance and Yield of Rice at 16WAP

Treatment	Plant height (cm)	No. of Leaves/plant	No. of Tillers/plant	Stalk yield (g/pot)	Grainyield (g/pot)
0	35.94b	30.11c	8.50c	28.32b	3.20b
2	38.91ab	32.47bc	8.67c	32.55ab	4.35ab
5	46.99a	33.58b	10.17b	33.93ab	5.20ab
10	40.14ab	52.25a	15.59a	38.67a	6.32a
SE	1.58	1.88	0.63	1.10	0.30
SIG.	*	*	*	*	*

Means followed by same letter (s) within same row are not significantly different at 5% level

* significant at 5% level.

The effect of different levels of poultry dropping on growth and yield parameters of rice is presented in Table 3. The result shows that, levels of poultry dropping had significant effect ($p < 0.05$) on plant height, number of leaves and number of tillers per plant.

Plant Height

Application of 5tha⁻¹ poultry dropping gave the highest (46.99cm) plant height at 16WAP. However, this was statistically similar with application of 10 and 2tha⁻¹, while the lowest plant height (35.94cm) was recorded in pots without poultry dropping (control). This could be due to the availability of more nutrients by the plant throughout the growing period as stated by Farhadet *al.* (2009) that, increase in plant height with poultry manure application was mainly due

tomore availability of nutrients by poultry manure throughout the growing season. These results are in accordance with the findings of Mitchell and Tu (2005) and Warren *et al.* (2006).

This result also corroborates with the report of Opara – Nadiet *al.*(2000) on increase in the height of maize treated with organic fertilizer. This indicates the significance of organic manure on this very important growth parameter of plant. Furthermore, Awotundunet *al.*(2000) observed a similar increase in height of maize plant that had cow dung application. Kwari (2003) observed that the height of millet increased when 7.5 t ha^{-1} cattle manure was added to soil relative the control plots. The positive influence of organic manure on plant height is also consistent with the report of Arunahet *al.* (2007) who also observe that, the height of two sorghum varieties had significantly increased due to amendment of soil with quantities of organic materials that was applied at $2 - 4 \text{ t ha}^{-1}$ in Zaria, Nigeria.

Number of Leaves and Tillers

Application of poultry dropping at 10tha^{-1} recorded the highest number of leaves and tillers per rice plant at 16WAP (52.25 and 15.59 respectively). While the lowest numbers of leaves (30.11) and tillers (8.50) were obtained from plants in the control pots. This could also be attributed to availability of the required plant nutrients in that treatment which helped in promoting the vegetative growth of the plants. This is in line with the findings of Akanni (2005) who reported that manure application improved organic matter, N, P and exchangeable cation concentration of soil that could benefit growing crops. In a similar way, Agbedeet *al.* (2008) reported that application of 7.5tha^{-1} poultry manure increased growth parameters (plant height, stem girth, leaf area) of sorghum in south-west Nigeria.

Stalk and Grain Yield

Effect of levels of poultry dropping on stalk and grain yield of rice is presented in Table 3. The result indicated that, the treatments had significant effect ($p < 0.05$) on stalk and grain yield. Application of 10 t ha^{-1} recorded the highest stalk and grain yield. This was similarly reported by Farhadet *al.* (2009) that, grain yield of spring maize was significantly affected by the application of different levels of poultry manure. These results are in accordance with the findings of Boateng et *al.* (2006), Deksissa et *al.* (2008) and Ezeibekweet et *al.* (2009) that poultry manure significantly increased the grain yield of maize in Ghana.

Conclusion

This research revealed that application of different levels of poultry dropping on soils under rice production had significant effect on all the soil and rice parameters measured. Application of poultry dropping at 10 t ha^{-1} gave significantly the highest value of organic carbon, available phosphorus, exchangeable bases and CEC status of the soil. It also gave the best growth and yield performance of rice in the study area except in plant height where 5 t ha^{-1} of poultry dropping was superior. Therefore application of 10 t ha^{-1} of poultry dropping incorporated at least two weeks before planting could be a satisfactory rate for soil fertility management under rice production in the study area.

References

- Adeboye, M. K. A., E. N. O. Iwuafor and Agbenin, J. (2006). Rotation Effect of Grain and Herbaceous Legume on Maize yield and Chemical Properties of an Alfisol in the Northern Guinea Savanna. *Nigerian Journal of Soil Science Research*, 6:22-31.
- Adeniyi, N. O. (2008). Evaluation of NPK Fertilizer and Poultry Manure for Soil Fertility Enhancement and Cassava Production in two agro-ecologies of southwest Nigeria. *Nigerian Journal of Soil Science*, 18: 48 – 53.
- Adeniyi, O.N. and Ojeniyi, S.O. (2003). Comparative effectiveness of different levels of poultry manure with NPK fertilizer on residual soil fertility nutrient uptake and yield of maize. *Moor Journal of Agricultural Research*, 4:191-197.

- Adesodun, J. K., J. S. C.MbagwuandOti, N. (2005).Distribution of Carbon, Nitrogen and Phosphorus in Water Stable Aggregates of an Organic Waste Amended Ultisol in Southern Nigeria.*Bio-resources Technology*, 96:509 – 516.
- Agbede, T.M., S.O. Ojeniyiand Adeyemo, A.J. (2008).Effect of poultry manure on soil physical and chemical properties, growth and grain yield of sorghum in southwest, Nigeria.*American-Eurasian Journal of Sustainable Agriculture*, 2(1):72-77.
- Agboola, A. A. and Obatolu, C. R.(1989).Problems and Prospects of Maintaining Continuous Arable Crops Production in the Humid Tropics through Soil Organic Matter.Proceedings of International Symposium UNSECO, UNDP, ICSU, Yamoussoukro pp. 641-653.
- Akanni, D.I. (2005).Response of Nutrient Composition and Yield Components of Tomato (*Lycopersiconesculentum*) to Livestock Manure.Ph.D. Thesis, Department of Crop, Soil and Pest Management, Federal University of Technology, Akure.Pp. 120.
- Arnborg, T. (1988).Where Savannah turns into Desert. International Rural Development Centre, Swedish University of Agriculture Sciences and Rural Development Studies No. 24: pp. 21-23.
- Aruna, U.L., U.F. Chiezey and Aliyu, L. (2007). Effect of Nitrogen Fertilizer and Poultry Manure on Growth Attributes of Two Sorghum Varieties. In: Uyovbisere, E.O., Raji, B.A., Yusuf, A.A., Ogunwole, J.O., Aliyu, L. and Ojiniyi, S.O. (eds.).*Soil and Water Management for Poverty Alleviation and Sustainable Environment*. Proceedings of the 31st Annual Conference of Soil Science Society of Nigeria, held at Ahmadu Bello University, Samaru Zaria. Nov. 13th – 17th, 2006. Pp. 299 – 304.
- Awotundun, J.S., A.O. Okuntade and Oyinloye, R. (2000).Comparative Effects of Organic and Inorganic Fertilizers on the Yield of an open Pollinated Maize. In: Babalola O.A., Ogunkunle, A.O., Olayinka A., Fagbemi, A.A.,Agboola, A.A., Ewenzor, W.O. (Eds.). Proceedings of the 26th Annual Conference of the Soil Science Society of Nigeria, held at Ibadan Oyo State Oct., 30th to Nov., 3rd 2000. Pp. 136 – 139.
- Balasubramania, V. and Singh A. (1978).Organic Recycling in Asian Small Holder Farms in Semi-arid Eastern Kenya.*Experimental Agricultural*, 31: 371-381.
- Boateng, S.A, J.Zickermann and Kornaharen,M. (2006).Effect of Poultry Manure on Growth and Yield of Maize.*West African J. App. Eco.*, 9:1 – 11.
- Dauda, S.N., F.A. Ajayi and Ndor,E. (2008).Growth and yield of water melon (*Citrulluslanatus*) as affected by poultry manure application.*Journal of Agricultural Science Society*, 4: 121–124.

- Deksissa T., I. Short and Allen, J. (2008). Effect of soil amendment with compost on growth and water use efficiency of Amaranth. In: Proceedings of the UCOWR/NIWR annual conference: International water resources: challenges for the 21st century and water resources education, July 22 – 24, 2008, Durham, NC.
- EIARD, (European Initiative on Agricultural Research for Development) (2013). Improving small-holder incomes through intensification of upland rice. Published 2013.
- Esu, I.E. (1991). Detailed soil survey of Nigerian Institute for Agricultural Research Farm at Bunkure, Kano state, Nigeria. p.72.
- Ezeibekwe, I.O., C.I. Ogbonnaya and Onuoha, C.I. (2009). Comparative Effect of Poultry manure and Urea on the Growth and Yield of Maize (*Zea mays* L.) Report and Opinion, (4) <http://www.sciencepub.net/> report.
- FAO (2004). (Food and Agricultural Organization of the United Nations). *Rice is Life*, 2004
- Farhad .W, M. F. Saleem, M. A. Cheema and Hammad, H. M. (2009). Effect of poultry manure levels on the productivity of spring maize (*Zea mays* L.). *Journal of Animal and Plant Sciences*, 19(3): 122-125.
- Gomez, K.A. and Gomez, A.A. (1984). *Statistical Procedures for Agricultural Research*. (2nd edition), John Wiley Sons, New York. 680 p.
- IRRI, (International Rice Research Institute) (2001). Rice Statistics. Accessed Online at <http://oryza.com/africa/nigeria/index.shtml> on 27th March, 2014.
- Kwari, J. D. (2003). Growth and Yield of Millet after four years of application of cattle manure, millet residue and green manure in sandy loam soils of Maiduguri, Nigeria. In: Ojeniyi, S. O., Ano, A. O., D. O. Awasalam and Chukwu, G. O. (Eds.). *Land Degradation, Agricultural Productivity and Rural Poverty Implications*. Proceedings of the 28th Annual Conference of the Soil Science Society of Nigeria, held at the National Root Crops Research Institute, Umudike, Abia State. Nov. 4th – 7th 2003. P.45-57.
- IITA (International Institute of Tropical Agriculture) (1989). Selected methods of soil and plant analysis. *Manual Series No. 1*. IITA, Ibadan, Nigeria, 70p.
- Lombin, L. G., J. A. Adepetu and Ayoyatade, K.A. (1991). Complementary Use of Organic and Inorganic Fertilizers. In: Proceedings of National Organic Fertilizers Seminar, Kaduna pp. 146-162.
- Maerere, A.P., G.G. Kimbi and Nonga, D.L.M. (2001). Comparative Effectiveness of Animal Manures on Soil Chemical Properties, Yield and Root Growth of Amaranthus (*Amaranthus cruentus* L.). *African Journal of Science and Technology*, 1(4):14-21

- Moabe, S. M., N. L. Kidula and Ondicho, A. R. (2000). Effect of Green Manure Residue Management Practices on Maize Yield in Western Kenya. Proceedings of the second scientific conference of soil management and legume research network project, Mombasa, Kenya. Pp.1-8
- Mitchell, C. C and Tu, S. (2005). Long term evaluation of poultry litter as a source of nitrogen for cotton and corn. *Agronomy Journal*, 97: 399-407.
- Naeem, M., J. Iqbal and Bakhsh, M. A. A. (2006). Comparative study of inorganic fertilizers and organic manures on yield and yield components of Mungbean (*Vigna radiata* L.). *Journal of Agricultural Science Society*, 2: 227-229.
- Obi M. E. and Ebo, P. O. (1995). The effects of different application rates of organic and inorganic fertilizers on soil physical properties and maize production in a severely degraded ultisol in southern Nigeria. *Bio-resource Technology*, 51(2-3): 117-123.
- Ogunwole, J. O., J. O., Atabo, D. T., Taro, A. B., Lawal, and Alabi, S. O. (2005). Effect of cow dung and poultry litter on soil amendment and extraction of garlic and pepper as insecticide on cotton production in the Nigerian savanna. *Journal of Agronomy*, 4(4): 267-272.
- Ojanuga, A. G. (2005). Agro-ecological zone map of Nigeria. National special programme for food security, FAO-UNESCO, p. 24.
- Ojeniyi, S. O. and Adeniyi, N. O. (1999). Effect of poultry manure and NPK fertilizer on soil fertility, nutrient content and yield of maize at Akure, Southwestern Nigeria. In: Babalola O., U. Omoti and A. E. Isenmila (Eds.). *Management of the Soil Resources of Nigeria for Sustainable Agricultural Production in the 21st century*. Proceedings of the 21st Annual Conference of the Soil Science Society of Nigeria held at Benin, 21st-25th Nov, 1999. Pp.185-191.
- Okafor, M. O. (1999). Effect of liming and manuring an Ultisol with locally available material for aluminium reduction and maize grain production in Ishaagu, Southeastern Nigeria. In: Babalola O., U. Omoti and A. E. Isenmila (Eds.). *Management of the soil resources of Nigeria for sustainable agricultural production in the 21st century*. Proceedings of the 21st Annual Conference of the Soil Science Society of Nigeria held at Benin, 21st-25th Nov, 1999. pp. 179-184.
- Opera-Nadi, O. A., A. A. Omenihu and Ifemedebe, S. N. (2000). Effect of Organic Waste, Fertilizers and Mulch on Productivity of an Ultisol. In: Babalola, O., Ogunkunle, A. O., Olayinka, O., Fagbami, A., Agboola, A. A. and Ewenzor, W. O. (Eds.). Proceedings of the

- 26th Annual Conference of Soil Science Society of Nigeria held at Ibadan Oyo state Nigeria, October 30th- Nov, 3rd 2000. Oyo State.pp.112-120.
- Pierre, C. and Morrean, R. (1997). Soil Fertility and Fertilization of Tropical Crops. John Wiley and Sons, New York 415 p.
- SAS (2003). Statistical Analysis System. SAS released 9.1 for windows, SAS Institute Inc. Cary, N.C., USA.
- Singh, B.R (1995). Soil management strategies for the semi-arid states. Africa Ecosystem in Nigeria. The Case of Sokoto and Kebbi States. *African Soils*, 28: 317-320.
- Sobulo, R. A. and Osiname, O. A. (1985). Fertilizer use in the Tropics, Nigerian Experience. In: Sobulo, R. A. and Udo, E. J. (Eds.). Soil fertility, tilth and degradation in the Tropics. 231 pp.
- Suresh, K.D., G.Sneh, K.K. Krishnand Mool, C.M. (2004). Microbial biomass, carbon and microbial activities of soils receiving chemical fertilizers and organic amendments. *Archive of Agronomic Soil Science*, 50: 641–647.
- Tisdale, S.L., W.L. Nelson and Beaton, J.D. (2003). Soil Fertility and Fertilizers. 4th Edition Macmillan Publishing Company, New York, 61-203p
- Usman, A., O. A. Osunde, A. Bala and Ezenwa, M.I.S. (2007). Carbon and Nitrogen Mineralization of Some Selected Legume Species. In: Uyovbisere E. O., Raji, B. A., Yusuf, A. A., Ogunwale, O., Aliyu, L. and Ojeniyi, S. O. (Eds.). *Soil and Water Management for Poverty Alleviation and Sustainable Environment*. Proceedings of the 31st Annual Conference of the Soil Science Society of Nigeria, held at Ahmadu Bello University, Samaru Zaria Nov 13th-17th 2006. pp. 489-495.
- Warren J. G., S. B Phillips, G. L. Mullins, D. Keahey and Penn, C. J. (2006). Environmental and production consequences of using alum-amended poultry litter as a nutrient source for corn. *Journal of Environment and Quality*, 35: 172-182.
- Wild, A. (1988). *Soils Condition and Plant Growth*. Longman group, London 116 p.