

Differences in Schistosomiasis knowledge among irrigation rice farming communities in different irrigation schemes in Tanzania.

Farida S. Salehe^{1*}, Amon Z. Mattee², Andrew K.P.R.Tarimo³, Jeroen H.J. Ensink⁴ and Madundo M.A. Mtambo⁵

^{1*}Development Studies Institute, Sokoine University of Agriculture,
P.O BOX 3024 Morogoro, Tanzania

² Department of Agricultural Education and Extension, Sokoine University of Agriculture,
P.O Box 3002 Morogoro, Tanzania

³ Department of Agricultural Engineering and Land Use Planning, Sokoine University of
Agriculture, P.O BOX 3003, Morogoro, Tanzania

⁴ Department of Infectious and Tropical Diseases, London School of Hygiene and
Tropical Medicine, Keppel Street, WC1E 7HT, London, United Kingdom

⁵ Department of Veterinary Medicine and Public Health, Sokoine University of
Agriculture P.O Box 3021, Morogoro, Tanzania

ABSTRACT

Aim: To assess differences in schistosomiasis knowledge in farmers working in traditional, improved traditional and modern irrigation schemes in Tanzania.

Study design: A cross-sectional survey among farmers practicing irrigation rice farming, in 2 different regions and 6 different irrigation systems.

Place and Duration of Study: The study was carried out between November and December 2011 in the Morogoro and Kilimanjaro Regions, Tanzania.

Methodology: Random sampling with equal number of irrigators in each scheme was employed. Irrigators were chosen by the researcher from the list provided by the village government in the six schemes of all farmers practicing irrigation rice farming. 120 samples were used, 120 from each region, i.e. farmers practicing irrigation rice farming in each scheme. Independent sample t-test was used for testing schistosomiasis knowledge differences among irrigators between schemes with the same design

and construction of their infrastructures between the two regions.

Results: More than 88% of irrigators surveyed in Kilimanjaro schemes had better knowledge regarding to all schistosomiasis items asked compared to those in Morogoro. There were significant difference on irrigators knowledge on schistosomiasis symptoms ($P<0.001$), and predisposing factors ($P<0.001$) between Morogoro and Kilimanjaro Modern schemes. Knowledge on predisposing factors differed significantly ($P<0.001$) between irrigators in Morogoro and Kilimanjaro improved traditional schemes. Moreover significant difference were noted on irrigators knowledge on schistosomiasis symptoms ($P<0.001$), predisposing factors ($P<0.004$) and schistosomiasis control measures ($P=0.003$) between irrigators in Morogoro and Kilimanjaro traditional schemes.

Conclusion: Schistosomiasis knowledge was found to be lower in Morogoro schemes, and particularly in traditional scheme, indicating that water management education deliveries are needed for sustainable control of schistosomiasis in these schemes, which have found to be successful in other irrigation systems world-wide.

Keywords: schistosomiasis, knowledge, different irrigation schemes, Morogoro, Kilimanjaro, Tanzania.

1. INTRODUCTION

It is estimated that 207 million people world-wide suffer from schistosomiasis, and the majority of these live in sub-Saharan Africa (Steinmann et al., 2006). Moreover, some estimates indicate that as many as 400 million people are affected (King, 2010). Schistosomiasis is one of the major causes of morbidity in endemic areas of Asia, Africa and South America (WHO, 1995; Hotez et al., 2008). Professions with a high water contact pattern like: house wives in charge of collecting drinking water, fisher men and farmers are at a high risk of contracting the infection (Steinmann et al. 2006). In Tanzania the disease affects the irrigated rice growing communities in the Morogoro and Kilimanjaro regions and causes a high disease burden in both health of the irrigation farming communities and socio-economy of the country (URT, 2008).

Several factors influence schistosomiasis transmission levels in irrigation schemes, these include: the design of irrigation infrastructures farming practices, like poor drainage and vegetation cover in canals. Changes in the management of irrigation systems can help control the intermediate host. Increases in the flow velocity in irrigation canals, through the construction of irrigation canals with an elevated proper slope, is a potential way to disperse snails, though this should be balanced with possible increased erosion of irrigation canals (Oomen, de Wolf et al. 1990), and might not always be possible. The lining of irrigation canals, regular removal of weeds and improving drainage are however expensive.

Schistosomiasis is claimed to be primarily a disease that results from lack of education, public health facilities and poverty (Oniya, et al., 2007; King et al., 2010). Although chemotherapy can reduce morbidity caused by the infection, rapid re-infection usually occurs requiring repeated treatment (N'Goran et al.,

2001). Therefore, improvements in hygiene, water supply and sanitation combined with health promotion and sustainable behavior change will need to be put on the agenda in case long term control of schistosomiasis can be achieved in Tanzania (Hamed, 2010; WHO, 2012). Understanding the levels of knowledge regarding schistosomiasis transmission will enable policy makers to help design control programmes. Integrating health education and medical intervention remain a high priority for the World Health Organization's (WHO) schistosomiasis control programs. However, in many endemic areas, control programs often do not incorporate public health promotion of the target groups.

In Tanzania there have been several studies assessing community perceptions and knowledge about schistosomiasis, though most of these studies focused on school children (Mazigo *et al.*, 2010; Clement *et al.*, 2008; Freudenthal *et al.*, 2006; Lansdown *et al.*, 2001; Lwambo *et al.*, 1999; Guyatt *et al.*, 1999), with the exception of one, which involved irrigated rice growers (Poggensee *et al.*, 2000). The level of knowledge demonstrated by school children does not necessarily represent the level of knowledge of their parents, nor those of specific communities at risk like farmers and fishermen and neither is it necessarily comparable between traditional, improved traditional and modern irrigation schemes. In this case, there is no certainty on this subject matter. Therefore, this study sought to assess schistosomiasis knowledge differences among the farming communities in terms of the level of knowledge related to schistosomiasis cause, symptoms, transmission, control and health-effects that may occur due to schistosomiasis infection between schemes in Morogoro and Kilimanjaro regions. Results were envisioned to contribute to the identification of knowledge gaps on schistosomiasis among irrigation rice farming communities thus provide entry/departure points towards future schistosomiasis control efforts in Tanzania.

2. MATERIALS AND METHODS

2.1 Study Areas

Six irrigation schemes which differed in terms of design and construction of their irrigation infrastructures were purposively selected from two Regions, Morogoro and Kilimanjaro. Three irrigation schemes in each region were selected. Two irrigation systems: Lower Moshi and Mwega schemes were categorized as modern because their intakes, main canals, secondary and tertiary canals, division and drop structures are concrete lined. The schemes Kikafu Chini and Mkindo were categorised as improved traditional systems because only the intakes, and part of their main canals are concrete lined, while the rest of the

infrastructure are earth lined. The final two: Njoro and Chabi were categorised as traditional schemes as from the intake, main, secondary canal and other infrastructures are earth lined and constructed using indigenous knowledge.

Njoro scheme is located in Kaloleni Ward, which is found in Moshi Municipal- Tanzania. It is the smallest ward in Moshi town. The major activity done in njoro is rice farming. There are about 232 rice farmers engaged in rice farming. Kaloleni ward has a population of about 6000 people with majority being poor farmers.

In 1999, Kikavu Chini had a population of nearly 3000, an average sized village in Tanzanian terms. The irrigation scheme is managed by about 760 farmers, including 430 land owners, 230 tenants, and 100 laborers. Kikavu is quite small, with 1600 acres. The irrigation scheme dominates the landscape taking up 1000 acres out of which only 300 acres are suitable for rice farming (Koopman et al. 2001).

Lower Moshi scheme is located 15 kilometers in the southern part of Moshi Town at the foot of the slopes of Mount Kilimanjaro (3°21'S, 37°21'E). The scheme lies at altitude ranging between 700 to 800 meters above sea level.

The detailed description of Chabi, Mkindo and Mwega schemes are given in Salehe and Hassan (2012).

The six schemes were selected due to the reason that they are among the main rice growing areas in Morogoro and Kilimanjaro. Additionally, nearly all the scheme inhabitants in the two regions use water from the canals for cooking, drinking, washing clothes and household utensils and for bathing.

2.2 Study design and sampling procedure

A cross sectional research design was used and data collection was carried out in six irrigation schemes between November and December, 2011. Data were collected at a single point in time. Random sampling with equal number of irrigators in each scheme was employed. Irrigators were chosen by the researcher from the list provided by the village government in the six schemes of all farmers practicing irrigation rice farming. The researcher chose a total of 240 irrigation rice farmers for the study, 120 from each Region and 40 from each irrigation scheme. A structured questionnaire with closed and open ended questions

was used as a method of collecting data. Information on what causes schistosomiasis, how schistosomiasis can be transmitted, effects that can be caused by the disease, symptoms, schistosomiasis predisposing factors and schistosomiasis control measures was sought from each irrigator interviewed.

122

123 **2.3 Determination of schistosomiasis knowledge index**

124 For better assessment of schistosomiasis knowledge status, a nine item index was used (Table 1).
125 However, the items were divided into 6 Schistosomiasis knowledge domains: (i) causes of
126 schistosomiasis, (ii) effects of schistomiasis, (iii) symptoms of schistomiasis, (iv) schistomiasis
127 predisposing factors, (v) transmission of schistosomiasis and (vi) schistomiasis control measures. Where
128 a domain had more than one item, the items were combined into one index for the domain. Respondents
129 were asked to respond to whether they understood each item concerning schistosomiasis or not. The
130 respondents scored 1 for a correct answer and 0 for incorrect answer. Respondents' responses were
131 added to form a schistosomiasis knowledge index which ranged from 0 to 6.

132

133 **Table 1: Schistosomiasis knowledge index**

Knowledge Domains and Items
Domain 1: Causes of schistomiasis -Schistosomiasis is caused by snails
Domain 2: Effects of schistomiasis - Schistosomiasis deteriorate people's health - Schistosomiasis patient can die if not well treated
Domain 3: Symptoms of schistomiasis The following are not schistosomiasis symptoms; -Blood in urine, blood in stool, pains during urinating, pains during passing stool, pains below abdomen, abdominal pains
Domain 4: Schistomiasis predisposing factors - One can acquire schistosomiasis through washing limbs, bathing and washing farming tools in irrigation infrastructures.

Domain 5: Schistosomiasis transmission

- Urinating and defecating in the scheme areas transmits schistosomiasis

Domain 6: Schistosomiasis control measures

- Construction of toilets in the scheme areas does not control schistosomiasis
- Proper canal cleaning to remove vegetation growth does not control schistosomiasis
- Draining excessive water in the plots and canals does not control schistosomiasis

134

135 **2.4 Data analysis**

136 The number of respondents who answered correctly to each knowledge item asked was expressed in
137 percentage. Independent sample t-test was used for testing schistosomiasis knowledge differences
138 among irrigators between schemes with the same design and construction of their irrigation infrastructure
139 between the two regions.

140

141 **3.0 RESULTS AND DISCUSSION**

142 **3.1 Results**

143 **3.1.1 Respondents' knowledge on Schistosomiasis among irrigation rice growers in**
144 **Modern, Improved traditional and Traditional schemes in Morogoro Region**

145 Table 2 presents responses related to schistosomiasis knowledge among irrigators interviewed in Modern
146 (Mwega), Improved traditional (Mkindo) and Traditional (Chabi) schemes in Morogoro. 90% of irrigators in
147 all three schemes had enough knowledge about schistosomiasis as a disease caused by snails. In
148 addition, 93% of irrigators were knowledgeable on the effects that can be caused by schistosomiasis.
149 Regarding irrigators' knowledge on schistosomiasis symptoms, only 58% in all three schemes had
150 knowledge on that item. Moreover, less than half (44.2%) had knowledge on schistosomiasis
151 predisposing factors. However, 81% of irrigators interviewed in the three schemes were aware of
152 schistosomiasis control measures and 92% had knowledge on how schistosomiasis can be transmitted.
153 Generally, irrigators in Morogoro traditional scheme (Chabi) had poor knowledge on schistosomiasis
154 symptoms and schistosomiasis predisposing factors while those in Morogoro improved traditional Scheme
155 (Mkindo) had also poor knowledge on schistosomiasis predisposing factors (Table 2).

156

157 **3.1.2 Respondents' knowledge on Schistosomiasis among irrigation rice growers in Modern,**
 158 **Improved traditional and Traditional schemes in Kilimanjaro Region**

159 Table 2 results reveal that more than 88% of irrigators interviewed in modern, improved traditional and
 160 traditional schemes in Kilimanjaro had better knowledge on what causes schistosomiasis, effects that can
 161 be caused by schistosomiasis, schistosomiasis predisposing factors, schistosomiasis symptoms, how the
 162 disease can be transmitted and controlled. Generally irrigators in Kilimanjaro traditional scheme (Njoro)
 163 had better schistosomiasis knowledge compared to their counterparts in Improved traditional and Modern.

164

165 **Table 2: Schistosomiasis Knowledge among surveyed irrigation rice growers in modern, improved**
 166 **traditional and traditional schemes in the two Regions.**

167

Variables	Morogoro schemes			Kilimanjaro schemes		
	Traditional (Chabi)	Improved tradition (Mkindo)	Modern (Mwega)	Traditional (Njoro)	Improved traditional (Kikavu chini)	Modern (Lower moshi)
Causes of schistomiasis	38 (95)	28 (70)	35 (87.5)	37 (92.5)	33 (82.5)	37(92.5)
Effects of schistomiasis	38 (95)	37 (92.5)	37 (92.5)	39 (97.5)	38 (95)	38 (95)
Symptoms of schistomiasis	13 (32.5)	33 (82.5)	24 (60)	40 (100)	36 (90)	40 (100)
Schist. predisposing factors:	14 (35)	18 (45)	21 (52.5)	40 (100)	38(95)	39(97.5)
Schistosomiasis transmission	38 (95)	33 (82.5)	39 (97.5)	39 (97.5)	37 (92.5)	39(97.5)

Schist. control measures	32 (80)	28 (70)	37 (92.5)	40 (100)	34 (85)	36 (90)
168	*Numbers in parenthesis refer to percentage of respondents that responded to schistosomiasis items.					
169						
170	<u>3.1.3 Schistosomiasis knowledge differences among irrigators in Modern irrigation schemes</u>					
171	<u>between Morogoro and Kilimanjaro Regions.</u>					
172						
173	An independent sample t-test indicates that there were significant differences on schistosomiasis					
174	symptoms and schistosomiasis predisposing factors among irrigator's between Morogoro modern scheme					
175	(Mwega) and Kilimanjaro modern scheme (Lowermoshi). Furthermore, results show that irrigators in					
176	Lower Moshi had more knowledge on schistosomiasis symptoms compared to those in Mwega (Table 3).					
177	Moreover irrigators in Lower Moshi had better understanding on schistosomiasis predisposing factors					
178	compared to their counterparts in Mwega.					
179						
180	<u>3.1.4 Schistosomiasis knowledge differences among irrigators in Improved traditional irrigation</u>					
181	<u>schemes between Morogoro and Kilimanjaro Regions.</u>					
182	Table 3 revealed significant difference on schistosomiasis predisposing factors among irrigators between					
183	Morogoro Improved traditional scheme (Mkindo) and Kilimanjaro Improved traditional scheme (Kikafu					
184	Chini). Irrigators in Kikafu Chini scheme had more schistosomiasis knowledge on the above mentioned					
185	schistosomiasis item compared to those in Mkindo scheme.					
186						
187	<u>3.1.5 Schistosomiasis knowledge differences among irrigators in Traditional irrigation schemes</u>					
188	<u>between Morogoro and Kilimanjaro Regions.</u>					
189	Table 3 results show significant differences on three schistosomiasis knowledge items between irrigators					
190	interviewed in Morogoro traditional scheme (Chabi) and those in the Kilimanjaro traditional scheme					
191	(Njoro). Irrigators' knowledge on schistosomiasis symptoms, schistosomiasis control measures and					
192	schistosomiasis predisposing factors differed significantly between the two schemes. Schistosomiasis					
193	knowledge on the mentioned items was widely understood by irrigators in Njoro than those in Chabi					
194	scheme (Table 5).					
195						

196 **Table 3: Schistosomiasis knowledge difference among irrigators in Modern, improved traditional**
197 **and traditional irrigation schemes between Morogoro and Kilimanjaro regions**

Variables	Modern Schemes			Improved Traditional Schemes			Traditional Schemes		
	Mwega	Lower	P-Value	Mkindo	Kikafu	P-Value	Chabi	Njoro	P-Value
		Moshi			Chini				
Causes of schistosomiasis	0.88	0.92	0.46	0.70	0.82	0.19	0.95	0.92	0.65
Effects of schistosomiasis	0.92	0.95	0.65	0.92	0.95	0.65	0.95	0.98	0.56
Symptoms of schistosomiasis	0.60	1.00	<0.001**	0.82	0.90	0.34	0.32	1.00	<0.001**
Schistosomiasis transmission	0.98	0.98	1.00	0.82	0.92	0.18	0.95	0.98	0.56
Schistosomiasis predisposing factors	0.52	0.98	< 0.001**	0.45	0.95	<0.001**	0.35	1.00	0.004*
Schistosomiasis control	0.92	0.90	0.69	0.70	0.85	0.11	0.80	1.00	0.003*

198 *Mean difference statistically significant at P<0.05 level

199 ** Mean difference statistically significant at <0.001 level

200

201 **4. DISCUSSION**

202 Schistosomiasis is one of the most widespread parasitic infections that affect humans in terms of socio-
203 economic and public health importance in the developing world. Yet majority of the people in these
204 countries are not aware of its symptoms and predisposing factors as highlighted in this study. Among
205 irrigators interviewed in the three schemes in Morogoro, those in traditional scheme (Chabi) had poor
206 knowledge on schistosomiasis symptoms and predisposing factors while those in Improved traditional
207 (Mkindo) scheme had low knowledge on schistosomiasis predisposing factors only. This might be due to
208 the reason that respondents in Chabi (Traditional) irrigation scheme had never received any training
209 regarding schistosomiasis as a disease since the village does not even have health facilities such as
210 dispensary. Therefore, our results tally with the study in Nigeria by Oniya (2007), that schistosomiasis
211 results from lack of education and public health facilities. The findings are also in line with the results in

212 Nigeria by Onyeneho *et al* (2010), that lack of knowledge is the main factor responsible for the high level
213 of schistosomiasis.

214

215 Rice irrigators in Mkindo scheme had low knowledge on schistosomiasis predisposing factors despite
216 existing evidence on provision of training on schistosomiasis from several institutions and the Mkindo
217 Farmer's Centre. Such training programmes include those offered by the Japan International Cooperation
218 Agency (JICA) whereby farmers in Morogoro schemes particularly Mwega and Mkindo schemes visited
219 Lower Moshi scheme for training on rice production and water management practices. The findings
220 correspond with the study done in Ghana by Wagatsuma *et al.* (2003), that children who had more
221 awareness about urinary schistosomiasis raised the question why awareness did not result in low
222 prevalence and avoidance of risk behaviour.

223

224 Irrigators interviewed in Modern, improved traditional and traditional schemes in Kilimanjaro had better
225 schistosomiasis knowledge compared to their fellow irrigators in Morogoro schemes. This might be
226 attributed to high literacy rate by Kilimanjaro farmers. This is supported by the URT (2005) report, that
227 Moshi Urban and Moshi Rural are known to have higher percentage of adult literacy rate of 96% and 89%
228 respectively compared to Morogoro Urban 85%. Njoro farmers' schistosomiasis knowledge was high may
229 be because they cultivate rice twice a year and therefore due to frequent contact with water they have
230 become more aware of the schistosomiasis symptoms. Moreover, multiple schistosomiasis infection by
231 farmers in Kilimanjaro traditional scheme (Njoro) is probably another reason which might have made them
232 understand better schistosomiasis symptoms compared to their counterparts in Morogoro traditional
233 scheme (Chabi).

234

235

236 **5. CONCLUSION AND RECOMMENDATION**

237 Generally schistosomiasis knowledge level was lower among irrigators interviewed in Morogoro schemes
238 particularly in traditional scheme (Chabi) and higher among respondents in Kilimanjaro schemes. From the
239 findings of this study it is concluded that schistosomiasis knowledge delivery is needed for sustainable
240 control of schistosomiasis in Morogoro schemes as this strategy has been successful in Niger and
241 Senegal.

242

243 Deliberate efforts are needed on the link between schistosomiasis and irrigation farming in modern,
244 improved traditional and traditional schemes in Morogoro and Kilimanjaro. Farmers' awareness and
245 provision of education on schistosomiasis, and how it relates with water need to be taken into serious
246 consideration in designing future schistosomiasis mitigation/control measures in all old and new irrigation
247 schemes.

248

249 **ACKNOWLEDGEMENTS**

250 Authors thank Hussein A, Swai G. and Msimbe L. for their assistance during data collection. Authors also
251 wish to express their appreciation to irrigators surveyed in all six schemes whose cooperation and
252 understanding was essential towards accomplishing this research. Funds for this work were provided by
253 the Tanzania Government through the Tanzania Commission for Science and Technology (COSTECH),
254 Dar Es Salaam -Tanzania.

255

256 **AUTHORS' CONTRIBUTIONS**

257 Farida S. Salehe designed the study, collected data, performed the statistical analysis and wrote the first
258 draft of the manuscript. Amon.Z. Mattee, Andrew K.P.R. Tarimo, Jeroen H.J. Ensink and Madundo M.A.
259 Mtambo gave comments. All authors read and approved the final manuscript.

260

261 **REFERENCES**

262 Clements ACA, Brooker S, Nyandindi U, Fenwick A, Blair L. Bayesian spatial analysis of a national
263 urinary schistosomiasis questionnaire to assist geographic targeting of schistosomiasis control in
264 Tanzania, East Africa . International Journal of Parasitology. 2008;38:401-415.
265 Engel D, Chitsulo L, Montresor A, Savioli L. The global epidemiological situation of schistosomiasis and
266 new approaches to control and research. Acta Tropica. 2002; 83:139-146.

267 Freudenthal S, Ahlberg BM, Mtweve S, Nyindo P, Poggensee G, Krantz I. School-based prevention of
 268 schistosomiasis: Initiating a participatory action research project in northern Tanzania. *Acta*
 269 *Tropica*. 2006;100: 79-87.

270 Gallup JL, Sachs JD. The economic burden of malaria. *The American Journal of Tropical Medicine and*
 271 *Hygiene*. 2001; 64: 85-96.

272 Garba A, Aboubacar A, Barkire A, Vera C, Sellin B, Chippaux J. Impact of health education campaigns on
 273 the control of urinary bilharziasis in Niger. *Cahiers Sante*. 2001;11: 35-42.
 274

275 Guyatt H, Brooker S, Lwambo NJS, Siza JE, Bundy DAP. The performance of school-based
 276 questionnaires of reported blood in urine in diagnosing schistosoma haematobium
 277 infection; patterns by age and sex. *Tropical medicine and International Health*. 1999; 4:751-
 278 757.

279 Gwathirira PR, Ndamba J, Nyazema, NZ. The impact of health education on the knowledge, attitudes,
 280 practices of a rural community with regard to schistosomiasis control using plant
 281 *molluscicide phytolacca dodecandra*. *Central African Journal of Medicine*. 1999; 45: 94-97.

282 Hamed MA. Strategic Control of Schistosome Intermediate Host. *Asian Journal of Epidemiology*. 2010;
 283 3: 123-140.

284 Hotez PJ, Bethony JM, Oliveira SC, Brindley PJ, Loukas A. Multivalent anthelmintic vaccine to prevent
 285 worm and schistosomiasis. *Expert Review of Vaccines*. 2008; 7:745-752.

286 King CH. Parasites and poverty: the case of schistosomiasis. *Acta Trop*. 2010; 113: 95–104.
 287

288 Kloos H, Ouma JH, Kariuki HC, Butterworth AE. Knowledge, perceptions and health behavior pertaining
 289 to schistosomiasis mansoni related illness in Machakos district, Kenya. *Tropical Medicine and*
 290 *Parasitology*. 1986; 37:171-175.

291 Koopman J, Kweka R, Mboya M, Wangwe S. Community Participation in Traditional Irrigation Scheme
 292 Rehabilitation Projects in Tanzania: Report of a Collaborative Research Project; 2001.

293 Lansdown R, Ledward A, Hall A, Issae W, Yona E, Matulu J, Mweta M, Kihamia C, Nyandindi U, Bundy
 294 D. Schistosomiasis, helminth infection and health education in Tanzania: achieving
 295 behavior change in primary schools. *Health Education Research*. 2001;17:425-433.

296 Lwambo NJS, Siza JE, Brooker S, Bundy DAP, Guyatt H. Patterns of concurrent infection with hookworm
 297 and schistosomiasis in school children in Tanzania. Transactions of the Royal Society of Tropical
 298 Medicine and Hygiene. 1999; 93:497–502.

299 Mazigo H, Waihenya R, Mkoji GM, Zinga M, Ambrose EE, Jahanpour OF, Bahemana E, Mnyone LL,
 300 Kweka EJ, Lwambo NJS. Intestinal Schistosomiasis: Prevalence, Knowledge, Attitude
 301 and Practices among School Children in an Endemic Area Of Norther Western Tanzania.
 302 Journal of Rural and Tropical Public Health. 2010; 9: 53 – 60.

303 Oniya MO. Socio-Cultural Practices Promoting the Transmission of Urinary Schistosomiasis among
 304 School Aged Pupils in a South Western Village in Nigeria. Research Journal of Biological
 305 Sciences. 2007; 2: 1- 4

306 Onyeneho NG, Yinkore P, Egwuage J, Emukah E. Perceptions, attitudes and practices on
 307 schistosomiasis in Delta State, Nigeria. Tanzania Journal of Health Research. 2010;12.

308 Poggensee G, Krantz I, Kiwelu I, Feldmeier H. Screening of Tanzanian women of childbearing age for
 309 urinary schistosomiasis: validity of urine reagent strip readings and self-reported symptoms.
 310 Bulletin of the World Health Organisation. 2000;78(4).

311 Salehe FS, Hassan SN. Socio-economic Effects of Schistosomiasis on Irrigation Rice Growers in
 312 Morogoro, Tanzania. American Journal of Experimental Agriculture. 2012; 2(3): 395-406.

313 Steinmann P, Keiser J, Bos R, Tanner M, Utzinger J. Schistosomiasis and water resources development:
 314 systematic review, meta-analysis, and estimates of people at risk. Lancet Infect Dis. 2006; 6: 411–
 315 25.

316 Tesfa-Michael T, Ayele T, Birrie H, Lo CT. Evaluation of Multidisciplinary control of schistosomiasis
 317 mansoni in a northern Ethiopian community. Institute of Pathobiology Research Report. 1987; 3:
 318 17-29.

319 United Republic of Tanzania. The National Schistosomiasis and Soil-transmitted helminth Control
 320 Programme (NSSCP). 2008. Dar es Salaam, Tanzania.

321 United Republic of Tanzania. Poverty at the District Level in mainland Tanzania, Poverty and Human
 322 Development Report. 2005.

323 Wagatsuma Y, Aryeetey MF, Nkrumah FK, Sack DA, Kojima S. Highly symptom-aware children were
 324 heavily infected with urinary schistosomiasis in southern Ghana. Central African Journal of
 325 Medicine. 2003; 49:16-19.

326 World Health Organisation. Schistosomiasis Fact Sheet N°115. 2012. Available online
327 <http://www.who.int/mediacentre/factsheets/fs115/en/index.html>

328 World Health Organisation. A picture of health? 1995. WHO, Geneva/UNICEF, New York

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

UNDER PEER REVIEW