

Research Paper

Rv2485c, a putative lipase of *M. tuberculosis*: expression, purification and biochemical characterization

ABSTRACT

Lipases have been demonstrated to have a role in virulence in several pathogens. Rv2485c gene product of *Mycobacterium tuberculosis* has been annotated as putative carboxyl esterase (LipQ) involved in cellular metabolism and respiration. The Rv2485c gene was cloned and expressed in *E. coli*. The LipQ enzyme was purified as a His-tagged protein from inclusion bodies and refolded with 37% protein yield. The specific activity of purified enzyme was calculated to be 93 U/mg with pNP-Palmitate as a preferred substrate. It showed optimum enzyme activity in the range of 40-50°C and pH 8.0. The Ser-249, Asp-344 and His-377, predicted as the member of the catalytic triad, were confirmed by site directed mutagenesis. The enzyme was inhibited in the presence of PMSF and DEPC while no effect on enzyme activity was observed with eserine. The apparent K_m and V_{max} were calculated to be 1.45 mM & 196.08 U/ml respectively. The turnover number (k_{cat}) of the enzyme was calculated to be 6.597 min⁻¹. The gene was expressed only in oxidative stress condition in *in vitro* culture of *M. tuberculosis* H37Ra as shown by Real Time PCR.

Keywords: *Mycobacterium tuberculosis*, lipase, pNP-palmitate, Real Time PCR, oxidative stress.

28

29

30

31 **Introduction**

32 *Mycobacterium tuberculosis* is an intracellular pathogen infecting primarily mononuclear
33 phagocytes. The bacterium has developed intricate strategies to evade killing mechanisms of
34 phagocytes[1]. A key to fight this disease is the identification and characterization of all these
35 different strategies[2]. Lipolytic enzymes demonstrated several effects on macrophages i.e.
36 inhibition of phagocytic functions well as modulation of the release of inflammatory
37 mediators[3].

38 Certain virulence factors are necessary for full pathogenicity regardless of the host. The
39 lipolytic enzymes are one of the known virulence factors in many bacteria such as
40 *Pseudomonas cepacia*, *Staphylococcus aureus*[4],[5] and in fungal species like *Alternaria*
41 *brassicicola*, *Candida albicans*, and *Fusarium graminearum*. [6],[7],[8] Many different
42 bacterial species produce lipases, which hydrolyze esters of glycerol with preferably long-
43 chain fatty acids. They act at the interface generated by a hydrophobic lipid substrate in a
44 hydrophilic aqueous medium.[9] Despite lipolytic activities being described in mycobacteria
45 many years ago, the associated enzymes, lipases/esterases and phospholipases, have only
46 been the focus of intense research for the last few years. Because these enzymes are shared
47 by several mycobacterial species, they represent a huge field of exploration not only for
48 tuberculosis but also for other mycobacterial diseases[10]. Recently reported immunogenicity
49 of mycobacterial lipases prompted its use as biomarkers for active tuberculosis. On the other
50 hand these enzymes are reported to be involved in the degradation of host lipids to provide
51 fatty acids to pathogens[11].

52 Lipolytic enzymes may hence be considered as potent therapeutic targets for developing
53 future anti-tuberculosis drugs. *M. tuberculosis* expresses a plethora of lipolytic enzymes, of
54 which only a limited number have been studied so far and much remains to be learnt about
55 the diversity of their biological functions. It has been shown that during infection, *M.*
56 *tuberculosis* accumulates intracellular lipid inclusions (ILI) that are probably derived from
57 lipids of the host cell membrane. LipY has been found to be responsible for the utilization of

58 accumulated TG during dormancy and reactivation of the pathogen[12]. Deletion mutant of
59 LipY, with compromised ability to mobilize the stored TG, but not the complemented mutant,
60 was unable to come out of dormancy upon treatment with anti-TNF α mAbs[13]. The
61 overproduction of LipY in mycobacteria resulted in a significant reduction in the pool of
62 TAGs, consistent with the lipase activity of this enzyme[14]. The gene product of Rv3487c
63 (LipF) has been cloned and characterized, but no long-chain TG hydrolysis was detected[15].
64 On the basis of structural genomic approach, cloning and biochemical characterization of
65 Rv1399c gene product, LipH was carried out[16]. Its biochemical behavior univocally
66 classifies Rv1399c as a non-lipolytic rather than a lipolytic hydrolase. However, at present
67 the physiological roles for many of these lipases have not been completely deciphered. As it
68 is already described that in latent state, the lipids are the main source of energy to the starving
69 pathogen, therefore, all the lipid hydrolyzing enzymes of *M. tuberculosis* should be
70 characterized to ascertain their role in survival/pathogenicity/virulence.

71 LipQ enzyme was predicted to be involved in intermediary metabolism and demonstrated
72 66% similarity with its counterpart in non-pathogenic strain *M. smegmatis*, while no change
73 in gene sequence was noticed in *M. tuberculosis* H37Ra strain. The effect of Rv2485c gene
74 product on activation of murine macrophages was investigated in lab (unpublished results).
75 No cytotoxicity was observed after treatment of cells with LipQ protein. The LipQ protein
76 inhibited LPS induced nitric oxide (NO) production by macrophages. Decreased NO
77 production could be directly correlated with inhibition of iNOS mRNA expression in murine
78 macrophage (unpublished results). These observations suggested the probable role of this
79 enzyme in survival of mycobacteria in the mature granulomas during the dormant phase.
80 Therefore in the present investigation Rv2485c gene from *M. tuberculosis* H37Rv was cloned
81 and expressed in *E. coli*. The detail biochemical characterization of LipQ protein was carried
82 out. The expression of Rv2485c gene was checked in normal *in vitro* culture of *M.*
83 *tuberculosis* H37Ra in different phases of growth and under various stress conditions.

84

85

86 **Materials and Methods**

87 Databases and software

88 The DNA sequences of the Rv2485c gene was retrieved from the NCBI GenBank. Online
89 BLAST (<http://blast.ncbi.nlm.nih.gov>) was employed for analyzing the homologous identity
90 of the Rv2485c of *M. tuberculosis* with other mycobacterial species[17]. The identical protein
91 sequences of the Rv2485c in different species were aligned with Clustal X1.83
92 (<http://www.clustal.org/>) and all the positions containing gaps were eliminated[18]. By using
93 software Phyre[19] (Imperial college, London), secondary structure was predicted. The *M.*
94 *tuberculosis* H37Rv chromosomal DNA was a kind gift from Dr V. M. Katoch, JALMA,
95 Agra, UP, India

96 Cloning of Rv2485c gene in *Escherichia coli*

97 The Rv2485c coding region from *M. tuberculosis* H37Rv chromosomal DNA was amplified
98 using the primers GSF2 and GSR2 (Table 1).

99 Table 1: Primer sequence used to clone the Rv2485c gene

Primer Name	Sequence
GSF2	5'-tgcacatcgccagcgtgactt-3'
GSR2	5'-gaacggccaagctcagct -3'

100

101 The thermocycler (Perkin Elmer Cetus, Foster City, CA, U.S.A.) was programmed for a hot
102 start of 94⁰C for 4 min followed by 30 cycles at 94⁰C for 30 sec, 60⁰C for 40 sec and 72⁰C for
103 2 min and a final cycle of 7 min at 72⁰C. The amplified and purified DNA was ligated to the
104 pQE30-UA vector (Qiagen, MD, USA) to generate construct pQE-LipQ. *E. coli* M15
105 (PREP4) was used for the transformation and expression of Rv2485c. The transferred *E. coli*
106 strain was grown in Luria-Bertani (LB) medium with kanamycin (30 µg/ml) and ampicillin
107 (100 µg/ml). The presence of Rv2485c gene was confirmed by plasmid isolation[20], colony
108 PCR and sequencing.

109 Expression and Purification of LipQ

110 For shake flask growth, 1 liter of Luria broth medium supplemented with kanamycin (30
111 µg/ml) and ampicillin (100 µg/ml) was inoculated with 10 ml of an overnight culture and

incubated at 37°C. The culture was grown upto midlog phase ($A_{600} \sim 0.6$), induced with 0.5 mM isopropyl thio- β -D- galactosidase (IPTG), and further incubated for 3 h at 37°C.

Cells were harvested at 4°C by centrifugation at 9000 g for 15 min and resuspended in cold lysis buffer (50 mM TrisCl, pH 8.0, 0.1% (v/v) triton X-100, 150 mM NaCl and 1 mM EDTA) at 5 ml/g wet weight and stored overnight at -80°C. The cell suspension was thawed on ice for 1 h and 20 mM MgSO₄ (final concentration) was added. Cells were disrupted by ultrasonication (10× with a 15 s cycle) using a Pharmacia Biotech Sonifier. Cell debris were separated from the cell extract by centrifugation at 14 000 g for 20 min. The pellet was solubilized by stirring at 4°C overnight in a 40 ml solution containing 10 mM TrisCl, pH 8.0, 100 mM sodium dihydrogen phosphate and 8 M urea. The inclusion bodies were separated from solubilized protein by centrifugation at 14 000 g for 20 min at 4°C. Recombinant LipQ was purified from solubilized inclusion bodies by metal affinity chromatography under denaturing conditions using a nickel nitrilotriacetic acid (NiNTA) column as described by the manufacturer (Qiagen). Solubilized inclusion bodies were loaded on a nickel nitrilotriacetic acid column previously equilibrated with equilibration buffer (10 mM TrisCl, pH 8.0, 100 mM sodium dihydrogen phosphate and 8 M urea). After that protein loading column was washed with equilibration buffer at pH 6.3, and bound protein was eluted using a pH gradient (pH 6.3 - 4.3) of equilibration buffer. The fractions of eluted peak containing purified LipQ were checked for presence of protein after dilutions by measuring absorption at 280 nm with prepared blank. The protein was also analyzed by SDS-PAGE[20] on a 12.5% gel that was subsequently stained with Coomassie brilliant blue.

Refolding LipQ by dialysis

Refolding was carried out with chaotropic agent for concentration gradient dialysis. The solution of denatured protein (1.5 mg/ml) was dialyzed against 2 L of freshly made 6, 4, 2, 1, 0.5, and 0 M urea, respectively, in 10 mM sodium phosphate buffer (pH 8.0). With each concentration, the protein was dialyzed 6 h at 4°C. Protein concentration was determined by Protein estimation kit (BCA method) procured from Bangalore GeNei.

Lipase assay

The enzyme activity was determined according to the modified method of Sigurgisladottir *et al.* 1993 [21] Hundred microliter enzyme (100 mg/ml) and 0.1 ml of 0.002 M pNP-palmitate was added in 0.8 ml of 0.05 M phosphate buffer (pH 8.0). The reaction was carried out at

37⁰C for 15 min, after which 0.25 ml 0.1 M Na₂CO₃ was added. The mixture was centrifuged and the activity was determined by measuring absorbance at 420 nm. One unit of enzyme activity is defined as the amount of enzyme, which liberates 1 μmole of *p*-nitrophenol from pNP-ester as substrate per min under standard assay conditions.

Biochemical properties of Rv2485c

Effect of pH and Temperature on enzyme activity/stability

Lipase activity was assayed in the buffers of different pH (2.0-10.5) at 37⁰C. The following buffers (50 mM) were used: Glycine/Hydrochloric acid, pH 2.0-3.0; acetate buffer, pH 4.0–5.5; phosphate buffer, pH 6.0–8.0; Tris/Cl buffer, pH 8.0–9.0 and carbonate bicarbonate buffer, pH 10.0–10.5. The pH stability of the lipase was determined by incubating the enzyme at different pH for 1 h at room temperature. The residual lipase activity was determined by standard assay method at pH 8.0.

The effect of temperature on enzyme activity was determined by carrying out the enzyme assay at different temperatures (20-90⁰C). Thermal stability of LipQ was determined by incubating enzyme at different temperatures (20-90⁰C) for 30 min and pH 8.0 followed by cooling of enzyme on ice for 15 min. Enzyme activity was measured as per standard assay conditions.

Substrate specificity

The substrate specificity of the purified recombinant protein was measured using pNP esters (Sigma, USA) with carbon chain length ranging from C₄ to C₁₈. The released pNP from the substrates *p*-nitrophenyl butyrate (pNPB; C₄), *p*-nitrophenyl-Decanoate (pNPD; C₁₀), *p*-nitrophenyl laurate (pNPL; C₁₂), *p*-nitrophenyl myristate (pNPM; C₁₄), *p*-nitrophenyl palmitate (pNPP; C₁₆) and *p*-nitrophenyl stearate (pNPS; C₁₈) were monitored using spectrophotometric assay.

Enzyme Kinetics

The effect of pNP-palmitate concentration (0.1- 5 mM) on the reaction rate of the enzyme was measured by using standard enzyme assay. The Michaelis–Menten constant (K_m) and

171 maximum velocity for the reaction ($V_{\max}$) were determined by Lineweaver–Burk plot and k_{cat}
172 (turnover number) & k_{cat}/K_m (specific constant) were calculated.

173 **Site-directed mutagenesis**

174 To confirm the predicted active site residues of this enzyme, site-directed mutagenesis of Ser-
175 249, Asp-289, Asp-344 and His-377 of LipQ was carried out using a pair of complementary
176 primers (Table 2) containing the desired mutations by a non-strand-displacing, high-fidelity
177 polymerase (New England Biolabs) in a linear amplification.

178 Table 2: Primer sequence used to clone site directed mutants of the Rv2485c gene

Primer Name	Sequence
t745g	5'-caccggcggggcccgggtgc-3'
t745g_antisense	5'-gcaccggcgccccgggtg-3'
a866c	5'-tacggcgtctacgccctaccaacgcc-3'
a866c_antisense	5'-ggcgttggtgaggcgtagacgccga-3'
a1031c	5'-tgttgacacggtgagaaggctccgatggtgc-3'
a1031c_antisense	5'-gcaccatcgagccttctaccgtgcaaca-3'
a1130c	5'-ccaacgcccaccccggttcgacct-3'
a1130c_antisense	5'-aggtcgacgcgggggtggcggttg-3'

179

180 The clones were sequenced to confirm the mutation. The mutant enzymes were expressed,
181 purified and refolded along with the native protein. The activity of enzyme was determined as
182 per standard assay method.

183 **Effect of various additives on enzyme activity:**

184 The effect of various lipase /esterase inhibitors phenylmethanesulfonylfluoride (PMSF),
185 chemical modifier diethylpyrocarbonate (DEPC) and beta mercaptoethanol (β -ME) (0.1 and
186 1.0 mM) were studied with reference to enzyme activity of LipQ. The enzyme and additives
187 were incubated at room temperature for 5 min prior to assay. The enzyme without any

additive served as 100%. Additionally in the case of PMSF and DEPC (1 mM concentration), the enzyme and PMSF or DEPC was incubated at 50°C for 5 min prior to assay. Then enzyme assay was performed according to standard assay method.

Growth of *M. tuberculosis* H37Ra

M. tuberculosis H37Ra culture was grown in middlebrook 7H9 broth base (Himedia) supplemented with 1 % glycerol and 0.05 % Tween-20. An additional 1 % (v/v) growth supplement OADC (BBL) was added. The cells were grown for 7 days upto mid log phase (A_{600} 0.7-0.8) and harvested by centrifugation. The bacterial pellet was resuspended in different media for exposure to stress conditions such as oxidative stress (7H9 growth media with 5 mM H_2O_2), acidic stress (7H9 growth media at pH 4.5) and nutrient stress (1X phosphate buffer saline). These three stressed cultures were grown for 6 h , alongwith control culture in middlebrook 7H9 broth base (normal pH, without any additive). Samples were also taken out at mid log phase and stationary phase(A_{600} 1.8-2). RNA was isolated from all the samples using TRI reagent from Sigma as per company's manual instructions.

Rv2485c expression analysis

The RNA samples were converted into complementary DNA (cDNA) using RevertAid™ cDNA kit (Fermentas). This cDNA was used as a template to check the relative expression of the Rv2485c gene. 16S RNA was used as an internal control (Table 3). Expression pattern of Rv2485c gene was analyzed in acidic, nutritional, and oxidative stress and mid log and stationary phase in *in vitro* culture of *M. tuberculosis* H37Ra using short primers, sGSF2 as forward and sGSR2 as reverse primer (Table 3). The specificity of the reactions was verified by melting curve analysis. Determination of expression level of the gene was carried out by using Real-Time Quantitative PCR and the $2^{-\Delta\Delta Ct}$ Method [22]. Ct values for each run were used to calculate mean Ct with SD (Standard Deviation). For each sample: ΔCt test= Ct test- Ct 16S rRNA, ΔCt control= Ct control- Ct 16S rRNA. $\Delta\Delta Ct$ values relative to control were calculated by following formula: $\Delta\Delta Ct = \Delta Ct$ test – ΔCt control. The fold change in test gene expression was determined by taking 2 to the power of $\Delta\Delta Ct$ values i.e $2^{-\Delta\Delta Ct}$.

219 Table 3: Short Rv2485c gene primer sequence and 16s RNA primer sequence used in the
220 expression analysis.

Primer Name	Sequence
sGSF2	5'- acctccgacatcccgtac-3'
sGSR2	5'- ttgtaccgccgtgtcggc- 3'
16sRNA fwd	5'-gaggaagggtggggatgacgt-3'
16sRNA rev	5'-aggcccggaacgtattcac-3'

221

222

223 Results

224 All the values represented are mean $\pm$ SD of three independent experiments carried out in
225 triplicate. The DNA sequences of the Rv2485c gene was compared with homologous
226 sequences of other species of mycobacteria. It showed least identity with β -lactamase of non-
227 virulent strain, *M. smegmatis* (Table 4).

228 Table 4. Percentage identity of Rv2485c nucleotide sequence with other species of
229 *Mycobacterium* by using BLAST.

Gene	<i>M.</i>	<i>M. marinum</i>	<i>M.</i>	<i>M.</i>	<i>M. bovis</i>
Product	<i>smegmatis</i>		<i>leprae</i>	<i>abscessus</i>	
Rv2485c	66% with	76% with LipQ	% with	-----	100% with putative
(LipQ)	peptidase		ML1243		carboxylesterase

230


231 Cloning of Rv2485c

232 Rv2485c gene of *M. tuberculosis* H37Rv, with its original start codon was amplified using
233 chromosomal DNA as template. The amplified product corresponded with the expected size

(1266 bp) of Rv2485c gene of *M. tuberculosis* H37Rv. The amplified fragments were purified and ligated into pQE30-UA vector and transformed into competent *E. coli* M15 cells carrying lacI plasmid pREP4. Transformants were confirmed for the presence of Rv2485c insert by colony PCR. The sequencing results showed 100% homology with Rv2485c nucleotide sequence present in the complete genome sequence of *M. tuberculosis* H37Rv[23]

Purification of LipQ

The enzyme could not be expressed as soluble protein by varying the induction conditions. Purification of the urea solubilized enzyme, using affinity chromatography on Ni-NTA column as major method, yielded purified enzyme with 37% protein yield. Approximately 6 mg of pure and active recombinant enzyme was recovered from 1 liter of culture. The protein migrated as a single band on SDS-PAGE with an approximate molecular mass of 47.35 kDa (Fig.1). The molecular mass was little higher than the calculated mass from amino acid sequence which agreed with the calculated molecular mass obtained from the sequence data plus the mass of the amino terminus-appended (His)₆ tag (2.1 kDa).



248

Fig. 1. SDS-PAGE analysis of purified LipQ. Protein samples were loaded on 12% SDS polyacrylamide gel under reducing conditions. Lane 1 - Lysate, Lane 2- purified LipQ and Lane 3- Protein Molecular Weight Marker.

252

Bioinformatics analysis

The theoretical *pI* of the LipQ protein was calculated to be 8.98 by using online tool ScanSite *pI*/Mw. The molecular weight of the protein was predicted to be 45249.5 daltons. The protein was rich in hydrophobic amino acids (31%). It has 49.69% negatively charge amino acids and

257 31.15% positive charged amino acids and net charge at pH 7.0 was calculated to be 6.29. The
258 secondary structure (SOPMA) predicted revealed that the protein consisted of 47.51% α -
259 Helix, 10.45% extended strand, 4.51% β -turn and 37.53% random coils. Multiple sequence
260 alignment of LipQ from *M. tuberculosis* and five other bacteria including *Bacillus* shows all
261 these three active site amino acid residues are conserved (Supplementary Fig 1).

M.tuberculosis	-----MHIAVTSRCSRAGAEALRQGAQLAADARDTCRAGALLLRGSPCAIG--WVAG	51
M.smegmatis	-----MAKPLLT-----ASAEVLNAVNAVKPFGRKGYSTVVAFAGWPTSENGPLITT	48
Streptomyces	MAAALGGGSIGQVKPDFVTRRAAQLGLAALAVAPPKPRSTSIPVFFAGWLTGELAPHLLA	60
Rhodococcus	-----MSISNRFLLRQVATAGLAANALRPIGMPPLSVPTFLAGWLTAEAPQLLA	50
Clostridium	-----	
Bacillus	-----MKHTTSWRPRGIIQWILTILTLVAFPIIGIISYYIMNPDSMD--KLMS	46
M.tuberculosis	WLSAEFPARVVTGHALSRSISPRSIGRFGTSWAAQRADQILHAALVDAFGPDFRDLVWHPT	111
M.smegmatis	VSALDALRRAVRGDYRTTPGRISLLKAVAWALLYIVHRRNRTSRFFEEPLRSALGADY	108
Streptomyces	LTAADTVQHLARHGVRRPSDRVGLALAGLSAAGFASLIGTSLKARLEVEAALVEALGEDY	120
Rhodococcus	AAVVDSAIHVARRGVRTKADVGLAAAGLS TAGLAAVIAAGRGARVEEAALTSALGSDY	110
Clostridium	-----MDRNDAGREVSNREALMLDVEKVFETY	29
Bacillus	SLAWATALFPVSLIVTLFIIVLLILAFWKKAKIAQVILVPLSLLLIFLTVQPITSMLSY	106
M.tuberculosis	GEQS-----EAARRSGLLNLPHIPGPHRRYAAQTS DIPYGGGRENLLDIWRRPDLAPGR	166
M.smegmatis	AVAR-----RARRPGGVIRTSWAR---HRYATTT--VRYGP-HRANLADIWMRPDLPRDG	157
Streptomyces	KDHL-ESPDSDFSTPWGRLLMPFFMH--EPDVVRDRNVAYAPGGKRFLLDVYRPRETVRD-	176
Rhodococcus	RSAAGESVPPVGVPRQLALPFFMR--RDDVVRVRLAYAPGGRRFRIDVYHRRDIPTD-	167
Clostridium	LDIS-----YANESKHQKMDIIPNEGEGP-	54
Bacillus	ANSK-----DATVSLSSHEFFNSQTISTKPMENVVYGKTTD GIELKMDVWPAKETS KNK	159
* : :		
M.tuberculosis	RAPVLIQVPGGAWTINGKRPQAYPLMSRMVELGWICVS INYSKSPRCTWPAHIVDVKRAI	226
M.smegmatis	KAPVLLQVPGGAWMIGMRPQSYPLMSHLAEQGWICVSIGYRISPKHSWPDHIVDVKQAL	217
Streptomyces	-APVLLQIHGGAWTIGNKEQQGLPLMRHLARRGWVCVAINYPLSPASRWPAPHIVAQAQAL	235
Rhodococcus	-APVLLQVHGGGWVIGSKDHQGVPLMLEMASRGWVCAAVNYPLSPRAVWPAHLVAVKQAV	226
Clostridium	-FPVVINIHGGAWFFGDKRNVHTRSSIQALFKGYAVATINYLRLSGDAKWPQAIYDCKAAV	113
Bacillus	LKPAVVLVHGGGWVSGDKG-EAPHWKQWLNLDLGYTVFDVQYRMPPPQAGWKDEVADIKSAL	218
*.: : **.* : : * : * . * .: **:		
M.tuberculosis	AWVRENIADYGGDPDFITITGGSAGGHLAALAALSANDPA---LQPGFESADTAVQAAAP	283
M.smegmatis	AWVKANIADFGGDPESVCIITGGSAGGHLTALAALTPNDPK---WQPGFEDADTSVVAAP	274
Streptomyces	AWIREHIAEYGGDPRFVAVTGGSAGGHLAALLALSQNDPA---LQPGFEEADTSVQACVP	292
Rhodococcus	AWLRDNVSSYGGDPGFVAVTGGSAGGHLAAMLALTANEPG---LQPGFEDADTSIQACAP	283
Clostridium	RFLRANAAYNIKTDKIGVIGNSSGGHLAGVMAATNGCPEYENLSMGNSEYSSDVQAVAT	173
Bacillus	GWVLQNADTYQIDPNKINVMGESAGGNLAMLAAYSMGEEQ---LPASTNVPEVHVNSVIN	275
: : : : .. : : * * : * : : * : . . . : :		
M.tuberculosis	YYGVYDLTNAEN-MHEMMMPFLEHFVMSRYVDNPG---FKAASPISYVHSEAPFFVL	339
M.smegmatis	VYGRYDWYSTTGPRTEFMEILERLIVKRPLDGNAL---YRDASPIMLVRPDAPFFVL	331
Streptomyces	HYGVYDIAASTGEAESRYR--LESLMARRVFAPEHRL-DDYLAASPMDRVSEDAPFFVI	349
Rhodococcus	HYGVYDFTGETGIKATRLR--VESGLMPMVLGKKARFPEDYEASPLAHLRPDAPFFV	341
Clostridium	WYGIFDFINMEN-----HKKQLVKVGET-----Y	197
Bacillus	MYGPADMTMFYNDNPSTNY--VHGVMKEYIGGTVSQFPERYKLLSPINYIEDNTPPTITL	333
** *		
M.tuberculosis	HGEKDPMPVSAQSRAFSAALRDAGAATVS YAE L PNAHHAFDLAA-TVRSRMVAEAVSDFL	398
M.smegmatis	HGTNDSLIPVVEGRDFVAALREVSNSPVVYAEIPHAQHAFDFFG-SPHGHTADAVEKFL	390
Streptomyces	HGKHDSLVPVAEAREFVARLRAKSAHPVAYAE LAGAQHAFDVFP-SIRSTHVVRGVEHFL	408
Rhodococcus	HGVSDSFIPVAEAREFVRLREVSRNPVAYAE LRGQHAFDIFP-SIRSAAVVQGVADFL	400
Clostridium	KGEDDGVLDRLMG-----	210
Bacillus	LGTGDRIVPEQG-EMLDKELTVKNVAHEFYLLPDVDHGFDINPGSLSTQFAKEKVTAFL	392
* * .:		
M.tuberculosis	GVIYGRRMGARKGSLALSSPPAS	421
M.smegmatis	NWVLAKRGVAQNR-----	403
Streptomyces	DWAYAQAFTP---VNEKNS---	424
Rhodococcus	GWARAAQRGEGGHVREDAG----	419
Clostridium	-----	
Bacillus	QRYNK-----	397

Supplementary Figure 1. The amino acid sequence alignment of Rv2485c (LipQ) of *M. tuberculosis* H37Rv, esterase/lipase (*Streptomyces* sp. AA4), alpha/beta hydrolase (*Rhodococcus equi* 103S), alpha/beta hydrolase (*M. smegmatis*), alpha/beta hydrolase fold protein (*C. carboxidivorans*) and lipase (*Bacillus cereus*). Identical amino acids are indicated by asterisks and similar by dots. Active site residues are presented in a box. Serine is part of pentapeptide.

Biochemical characterization of Rv2485c

Substrate specificity

The enzyme was active on broad range of substrates (C₁₀-C₁₈). The maximum enzyme activity (100%) was observed with pNP palmitate. The specific activity of enzyme was calculated to be 93 U/mg with pNP-Palmitate as a preferred substrate. The result demonstrated the substrate choice for the recombinant lipase was in the order C₁₆>C₁₈>C₁₄>C₁₂>C₁₀>C₄ (Fig. 2).




Figure 2. Substrate specificity of recombinant LipQ. The reaction mix without enzyme served as blank. Values are the average results from 3 independent assays in triplicates, with error bars indicating standard deviations.

Effect of temperature and pH on enzyme activity

The enzyme was active over wide range of temperature i.e. 30-50°C having temperature optimum at 40°C with pNP-palmitate as substrate. Nearly 45% activity was observed at 20°C and 60°C. The enzyme was stable from 20°C to 30°C for 30 min; however it could retain nearly 70 & 60% activity at 40 & 50°C respectively (Fig. 3A). The enzyme was completely stable at 50°C for 15 minutes; however the activity was reduced to 50% after 45 min incubation. The enzyme displayed maximum activity at pH 8.0 and was completely stable at pH 7.0 and nearly 90% stable at pH 6.0 & 8.0 at room temperature for 1h (Fig. 3B).




Figure 3. Temperature (A) and pH (B) dependence activity and stability of LipQ. Values are the average results from 3 independent assays in triplicates, with error bars indicating standard deviations.

Effect of different additives on Rv2485c activity

PMSF and DEPC (1.0 mM v/v) inhibited the enzyme activity to nearly 50% and 60% loss in enzyme activity respectively. However the enzyme did not show any activity with 1.0 mM of PMSF and DEPC after incubation at 50°C for 5 min (data not shown). Eserine did not affect the enzyme activity. The decline in enzyme activity was observed on treatment with β -ME (Fig. 4).




Figure 4. Effect of various additives/chemical modifiers on Rv2485c activity. The experiment was repeated three times in triplicate with error bars indicating standard deviations.


Kinetic Properties

The reaction catalyzed by LipQ (100 μ l of 190 U/ml enzyme stock) was studied by monitoring the increase in pNP absorbance at 420 nm. From the kinetic analysis, it was determined that the enzyme displayed maximum velocity of reaction at 1.0 mM concentration after which no further increase in velocity of the enzyme reaction was observed. The double-reciprocal plot (Lineweaver-Burk plot) of reaction rate was used to evaluate the Michaelis-Menten constant, K_m and maximum velocity, V_{max} . The apparent K_m and V_{max} values for substrate, pNP-palmitate was 1.45 mM & 196.08 U/ml. The turnover number (k_{cat}) of the enzyme was calculated to be 6.597 min^{-1} .

Structure Prediction and identification of active site residues

The Ser-249, Asp-344 and His-377 were predicted as the member of the catalytic triad. There were two Asp (one at 289 and another at 344 positions) residues found to be conserved in multiple sequence alignment (Supplementary Fig 1). The protein structure of LipQ was predicted by homology modeling by using online software, Phyre with template, esterase EstE1 cloned by metagenomics[24] and image was produced and labeled by PyMol[25] (Supplementary Fig 2). Serine residue was found to be buried in the hydrophobic core. It suggested the chances of Asp 344 to be the active site residue. The active site residues of

LipQ were confirmed by site-directed mutagenesis. For each of the altered proteins of Serine and Histidine residues, the enzyme activity was lost (~95%) with pNP-palmitate as substrate. Nearly 75% loss of enzyme activity was observed for Asp-344 mutant. More than 85% enzyme activity was retained with Asp-289 mutant.



Supplementary Figure 2. Cartoon diagram of LipQ model showing position of probable active site residues Ser249, Asp344 and His377. α -helices are shown in blue color, β turns are shown in green color and random loops are shown in orange color. Probability of Asp344 to be an active site residue than Asp-289 was predicted.

Rv2485c expression analysis:

Real Time PCR:

The expression of Rv2485c gene of *M. tuberculosis* H37Ra was analyzed in mid log and stationary phase samples and also in acidic stress, oxidative stress and nutrient depletion by real-time PCR. RNA of the control culture was used as a positive calibrator for real-time quantitative PCR. The specificity of the reactions was verified by melting curve analysis. No expression of Rv2485c was observed in log phase or stationary phase of bacterial growth by semi quantitative RT-PCR (data not shown). Out of all the three stress condition, there was upregulation of expression of Rv2485c in oxidative stress only. The expression of Rv2485c was increased about 3.14 fold during oxidative stress and it was slightly downregulated in

acidic sample in comparison to control samples. In case of nutrient sample there was not much difference than the control samples (Table 5).

Table 5: The expression analysis of Rv2485c using RT- PCR

Type	Ct Value (test)	Ct Value (16S RNA)	Δ Ct (test – 16 S)	$\Delta \Delta$ Ct (Δ Ct test – Δ Ct control)	Fold Change ($2^{-\Delta \Delta Ct}$)
Control	14.00	11.34	2.66	0	1
Acidic	16.00	11.34	4.66	2.00	0.25
Oxidative	13.50	12.50	1.00	-1.66	3.14
Nutrient	10.31	12.50	2.19	-0.47	1.386

Discussion

Pathogenic bacteria utilize a number of mechanisms to cause disease in human hosts. Bacterial pathogens express a wide range of molecules that bind host cell targets to facilitate a variety of different host responses. The development of novel drugs against *M. tuberculosis* has numerous hurdles like identification of therapeutically relevant metabolic targets and introduction of favorable physiochemical characteristics.

The previous studies demonstrated that bacteria grown *in vitro* had a preference for carbohydrates, whereas bacteria grown *in vivo* preferred fatty acids as their carbon source giving due importance to lipolytic enzymes. The bioinformatics study revealed the presence of lip family genes i.e. 24 putative lipases in *M. tuberculosis* H37Rv[26]. Present knowledge about the structure, biochemical properties, regulation and physiological role of all lipase enzymes of mycobacterium is still low. In some recent studies, few lipolytic enzymes of *M. tuberculosis* were characterized both biochemically and enzymatically[15][16][27] allowing us to propose that these enzymes, are very likely to participate in lipid metabolism during dormancy and/or during reactivation[28]. This view is supported by a recent study conducted

by Low *et al.* in 2009 demonstrating extensive accumulation and degradation of TAGs in the bacilli during entry into and exit from hypoxia-induced dormancy, respectively[29]. A lip family gene, *LipF* has already been shown its role in pathogenesis[15]. Still lots of putative lipases/esterases remained to be studied in detail. Therefore, this study aims to characterize LipQ that was predicted to play some role in metabolism and respiration. Till date no reports were available on biochemical properties of LipQ as catalyst and their effects on host cells.

After purification and refolding only 37% of protein could be recovered as active protein due to aggregation problem. It was observed that LipQ exhibited more activity towards the long carbon chain substrates in comparison to short carbon chain substrates. It displayed maximum activity towards pNP-palmitate (C₁₆), at pH 8.0. It might be hypothesized that this enzyme might hydrolyze the palmitate present in mammalian cells to provide energy and might allow mycobacteria to survive for a long period of time. However in present investigation LipQ did not trigger membrane lysis when incubated with macrophage cells, thus displaying no cytotoxic effects on RAW 264.7 cells. In contrast, Rv3452, a phospholipase A2, was reported to induce macrophage lysis[30].

The lip Q was demonstrated to show very low activity with long chain TG hydrolase[12]. In similar studies with other putative lipases, no long-chain TG hydrolysis was detected with LipF[15]. On the other hand LipH, previously annotated as a putative lipase, efficiently hydrolyzed soluble triacylglycerols and vinyl esters but was inactive against emulsified substrate[16].

The possible catalytic triad of the lipase was predicted by multiple sequence alignment (Supplementary Fig 1). LipQ showed presence of conserved pentapeptide, G-X-S-X-G motif (GGSAG) at positions 247-251, a conserved His at position 377 and two conserved aspartic acids at position 289 and 344. To further shed lights on the active site residues of this lipase, the enzyme was treated with PMSF and DEPC. The partial inhibition of enzyme with PMSF and DEPC at room temperature suggested that the active site residues appeared to be buried. Therefore the enzyme was preincubated with PMSF and DEPC at optimum temperature for LipQ activity. The enzyme activity was completely inhibited in this case. It appeared that at low temperature the substrate accessibility to the partially opened active site pocket hindered the inhibition process by PMSF and DEPC as serine was found to be buried in hydrophobic core. While at higher temperature the inhibitor competes with substrate molecules to bind with the active site residue. The inhibition of enzyme activity with PMSF and DEPC suggested the involvement of Ser and His residues in catalytic site. LipQ had five cysteine

residues and predicted to have two disulphide bonds. Inhibition of LipQ enzyme activity with β -ME suggested the importance of S-S bonds in maintaining the proper conformation of enzyme.

It was reported earlier that the *M. tuberculosis* resides in macrophage compartment which has pH range from 6.2 to 4.5. This depends on the activation state of the macrophage[31] *In vitro* assays carried out for the purified lipases showed significant activity in acidic pH range as well as enzyme could retain approximately 50% of enzyme activity at pH 5.0. However no noticeable up-regulation of Rv2485c transcription was observed during *in vitro* culture of *M. tuberculosis* H37Ra during acidic stress. It suggested that the promoter of this gene was not induced by acidic pH and therefore the protein was not specifically synthesized during acidic stress. On the other hand irrespective of the mode of synthesis, the enzyme might work during acidic stress condition as it was stable (retaining 90% enzyme activity) at pH 6.0. While the mechanisms by which *M. tuberculosis* responds to pH are poorly characterized, the global transcriptional response of *M. tuberculosis* to acid stress points to several regulators. In our study the enzyme was specifically induced during oxidative stress conditions only suggesting its role in dormant stage.

During initial infection the internal temperature of the host will be slightly higher than healthy patients. It is due to the defense mechanism of the host against the pathogen[32]. During this condition, the proteins, active at high temperature were predicted to play important role for the survival of pathogens[33]. During initial infection, temperature of the host is unlikely to soar beyond 45⁰C, activity of LipQ at 50⁰C might help not only to prevents its inactivation but ensures its functionality at this temperature. It was reported that the thermostability associated with Rv2416c protein play significant role in intracellular survival within human macrophages[34].

The idea that lipases and phospholipases could be involved in pathogenesis has been suggested by the results of several studies[35][36] The purified lipases/phospholipases from these organisms demonstrated various response in eukaryotic cells *in vitro*. They inhibited the phagocytic function of alveolar macrophages and modulate the release of inflammatory mediators by various kinds of cells in the immune system[35][36][37].

Conclusion

In the present study LipQ, a putative lipase, was demonstrated to be a lipase hydrolyzing long chain esters while eserine, an esterase inhibitor, did not show any effect on enzyme activity. Biochemical parameters provide a clue about the environment where it may work efficiently. Rv2485c gene was expressed only in oxidative stress conditions and might help in fighting the oxidative damage caused by reactive oxidative species (ROS) during the dormant stage and also during the reactivation of the disease. Although the nature of LipQ substrate in *in-vivo* conditions and the role of this enzyme in the life cycle of *M. tuberculosis* remain to be investigated, this study represents some important clues towards the elucidation of the biological function of LipQ.

Conflict of interest statement

Authors have no conflict of interest.

Acknowledgement

The Senior Research Fellowship to GS from CSIR, New Delhi and financial support to JK from DBT, New Delhi is acknowledged.

References

1. Ting LM, Kim AC, Cattamanchi A, Ernst JD. *Mycobacterium tuberculosis* inhibits IFN-gamma transcriptional responses without inhibiting activation of STAT1. J Immunol 1999;**163**:3898-3906.
2. Wilson J, Schurr M, LeBlanc C, Ramamurthy R, Buchanan K, Nickerson C. Mechanisms of bacterial pathogenicity. Postgrad Med J 2002;**78**:216-224.
3. Flynn JL, Chan J. Immunology of tuberculosis. Annu Rev Immunol 2001;**19**:93-129.
4. Lonon MK, Woods DE, Straus DC. Production of lipase by clinical isolates of *Pseudomonas cepacia*. J Clin Microbiol 1988;**26**:979-984.
5. Rollof J, Braconier JH, Soderstrom C, Ehle PN. Interference of *Staphylococcus aureus* lipase with human granulocyte function. Euro J Clin Microbiol Infect Dis 1988;**7**:505-510.

- 458 6. Berto P, Commenil P, Belingheri L, Dehorter B. Occurrence of a lipase in spores of
459 *Alternaria brassicicola* with a crucial role in the infection of cauliflower leaves.
460 FEMS Microbiol Lett 1999;**180**:183-189.
- 461 7. Tsuboi R, Komatsuzaki H, Ogawa H. Induction of an extracellular esterase from
462 *Candida albicans* and some of its properties. Infect Immun 1996;**64**:2936-2940.
- 463 8. Voigt CA, Schafer W, Salomon S. A secreted lipase of *Fusarium graminearum* is a
464 virulence factor required for infection of cereals. Plant J 2005;**42**:364-375.
- 465 9. A. Andrejew, J. Desbordes. Hydrolysis of Fatty Acids Esters by *Mycobacterium phlei*,
466 vol. 117 Ann Inst Pasteur, Paris 1969 pp. 486–500.
- 467 10. Dedieu L, Serveau-Avesque C, Kremer L, Canaan S (2012) Mycobacterial Lipolytic
468 Enzymes: a Gold Mine for Tuberculosis Research. Biochimie in press. doi:
469 10.1016/j.biochi.2012.07.008.
- 470 11. B. Brust, M. Lecoufle, E. Tuillon, L. Dedieu, S. Canaan, V. Valverde, L. Kremer.
471 *Mycobacterium tuberculosis* lipolytic enzymes as potential biomarkers for the
472 diagnosis of active tuberculosis. PLoS ONE, 6 (2011), p. e25078.
- 473 12. Deb C, Daniel J, Sirakova TD, Abomoelak B, Dubey VS, Kolattukudy PE. A novel
474 lipase belonging to the hormone-sensitive lipase family induced under starvation to
475 utilize stored triacylglycerol in *Mycobacterium tuberculosis*. J Biol Chem
476 2006;**281**:3866-3875.
- 477 13. Nidhi Kapoor, Santosh Pawar, Tatiana D. Sirakova, Chirajyoti Deb, William L.
478 Warren, and Pappachan E. Kolattukudy. Human Granuloma *In Vitro* Model, for TB
479 Dormancy and Resuscitation PLoS One. 2013; 8(1): e53657.
- 480 14. Mishra KC, Chastellier CD, Narayana Y, Bifani P, Brown AK, Besra GS, Katoch
481 VM, Joshi B, Balaji KN, Kremer L. Functional Role of the PE Domain and
482 Immunogenicity of the *Mycobacterium tuberculosis* Triacylglycerol Hydrolase LipY.
483 *Infect Immun* 2008;**76**(1):127-140.

- 484 15. Zhang M, Wang JD, Li ZF, Xie J, Yang YP, Zhong Y, Wang HH. Expression and
 485 characterization of the carboxyl esterase Rv3487c from *Mycobacterium tuberculosis*.
 486 Protein Expr Purif 2005;**42**:59-66.
- 487 16. Canaan S, Maurin D, Chahinian H, Pouilly B, Dourousseau C, Frassinetti F, Calvo LS,
 488 Cambillau C, Bourne Y. Expression and characterization of the protein Rv1399c from
 489 *Mycobacterium tuberculosis*. A novel carboxyl esterase structurally related to the
 490 HSL family. Eur J Biochem 2004;**271**:3953-3961.
- 491 17. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search
 492 tool. J Mol Biol 1990;**215**:403-10.
- 493 18. Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of
 494 progressive multiple sequence alignment through sequence weighting, position-
 495 specific gap penalties and weight matrix choice. Nucleic Acids Res
 496 1994;**22**(22):4673-4680.
- 497 19. Kelley LA and Sternberg MJE. Protein structure prediction on the web: a case study
 498 using the Phyre server. Nature Protocols 2009;**4**: 363-371
- 499 20. Sambrook J, Fritsch EF, Maniatis T. Molecular Cloning: A Laboratory Manual. Cold
 500 Spring Harbor Laboratory. Cold Spring Harbor, NY:1989.
- 501 21. Sigurgisladottir S, Konraosdottir M, Jonsson A, Kristjansson JK, Matthiasson E.
 502 Lipase activity of thermophilic bacteria from Icelandic hot springs. Biotechnol Lett
 503 1993;**15**(4):361-366.
- 504 22. K.J. Livak, T.D. Schmittgen, Analysis of relative gene expression data using realtime
 505 quantitative PCR and the $2\Delta\Delta C(T)$ Method. Methods 25 (2001) 402–408.
 506
- 507 23. Cole ST, Brosch R, Parkhill J, Garnier T, Churcher C, Harris D, Gordon SV,
 508 Eiglmeier K, Gas S, Barry CE III, Tekaia F, Badcock K, Basham D, Brown D,
 509 Chillingworth T, Connor R, Davies R, Devlin K, Feltwell T, Gentles S, Hamlin N,
 510 Holroyd S, Hornsby T, Jagels K, Krogh A, McLean J, Moule S, Murphy L, Oliver K,
 511 Osborne J, Quail MA, Rajandream MA, Rogers J, Rutter S, Seeger K, Skelton J,
 512 Squares R, Squares S, Sulston JE, Taylor K, Whitehead S, Barrell BG. Deciphering

- 513 the biology of *Mycobacterium tuberculosis* from the complete genome sequence. Nat
514 (London) 1998;**393**:537-544.
- 515 24. Byun JS, Rhee JK, Kim ND, Yoon J, Kim DU, Koh E, Oh JW, Cho HS. Crystal
516 structure of hyperthermophilic esterase EstE1 and the relationship between its
517 dimerization and thermostability properties. Bmc Struct Biol 2007;**7**:47.
- 518 25. <http://www.pymol.org/>
- 519 26. Cotes K, N’Goma JCB, Dhouib R, Douchet I, Maurin D, Carriere F, Canaan S.
520 Lipolytic enzymes in *Mycobacterium tuberculosis*. Appl Microbiol Biotechnol
521 2008;**78**:741-749.
- 522
- 523 27. Cotes K, Dhouib R, Douchet I, Chahinian H, de Caro A, Carriere F, Canaan S.
524 Characterization of an exported monoglyceride lipase from *Mycobacterium*
525 *tuberculosis* possibly involved in the metabolism of host cell membrane lipids.
526 Biochem J 2007;**408**:417-427.
- 527 28. Deb C, Lee CM, Dubey VS, Daniel J, Abomoelak B, et al. (2009) A novel *in vitro*
528 multiple-stress dormancy model for *Mycobacterium tuberculosis* generates a lipid-
529 loaded, drug-tolerant, dormant pathogen. PLoS One 4: e6077.
530 doi:6010.1371/journal.pone.0006077.
- 531 29. Low KL, Rao PS, Shui G, Bendt AK, Pethe K, Dick T, Wenk MR. Triacylglycerol
532 utilization is required for regrowth of *in vitro* hypoxic nonreplicating *Mycobacterium*
533 *bovis* bacillus Calmette-Guerin. J Bacteriol 2009;**191**:5037-5043.
- 534 30. Schue M, Maurin D, Dhouib R, N’Goma JCB, Delorme V, Lambeau G, Carriere F,
535 Canaan S. Two cutinase-like proteins secreted by *Mycobacterium tuberculosis* show
536 very different lipolytic activities reflecting their physiological function. FASEB J
537 2010;**24**:1893-1903.
- 538 31. Vandal OH, Nathan CF, Ehrt S. Acid Resistance in *Mycobacterium tuberculosis*. J
539 Bacteriol. 2009 **191**(15): 4714–4721.

- 540 32. Jones CH. Case of Low Fever: Delirium: Incomplete Dementia: Convulsions: Death:
541 Tuberculosis. Br Med J 1867;**344**:84-85.
- 542 33. Lack NA, Kawamura A, Fullam E, Laurieri N, Beard S, Russell AJ, Evangelopoulos
543 D, Westwood I, Sim E. Temperature stability of proteins essential for the intracellular
544 survival of *Mycobacterium tuberculosis*. Biochemistry J 2009;**418**:369-378.
- 545 34. Ganaie AA, Lella RK, Solanki R, Sharma C. Thermostable hexameric form of Eis
546 (Rv2416c) protein of *M. tuberculosis* plays an important role for enhanced
547 intracellular survival within macrophages. PLoS ONE 2011;**6**:e27590. doi:10.1371/
548 journal.pone.0027590.
- 549 35. G. Shen, K. Singh, D. Chandra, C. Serveau-Avesque, D. Maurin, S. Canaan, R.
550 Singla, D. Behera, S. Laal. LipC (Rv0220) is an immunogenic cell surface esterase of
551 *Mycobacterium tuberculosis* Infect. Immun. 2012;**80**:243–253
- 552 36. J.C. Bakala N'goma, M. Schue, F. Carriere, A. Geerlof, S. Canaan. Evidence for the
553 cytotoxic effects of *Mycobacterium tuberculosis* phospholipase C towards
554 macrophages. Biochim. Biophys. Acta. 2010; **1801**:1305–1313.
- 555 37. Singh G, Singh G, Jadeja D, Kaur J. Lipid hydrolizing enzymes in virulence:
556 *Mycobacterium tuberculosis* as a model system. Crit Rev Microbiol 2010;**36(3)**:259-
557 269.
- 558
- 559
- 560
- 561
- 562
- 563