

MODS culture for primary diagnosis of tuberculous meningitis and HIV-associated pulmonary tuberculosis in Indonesia

Lidya Chaidir^{1*}, Jessi Annisa¹, Sofiati Dian², David A.J. Moore³, Soni Muhsinin¹, Ida Parwati⁴, Ahmad Rizal Ganiem², Merry Lestari¹, Tina Kusumaningrum¹, Bacht Alisjahbana¹, and Reinout van Crevel⁵

¹ Laboratory for Health Research and Community Service, Faculty of Medicine, Universitas Padjadjaran / Hasan Sadikin Hospital, Bandung 40161, Indonesia

² Department of Neurology, Faculty of Medicine, Universitas Padjadjaran / Hasan Sadikin Hospital, Bandung 40161, Indonesia

³ LSHTM TB Centre and Department of Clinical Research, London School of Hygiene and Tropical Medicine, London WC1E 7HT, England

⁴ Department of Clinical Pathology, Faculty of Medicine, Universitas Padjadjaran / Hasan Sadikin Hospital, Bandung 40161, Indonesia

⁵ Department of Medicine, Radboud University Medical Centre, Nijmegen 6500 HB, The Netherlands

ABSTRACT

Aims: To compare the microscopic observation drug susceptibility (MODS) culture with microscopy and solid culture for diagnosis of TB meningitis and HIV-associated pulmonary tuberculosis (TB).

Study design: Comparative study

Place and Duration of Study: Department of Clinical Pathology, Hasan Sadikin Hospital, Bandung, Indonesia, between 2010 and 2012.

Methodology: Two groups of patients were included. The first group consisted of 167 consecutive HIV-infected patients presenting with suspected pulmonary TB. The second group consisted of 88 patients with clinical suspicion of TB meningitis. Sputum samples from HIV-associated pulmonary TB patients and cerebrospinal fluid (CSF) from patients with TB meningitis were analyzed using microscopy of Ziehl-Neelsen (ZN) stained smears, culture on solid medium (Ogawa), and MODS culture.

Results: MODS showed the highest detection rate in both patient groups. Among HIV-associated pulmonary TB patients, positivity of MODS was 31.2% compared with 26.9% for Ogawa and 20.6% for ZN. Among TB meningitis patients, positivity of MODS was 41.2% compared with 38.8% for Ogawa and 8.3% for ZN. The median time to culture positivity was significantly shorter for MODS compared to Ogawa, both for sputum (median 11 vs 21 days) and CSF (14 vs 33 days). In 14 days, MODS detected significantly more cases compared with Ogawa in both patients group (79.2% vs 2.4% and 68.6% vs 0%, respectively). Laboratory staff readily used MODS after two weeks of training.

Conclusion: We were able to implement MODS culture as a robust, sensitive, and rapid method for diagnosis of HIV-associated pulmonary TB and TB meningitis in a hospital setting in Indonesia. Further studies may be needed to assess the feasibility of MODS culture in other settings and assess its impact on case detection and timely treatment of both forms of TB.

Keywords: MODS culture, primary diagnosis, HIV-associated pulmonary TB, TB meningitis, Indonesia.

1. INTRODUCTION

Diagnosis of tuberculosis (TB) is often difficult, especially among HIV-infected patients [1] and among patients with extrapulmonary TB [2]. Microscopy is insensitive and conventional culture is more sensitive but slow. Commercial liquid culture systems which can shorten the time to detect growth of *Mycobacterium tuberculosis* require sophisticated instrumentation and are more costly [1]. The World Health Organization has recently endorsed the Xpert MTB/RIF which is a nucleic acid amplification-based test. The Xpert test is a rapid and simple method for TB diagnosis and detection of rifampicin resistance. However, it is less sensitive for diagnosis of paucibacillary disease [2], it is relatively expensive, and cannot evaluate treatment response. In addition, culture is still needed to confirm the results of drug susceptibility testing (DST). Therefore, there is a need for rapid, efficient, and inexpensive culture-based diagnostic tools.

One of several recently developed methods for diagnosis and drug resistance testing of TB is the microscopic observation drug susceptibility (MODS) culture. MODS uses broth culture and inverted light microscope to rapidly detect specific growth of *M. tuberculosis*. MODS can be used for primary diagnosis as well as DST. With regard to primary diagnosis, a few studies have examined MODS for diagnosis of pulmonary TB [1, 3, 4], and one study has evaluated MODS for diagnosis of TB meningitis [5]. These studies suggest that MODS culture, with its high sensitivity, speed and low price, and its potential to add DST, may be a suitable alternative for diagnosis of paucibacillary forms of TB, and for use in low-resource settings.

However, for successful implementation more studies are needed, in different patient groups and settings. We compared MODS culture with microscopy and solid culture for diagnosis of TB meningitis and HIV-associated pulmonary TB in Indonesia, which has a high burden of TB and a rapidly growing HIV epidemic. Like in other settings, bacteriological confirmation of HIV-associated pulmonary TB and TB meningitis is difficult, and mortality is very high [6, 7]. More rapid diagnosis would help timely treatment to improve patient outcome and reduce TB transmission.

2. MATERIAL AND METHODS

This study was conducted between February 2010 and January 2012 at Dr. Hasan Sadikin Hospital, the referral hospital for West Java Province, Indonesia. Written informed consent was obtained from all participants and the study was approved by the ethical committee of the Faculty of Medicine, Padjadjaran University/Hasan Sadikin Hospital, Bandung, Indonesia.

We included two groups of patients with suspected TB. The first group consisted of HIV-infected patients, either from the outpatient clinic or the HIV clinic in the hospital, who had symptoms or chest X-ray (CXR) abnormalities suggesting pulmonary TB. From this group one to three consecutive sputum samples were collected per patient prior to initiation of TB treatment. Direct smears were prepared from each sputum specimen for Ziehl-Neelsen (ZN) staining. The remaining sputum specimens were decontaminated by a standard N-acetyl-L-cysteine (NALC)-NaOH method and concentrated by centrifugation at 3000 x g for 15 minutes. For each patient, one sputum culture was made. Sputum specimens from patients who had provided ≥ 1 specimen were mixed before decontamination. The resulting

76 sediments were then resuspended and inoculated into MODS medium as described
77 previously [8, 9] and onto two slants of Ogawa solid medium.

78 The second group consisted of adult patients (≥ 18 years old) with suspected TB meningitis.
79 A clinical diagnosis of meningitis was based on clinical findings, CSF criteria or both [10].
80 Five to ten milliliters of CSF was obtained and divided into two tubes. The first tube (0.5–1
81 mL) was used for CSF cells, protein and glucose. The second tube (4-10 ml) was
82 concentrated by centrifugation at 3000 x g for 15 minutes. The CSF sediment was used to
83 prepare smears for ZN-staining and was inoculated into MODS medium [5] and onto two
84 slants of Ogawa solid medium.

85 For this study, MODS was not performed on a daily basis to simplify the technical work,
86 reduce the high work load in our laboratory, and to standardize the time for microscopic
87 observation. After preparation of smears, specimens were stored in 4°C before they were
88 processed for culture. Positive MODS cultures due to cross-contamination seems very
89 unlikely, as previous studies have shown that cross-contamination is extremely rare in
90 MODS [11]. To further reduce the possibility of cross-contamination and to improve
91 biosafety, MODS plates were sealed in ziplock bags after inoculation.

92 ZN-staining and microscopic reading was done according to standard guidelines. Ogawa
93 cultures, considered positive when mycobacterial growth > 1 colony forming unit (CFU) was
94 observed were read twice weekly until 8 weeks after incubation. Mycobacterial growth in
95 MODS plates was examined twice weekly under an inverted light microscope at 40x
96 magnification from day 4 until day 40 after inoculation. Positive MODS cultures were
97 identified by cord formation which is characteristic of *M. tuberculosis* growth.

98 For quality control of microscopy positive (*M. tuberculosis* H37Rv) and negative (sterile
99 distilled water) controls are included daily. Our laboratory also participates and qualifies for
100 the external quality assurance (EQA) program for smear microscopy provided by the WHO.
101 For mycobacterial culture, we monitor the contamination rate on a monthly basis, as a
102 quality control for the decontamination process. For MODS, negative controls (sample-free
103 liquid medium) were included in each plate, and positive controls (H37Rv) for each run. To
104 minimize the risk of cross-contamination, the positive controls were set up in a separate
105 plate, after all plates with patients samples had been sealed.


106 The positivity rate of microscopy and mycobacterial culture using Ogawa and MODS was
107 expressed as percentage. For HIV-infected individuals suspected of having pulmonary TB,
108 more than one sputum sample was often available. However, analysis in this group was
109 done per patient, not per individual sample. Time to culture positivity was expressed as
110 median days (interquartile range, IQR). Differences between groups were compared using
111 Chi-square test for proportions with $P < .05$ considered statistically significant.

112 113 3. RESULTS

114
115 We included 167 HIV-infected patients screened for pulmonary TB. All patients were ART-
116 naive at the time they were screened for TB. Most patients came with advanced HIV
117 infection, with a median CD4 cell count of 51/ μ l (IQR 10-252). As a second group, we
118 included 88 patients with suspected TB meningitis. Most patients presented with lowered
119 consciousness and focal neurological signs, and with typical CSF findings of moderate
120 pleiocytosis, elevated protein and lowered glucose, 9 patients (11%) were HIV-infected.

121 MODS culture had the highest detection rate among both groups (Figure 1). In HIV-infected
122 pulmonary TB suspects, MODS culture of 7 patients (4.2%) were contaminated and were

123 excluded from the calculation of positivity rate. Of 160 patients left, MODS culture was found
 124 positive in 50 patients (31.2%) compared with 43 patients (26.9%) for Ogawa culture and 33
 125 patients (20.6%) for ZN, the difference between MODS culture and ZN was statistically
 126 significant ($P = .02$). None of the patients whose MODS cultures were contaminated had a
 127 positive Ogawa culture. Among 127 ZN-smear negative patients, MODS was positive in 20
 128 patients (15.8%) compared with 15 (11.8%) using Ogawa. Among TB meningitis suspects,
 129 MODS cultures of 3 patients (3.4%) were contaminated and were excluded from the
 130 analysis. Of 85 patients left, MODS culture also gave the highest detection rate, in 35
 131 patients (41.2%) compared with 33 patients (38.8%) for Ogawa and 7 patients (8.3%) for ZN.
 132 Both MODS and Ogawa culture were significantly more sensitive compared to ZN ($P <$
 133 $.001$). Of patients whose MODS cultures were contaminated, one patient was positive for the
 134 Ogawa culture. Among 77 smear-negative patients, MODS was positive in 30 (39.0%) and
 135 Ogawa in 27 (35.1%) patients. Ten patients in both patient groups were MODS-
 136 positive/Ogawa-negative, and 3 and 8 patients, respectively, were MODS-negative/Ogawa-
 137 positive. This was thought to be due to the low numbers and clumping of mycobacteria
 138 which makes that bacilli are not always uniformly distributed in sample aliquots.



139

140 Figure 1. Positivity rate of microscopy of Ziehl-Neelsen-stained smears, Ogawa solid culture,
 141 and MODS culture for suspected pulmonary TB among 160 HIV-infected patients and
 142 suspected TB meningitis among 85 patients. Seven patients in HIV-infected pulmonary
 143 group and three patients in TB meningitis group were excluded due to contamination in
 144 MODS cultures.

145 The median time to culture positivity was shorter for MODS culture than for Ogawa culture in
 146 both groups of patients (figure 2). Using sputum from HIV-infected patients, positive results
 147 for MODS were obtained after a median 11 days (IQR 7-14 days) compared with 21 days
 148 (IQR 18-28 days; $P < .001$) for Ogawa. Within 14 days, MODS detected 79.2% of positive
 149 samples, compared to only 2.4% for Ogawa ($P < .001$). Among suspected TB meningitis
 150 patients, the median time to culture positivity of CSF samples was 14 days (IQR 11-18 days)
 151 for MODS, compared to 33 days (IQR 23-39 days, $P < .001$) for Ogawa. In this group,

MODS detected 68.6% of positive cultures within 14 days, while all Ogawa cultures were still negative by this time.




Figure 2 Positive culture results (n) in time for MODS (Δ) and Ogawa (□) among patients with suspected HIV-associated pulmonary TB [A] and suspected TB meningitis [B]

4. DISCUSSION

Bacteriological confirmation of certain paucibacillary forms of TB, like TB meningitis, is problematic. We examined the performance of MODS as a rapid culture-based method for diagnosis of TB meningitis and HIV-associated pulmonary TB in a hospital setting in Indonesia. As comparative tests we used solid culture and microscopy of Ziehl-Neelsen stained smears, the two most widely used methods for TB diagnosis in low-resource settings. For both study groups, MODS culture had the highest positivity rate. In addition, MODS culture was much quicker compared with Ogawa solid culture.

The high positivity rate of MODS culture, which is due to the use of liquid medium, was in line with previous studies [8, 12, 13]. One previous study reported that MODS was more sensitive than automatic mycobacterial culture for diagnosis of pulmonary TB [3], but this was not confirmed in another study [1]. To our knowledge, only one study evaluated MODS for suspected TB meningitis [5]. This study from Vietnam showed that MODS was a sensitive and simple culture-based method for diagnosis of TB meningitis. Mycobacterial growth Indicator tube (MGIT, Becton Dickinson) showed a slightly higher sensitivity in the Vietnam study, but the difference was not significant. A smaller volume of deposit was inoculated into the MODS culture than the MGIT culture, possibly contributing to the slight difference in sensitivity.

In our study MODS culture significantly reduced the time to both positive and negative results. After two weeks, 79.2% of HIV-associated pulmonary TB had been diagnosed with MODS, and 68.6% of TB meningitis, compared with 2.4% and 0%, respectively, using Ogawa solid culture. These findings are in line with previous studies in HIV-associated pulmonary TB [1, 3] and TB meningitis patients [5]. With the high sensitivity and rapidity, MODS culture is suitable for the TB screening of HIV-infected patients, as shown by a study in Peru [4].

Among HIV-infected patients examined for pulmonary TB, the positivity rate of ZN was relatively high compared to culture. This is probably caused by the fact that ZN microscopy

187 was performed on all available sputum samples for each patient, while only single Ogawa
188 and MODS culture were performed for every patient.

189 The contamination rate of MODS culture was around 3-4%. A contamination rate of 2%-3%
190 is acceptable for fresh specimens, but it may be as high as 5%-10% for stored samples [14].

191 MODS culture is a simple test. Sample preparation for MODS is similar to that required for
192 other culture methods. The microscopic detection of mycobacteria by MODS is technically
193 similar to that of a smear. Our laboratory technicians were able to perform the MODS assay
194 and accurately read the results after two weeks of training. The cord formation, which is
195 characteristic of *M. tuberculosis* in liquid culture, is easy to recognize using inverted
196 microscopy. Certain species of non-tuberculous mycobacteria (NTM), for example *M.*
197 *chelonae*, may also form cording in liquid culture but this can be recognized by rapid
198 overgrowth by day 5 [3]. Of note, genotyping of > 800 cultures isolated in our laboratory has
199 only revealed a very low number of NTM (<3%)[15].

200 A possible limitation of this study is the fact that we did not include automated liquid culture
201 (MGIT), which is recommended as diagnostic standard for culture, as the comparative test.
202 The few studies comparing MODS culture with MGIT showed slightly different results [1, 3,
203 5]. Furthermore, our study was performed in a single laboratory. However, as this is a busy
204 hospital laboratory of a large hospital, our results suggest that MODS can be implemented in
205 high-burden settings. Finally, we did not evaluate MODS for DST. Therefore, further studies
206 may be needed to assess the feasibility of MODS culture in other settings, and assess its
207 impact on case detection and timely treatment of paucibacillary forms of TB like TB
208 meningitis.

209

210 **5. CONCLUSION**

211

212 MODS culture showed a high sensitivity and rapidity (compared with solid culture), simplicity,
213 and potential low cost (compared with automated liquid systems) for diagnosis of HIV-
214 associated pulmonary TB and TB meningitis in a hospital setting in Indonesia. As such, this
215 seems a reliable method to help implement *M. tuberculosis* culture more widely and speed
216 up TB diagnosis in TB endemic settings.

217

218

219 **ACKNOWLEDGEMENTS**

220

221 We thank Dr. Bayu Wahyudi, Director of Hasan Sadikin hospital; Prof. Tri Hanggono
222 Achmad, Dean of the Faculty of Medicine, Universitas Padjadjaran; and dr. Tintin Gartinah,
223 the Head of West Java Province Reference Laboratory, for encouraging and accommodating
224 research at their institutions. We also thank dr. Ni Sayu Dewi and dr. Nieke Dewi for their
225 help in sample collection.

226

227 **COMPETING INTERESTS**

228

229 Authors have declared that no competing interests exist.

230

231 **AUTHORS' CONTRIBUTIONS**

232

233 Lidya Chaidir designed the study, performed the statistical analysis, wrote the protocol, and
234 wrote the first draft of the manuscript. Jessi Annisa managed the specimens and data
235 collection in laboratory. Sofiati Dian and Ahmad Rizal Ganiem managed the inclusion of TB

meningitis patients in the hospital as well as the patient information. David A. J. Moore reviewed the manuscript before submission. Soni Muhsinin and Tina Kusumaningrum performed the study in the laboratory. Ida Parwati coordinated the laboratory work. Merry Lestari managed the inclusion of HIV-infected pulmonary TB in the hospital as well as the patient information. Bacht Alisjahbana and Reinout van Crevel designed the study and reviewed the data analysis and the manuscript. All authors read and approved the final manuscript.”

CONSENT

All authors declare that written informed consent was obtained from the patient.

ETHICAL APPROVAL

This study was approved by the ethical committee of the Faculty of Medicine, Padjadjaran University/Hasan Sadikin Hospital, Bandung, Indonesia.

REFERENCES

1. Ha DT, Lan NT, Kiet VS, Wolbers M, Hang HT, Day J, et al. Diagnosis of pulmonary tuberculosis in HIV-positive patients by microscopic observation drug susceptibility assay. *J Clin Microbiol.* 2010;48(12):4573-9.
2. Armand S, Vanhuls P, Delcroix G, Courcol R, Lemaitre N. Comparison of the Xpert MTB/RIF test with an IS6110-TaqMan real-time PCR assay for direct detection of *Mycobacterium tuberculosis* in respiratory and nonrespiratory specimens. *J Clin Microbiol.* 2011;49(5):1772-6.
3. Moore DA, Evans CA, Gilman RH, Caviedes L, Coronel J, Vivar A, et al. Microscopic-observation drug-susceptibility assay for the diagnosis of TB. *N Engl J Med.* 2006;355(15):1539-50.
4. Reddy KP, Brady MF, Gilman RH, Coronel J, Navincopa M, Ticona E, et al. Microscopic observation drug susceptibility assay for tuberculosis screening before isoniazid preventive therapy in HIV-infected persons. *Clin Infect Dis.* 2010;50(7):988-96.
5. Caws M, Ha DT, Torok E, Campbell J, Thu do DA, Chau TT, et al. Evaluation of the MODS Culture Technique for the Diagnosis of Tuberculous Meningitis. *PLoS ONE.* 2007;2(11):e1173.
6. Wisaksana R, Indrati AK, Fibriani A, Rogayah E, Sudjana P, Djajakusumah TS, et al. Response to first-line antiretroviral treatment among human immunodeficiency virus-infected patients with and without a history of injecting drug use in Indonesia. *Addiction.* 2010;105(6):1055-61.
7. Ruslami R, Ganiem AR, Dian S, Apriani L, Achmad TH, van der Ven AJ, et al. Intensified regimen containing rifampicin and moxifloxacin for tuberculous meningitis: an open-label, randomised controlled phase 2 trial. *Lancet Infect Dis.* 2012;13(1):27-35.
8. Caviedes L, Lee TS, Gilman RH, Sheen P, Spellman E, Lee EH, et al. Rapid, efficient detection and drug susceptibility testing of *Mycobacterium tuberculosis* in sputum by microscopic observation of broth cultures. The Tuberculosis Working Group in Peru. *J Clin Microbiol.* 2000;38(3):1203-8.
9. Moore DA, Mendoza D, Gilman RH, Evans CA, Holm Delgado MG, Guerra J, et al. Microscopic observation drug susceptibility assay, a rapid, reliable diagnostic test for multidrug-resistant tuberculosis suitable for use in resource-poor settings. *J Clin Microbiol.* 2004;42(10):4432-7.
10. Ganiem AR, Parwati I, Wisaksana R, van der Zanden A, van de Beek D, Sturm P, et al. The effect of HIV infection on adult meningitis in Indonesia: a prospective cohort study. *AIDS.* 2009;23(17):2309-16.

- 289 11. Moore DA, Caviedes L, Gilman RH, Coronel J, Arenas F, LaChira D, et al. Infrequent
290 MODS TB culture cross-contamination in a high-burden resource-poor setting. *Diagn*
291 *Microbiol Infect Dis*. 2006;56(1):35-43.
- 292 12. Shiferaw G, Woldeamanuel Y, Gebeyehu M, Girmachew F, Demessie D, Lemma E.
293 Evaluation of microscopic observation drug susceptibility assay for detection of multidrug-
294 resistant *Mycobacterium tuberculosis*. *J Clin Microbiol*. 2007;45(4):1093-7.
- 295 13. Arias M, Mello FC, Pavon A, Marsico AG, Alvarado-Galvez C, Rosales S, et al. Clinical
296 evaluation of the microscopic-observation drug-susceptibility assay for detection of
297 tuberculosis. *Clin Infect Dis*. 2007;44(5):674-80.
- 298 14. World Health Organization. Laboratory services in tuberculosis control; Part 3 : Culture.
299 1998. Accessed 28 June 2011. Available:
300 http://whqlibdoc.who.int/hq/1998/WHO_TB_98.258_%28part3%29.pdf
- 301 15. Parwati I, Alisjahbana B, Apriani L, Soetikno RD, Ottenhoff TH, van der Zanden AG, et al.
302 *Mycobacterium tuberculosis* Beijing genotype is an independent risk factor for
303 tuberculosis treatment failure in Indonesia. *J Infect Dis*. 2010;201(4):553-7.
304