

Original Research Article

Dietary Pattern to Antioxidants Levels in Patients with Simple Goiter and Thyroid Cancer

ABSTRACT

Background /Aims: Oxidative stress has been implicated in the pathogenesis of thyroid cancer and simple goiter among other diseases however; there has not been a direct comparison of the severity of the stress factor in these two conditions in relation to their dietary pattern in our environment.

Objective: This study assessed oxidative stress indices, antioxidant status and dietary pattern in thyroid cancer and simple goiter compared with the controls.

Study design: This is a case-control study

Place and Duration of Study: Study groups were recruited from Departments of Surgery and Nuclear medicine of University of Ilorin teaching hospital, Ilorin, Kwara State and University College Hospital, Ibadan, Oyo State respectively between March 2013 and September 2013.

Materials and Methodology: A total number of one hundred and five (105) age matched participants consisting of 88 females and 17 males were recruited for this study. They were divided into 3 groups; Group one 35 participants with thyroid cancer and group two, 35 participants with simple goiter and group three, 35 control participants. Plasma levels malondialdehyde (MDA), hydrogen peroxide (H_2O_2) and total plasma peroxide (TPP) were determined by colorimetric methods. Antioxidant status was determined by measuring total antioxidant potential (TAP), glutathione peroxidase (GPx), Superoxide dismutase (SOD), reduced glutathione (GSH), glutathione-s-transferase (GST) by colorimetric methods. Selenium (Se) was determined by atomic absorption spectrophotometry (AAS). Oxidative stress index (OSI) was measured and all values expressed as mean $\pm$ SD, while frequency of intake of various dietary sources of the participants were collated.

Results: TAP, SOD, GPx, GSH and GST (antioxidants) levels in group 1 (552.17 ± 74.67 ; 1.30 ± 0.32 ; 1.91 ± 0.23 ; 4.47 ± 0.59 ; 0.92 ± 0.25 ; 1.28 ± 0.04 respectively) were significantly lower than group 3 (933.51 ± 80.15 ; 2.85 ± 0.39 ; 5.40 ± 0.76 ; 8.34 ± 1.12 ; 2.11 ± 0.31). While the mean plasma levels of antioxidants in group 2 (704.74 ± 62.22 ; 2.01 ± 0.27 ; 4.62 ± 0.89 ; 6.19 ± 0.56 ; 1.47 ± 0.14 and 1.40 ± 0.04) were significantly higher than that of group 1. However, significant increase was observed in mean levels of oxidative stress markers; TPP, MDA, H_2O_2 and OSI, in group 1 (16.24 ± 2.52 , 13.88 ± 2.95 , 13.52 ± 6.91 and 2.93 ± 1.40 respectively) compared to group 3 (4.57 ± 1.32 , 3.22 ± 1.20 , 3.01 ± 0.86 and 0.48 ± 0.14). Oxidative stress markers in group 2 (9.79 ± 1.40 , 5.90 ± 1.13 , 3.81 ± 1.40 and 1.62 ± 0.23) were significantly lower than group 1. A P -value < 0.05 was considered significant.

Conclusion: The study reported higher oxidative stress markers, oxidative stress index and lower antioxidant status. Oxidative stress appeared more marked in thyroid malignancy than the benign state. Adequate intake of fresh fruits and vegetables is recommended for thyroid cancer patients.

Key Words: Thyroid cancer, simple goiter, enzymatic antioxidant, oxidative stress index.

1. INTRODUCTION

Thyroid cancer is the most common endocrine malignancy with more deaths annually than other endocrine cancers combined [1,2,3]. Free radicals have been implicated in a wide variety of degenerative diseases including thyroid cancer. It is believed that oxidative stress plays a major role in the pathogenesis of thyroid cancer [4]. This may occur when the cellular antioxidant defense is unable to eliminate the detoxification of the free radicals. The imbalance between the oxidant (free radicals) and antioxidant system in favour of oxidant generation has carcinogenic effect [5,6].

Thyroid hormones regulate oxidative metabolism and thus play an important role in production of free radical such as hydrogen peroxide [7,8,9]. Hydrogen peroxide is abundantly present in the thyroid gland

as a substrate for the thyroperoxidase that catalyzes iodine oxidation, binding to thyroglobulin and oxidative coupling of iodotyrosines into iodothyronine. Therefore reduced detoxification of hydrogen peroxide may lead to thyroid cell death. Thyroid hormones regulate the synthesis and degradation of antioxidants enzymes such as superoxide dismutase (SOD), catalase, glutathione peroxidase (GPx) and non-enzymatic antioxidants such as vitamin E and C, uric acid, ferritin, transferrin and ceruloplasmin [10,11]. Vitamin A, C and E have been proved to have antioxidant efficacy *invivo*. Many common foods are good sources of antioxidants. Spices, herbs, fibers and essential oils are rich in antioxidants properties in the plant. Fruits like oranges, guava, mango, grape juice; spices such as ginger, pepper, garlic, onion are also good sources of antioxidants.

This study assessed the degree of imbalance between the pro-oxidants and antioxidants in thyroid cancer and simple goiter in relation to their diet. The findings may help in understanding the role of diet in ameliorating oxidative stress and the development of thyroid cancer in our environment.

2. MATERIALS AND METHODS

Ethical approval was obtained from University of Ibadan/University College Hospital and University of Ilorin Teaching Hospital ethical review committee respectively and written informed consent (attached to every questionnaire completed by each participant) was obtained from individual patients prior to recruitment into the study.

2.1 Study Design

A total of 105 participants matched by age and sex were recruited for this study from Nuclear Medicine and Surgery department of University College Hospital, Ibadan and University of Ilorin Teaching, Ilorin respectively. All subjects were examined and classified into 3 groups. Participants without any form of cancer or chronic illnesses constituted the controls (group1), participants with thyroid cancer constituted the second while group 3 comprised of participants with simple goiter. Each group consisted of 35 participants. Their medical history and physical examination were documented. Diagnosis of simple goiter was by clinical examinations: thyroid swelling and biochemical result of T3, T4 and TSH while participants with thyroid cancer were diagnosed with thyroid swelling, thyroid ultrasound and histological diagnosis either by Fine Needle Aspiration Cytology (FNAC) or excision biopsy for confirmation. A questionnaire was used to capture the participants' dietary recall and the frequency of consumption of major food items. Exclusion criteria included Individuals with other chronic illness or combination of any other forms of cancer. Participants on multivitamins or food supplements were also excluded from the study. Ethical approval and informed consent were obtained from UI/UCH and UITH ethical review committee respectively.

2.2 Blood Sample Collection

Five milliliters of venous blood was collected into lithium heparin anticoagulant bottles. This was gently mixed to avoid coagulation and hemolysis. The plasma samples were frozen at -20° until samples were analyzed. A structured questionnaire was administered detailing the dietary intake of the study participants.

2.3 Method of analysis

Plasma antioxidants such as SOD, GPx, GSH, GST were measured based on standard colorimetric method respectively [12,13,14,15]. Total Antioxidant Potential (TAP) was measured by colorimetric method [16]. Total plasma peroxide (TPP), MDA and H₂O₂ levels were measured based on colorimetric method of [17,18,19] method respectively. Selenium was determined by Atomic Absorption Spectrophotometry (ASS) with appropriate control [20]. The ratio of TPP level to TAP level was calculated as oxidative stress index (OSI).

2.4 Statistical analysis

Statistically analysis was performed using SPSS software version 17.0. All data were expressed as mean $\pm$ SD (standard deviation). ANOVA was also used to compare the 3 groups (thyroid cancer, simple goiter and control) while Post Hoc test was used to compare the difference between the means of the participant's groups. The Pearson's correlation was used to determine the relationship between the oxidative stress index (OSI) and other biochemical parameters. Data were analyzed at 95% confidence interval and *P*-values 0.05 (*P* < .05) were considered statistically significant.

3. RESULTS

The anthropometric data of the study participants showed no significant differences among the three groups (*P* < 0.05) (Table 1).

Consumption of carbohydrate, animal protein, smoked food and sea foods was similar among the three groups. Larger percentages of the control participants consumed fruits and vegetables (sources of exogenous antioxidants) regularly more than the test groups (Table 2).

Table 1: Anthropometric indices in thyroid cancer, simple goiter and control participants.

Variables	Thyroid cancer n=35	Simple goiter n=35	Control n=35	F-value	p-value
Age (years)	41.83 $\pm$ 6.11	41.97 $\pm$ 6.14	44.51 $\pm$ 7.12	1.908	.15
Weight (Kg)	64.40 $\pm$ 4.29	65.03 $\pm$ 3.88	64.11 $\pm$ 4.57	0.422	.66
Height (M)	1.60 $\pm$ 0.03	1.60 $\pm$ 0.05	1.59 $\pm$ 0.03	1.230	.30
BMI (kg/m ²)	25.33 $\pm$ 1.93	25.54 $\pm$ 1.59	25.53 $\pm$ 2.09	0.142	.86
Systolic BP (mmHg)	136.97 $\pm$ 7.55	134.29 $\pm$ 9.67	120.41 $\pm$ 4.24	29.206	.30
Diastolic BP (mmHg)	80.49 $\pm$ 5.99	76.14 $\pm$ 4.75	74.26 $\pm$ 3.25	0.843	.43

Table 2: Percentage frequency of diet style in the test groups and control group.

Variables	Frequency intake	of Thyroid cancer (%) n=35	Simple goiter (%) n=35	Control (%) n=35
Carbohydrate	Regular	86	83	84
	Not regular	14	17	16
Protein (Animals)	Regular	82	76	71
	Not regular	18	24	29
	(Plants)	Regular	62	70
	Not regular	56	38	30
Smoked food	Regular	77	74	66

	Not regular	23	26	34
Sea foods	Regular	62	66	69
	Not regular	38	34	31
Fruits/vegetables	Regular	59	63	70
	Not regular	41	37	30

Table 3 The mean values of biochemical parameters in the study participants (thyroid cancer and simple goiter) and control participants.

	Thyroid cancer	Simple goiter	Controls	F-value	P-value
MDA	13.88 ± 2.95 ^{a,b}	5.90 ± 1.13 ^a	3.22±1.20	283.25	.00*
H ₂ O ₂	13.52 ± 6.91 ^{a,b}	3.81 ± 1.40 ^a	3.01±0.86	220.51	.00*
TPP	16.24 ± 2.52 ^{a,b}	9.79 ± 1.40 ^a	4.57±1.32	356.82	.00*
TAP	552.17 ± 74.67 ^{a,b}	704.74 ± 62.22 ^a	933.51±80.15	243.72	.00*
SOD	1.30 ± 0.32 ^{a,b}	2.01 ± 0.27 ^a	2.85±0.39	193.56	.00*
GPX	1.91 ± 0.23 ^{a,b}	4.62 ± 0.89 ^a	5.40±0.76	247.48	.00*
GSH	4.47 ± 0.59 ^{a,b}	6.19 ± 0.56 ^a	8.34±1.12	206.30	.00*
GST	0.92 ± 0.25 ^{a,b}	1.47 ± 0.14 ^a	2.11±0.31	199.59	.00*
Se	1.28 ± 0.04 ^{a,b}	1.40 ± 0.04 ^a	1.60±0.05	450.20	.00*
OSI	2.93 ± 1.40 ^{a,b}	1.62 ± 0.23 ^a	0.48±0.14	348.75	.00*

* Statistically significant at p< 0.05 (2-tailed).

^a Value significantly different (p< 0.05) from control value (Post hoc test)

^b Value significantly different (p< 0.05) from simple goiter value(post hoc test)

3.1 Oxidative stress markers

The mean plasma oxidative stress markers were significantly different between the study participants and the controls. The plasma levels of MDA (μmol/L), H₂O₂ (μmoles) and TPP (μmolH₂O₂/L) in group 1 (thyroid cancer group) were significantly higher than group 3 (controls) (13.88±2.95, 13.52±6.91; 16.24±2.52) vs (3.22±1.20; 3.01±0.86; 4.57±1.32) respectively (Table 3). While, the mean plasma levels of MDA, H₂O₂, TPP in group 2 (simple goiter) were also significantly lower in group 1(5.90±1.13; 3.81±1.40; 9.79±1.40)vs(13.88±2.95, 13.52±6.91; 16.24±2.52) respectively. The plasma means levels of oxidative stress markers were significantly higher in group 2 than group 3.

3.2 Antioxidants

The mean plasma levels of total antioxidant potential (TAP), superoxide dismutase (SOD), glutathione peroxidase (GPx), reduced glutathione (GSH), glutathione-s-transferase (GST) in group 1 were significantly lower than group 3 (552.17 ± 74.67 ; 1.30 ± 0.32 ; 1.91 ± 0.23 ; 4.47 ± 0.59 ; 0.92 ± 0.25 ; 1.28 ± 0.04) vs (933.51 ± 80.15 ; 2.85 ± 0.39 ; 5.40 ± 0.76 ; 8.34 ± 1.12 ; 2.11 ± 0.31) respectively (Table 3). Meanwhile, the mean plasma levels of TAP, SOD, GPx, GSH, GST in group 2 were significantly higher than that of group 1 (704.74 ± 62.22 ; 2.01 ± 0.27 ; 4.62 ± 0.89 ; 6.19 ± 0.56 ; 1.47 ± 0.14 ; 1.40 ± 0.04) vs (552.17 ± 74.67 ; 1.30 ± 0.32 ; 1.91 ± 0.23 ; 4.47 ± 0.59 ; 0.92 ± 0.25 ; 1.28 ± 0.04) respectively. The plasma mean levels of TAP, SOD, GPx, GSH, GST were significantly lower in group 2 than that of controls (group 1).

3.3 Selenium

The mean plasma selenium in the thyroid cancer group (group 1) (1.28 ± 0.04) was significantly lower compared with the control group (3). While, the mean plasma selenium in group 2 (1.40 ± 0.04) was significantly higher than group 1. The selenium level in group 2 was lower compared to that of controls (group 3) (Table 3).

3.4 Oxidative stress index (OSI)

The mean plasma OSI in group 1 was significantly higher compared to that of group 2 (2.93 ± 1.40) vs (1.62 ± 0.23). The OSI level in group 2 was also significantly lower than group 1 but was significantly higher than the control (group 3) (0.48 ± 0.14) (Table 3) respectively.

Table 4: Correlation studies between oxidative stress index (OSI) and biochemical parameters in the study participants and controls.

	Group 1 (Thyroid cancer) n=35	Group 2 (Simple goiter) n=35	Group 3 (Control) n=35
MDA ($\mu\text{mol/L}$)	$r = 0.554^*$ $P = .00$	$r = 0.192$ $P = .27$	$r = -0.345^*$ $P = .04$
H_2O_2 (μmoles)	$r = -0.420^*$ $P = .012$	$r = 0.099$ $P = .57$	$r = -0.187$ $P = .28$
TPP ($\mu\text{molH}_2\text{O}_2/\text{L}$)	$r = 0.775^*$ $P = .00$	$r = 0.844^*$ $P = .00$	$r = -0.898^*$ $P = .00$
TAP	$r = -0.762^*$ $P = .00$	$r = -0.610^*$ $P = .00$	$r = -0.139$ $P = .43$
SOD	$r = -0.375^*$ $P = .03$	$r = 0.842$ $P = .64$	$r = -0.190$ $P = .27$
GPx	$r = -0.074$ $P = .69$	$r = -0.051$ $P = .75$	$r = -0.017$ $P = .93$
GSH	$r = 0.277$ $P = .11$	$r = 0.009$ $P = .96$	$r = -0.271$ $P = .12$

GST	$r = -0.368^*$ $P = .03$	$r = 0.245$ $P = .16$	$r = 0.370^*$ $P = .03$
Se	$r = -0.401$ $p = .17$	$r = -0.149$ $p = .39$	$r = 0.012$ $p = .95$

* Correlation is significant at the 0.05 level.

The correlation studies showed a strong positive significant association between OSI and oxidative markers TPP, MDA and H_2O_2 in thyroid cancer participants. Also there was a negative significant association between OSI and TAP ; SOD GST in thyroid cancer group. Table 4

There was also a positive significant association between OSI and TPP in simple goiter group and GST in the controls respectively.

TAP in the simple goiter group, MDA, and TPP in controls showed negative significant association with OSI (Table 4).

4. Discussion

An imbalance between the oxidant and antioxidant system has been reported to play a major role in development of thyroid cancer and simple goiter [6]. The observed higher levels of pro oxidants and lower levels of antioxidants in thyroid cancer participants compared with the controls supports the reported imbalance between the oxidants and antioxidant defense system.

Observed increases in level of hydrogen peroxide (H_2O_2) a pro-oxidant generated in the synthesis of thyroxine by the thyroglobulin-iodine-tyrosine complex in the thyroid gland may be the basis of the free radical generation as earlier reported by other authors [6,21]. Oxidative stress in thyroid gland is very common because hydrogen peroxide (H_2O_2) is generated during thyroid synthesis.

In this study, there were significantly increased levels of total plasma peroxide (TPP), malondialdehyde (MDA) and hydrogen peroxide (H_2O_2) in thyroid cancer and to a lesser extent in simple goiter compared with the controls. TPP is a measure of increased ROS in the test group. This further confirms the increase in generation of free radicals which appears to be progressive or worsened in the thyroid cancer group. It has also been observed that as age increases the risk of development of thyroid cancer increases. This may not be unexpected from increase in the duration of exposure to environmental factors and toxic chemicals as reported by Jemal *et al*, 2006 [22].

Significantly decreased levels of antioxidants (superoxide dismutase, glutathione peroxidase, reduced glutathione, glutathione-s-transferase) were observed in thyroid cancer and simple goiter subjects when compared with controls. The significant decrease observed in the test group could be due to their involvement in scavenging free radicals generated in thyroid gland. Several authors[6,23] reported a significant correlation between antioxidant enzymes and oxidants in patients with thyroid cancer and simple goiter. Therefore as the oxidant level increases, the antioxidants level decreases in thyroid cancer and simple goiter patients. It was suggested that a decrease in total antioxidant potential (TAP) of the body may increase the oxidative damage in the tissue including the thyroid gland.

The significant decrease in TAP in the thyroid cancer and simple goiter compared with controls confirmed the reduced plasma antioxidant defense system. Similar findings by Senthil and Manoharan 2004; Akinci *et al*, 2008; Wang *et al*, 2010 confirmed the significant difference between the total antioxidant status of thyroid cancer patients and controls. Hence the progression of oxidative stress contributes to the development of thyroid cancer in the absence of adequate consumption of antioxidant vitamins and failure of the endogenous antioxidants to detoxify the free radicals.

The effects of dietary antioxidants on the overall antioxidant status of test groups (thyroid cancer and simple goiter), indicated that more than 70% of the thyroid cancer participants consumed food rich in carbohydrate, animal proteins and smoked foods as against the controls who regularly took more of plant

protein and less of the smoked food. About 59% of thyroid cancer participants ingest fruits and vegetables regularly as against 70% of the controls that consumed fruits and vegetables more than four days in a week. Therefore, dietary antioxidants seem to be associated with an improved antioxidant status, this finding has been earlier reported [24].

Diet is one of the lifestyle factors that influence the risk of developing cancer. The foods that we eat affect the risk of developing certain types of cancer; Diet high in energy, fat and low in fruits, vegetables and fibers are more likely to influence cancer development [25]. There is convincing evidence that smoked foods and meat cooked at high temperature increase risk of developing cancer. Smoking is a known source of food contamination caused by carcinogenic polycyclic aromatic hydrocarbons while meat cooked at high temperature contains heterocyclic amines (HCA) also a carcinogenic compound [26]. Studies indicate a correlation between the increase occurrence of cancer and the frequent intake of smoked foods [26]. Smoked foods absorb large amount of tars that arise from incomplete combustion of wood or charcoal fire which are known to contain numerous carcinogens [27].

Plants based diet (fresh fruits, legumes and vegetables) contain many beneficial nutrients such as vitamins, minerals, antioxidants, phytochemicals and fiber which may reduce risk of developing cancer [28]. Carotenoids present in significant amounts in fruits, as well as vitamin E in oil are probably responsible for protection against cancer [29]. Lycopene is a potent antioxidant found in tomatoes, watermelon and strawberries that may lower the risk of cancer development [29]. While fatty red meats, processed meats, processed foods that are low in fiber, antioxidants, phytochemicals, heavily salted and pickled foods may increase risk of cancer development [30]. Therefore, these dietary antioxidants may increase the level of antioxidant defense system. A low –fat, high fiber diet is recommended with plenty of fresh fruits and vegetables that are rich in vitamins A, B, C, D, E and trace elements such as zinc, selenium can help to reduce the risk of complication [31]. The effect of free radicals generated in these subjects (thyroid cancer & simple goiter) could or possibly be ameliorated by dietary antioxidants thereby reducing the risk of oxidative stress.

Significant decreased level of selenium was observed in thyroid cancer and simple goiter subjects compared with the control. Selenium is an essential trace element for the biosynthesis and function of selenocysteine-containing selenoproteins implicated in thyroid hormone metabolism and thyroid gland function. However, generation of hydrogen peroxide and oxidative damage are decreased by selenoenzyme systems in the thyrocytes involved in regulating hormone synthesis. Low selenium levels and lower selenoenzyme activity levels have been documented in thyroid cancer and simple goitre [29].

5. Conclusion

The findings in this study support the hypothesis that oxidative stress contributes to the development of thyroid cancer and simple goiter in our environment. The study has established the fact that inadequate level of antioxidants (both endogenous and exogenous) as well as selenium is associated with the development of thyroid cancer and simple goiter. Diets including fruits and vegetables which contain vitamins could increase dietary antioxidant status and mitigate the harmful effect of increased oxidative stress in the thyroid gland.

CONSENT

All authors declare that informed consent was obtained from the participants for publication.

ETHICAL APPROVAL

All authors declare that all experiments have been examined and approved by the appropriate ethics committee.

References

1. Davies L. and Welch H.G. The increasing incidence of thyroid cancer in the United States, 1973-2002'. *JAMA*. 2006; 295(18):2164-2168.
2. Kulldorff. Cancer map patterns: are they random or not? *American Journal of Preventive Medicine*.
3. Huang L., Pickle L.W., and Das B. 2008. Evaluating spatial methods for investigating global clustering and cluster detection of cancer cases. *Statistical Medicine* 27(25):5111-5142.
4. Wang D, Feng J, ZengP, Yang Y, Luo J and Yang Y. 2011. Total oxidant/antioxidant status in sera of patients with thyroid cancers'. *Endocrine-Related Cancer* 18 773-782.
5. Ames B.N. 1988. International Agency for Research on Cancer, *SciPubl* 89,407-416.
6. Andryskowski G and Owczarek T. 2007. The evaluation of selected oxidative stress parameters in patients with hyperthyroidism'. *Polish Archives of Internal Medicine* 117, 285-289.
7. Aslan M, Cosar N, Celik H, Akosoy N, DulgerAC, Soyoral YU, Kucukoglu ME and Selek S. 2011. Evaluation of oxidative status in patients with hyperthyroidism'. *Endocrine* 40, 285-289.
8. Erdamar H, Demirci H, Yaman H, Erbil MK, Yakar T, Sancak B, Elbeg S, Biberoğlu G and Yetkin I. 2008. The effect of hypothyroidism, hyperthyroidism, and their treatment on parameters of oxidative stress and antioxidant status'. *Clinical Chemistry and Laboratory Medicine* 46:1004-1010.
9. Lu H., Ouyang W. and Huang C. 2006. 'Inflammation, a key event in cancer development', *Molecular Cancer Research* 4: 221-233.
10. Spencer JP (2005). Interactions of flavonoids and their metabolites with cell signaling cascades. In *Nutrigenomics* 353-377.
11. Saad-Hussein A, Hamdy H, Aziz HM and Mahdy-Abdallah H. 2011. Thyroid functions in paints production workers and the mechanism of oxidative-antioxidants status. *Toxicology and Industrial Health* 27:257-263.
12. Misra HP and Fridovich I. 1972. The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase. *J BiolChem* 247: 3170-5.
13. Rotruck JT, Pope AL, Ganther HE, Swanson AB, Hafeman DG and Hoekstra W.G. 1973. Selenium: biochemical role as a component of glutathione peroxidase.
14. Beutler E, Duron O and Kellin B.M. 1963. Improved method for the determination of blood glutathione. *J. Lab.Clin. Med* 61: 882-888.
15. Habig WH, Pabst MJ, Jakoby W.B. 1974. Glutathione S-transferase. The first systemic step in mercapturic acid formation. *J BiolChem* 246: 7130-7139.
16. Benzie, F.F. and Strain, J.J. 1999. Ferric Reducing/ Antioxidant Power Assay: Direct Measure of Total antioxidant Activity of Biological Fluids and Modified Version for Simultaneous Measurement of Total Antioxidant Power and Ascorbic Acid Concentration. *Methods in enzymology* 299:15-23.
17. Harma M, Harma M and Erel O. 2005. Measurement of the total antioxidant response in preeclampsia with a novel automated method. *Eur J. Obstet. Gynaecol. Reprod. Biol.*, 10: 47-51.
18. Varsheny R and Kale R.K. 1990. Effect of calmodulin Antagonist. *Int. J. Rad. Biol*, 58:733-743.
19. Wolff 1999. Spectrophotometric measurement of hydroperoxides at increased [sensitivity by oxidation of Fe²⁺ in the presence of xylenol orange.
20. Kaneko J.J. 1999. *Clinbiochem of animals*. 4th ed. New York; Academic Press Inc; p.932.
21. Erdamar H, Cimen B, Gülcemal H, Saraymen R, Yerer B, Demirci H. 2010. Increased lipid peroxidation and impaired enzymatic antioxidant defense mechanism in thyroid tissue with multinodular goiter and papillary carcinoma. *Clinical Biochemistry* 43, 650-654.
22. Jemal A, Siegel R and Ward L. 2006. Cancer Statistics, *Cancer J Clin* 56: 106-130.
23. Lassoued S, Mseddi M, Mnif F, Abid M, Guermazi F, Masmoudi H, ElFeki A and Attia H. 2010. A comparative study of the oxidative profile in Graves' disease, Hashimoto's thyroiditis, and papillary thyroid cancer. *Biological Trace Element Research* 138:107-115.
24. Anlasik T, Sies H, Griffiths HR, Mecocci P, Stahl W, and Polidori M.C. 2005. Dietary habits are major determinants of the plasma antioxidant status in healthy elderly subjects. *Brit J Nutr*, 94:639-42.
25. Ross SA. 2010. Evidence for the relationship between diet and cancer. *Exp. Oncology*, 32(3): 137-42.

26. Barnard ND, Nicholson A, Howard JL. 1995. The medical costs attributable to meat consumption. *Prev Med*; 24:646-655.
27. Shihadeh A and Saleh R. 2005. Polycyclic aromatic hydrocarbons, carbon monoxide, "tar" and nicotine in the mainstream smoke aerosol of the narghade water pipe.
28. Diplock AT, Charleux JL, Crozier WG. 1998. Functional food science and defence against reactive oxygen species. *British Journal of Nutrition*; (80 Suppl 1): S77- S112.
29. Davies CD, Tsuji PA, Milner JA. 2012. Selenoproteins and cancer prevention. *Annual Review of Nutrition*; 32: 73-95.
30. Thorogood M, Mann J, Appleby P, Mipherson K. 1994. Risk of death from cancer and ischaemic heart disease in meat and non meat eaters. *B.Med J*.308: 1667-1670.
31. World Cancer Research Fund 2007. Food, nutrition, physical activity and the prevention of cancer, A global perspective. American Institute of cancer Research Washington DC