

THE PATTERN OF FEMALE GENITAL MUTILATION IN PORT HARCOURT, SOUTHERN NIGERIA.

Abstract:

Background:

Female genital mutilation is a harmful traditional practice which is an infringement on the sexual and reproductive rights of women and girls and has profound psychosocial as well as reproductive health morbidities. This study sought to determine the pattern of female genital mutilation in Port Harcourt, South-South Nigeria.

Methods:

A cross-sectional study was carried out on clients seen at the Department of ObstetricGynaecology of the University Port Harcourt Teaching Hospital (UPTH) between 1st January and 31st January 2009. Five hundred clients were selected randomly and agreed to participate in the study. They were interviewed using a structured questionnaire and examined clinically. Data management was with SPSS 15.0 for Windows statistical software.

Results:

The prevalence of female genital mutilation (FGM) was 34%. Four hundred and eighty one (96.2%) were aware of female circumcision generally. Ninety five

(55.8%) of those circumcised, had it done in infancy. Thirty seven percent of the practitioners of female circumcision were traditional birth attendants while 14.7% were trained health professionals. The commonest reason for female circumcision included reduction sexual passion/promiscuity and conformity with tradition. Type 1 female genital mutilation was the commonest (58.2%). Among those clients aged 50 years and above, 78.8% had FGM while 9.1% of those aged 10 – 19 years had FGM. This showed a downward trend.

Conclusion:

Female circumcision is a harmful traditional practice which has remained a serious health problem. Its prevalence is still high in our environment. This study suggests that it is on a downward trend.

Keywords: Female genital mutilation, patterns, trends.

INTRODUCTION:

Female Genital Mutilation (FGM) also known as female circumcision or female genital cutting comprises all procedures involving partial or total removal of the external female genitalia or other injury to the female genital organs whether for cultural, religious or other non-therapeutic reasons¹⁻³. It is a harmful traditional practice which is an infringement on the sexual and reproductive rights of women and girls⁴.

43 About 130million women and girls worldwide have undergone FGM, and
44 about 2 million girls are at risk each year^{1,2,5}. It crosses all religious, racial and
45 social boundaries. In Africa, particularly in countries such as Nigeria, Ethiopia,
46 Sudan, Egypt and Somalia, FGM is widespread and a highly valued ritual ⁶. In
47 Nigeria, national prevalence rate of 50-60% has been reported and the practice
48 is said to be commoner in the South -East, South-West and South-South regions
49 of the country ^{7,8}.

50 Various reasons have been given for the perpetuation of the practice of
51 FGM. These have been categorized into psychosexual, sociological, hygiene
52 and aesthetic, mythical and religious reasons. These include prevention of sexual
53 promiscuity/infidelity in marriage, identification with cultural heritage, initiation of
54 girls into womanhood, promotion of hygiene and aesthetic appeals of the female
55 external genitalia and enhancement of fertility and child survival^{1,9,10}.

56 Female genital mutilation is associated with profound reproductive health
57 morbidities and mortality. Immediate complications include severe pain,
58 haemorrhage, shock, urinary retention, ulceration of the genital region, injury to
59 adjacent tissues, tetanus and death. Long term complications include cysts and
60 abscesses, keloid formation, urinary incontinence, gynaetresia, sexual
61 dysfunction, infertility, prolonged obstructed labour and higher rate of episiotomy
62 and caesarean sections ¹¹⁻¹⁴.

63 This study sought to determine the pattern of female genital mutilation in
64 Port Harcourt, South-South Nigeria. It is a hospital based study and offers
65 indispensable opportunities for enlightenment and education and intervention

strategy to change the concept and behaviour of those who were hitherto naïve or misinformed on the subject. Therefore it seeks to reduce the subsequent rate of female circumcision. Studies that are community based are punctuated by scepticism on candid or true responses to avoid shame and condemnation. This study has physically corroborated interview responses with clinical examination findings.

SUBJECTS AND METHODS

This was a cross-sectional study carried out among clients seen at the Department of Obstetrics and Gynaecology (antenatal clinic, gynaecology clinic, family planning clinic, labour ward, accident and emergency unit) of the University of Port Harcourt Teaching Hospital, South-South Nigeria. The required sample size was derived using the formula: $n = z^2pq / d^2$ where z is the standard normal deviate, usually set at 1.96 (or more simply at 2.0), which corresponds to the 95% confidence level; p represents the prevalence of FGM set at 50%, q is $1.0 - p$ and d is the margin of error tolerable (or degree of accuracy desired) which is set at 5%. This gave a sample size of 384. The participants were selected randomly and informed of the study, and all agreed to participate.

Data was collected using an interviewer-administered structured questionnaire between 1st January and 31st December 2009. Information obtained included socio-demographic characteristics, awareness, attitude and practice of FGM. This study had a team who physically examined and reviewed all the clients with a view to corroborating the interview responses with clinical

examination findings. The data collected were coded and entered into a computer spreadsheet using SPSS 15.0 for Windows statistical software. Results were presented as means with standard deviations, rates and proportions, graph, tables and figures. Chi-square tests were carried out where necessary. Cross tabulation was performed to establish relationships among variables. These were reported when statistically significant at p value of ≤ 0.05 .

RESULTS:

The sociodemographic characteristics of the respondents were shown in table 1. The mean age in years was 31.57 ± 9.419 . The minimum age was 10 years and the maximum was 76 years. Thirty percent of the respondents were housewives while 79 (15.8%), 56 (11.2%) 101 (20.2%), and 99 (19.8%) were civil servants, teachers, traders and students respectively. Clients who had tertiary education constituted 66.2%. Respondents who were married made up 90.6% while 43% of the respondents were nulliparae.

Out of the 500 clients who participated in this study, 170 had female genital mutilation this gave a prevalence of 34% as shown in figure 1, 0.4% of respondents were circumcised but did not know about it.

Figure 2 showed that 96.2% of the respondents were aware of FGM. It was observed that 55.8% of those who were circumcised had it done during infancy. Thirty seven percent of the practitioners (perpetuators) of FGM were

traditional birth attendants while 14.7% were trained health professionals (doctors, nurses and midwives) as shown in figure 3.

The commonest reasons for FGM were reduction of sexual passion and promiscuity, and conformity with tradition. Other reasons included rite of passage to womanhood, ensuring virginity and chastity in marriage and narrowing/tightening of the vaginal introitus for maximal sexual performance. This was shown in figure 4. The commonest complications following FGM were decrease in sexual desire or arousal, dyspareunia, dysmenorrhoea, episiotomies/perineal tears especially during their first deliveries. Some of them had caesarian section for obstructed labour/rigid perineum.

Type 1 (58.2%) was the commonest type of FGM, followed by type 2 (40.0%). None of the respondents had type 3, while 1.8% had type 4.

Majority of the respondents (99%) would not recommend FGM for their daughters or wards. The reasons for not recommending FGM included complications of FGM, knowledge of the efforts of governments and various bodies including NGOs to ban FGM, and the practice not being done in some communities/cultural groups.

The issue of sanctions / benefits from FGM was also studied. Out of the 500 clients, 481 (96.2%) responded that they were not aware of sanctions in their communities against those who refused circumcision, while 19 (3.8%) responded that they were aware of sanctions. The sanctions included payment of fines (money, goats, yam), ostracism and denial of burial rights. Similarly 487 (97.4%) responded that they were not aware of any benefits in their communities for

those who were circumcised, while 13 (2.6%) responded that benefits were accruable to these who were circumcised. The benefits included provision of crops, allocation of land and special burial rights.

It was observed that 48% of the respondents were aware of the efforts of governments and NGOs to eradicate FGM. Also 62.4% of the respondents had other means of livelihood.

Figure 5 showed the trends in circumcision, where 78.8% of those aged 50 years and above had FGM while 9.1% of those aged 10-19 years had FGM. This showed that the trend is on a decline.

DISCUSSION

The prevalence of FGM was 34%. This is different from other studies^{2,9} with higher prevalence. While 78.8% of those aged 50 years and above had FGM, only about 9.1% of those in the 10-19 years -age group had FGM. The trend here showed that there is a decline in FGM- This is similar to a study in South West Nigeria ⁶. Majority of the respondents had tertiary education. This is similar to other studies ^{7,11}. This high level of education will impact on their knowledge, attitude and practice of FGM. It is not surprising then that 99% of them would not recommend FGM for their daughters or wards. These were probably empowered to resist the cultural norms/pressure to circumcise their children.

Out of the 500 respondents 96% were aware of FGM. This was similar to a study in Kwara State ¹⁵. Among those who were circumcised 55 % had it done

as infants. This is the commonest age group in which it is done in this part of the country. Traditional birth attendants constituted 37% of the practitioners of FGM. However, 14.7% of the practitioners were trained health professionals who included doctors, nurses and midwives. This showed that there was a high level of medicalisation of FGM as was found in a similar study¹⁶. The world Health Organization has banned medicalisation of FGM¹.

Reduction of sexual passion and promiscuity, and conformity with tradition were the commonest reasons for FGM. However, some studies showed that FGM does not significantly affect women's sexual behavior^{17,18}.

The common gynecological complications of FGM observed in this study were dyspareunia, dysmenorrhoea and decrease in sexual desire or arousal. However other studies^{12, 14} showed other complications like chronic pelvic pain, pelvic inflammatory disease, infertility, urinary retention or urinary tract infection, acquired gynaetresia. This may be explained by the fact that only types I and II were mostly performed here. Type 3, which is implicated in most of the severe gynaecological complications was not observed in this study.

Only 48% of the respondents were aware of the efforts of the various governments, organs of the United Nations, International federation of Gynaecology and Obstetrics (FIGO), women organizations and the NGO's to eradicate FGM. This showed that a lot of education and enlightenment of the people have to be done. The religious bodies, traditional rulers, health providers, women organizations, the government and mass media are all stake holders for the mobilisation of the people against FGM. FGM was found to be commoner

among the Ibos, Yorubas and Ijaws than the Hausa/Fulani. This was similar to some studies in Northern Nigeria^{8,15}. Majority of the practitioners of FGM had other means of livelihood. This is a positive development towards the efforts to reduce the practice as these practitioners do not need to find new jobs when they eventually stop performing FGM. Medical practitioners should be reminded of the ban on medicalisation of FGM by the World Health Organization^{1, 19}.

CONCLUSION

Female genital mutilation is a harmful traditional practice, which has remained a serious public health problem. This study suggests that it is on the decline. However, more efforts still have to be made by the governments and other bodies involved in the fight against the practice. Various governments should pass legislation and enforce them. Religious bodies, traditional rulers, community development committees, health providers and the mass media still have big roles to play in this fight. Medicalisation of FGM is an emerging problem which needs to be curbed. Appropriate sensitization and advocacy through faith based religious organizations may reverse the concept and practice.

REFERENCES

1. World Health Organization (WHO). Female genital mutilation. Fact sheets No. 241 June 2000.
2. World Health Organization (WHO). Gender and Women's Health Department. Female genital mutilation: report of a WHO technical working group Geneva; 17-19 July 1995.

- 200 3. Sakeah E, Beke A, Doctor HV, Hodgson AV. Males' preference for
201 circumcised women in Northern Ghana. *Afr Reprod Health* 2006; 10 (2):
202 37-47.
- 203 4. Adinma JIB. Overview of the global policy consensus on women's sexual
204 and reproductive rights: the Nigerian perspective. *Trop. J. Obstet.*
205 *Gynaecol.* 2002; 19(suppl. 1): S9
- 206 5. Braddy CM, Files JA. Female genital mutilation, Cultural Awareness and
207 Clinical considerations. *J Midwifery Women's Health* 2007; 52: 158 –163.
- 208 6. Adeokun LA, Oduwole M, Oronsaye F, Gbogboade A. O, Aliyu N,
209 Adekunle W. Sadiq G, Sutton I, Taiwo M. Trends in female circumcision
210 between 1933 and 2003 in Osun and Ogun States, Nigeria: a cohort
211 analysis. *Afr J Reprod Health* 2006; 10(2): 48 – 56.
- 212 7. Sule ST. Female genital mutilation: the enormity of the problem. *Nig J*
213 *Med* 1997; 6:54-56.
- 214 8. Abubakar 1, Iliyasu Z, Kabir M, Uzoho CC, Abdulkadir MB. Knowledge,
215 attitude and practice of female genital cutting among antenatal patients in
216 Amino Kano Teaching Hospital, Kano. *Nig J Med* 2004 Jul-Sep;13 (3):250
217 -3.
- 218 9. Mandara MU. Female genital mutilation in Nigeria. *Int J Gynaecol Obstet*
219 2004; 84:291 – 298.
- 220 10. Oguntuyi A. History of Ekiti. Ibadan: Bisi Books, 1979. pp 28-29.
- 221 11. De Silva S. Obstetric sequelae of female circumcision. *Eur J Obstet*
222 *Gynaecol Reprod Biol* 1989; 32:233-240.
- 223 12. Ozumba BC. Acquired Gynaetresia in Eastern Nigeria. *Int J Gynaecol*
224 *Obstet* 1992; 37:105 –109.
- 225 13. Tindall VR. Injuries. In: Jeffcoate's Principles of Gynaecology 1987. Fifth
226 Edition. Butterworth – Heinemann pp. 238-259.
- 227 14. Larsen U, Okonofua FE. Female circumcision and obstetric complications.
228 *Int J Gynaecol Obstet* 2002; 77: 255 – 65.

- 229 15. Aboyeji AP, Abubakar D, Ijaiya MA, Nwachukwu D. Female genital
230 mutilation in rural areas of Kwara State. Trop J Obstet Gynaecol 2005; 22
231 (supp 1); S30.
- 232 16. Ugboma HA, Akanni CI, Babatunde S. Prevalence and medicalisation of
233 female genital mutilation. Nig J Med 2004 Jul-Sep; 13(3):254-8.
- 234 17. Msuya SE, Mbizvo E, Hussain A, Sundby J, Sam, NE. Stray – Pedersen
235 B. Female genital cutting in Kilimanjaro, Tanzania: changing attitudes?
236 Trop Med Int Health 2002; 7:159-65.
- 237 18. Elmusharaf S, Elkhidir I, Hoffman S, Almroth L. A case control study on
238 the association between female genital mutilation and sexually transmitted
239 infections in Sudan. BJOG 2006; 113:469 – 74.
- 240 19. Adinma JIB, Agbai AO. Practice and perceptions of female genital
241 mutilation among Nigerian Igbo women. J. Obstet. Gynaecol. 1999; 19(1):
242 44-48

Table 1: Socio-demographic characteristics

Characteristics	Number	Percentage %
Age (mean + SD)	31.57years $\pm$ 9.42	
10-19	33	6.6
20-29	199	39.8
30-39	192	38.4
40-49	43	8.6
≥ 40	33	6.6
Parity		
0	218	54.25
1	90	23.50
2	73	13.25
3	51	5.50
4	39	2.75
≥ 5	29	0.75
Educational status		
No formal education	1	0.2
Primary	18	3.6
Secondary	150	30.0
Tertiary	331	66.2
Occupation		
Housewives	150	30.0
Civil servants	79	15.8
Traders/Business women	101	20.2
Student	99	19.8
Doctor /Nurse / Pharmacist	15	3.0
Teacher	56	11.2
Religion		
Christian	497	99.4
Islam	3	0.6
Marital status		
Married	453	90.6
Single	41	8.2
Divorced	3	0.6
Widowed	3	0.6

Table 2: Tribe / Ethnic group.

Tribe/Ethnic group	No.	Percentage %
Ijaw	66	13.2
Igbo	252	50.4
Ikwerre/Etche	65	13.0
Yoruba	20	4.0
Hausa/Fulani	15	3.0
Bini	15	3.0
Others	67	13.4
Total	500	100

Figure 1: Proportion Circumcised




Figure 2: Awareness of circumcision




Figure 3: Practitioner




Figure 4: Reason for circumcision





Figure 5: Trends in circumcision



464
465
466

UNDER PEER REVIEW