

MALARIA PARASITAEMIA ASSOCIATION WITH ABO BLOOD GROUPS AMONG STUDENTS OF A PRIVATE UNIVERSITY IN WESTERN DELTA, NIGERIA

ABSTRACT

The focus of this work was to investigate any relationship between ABO blood types and malaria parasitaemia among students of a private University based in the western part of Delta state, Nigeria. Venous blood of 2ml volume was obtained by venepuncture from 360 students made up of 150 (41.7%) males and 210 (58.3%) females of 28 years average and who were both symptomatic and asymptomatic for malaria due to *Plasmodium falciparum*. Malaria parasite screening was done by both *P. falciparum* antigen rapid (Micropoint, USA) test and Giemsa staining. ABO blood typing was done using Monoclonal Antisera A, B and D. One hundred and forty one (41.6%) male and 198(58.4%) female samples was rhesus positive. Nine (42.9%) and 12 (57.1%) males and females were rhesus negative. ABO blood group frequency occurrence was 55.8% (O), 22.5 % (A) 18.3 % (B) and 3.4 % (AB). A total of 255 (70.8%) students were infected with *P.falciparum* parasites of which 55.3% and 44.7% were females and males respectively. ABO blood group malaria parasitaemia frequencies were 76.4% (O), 56.3% (B), 52.4% (A) and 0.0% (AB) for non-severe malaria and 70.5% (O), 58.3% (A), 55.6% (AB) and 50.0 % (B) for severe plasmodiasis. Whereas there was significant association between malaria infection and gender ($P < 0.05$) there was no significant association between severe and non-severe malaria parasitaemia and ABO blood types ($P > 0.05$). The implications of these findings are discussed.

32

33

34

35

36

37

Introduction

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

Malaria (which is caused by *Plasmodium* species transmitted by anopheles mosquitoes) is the second leading disease following Acquired Immunodeficiency Syndrome-AIDS (Uneke, 2006) associated with high morbidity and mortality through anaemia, cerebral complications and other mechanisms (Uneke, 2006).

Despite the high morbidity and mortality, certain individuals living in malaria endemic regions appear relatively protected compared to those who suffer frequent severe malaria attacks. Resistance to malaria infection is dependent on the development of an immune response by the host and to a varying extent, a certain innate characteristic possessing protective value against infection (Cserti and Dzik, 2007). These factors include Sickle cell trait (HbAS) and Sickle cell disease –HbSS (Alouch, 1997), ABO blood types (Agbonlahor *et al.*, 1993) and the level of G-6-P-dehydrogenase (Ruwende and Hill, 1998).

It is thought that an understanding of the nature of relationship (if any) between ABO blood groups and malaria parasitaemia should provide an invaluable window in the scourge and that studies of malaria parasitaemia from that stand point in populations of malaria endemic regions will be helpful in elucidating any such relationship (Ilozumba and Uzozie, 2009)

56 A high percentage of severe malaria cases have been reported among blood
57 group A individuals (Fischer and Boone, 1998; Lell *et al.*, 1999). Furthermore, Migot-
58 Nabias (2000) and Pathirana *et al.* (2005) observed low paraitaemia and
59 uncomplicated malaria cases among blood group O individuals. Similarly, a
60 significant advantage of *Plasmodium falciparum* with ABO blood groups with respect
61 to group O has been reported by some authors (Cserti – Gazdewich *et al.*,
62 2012; Igbineghu *et al.*, 2012; Nasr, 2013). Other studies have shown high frequency of
63 malaria episodes among blood group A individuals as compared with other blood
64 group individuals (Beigulman *et al.*, 2003; Akhigbe *et al.*, 2011) reported that blood
65 group AB persons had the lowest malaria attack while blood group A persons had the
66 highest attack.

67 Studies to investigate any possible association between ABO blood group
68 system and some disease conditions have been carried out by some authors (Tursen *et*
69 *al.*, 2005; Kassim and Ejezie, 1982; Opera, 2007; Abdulazeez *et al.*, 2008; Ndambaa
70 *et al.*, 1997; Blackwell, 2002). Some of these studies reported significant associations
71 thereby suggesting that ABO blood groups have an impact on the infection status of
72 the individuals possessing a particular ABO blood group (Opera, 2007; Abdulazeez *et*
73 *al.*, 2008; Ndambaa *et al.*, 1997, Blackwell, 2002).

74 The absence of significant association between *P. falciparum* prevalence
75 and ABO antigens has also been reported by some other studies (Uneke *et al.*,
76 2006; Herrera *et al.*, 2009; Boel *et al.*, 2012; Igbineghu *et al.*, 2012). Other authors are
77 less equivocal in relating malaria parasitaemia to ABO blood group (Senga *et al.*,
78 2007; Rowe *et al.*, 2008). Otajevwo (1997) in a study associating malaria parasitaemia

79 with ABO blood types among residents of Warri, Nigeria, reported that 6.9%, 19.0%,
 80 20.7% and 53.3% of a total of 174 whole blood samples processed belonged to blood
 81 types AB, B, A and O respectively and 138(79.3%) of total sample size were infected
 82 with malaria parasites of which *P. falciparum* was the predominant species. In the
 83 report, the highest malaria parasite load was observed among group O (52.2%)
 84 individuals while the least was noticed among blood group AB (8.7%) individuals.
 85 Malaria parasitaemia was higher among the males (83.3%) than females (75.0%).
 86 Otajevwo (2013) also reported ABO blood types among students of Igbinedion
 87 University, Okada, Nigeria of which out of 104 samples analysed, 44(3.9%),
 88 16(15.4%), 32(30.8%) and 52(50.0%) students occurred in AB, A, B and O groups
 89 respectively. On the whole, 80(76.8%) of total samples processed, were positive for
 90 malaria parasitaemia.

91 Nkuo-Akenji (2004) reported that blood group O individuals maybe more
 92 susceptible to malaria attack. A study conducted in Edo State University, Nigeria
 93 reported that blood groups O and B male individuals were the most and least
 94 susceptible to malaria attack respectively (Agbonlahor *et al.*, 1993). In a study
 95 conducted on inhabitants of Odakpu, Anambra State ,Ilozumba and Uzozie (2009)
 96 reported ABO blood group prevalence of 2.63%,12.05%,21.1% and 63.8% for groups
 97 AB,B,A, and O respectively .Zerihun *et al.*(2011) reported that A, B and AB blood
 98 group individuals are more vulnerable to *P. falciparum* infection compared to blood
 99 group O individuals in a study they carried among 269 febrile Ethiopian out patients
 100 who visited a health care .

Despite the above researches, a consensus on possible association between ABO blood group genes and malaria infection is still lacking (Omotade *et al.*, 1999). This might be due to limited data or unreported data on the cell antigens (Nkuko-Akenji, 2004), peculiar demographic distributions and characteristics of study areas (Herrera *et al.*, 2009). A definite statement on the apparent trend can be made therefore, if results of more studies on ABO blood groups associated with malaria infection in different parts of Nigeria and other neighbouring countries are made available and accessible (Ilozumba and Uzozie, 2009).

This work is a response to the call for further research in the area is aimed at studying ABO blood type association with malaria parasitaemia among students of a private University based in Western Delta, Nigeria with the following objectives:

1. Determine the sex distribution of malaria parasitaemia among students recruited for the study.
2. Determine the distribution of malaria parasitaemia among ABO blood group types of male students
3. Determine the distribution of malaria parasitaemia among ABO blood types of female students.
4. Determine the distribution of severe malaria parasitaemia among ABO blood types of male students.
5. Determine the distribution of severe malaria parasitaemia among ABO blood types of female students.

124

125

126

127

128

129

130 **Materials and Methods**

131

132 **Ethical Clearance**

133 Informed consent was given by students recruited for the study .A consent
134 form (in form of a questionnaire) was given to students who were all adults to fill and
135 sign. Participating students also gave their verbal informed consent by presenting
136 themselves for blood sample collection

137 **Sampling**

138 Two millilitres (2ml) of venous blood was collected from 360 randomly
139 selected Western Delta University students made of 150(41.8%) males and
140 210(58.2%) females from various departments of the University .Blood samples were
141 collected with the aid of 5ml sterile needles and syringes by venipuncture (by tying a
142 tourniquet around the upper arm and sterilizing it with 70% ethanol to increase blood
143 pressure in the veins) and dispensed into ethylene diamine tetra acetic acid (EDTA)
144 anticoagulant bottle containers. Blood and blood containers were properly mixed by
145 standard method and labelled appropriately. Both symptomatic and asymptomatic
146 subjects were used for the study. Symptomatic subjects were those malaria parasite

test were positive and showed obvious signs of illness .Asymptomatic subjects were those who were apparently healthy although positive for malaria parasite test. For clarity, symptomatic subjects were grouped as severe malaria parasitaemia while asymptomatic as non-severe malaria parasitaemia.

Malaria Parasite Detection Tests

The Malaria *Plasmodium falciparum* rapid test Device (whole blood) Micropoint package insert kit by Cook et al. (1999) was used to test for the presence of specific antigens (proteins) produced by *falciparum malaria* parasites (Murray, 2008; WHO,2011). Antigen targeted is *P. falciparum* specific histidine rich protein -2 (i.e. PfHRP -2) and aldolase (Wongsrichanalai, 2007; Murray, 2008).

Rapid (Micropoint, USA) Test Procedure

The test device, specimens, buffer and /or controls were allowed to equilibrate to 15-30⁰ C (room temperature) prior to testing. The test device cassette was removed from the foil pouch and used immediately. Using the sample loop (or rubber pipette) provided, 5ul (0.05ml) of whole blood was dropped into sample well (A). Two drops (60ul or 0.06ml) of assay buffer were put in well B and the set up was left to stand for 20minutes after which result was read. A positive result was one in which a pink or purple coloured control(C) band appeared in addition to a distinct pink coloured band in the test (P.f) region. In a negative result, only one pink or purple band appeared on the control (C) region while no apparent band appeared on the test (P.f) region. The

quality control region was to confirm if sufficient specimen volume was used and if the correct procedural technique was applied.

Sensitivity of Test Device

The Malaria P.f Rapid (Micropoint) test device (whole blood) has been tested in comparison with the thick blood smears of clinical samples. The results show that the sensitivity of the test device (whole blood) is > 99.0% relative to blood smears.

Specificity

The test device uses an antibody that is highly specific for Malaria *Plasmodium falciparum* antigen in the blood. The test device has a specificity of >99.0% relative to blood smears

Expected

The test device has been compared with traditional thick or thin blood smears microscopic analysis. The correlation between the two systems is >99.0%.

Limitation

The test device is for in-vitro diagnostic test use only. Neither the qualitative value nor the rate of increase in Pf antigen concentration can be determined by its qualitative test. The test Device will only indicate the presence of *Plasmodium falciparum* antigen and hence, the specimen should not be used as the sole criteria for the diagnosis of malaria infection. All results must be interpreted together with other clinical information available to the physician. A negative result does any time prelude the possibility of malaria infection.

Giemsa Staining

This manual staining was carried out as quality control of the rapid test method. Thick blood films were made on grease free microscope slides (after appropriate labelling and allowed to air dry on laboratory working bench. Slides were arranged on a staining rack and flooded with 10% (v/v) Giemsa stain solution for 15minutes (Brooks *et al.*, 2004). The plus system was used for the determination of parasite density (WHO, 1990). Only positive results with the rapid test device and Giemsa staining method were recorded and used for ABO group typing.

Typing Blood Samples for Blood Group Antigens

The ABO blood group of each subject was determined using cell grouping Antisera according to methods described by Rosenfield (1976) and Cheesbrough (2000). Monoclonal Antisera A, B and D (Agape Diagnostics, Ltd, India) were used.

Statistical Analysis of Data

Chi square (χ^2) analysis using test of independence of two characters or associations using a 4 x 3 contingency table at 95% confidence interval was used. The software used was the statistical package for social sciences (SPSS) version 17.0. Confidence interval at 95% (0.05) was calculated using mean + or - 0.05 S_x where 6 refers to degree of freedom calculated using (r-1) (c-1)

216

217

218

219

220

221

222

223

224

225

Results

226

227

228

229

230

231

232

233

234

235

236

237

238

239

In **Table 1**, the sex, department and age group distribution of students recruited for the study are shown. Students of fifteen (15) departments gave their informed consent and volunteered their blood samples as shown. Seventy two (20.0%) ,66(18.3%), 54(15.0%) ,33(9.2%), 30(8.3%) ,30(8.3%) ,18(5.0%) ,18(5.0%) ,12(3.3%) ,6(1.7%) ,6(1.7%) ,6(1.7%) ,3(0.8%) ,3(0.8%) and 3(0.8%) volunteering students of Microbiology/Biotechnology ,Accounting ,Geology, Economics, Biochemistry , Political Science , Mass Communication, Business Administration ,Computer Science , Chemistry ,Physics, Library science , Philosophy , Hotel and Tourism and Environmental Science departments respectively engaged in this study are also shown in the descending order . Out of the 360 sample size, 150(41.7%) and 210(58.3%) were males and females respectively. Table 1 also shows the age brackets of students used for the study. In decreasing order, 198(55.0%) , 111(30.8%) , 39 (10.8%) , 6(1.7%) , 3(0.8%) and 3(0.8%) number of students occurred in 21-25 (average age 23years) ,15-20(average age 19 years) , 26-30(average age 28 years),31-

240 35(average age 33 years) , 36-40(average age 40) and 41-45 (average age 44 years)
 241 Again , out of the total recruited in each age group, 90(45.5%) , 33(29.7%), 15
 242 (38.5%) , 6(100.0%),3(100.0%) and 3(100.0%) represented male students in 21-25,
 243 15-20, 26-30, 31-35, 36-40 and 41-45 age brackets respectively in that descending
 244 order while 108(54.5%) ,78(70.3%), 24(61.5%) ,0(0.0%), 0(0.0%) and 0(0.0%)
 245 represented female students in the same age group respectively

246 **Table 1:** Sex and Age Distribution OF Western Delta University Recruited For the Study

Department	Sex	15-20 Average=19 N=111(30.8%)	21-25 Average=23yrs N=198(55.0%)	26-30 Average=28 N=39(10.8%)	31-35 Average=33 N=06(1.7%)	36-40 Average=40 N=03(0.8%)	41-47 Average=44 N=03(0.8%)
Geology N=54(15.0%)	M=36 F=18	06 12	27 03	00 03	03 00	00 00	00 00
Microbiology N=72(20.0%)	M=21 F=51	12 09	09 33	00 09	00 00	00 00	00 00
Accounting N=66(18.3%)	M=24 F=42	09 15	12 24	03 03	00 00	00 00	00 00
Mass comm. N=18(5.0%)	M=03 F=15	00 00	00 06	00 00	00 00	03 00	00 00
Computer.Sci N=12(3.3%)	M=06 F=06	00 03	03 03	03 00	00 00	00 00	00 00
Bus.Admin N=18(5.0%)	M=06 F=12	00 00	03 12	03 00	00 00	00 00	00 00
Biochemistry N=30(8.3%)	M=03 F=27	00 21	03 06	00 00	00 00	00 00	00 00
Economics N=33(9.2%)	M=18 F=15	03 03	15 09	00 03	00 00	00 00	00 00
Pol. Science N=30(8.3%)	M=21 F=09	03 06	12 03	03 00	00 00	00 00	03 00
Env. Science N=03(0.8%)	M=03 F=00	00 00	03 00	00 00	00 00	00 00	00 00
Chemistry N=06(1.7%)	M=03 F=03	00 00	03 03	00 00	00 00	00 00	00 00
Physics N=06(1.7%)	M=03 F=03	00 00	00 00	03 03	00 00	00 00	00 00
Hotel&tourism N=03(0.8%)	M=00 F=03	00 00	00 03	00 00	00 00	00 00	00 00
Philosophy N=03(0.8%)	M=03 F=00	00 00	00 00	00 00	03 00	00 00	00 00
Library sci. N=06(1.7%)	M=00 F=06	00 00	00 03	00 03	00 00	00 00	00 00
TOTAL	M=150(41.7%) F=210(58.3%) 360	33(29.7%) 78(70.3%) 111(30.8%)	90(45.5%) 108(54.5%) 198(55.0%)	15(38.5%) 24(61.5%) 39(10.8%)	06(100.0%)0(0.0%) 06(1.7%)	03(100.0%) 0(0.0%) 03(0.8%)	03(100.0%) 0(0.0%) 03(0.8%)

Table 2 shows data on ABO blood groups and Rhesus factor frequency distribution among students studied. A total of 360 students' blood samples made up of 150(41.5%) and 210 (58.3%) males and females respectively were processed for ABO blood typing. Out of this sample size, 141(41.6%) male samples were rhesus positive. Nine (42.9%) and 12(57.1%) male and female samples respectively were rhesus negative. In terms of male ABO blood grouping, 6(4.0%), 18(12.0%), 36(24.0%) and 90(60.0%) male students were grouped into blood groups AB, B, A and O respectively in that ascending order. With regards to the female students, 6(2.9%), 45(21.4%), 48(22.9%) and 111(52.9%) female students were grouped into blood groups AB, A, B and O respectively in that ascending order.

Table 2: ABO blood groups and rhesus frequency distribution among students

Recruited for study

ABO blood types	Sex	Rhesus Positive (%)	Rhesus Negative (%)	Total
A	M	33(23.4%)	03(33.3)	36(24.0)
	F	39(19.7%)	06(66.7)	45(21.4)
B	M	18(12.8)	00(0.0)	18(12.0)
	F	45(22.7)	03(33.3)	48(22.9)
AB	M	06(4.3)	00(0.0)	06(4.0)
	F	06(3.0)	00(0.0)	06(2.9)
O	M	84(59.6)	06(66.7)	90(60.0)
	F	108(54.6)	03(33.3)	111(52.9)
Total	M	141(41.6)	09(42.9)	150(41.7)
	F	198(58.3)	12(57.1)	210(58.3)
		339(94.2)	21(5.8)	360(100.0)

Table 3 shows the sex (gender) distribution of malaria parasitaemia among students engaged in the study. Out of the 360 students recruited, 150(41.7%) and 210(58.3%) were males and females respectively. Out of the same sample size, 255(70.8%) students were infected (positive) with *P. falciparum* parasites in their peripheral blood circulation of which 114(44.7%) and 141(55.3%) were infected male and female students respectively. Table 1 also shows that 105 (29.2%) students were negative for malaria parasites in their blood system of which 36(34.3%) and 69(65.7%) were uninfected male and female students respectively. Malaria parasitaemia was significantly associated with female gender and therefore with sex (χ^2 p-value =3.841, cal. Value= 3.958, $p < 0.05$)

Table 3: Sex distribution of malaria parasitaemia among students recruited for Study

Sex	Positive Malaria parasite (%)	Negative malaria parasite (%)	Total
Males	114(44.7)	36(34.3)	150(41.7)
Female	141(55.3)	69(65.7)	210(58.3)
Total	255(70.8)	105(29.2)	360(100.0)

$$X^2 \text{ df } (1)_{0.05} = 3.841 (\text{P-value})$$

$$\text{Calculated } X^2 = 3.958$$

$$P < 0.05$$

The distribution of malaria parasitaemia and severe malaria parasitaemia among ABO blood types of students data are shown in **Table 4 and 5**. Out of a total sample size of 360 255(70.8%), students were infected (Table 3) of which 206 (67.5%) had non-severe form of malaria parasitaemia and 49(63.6%) students recorded severe form of malaria parasitaemia. Blood group O students recorded the highest non-severe malaria infection rate (76.4%) as well as the highest severe malaria infection rate (70.5%). Although blood type B students recorded the next highest non-severe malaria infection rate (56.3%) blood type, A students recorded the next highest severe malaria infection rate (58.3%) after group O. None of the AB blood type students recorded non-severe malaria infection but 5(55.6%) AB blood group subjects had severe malaria infection rate (50.0%) was recorded for blood group B subjects. Also, whereas 99(32.5%) students cutting across all blood types were uninfected with non-severe form of *P.falciparum* malaria parasites, 28(36.4%) students were uninfected with the severe form. Statistically, there was no significant association of blood types with non-severe form of *P.f* malaria infection (X^2 df (3)_{0.05}=7.815, cal . X^2 =3.408, $P > 0.05$) and there was also no significant association of the blood types with severe form of *P. f* malaria infection (X^2 df (3)_{0.05} = 7.815, cal x^2 =5.652, $P >_{0.05}$

Table 4: Non-severe malaria parasitaemia distribution among ABO blood types
Of students distribution.

Malaria status	A	B	AB	O	Total
Infected (%)	33(52.4)	27(56.3)	0(0.0)	146(76.4)	206(67.5)
Uninfected (%)	30(47.6)	21(43.7)	3(100.0)	45(23.6)	99(32.5)
Total	63(20.7)	48(15.7)	3(1.0)	191(62.6)	305(100.0)

$$X^2 \text{ df } (3)_{0.05} = 7.815 (\text{P-value})$$

$$\text{Calculated } X^2 = 3.408$$

$$P > 0.05$$

Table 5: Severe Malaria Parasitaemia Distribution among ABO blood types of
Students Distribution

Malaria status	A	B	AB	O	Total
Infected (%)	7(58.3)	6(50.0)	5(55.6)	31(70.5)	49(63.6)
Uninfected (%)	5(41.7)	6(50.0)	4(44.4)	13(29.5)	28(36.4)
Total	12(15.6)	12(15.6)	9(11.7)	44(57.1)	77(100.0)

$$X^2 \text{ df } (3)_{0.05} = 7.815 (\text{p-value})$$

$$\text{Calculated } X^2 = 5.652$$

$$P > 0.05$$

Discussion

A total of 360 students from fifteen departments of the University were engaged in the study. Of this sample size, 198 (55.0%) were of the 21-25 age category and 23 years average age. This somewhat suggests that the average age of majority of students in the institution under is 23 years. This is followed by an average age of 19 years as represented by 111 (30.8%) students who occurred in the 15-20 age brackets. This finding implies that the subjects recruited for this study were mature enough to make informed consent to be enlisted for the research.

Findings in this study showed that 339(94.2%) and 21(5.8%) students were rhesus and rhesus negative respectively of which 141(41.6%) and 9(42.9%) male students were rhesus positive and rhesus negative respectively. That there were more rhesus positive and rhesus negative females than males may have been a factor of chance. This finding is however in consonance with the report of a previous study (Otajevwo, 2013). It is to be noted that the 5.8% prevalence of rhesus negative factor appears small in view of the total sample size. This should however not be undermined in view of the far reaching medical implication in terms of child birth and still birth arising from haemolytic disease of newborn (HDN). It also suggests that rhesus negative factor is occurring increasing frequencies from population to population and the need for relevant health care providers as well as Ministry of health to track down people with this factor through compulsory blood typing .Results in this work also reveal that the frequency occurrence of the ABO blood types were 55.8%, 22.5%, 18.3% and 3.4% for group O, A, B and AB respectively

(Table 2). This therefore, confirms that blood group O is the most prevalent while blood AB is the least prevalent and this agrees with the reports of previous studies (Uzoegwu and Onwurah, 2003; Anees and Mirza, 2005; Jeremiah, 2006; Bakare *et al.*, 2004; Odokuma *et al.*, 2007; Adulazeez *et al.*, 2008; Epid *et al.*, 2008; Akhigbe *et al.*, *et al* 2009, 2011; findings in agreement with report of earlier authors Zerihun, *et al*, 2011; Ilozumba and Uzozie, 2009 Otajevwo, 2013) .

These findings are not consistent with an earlier study by Agbonlahor *et al.* (1993) which reported groups O and A as the highest and least occurring respectively. These differences may be due to ethnic, racial and geographical differences of various populations studied.

The findings in this work further revealed that 255 (70.8%) of the 360 students sampled were infected with *Plasmodium falciparum* parasites of various densities .This rather high parasitaemia rate suggests that Oghara town maybe hyper endemic for malaria. This parasitaemia rate appears to be low compared to 93.4% obtained in Odoakpu, Onitsha South by Ilozumba and Uzozie (2009) and 76.8% obtained in okada , Edo state by Otajevwo (2013) as well as 79.3% and 77.4% parasitaemia rates obtained in warri by Otajevwo (1997) and in Owerri by Mbanugo and Emenalo, (2004) respectively. Parasitaemia rate in this study appears high however, when compared to prevalence rates of 66.9%, 58.3%, 43.2%, 10.0% and 6.0% obtained respectively for subjects in Ogbomosho by Akhigbe *et al.* (2011), children in Awka by Mbanugo and Ejins (2002), coastal dwellers in Lagos by Nebe *et al.* (2002), blood donors in Ibadan by Edington *et al.* (2001) and blood donors in Maiduguri by Ahmed *et al.* (2001).

These results may suggest existence of regional differences in Malaria parasitaemia in Nigeria with the western, mid western and eastern areas (represented by Ogbomosho, Warri, Onitsha and Owerri) ranking highest in prevalence rating and northern area (represented by Maiduguri occupying the lowest position while Lagos and Ibadan take a middle position. Studies in this area should be undertaken by other authors in other parts of the country before a more definite statement on the apparent trends could be made.

Statistical analysis showed that at 95% C.I, the female students were significantly ($\chi^2 = 3.841$ p-value, Cal. $\chi^2 = 3.958$) $p < 0.05$.More infected than the males ($P < 0.05$) (Table 3). The reason why the females are more prone to malaria attack could not be fathomed immediately but it may be because of the (According to Portilo and Sullivan, (1943), Genetic factors could play a role by endowing females with immuno-regulatory potentials to cope better with some diseases) nature of the peculiar dressing of the females , most parts of their bodies (such as hands and legs) are exposed to mosquito bites most of the time compared to the males . This finding was in agreement with the of some earlier authors (Akhigbe *et al.*, 2011) but inconsistent with report of some previous studies (Agbonlahor *et al.*, 1993; Ilozumba and Uzozie, 2009; Mbanugo and Emenalo, 2004) .

ABO blood type frequencies of non-severe infection were 76.4%, 53.6%, 52.4% and 0.0% for blood groups O,B,A,AB respectively as obtained in this study. Clearly, blood group O students were the highest infected with non-severe malaria and the least were group A subjects. No AB subjects were infected. High prevalence of malaria

infection among women with blood group O has been reported (Uneke, 2007). Finding in this study is not tandem with the report of Akhigbe *et al.* (2011) which stated non malaria frequencies of 85.1%,46.1%, 40.0% and 33.5% for blood groups O,A,B,AB respectively . The present finding stating that group O subjects were the most infected agrees with the reports of earlier authors (Fischer and Boone, 1998; Cserti and Dzik, 2007; Adam *et al.*, 2007; A moo *et al.*, 2008, Akhigbe *et al.*, 2011; Pathirana *et al.*, 2007; Migot-Nabias *et al.*, 2000). Present finding therefore disagrees with the reports of Barragan *et al.* (2000), Carlson and Wahlgreen (1992), Rowe *et al.*, (1995), Udumsangpetch *et al.* (1993) which stated that blood group O individuals usually have low malaria attack because parasitized red blood cells have stronger tendency to form rosettes with uninfected erythrocytes of the A, B, AB blood groups than with those of blood group O.

In this study malaria infection was divided into severe and non-severe malaria .Infected students whose thick film shows the presence of *P. falciparum* asexual forms such as schizonts, merozoites and who showed obvious signs of malaria infection were grouped as severe. It is well known that pathogenesis of severe malaria are due to the presence of schizonts, merozoites and trophozoites of *Plasmodium falciparum* (Beng-Ong *et al.*, 2011). The frequency occurrence of severe malaria therefore among the blood types were 70.5%,58.3% ,55.6%and 50.0% for groups O,A,AB ,and O respectively(Table 5). Again group O subjects were the most infected with severe malaria while group B were the least infected. On the whole, group O subjects were the most prone to non –severe and severe forms of malaria. This is not consistent with the finding

of Akhigbe *et al.* (2011) which stated that blood group O persons are associated with the highest prevalence of non-severe malaria and a lower prevalence of severe malaria. Whereas group AB persons recorded no single non-severe malaria case, 55.6% of AB individuals were infected with the severe form of malaria. The finding by some authors stating that blood group O seems to confer a certain degree of protection against severe cases of malaria disease is therefore, not in consonance with present finding in that regard (Fischer and Boone, 1998; Lell *et al.*, 1999 Pathirana *et al.*, 2005; Migot-Nabias *et al.*, 2000). Also, according to Athreya and Coriell (1967), malaria is a disease for which an association with ABO blood group distribution seems to exist and which may have an important role in shaping the prevalence in favour of group O.

Present study reported that where as AB students were infected with 55.6% cases of severe malaria , no one single person was infected with the non-severe malaria .This is in agreement with report of previous studies (Akhigbe *et al.*, 2011) and inconsistent with that of Ilozumba and Uzozie (2009) .

The pathophysiological plausibility of an interaction between malaria parasite and red blood cells as well as the potential role erythrocyte surface antigens may play in cytoadhesion of infected erythrocytes and parasite invasion may give credence to possible association between ABO blood groups and malaria infection (Cserti and Dzik, 2007). Numerous studies have been carried out on the association between ABO blood groups and malaria (Uneke, 2007) and findings from such studies evaluating possible relationship have been contradictory (Uneke, 2007). Herrera *et al.* (2009) reported that there was no association between ABO blood types and malaria parasitaemia and that the

controversial association of these variables previously found in other populations may be due to their peculiar demographic distributions and characteristics.

In this study , attempt was made to associate non-severe malaria and severe malaria parasitaemia with ABO blood types (Tables 4 and 5) chi-square (χ^2) p value (book value) at degree of freedom (3) and 95% confidence interval was 7.815 as against calculated $\chi^2 = 3.408$ for non-severe malaria association with ABO blood groups (Table 4) . Hence $P < 0.05$ and therefore, no significant association or relationship was established. As for the severe malaria association with ABO blood group, $\chi^2 0.05(3) = 7.815$ (p value) and Cal. $X^2 = 5.652$ and hence, $p > 0.05$ (Table 5). This implied that there was also no statistically significant association existing the variables. This lack of significant association in both cases is disagreement with reports of previous studies which recorded significant associations (Zerihun *et al.*, (2011) and consistent with reports of some authors (Nkwo-Akenji, 2004); Kassim *et al.*, 1982; Thakur and Verman, 1992; Montoya *et al.*, 1994 and Uneke *et al.*, 2007).

Conclusion

This study showed that there are more rhesus positive female students than male students in the institution under study. The presence of rhesus negative factor up to 5.8% suggested a gradual and steady rise in the frequency of occurrence of the factor when compared to reports of earlier studies. This has far reaching medical implication in terms of erythroblastosis fetalis (or haemolytic disease of newborn). ABO blood groups O and AB recorded the highest and lowest frequencies respectively. Hence, this finding, lends further credence to earlier related studies that in any given population, the highest number of subjects belong to group O while the least number belong to group AB. A striking revelation that followed was that across all ABO blood groups screened for, there were more females than males. This may not have been due to chance as it is similar to reports of earlier studies.

The highest parasitaemia rate was observed among group O individuals and also among female group O individuals compared to group O male students. More female than male students suffered from both severe and non-severe forms of plasmodiasis. There was a significant statistical relationship between sex (gender) and malaria parasitaemia. Also, blood group O subjects had the highest prevalence of non-severe and severe malaria infection and whereas no single AB student recorded non-severe malaria infection, up to 50.0% AB students had severe form of malaria attack. It was not clear which of blood groups A and or B recorded the least of severe and non-severe forms of malaria. Statistically, there was however, no significant association of all ABO blood

521 types with severe non-severe malaria parasitaemia. This ostensibly implies that all ABO
 522 blood groups are equally at risk and therefore, available malaria prophylactic and
 523 therapeutic strategies by health agencies should be directed at individual of all groups.

524

525

526

527

528

529

530

531

532

533

534

535

536

537

REFERENCES

- 538 Abdulazeez, A. A., Alo, E. B. and Rebecca, S.N.(2008). Carriage rate of human
539 immunodeficiency Virus (HIV) infection among different ABO and Rhesus blood groups
540 in Adamawa state Nigeria. *Bio.Med.Res.* **19**:41-44
- 541 Adam, I., Babiker, S., Mohammed, A.A., Salih, M.M., Prins, M.H and Zaki, Z.M. (2007). ABO
542 Blood group system and placental malaria in an area of unstable malaria transmission In
543 eastern Sudan. *Malar J.*; **6**:110-120.
- 544 Agbonlahor, De, Obi, Fl. Esumeh, o, Ajanaku, A.A. and Igumbor, E.O.(1993). Association of
545 ABO blood Groups and malaria parasitaemia among students of Edo state university
546 Ekpoma, Nigeria. *Journ. Med. Lab. Scit.* **4** (2):12-19.
- 547 Ahmed, S. G., Ibrahim, U. A. and Ibrahim, G. (2001). Prevalence and clinical significance of
548 parasitaemia On blood donors in Maiduguri, Nigeria. *Nigeria Journal of*
549 *Parasitology* **22**:29-34
- 550 Aidoo, M., Terlouw, D.J., Kolczak, M.S. and *et al.* (2002). Protective effects of the sickle cell
551 gene Against malaria morbidity and mortality. *Lancet*; **359**:1311-1312
- 552 Aird, I., Bentall, H.H. and Roberts, J.A. (1953). A relationship between cancer of stomach and
553 the ABO blood groups. *Br Med J*; **1**:799-801.
- 554 Aird, I., Bentall, H.H., Mehigan, J.A., and Roberts, J.A. (1954). the blood groups in relation to
555 Peptic ulceration and carcinoma of colon, rectum, breast, and bronchus; an
556 association Between the ABO groups and peptic ulceration. *Br Med J*; **2**:315-321.
- 557 Akhigbe, R. E., S. F., Ige, A. O., Afolabi, O. M., Azeez, G. J. Adegunlola and J.O. Bamidele.
558 (2009). Prevalence of haemoglobin variants, ABO and rhesus blood groups in Ladoke
559 Akintola University of Technology, Ogbomoso, Nigeria. *Trends Med.Res.* **4**:24-29
- 560 Akide-Ndunge, O.B., Ayi, K., and Arese, P. (2003). The Haldane malaria hypothesis: facts,
561 Artifacts and a prophecy. *Redox Rep.* ; **8**:311-316
- 562 Akinboye, D.O and Ogunrinade, A.F. (1987). Malaria and loasis among blood donors at
563 Ibadan, Nigeria. *Trans R Soc Trop Med Hyg.* **81**:398-399
- 564 Allen, S.J., O'Donnell, A., Alexander, N.D., and *et al.* (1997). Alpha+-Thalassemia protects
565 children Against disease caused by other infections as well as malaria. *Proc Natl Acad Sci*
566 *U S A.*; **94**:14736-14741.
- 567 Allison, A.C. (1954). The distribution of the sickle-cell trait in East Africa and elsewhere, and
568 its Apparent relationship to the incidence of sub tertian malaria. *Trans R Soc Trop Med* ;
569 **48**:312-318.
- 570 Alouch, A. L. (1997). Sickle cell disease and Malaria parasitaemia in a tropical setting *Journ*
571 *Biol. Chem.* **12**:23-30.
- 572 Amer, J., Ghoti, H., Rachmilewitz, E., Koren, A., Levin, C. and Fibach, E. (2006). Red blood
573 cells, Platelets and polymorphonuclear neutrophils of patients with sickle cell disease
574 exhibit Oxidative stress that can be ameliorated by antioxidants. *Br J Haematol*; **132**:108-
575 113.

- 576 Anastasi, J. (1984) .Hemoglobin S-mediated membrane oxidant injury: protection from malaria
577 And pathology in sickle cell disease. *Med Hypotheses*; **14**: 311-320.
- 578 Anees, M. and M.S. Mirza. (2005). Distribution of ABO and Rh blood group alleles in Gujarat
579 region of Punjab, Pakistan. *Proc. Pak.Acad. Sci.***42**:233-238.
- 580 Arese, P., Turrini, F., and Schwarzer, E. (2005). Band 3/complement-mediated recognition and
581 Removal of normally senescent and pathological human erythrocytes. *Cell Physiol*
582 *Biochem*; **16**:133- 146.
- 583 Atamna, H. and Ginsburg, H. (1997).The malaria parasite supplies glutathione to its host cell—
584 Investigation of glutathione transport and metabolism in human erythrocytes infected
585 With *Plasmodium falciparum*. *Eur J Biochem.*; **250**:670-679.
- 586 Athreya, B.H. and Coriell, L.L. (1967). Relation of blood groups to infection. I. A survey and
587 Review of data suggesting possible relationship between malaria and blood groups. *Am J*
588 *Epidemiol*; **86**:292-304.
- 589 Ayi, K., Min-Oo, G., Serghides, L. and *et al.* (2008). Pyruvate kinase deficiency and malaria. *N*
590 *Engl J Med*; **358**:1805-1810.
- 591 Ayi, K., Patel, S.N., Serghides, L., Smith, T.G and Kain, K.C. (2005). Nonopsonic phagocytosis
592 of Erythrocytes infected with ring-stage *Plasmodium falciparum*. *Infect Immun*.
593 **73**:2559-2563.
- 594 Ayi, K., Turrini, F., Piga. A. and Arese, P. (2004) Enhanced phagocytosis of ring-parasitized
595 Mutant erythrocytes: a common mechanism that may explain protection against *falciparum*
596 malaria in sickle trait and beta-thalassemia trait. *Blood*; **104**:3364-3371
- 597 Bakare, A, A., M, A. Azeez and J.O. Agbolade. (2004). Gene frequencies of ABO and Rhesus
598 blood Groups and haemoglobin variants in Ogbomosho, South-west, Nigeria global. *J.*
599 *Med.Sci.***3**:17-22
- 600 Barnwell, J, W. and Galinski, M, R. (1995). Plasmodium vivax: a glimpse into the unique and
601 Shared biology of the merozoite. *Ann Trop Med Parasitol.* **89**:113-120.
- 602 Barnwell, J.W., Nichols, M.E., and Rubinstein, P.(1989;) In vitro evaluation of the role of the
603 Duffy blood group in erythrocyte invasion by *Plasmodium vivax*. *J Exp Med.*; **169**:1795-
604 1802.
- 605 Barragan, A., Kremsner, P.G., Wahlgren, M. and Carlson, J. (2000). Blood group A antigen is a
606 Co receptor in *Plasmodium falciparum* rosetting. *Infect Immun*; **68**:2971-2975.
- 607 Baum, J., Maier, A.G., Good, R, T., Simpson, K.M. and Cowman, A, F. (2005). Invasion by
608 *P.falciparum* merozoites suggest a hierarchy of molecular interactions. *PLoS Pathog*;
609 1:e37. Beckman L. Racial and Ethnic distribution of ABO Blood Types
- 610 Bayoumi, R.A., Bashir, A, H. and Abdulhadi, N.H. (1986). Resistance to *falciparum* malaria
611 Among adults in central Sudan. *Am J Trop Med Hyg*; **35**:45-55.
- 612 Beeson, J, G. and Brown, G, V. (2002). Pathogenesis of *Plasmodium falciparum* malaria: the
613 roles of parasite adhesion and antigenic variation. *Cell Mol Life Sci.*; **59**:258-271

- 614 Beiguelman, B.,Alves ,F.P., Moura,M.M.,Engracia, V.,Nunes,A.C.S.,Heckman,M.I.O. and *et al.*
615 (2003).The Association of Genetic Markers and Malaria infection in the Brazilian Western
616 Amazonian region. *Mern Inst Oswaldo Cruv.***98**:455-460
- 617 Berendt, A.R., Tumer, G.D. and Newbold, C.I.(1994);Cerebral malaria: the sequestration
618 Hypothesis. *Parasitol Today.* **10**:412-414.
- 619 Berger, S.A., Young, N.A. and Edberg, S.C. (1989). Relationship between infectious diseases
620 and Human blood type. *Eur J Clin Microbiol Infect Dis.*; **8**:681-689.
- 621 Beutler E. (1994). G6PD deficiency. *Blood.* **84**:3613-3636.
- 622 Blackwell,C.C.,Dundan, S.,James,V,S.,Mackenzie,A,C.,Braun,J,M.,Alkrout,A,M. and *et*
623 *al.*(2002).Blood Group and susceptibility to disease caused by *Escherichia coli* 0157. *J*
624 *infect Dis* **185**:393-396.
- 625 Boel,M,E., Rijken,M,J.,Pimanpanarak,M.,Proux, S.,Nosten,F. and McGready ,R.(2012). No
626 association of phenotypic ABO blood group and malaria during pregnancy. *American*
627 *Society of Tropical Medicine and Hygiene* .**8**:447-45
- 628 Bookchin, R.M. and Lew, V.L. (2002). Sickle red cell dehydration: mechanisms and
629 Interventions. *Curr Opin Hematol*; **9**:107-110.
- 630 Boren, T., Falk, P., Roth, K.A., Larson, G. and Normark, S. (1993). Attachment of Helicobacter
631 Pylori to human gastric epithelium mediated by blood group antigens. *Science.* **262**:1892-
632 1895.
- 633 Bowen, D.J. (2003). An influence of ABO blood group on the rate of proteolysis of von
634 Willebrand factor by ADAMTS13. *J Thromb Haemost.*; **1**:33-40.
- 635 Bridges, K.R. and Pearson, H.A. (2007). Anaemia and other red cell disorders (Ed 1st). New
636 York:McGraw-Hill.
- 637 Brockelman, C.R., Wongsattayanont, B., Tan-ariya, P. and Fucharoen, S. (1987). Thalassemic
638 Erythrocytes inhibit in vitro growth of *Plasmodium falciparum*. *J Clin Microbiol*; **25**:56-
639 60.
- 640 Brooks, G, F., Butel, J, S. and Morse, S.A. (2004).Medical parasitology 23rd edition. McGraw
641 Hill Companies. Inc.Boston; 818p
- 642 Brown, A.E., Webster, H.K., Teja-Isavadharm, P. and Keeratithakul, D. (1990). Macrophage
643 Activation in *falciparum* malaria as measured by neopterin and interferon-gamma. *Clin*
644 *Exp Immunol.*; **82**:97-101.
- 645 Brown, K.N., and Hills, L.A. (1971). Macrophage "activation" and the action of opsonising
646 Antibodies in malaria. *Trans R Soc Trop Med Hyg.* **65**:6-12
- 647 Bryan, J.H. and Smalley, M.E. (1978). The use of ABO blood groups as markers for mosquito
648 Biting studies. *Trans R Soc Trop Med Hyg*; **72**:357-360
- 649 Buchanan, G.R., DeBaun, M.R., Quinn, C.T., Steinberg, M.H. (2004): Sickle cell
650 disease.Haematology *Am Soc Hematol Educ Program.***9**: 35-47
- 651 Bulai, T., Bratosin, D., Pons, A., Montreuil, J. and Zanetta, J.P. (2003). Diversity of the human
652 Erythrocyte membrane sialic acids in relation with blood groups. *FEBS Lett*; **534**:185-189

- 653 Bunyaratvej, A., Butthep, P., Yuthavong, Y., and *et al.* (1986) Increased phagocytosis of
654 *Plasmodium falciparum*-infected erythrocytes with haemoglobin E by peripheral Blood
655 monocytes. *Acta Haematol.* **76**:155-158.
- 656 Burton, N.M. and Anstee, D.J. (2008). Structure, function and significance of Rh proteins in red
657 Cells. *Curr Opin Hematol*; **15**:625-630.
- 658 Cappadoro, M., Giribaldi, G., O'Brien E. and *et al.* (1998). Early phagocytosis of glucose-6
659 Phosphate dehydrogenase (G6PD)-deficient erythrocytes parasitized by *Plasmodium*
660 *falciparum* may explain malaria protection in G6PD deficiency. *Blood*; **92**:2527-2534.
- 661 Carlson, J., and Wahlgren M. (1992). *Plasmodium falciparum* erythrocyte rosetting is mediated
662 by Promiscuous lectin-like interactions. *J Exp Med.*; **176**:1311-1317.
- 663 Carter, R. and Mendis, K.N. (2002). Evolutionary and historical aspects of the burden of
664 malaria. *Clin Microbiol Rev*; **15**:564-594.
- 665 Cartron, J.P. and Rouger, P. (1995). Molecular basis of human blood group antigens. New York:
666 Plenum Press.
- 667 Castro, O. and Scott, R.B. (1985). Red blood cell counts and indices in sickle cell trait in a black
668 American population. *Hemoglobin*; **9**:65-67.
- 669 Cattani, J.A., Gibson, F.D., Alpers, M.P. and Crane, G.G. (1987). Hereditary ovalocytosis and
670 Reduced susceptibility to malaria in Papua New Guinea. *Trans R Soc Trop Med Hyg*;
671 **81**:705-709.
- 672 Cavasini, C.E., De Mattos, L.C., Alves, R.T., and *et al.* (2006). Frequencies of ABO, MNSs, and
673 Duffy phenotypes among blood donors and malaria patients from four Brazilian Amazon
674 areas. *Hum Biol.*; **78**:215-211.
- 675 Chaibaroj, S.C., Rutta, A.S., Muenthaisong, K., Watkins, P., Na Ubol, M. and Looareesuwan, S.
676 (2004). Reduced levels of transforming growth factor-beta1, interleukin-12 and Increased
677 migration inhibitory factor are associated with severe malaria. *Acta Trop*. **89**:319-327
- 678 Chang, K.H. and Stevenson, M.M. (2004). Malarial anaemia: mechanisms and implications of
679 Insufficient erythropoietin during blood-stage malaria. *Int J Parasitol*; **34**:1501-1516.
- 680 Cheesbrough, M. (2000). Parasitological tests. District laboratory practices in tropic Countries,
681 Cambridge University Press. **221**:23-45
- 682 Chen, Q., Schlichtherle, M. and Wahlgren, M. (2000). Molecular aspects of severe malaria. *Clin*
683 *Microbiol Rev*; **13**:439-450.
- 684 Cholera, R. B.N., Gillrie, M.R., Lopera-Mesa, T.M., Diakit , S.A., Arie, T., Krause, M.A.,
685 Guindo, A., Tubman, A., Fujioka, H., Diallo, D.A., Doumbo, O.K., Ho M, Wellems, T.E.
686 and Fairhurst, R.M (2007). Impaired cytoadherence of *Plasmodium falciparum*-infected
687 Erythrocytes containing sickle haemoglobin. *PNAS*. **105**:991-996.
- 688 Chotivanich, K.T., Udomsangpetch, R., Pipitaporn, B. and *et al.* (1998). Rosetting characteristics
689 of uninfected erythrocytes from healthy individuals and malaria patients. *Ann Trop Med*
690 *Parasitol*; **92**:45-56.
- 691 Chuluyan, H.E., and Issekutz, A.C. (1993). VLA-4 integrin can mediate CD11/CD18-
692 independent Transendothelial migration of human monocytes. *J Clin Invest.* **92**:2768-2777

- 693 Chung, W, Y., Gardiner, D.L., Hyland, C., Gatton, M., Kemp, D.J. and Trenholme, K.R.
694 (2005)Enhanced invasion of blood group A1 erythrocytes by *Plasmodium falciparum*. *Mol*
695 *Biochem Parasitol.*; **144**:128-130
- 696 Clark, I.A., Rockett, K.A., and Cowden, W.B. (1992). Possible central role of nitric oxide in
697 Conditions clinically similar to cerebral malaria. *Lancet*; **340**:894-896.
- 698 Conran, N., Franco-Penteado, C.F. and Costa, F.F. (2009); Newer aspects of the
699 Pathophysiology of sickle cell disease vaso-occlusion. *Haemoglobin*. **33**:1-16.
- 700 Cooke, A, H., Chiodini, P, L. and Doherty, T. (1999).Comparison of a parasite lactate
701 dehydrogenase Immunochromatographic antigen detection assay with microscopy for the
702 of malaria Parasite in human blood samples .*Am.Journ.Trop.Med.Hyp*.**60**:173-200.
- 703 Cortes, A., Benet, A., Cooke, B.M., Barnwell, J.W. and Reeder, J.C. (2004). Ability of
704 *Plasmodium falciparum* to invade Southeast Asian ovalocytes varies between parasites
705 Lines. *Blood*; **104**:2961-2966.
- 706 Cowman, A.F. and Crabb, B.S. (2006) Invasion of red blood cells by malaria parasites. *Cell*;
707 **124**:755-766.
- 708 Cserti, C.M. and Dzik, W.H. (2007) The ABO blood group system and *Plasmodium falciparum*
709 Malaria. *Blood*; **110**:2250-2258.
- 710 Cserti-Gazdewich, C, M., Dhabangi, A., Musoke, C., Sewanyana, I., Mpimbaza, A.and Dzik,
711 W.H. (2012). Cytoadherence in paediatric malaria: ABO blood groups and severe
712 *Plasmodium falciparum* infection. *British journal of haematology*. **159**: 223-236
- 713 Daniels, G. (2002). Human blood groups (ed 2nd). Oxford; Malden, MA: Blackwell Science;
714 Daniels, G. (2005). The molecular genetics of blood group polymorphism. *Transpl Immunol*;
715 **14**:143-153.
- 716 Day, N.P., Hien, T.T., Schollaardt, T. and *et al.* (1999).The prognostic and pathophysiologic role
717 of pro- and anti-inflammatory cytokines in severe malaria. *J Infect Dis.*; **180**:1288-1297
- 718 Dean, L. (2005) Blood Groups and red cell antigens. In: Beck B Ed.
- 719 Desimone, T.M., Jennings, C.V., Bei, A.K. and *et al.* (2009). Cooperativity between *Plasmodium*
720 *falciparum* adhesive proteins for invasion into erythrocytes. *Mol Microbiol*.
- 721 Domarle, O., Migot-Nabias, F., Mvoukani, J.L. and *et al.* (1999). Factors influencing resistance
722 to reinfection with *Plasmodium falciparum*. *Am J Trop Med Hyg*; **61**:926-931.
- 723 Downie, A.W, Meiklejohn, G., St Vincent, L., Rao, AR., Sundara Babu, B.V. and Kempe,
724 C.H.(1965). Smallpox frequency and severity in relation to A, B and O blood groups. *Bull*
725 *World Health Organ*; **33**:623-625.
- 726 Dronamraju, K.R., Arese, P. and Haldane, J.B.S. (2006). Malaria: genetic and evolutionary
727 Aspects. New York: Springer.
- 728 Duffy, P.E. and Fried, M. (2006). Red blood cells that do and red blood cells that don't: how to
729 Resist a persistent parasite. *Trends Parasitol*. **22**:99-101.
- 730 Dunstan, R.A. (1986) Status of major red cell blood group antigens on neutrophils, lymphocytes
731 and Monocytes. *Br J Haematol*. **62**:301-309
- 732 Edington, G, M. and Gillies, H.M. (2001).Disease In: Pathology in the tropics. **121**:234-245

- 733 Ejins, D. O. and Mbanugo, J.I. (2002) Plasmodium infections in children aged 0-5 years in Awka
734 Metropolis, Anambra state, Nigeria. *Nigerian Journal of Parasitology*.**21**:55-59
- 735 Epid, T. T., C.D. Nwani and N.P. Ugorji.(2008).Prevalence of malaria in blood donors in.
736 Abakaliki Metropolis, Nigeria. *Sci. Res. Essays*. **3**:162-164
- 737 F, D. Otajevwo. (2013). Prevalence of Malaria Parasitaemia and its association with ABO blood
738 Grouping among students of Igbinedion University okada, Nigeria. *Science*
739 *domain*.**3**(4):1164-1177
- 740 Facer, C.A., and Brown, J. (1979) ABO blood groups and *falciparum* malaria*Trans R Soc*
741 *TropMed Hyg*; **73**:599-600.
- 742 Febbraio, M., Hajjar, D.P. and Silverstein, R.L. (2001) CD36: a class B scavenger receptor
743 involved in Angiogenesis, atherosclerosis, inflammation, and lipid metabolism. *J Clin*
744 *Invest*. **108**:785-791.
- 745 Fischer, P.R., and Boone, P. (1998). Short report: severe malaria associated with blood group.
746 *Am J Trop Med Hyg*; **58**:122-123.
- 747 Franco, R.S., Lohmann, J., Silberstein, E.B. and *et al.* (1998). Time-dependent changes in the
748 Density and haemoglobin F content of biotin-labelled sickle cells. *J Clin Invest*.;
749 **101**:2730-2740.
- 750 Friedman, M.J. (1979). Oxidant damage mediates variant red cell resistance to malaria.Nature;
751 **280**:245-247
- 752 Fry, A.E., Griffiths, M.J., Auburn, S. and *et al.* (2008). Common variation in the ABO
753 Glycosyltransferase is associated with susceptibility to severe *Plasmodium*
754 *falciparum*Malaria. *Hum Mol Genet*; **17**:567-576.
- 755 Garratty, G. (2000). Blood groups and disease: a historical perspective. *Transfus Med Rev*.;
756 **14**:291-301.
- 757 Gaur, D., Furuya, T., Mu, J., Jiang, L.B., Su, X.Z. and Miller, L.H. (2006). Up regulation of
758 Expression of the reticulocyte homology gene 4 in the *Plasmodium falciparum* clone Dd2
759 is associated with a switch in the erythrocyte invasion pathway. *Mol Biochem Parasitol*;
760 **145**:205-215.
- 761 Genton, B., al-Yaman, F., Mgone, C.S., and *et al.* (1995). Ovalocytosis and cerebral malaria.
762 *Nature*. **378**:564-565
- 763 Gibbs, F.L. and Wallace. M.E. (1986). American Association of Blood Banks. Blood group
764 Systems, ABH and Lewis. Arlington, Va.: American Association of Blood Banks;
- 765 Gilles, H.M., Fletcher, K.A., Hendricks, R.G., Lindner, R., Reddy, S. and Allan, N. (1967).
766 Glucose-6- phosphate-dehydrogenase deficiency, sickling, and malaria in African children
767 In South Western Nigeria. *Lancet*. **1**:138-140.
- 768 Giribaldi, G., Ulliers, D., Manu, F., Arese, P. and Turrini, F. (2001). Growth of *Plasmodium*
769 *falciparum* induces stage-dependent haemachrome formation, oxidative aggregation of
770 Band 3, membrane deposition of complement and antibodies, and phagocytosis of
771 Parasitized erythrocytes. *Br J Haematol*. **113**:492-499.

- 772 Gowda, D.C. (2007). TLR-mediated cell signalling by malaria GPIs. *Trends Parasitol*; **23**:596-
773 604.
- 774 Grau, G.E., Piguet, P.F., Vassalli, P., and Lambert, P.H.O. (1989). Involvement of tumour
775 Necrosis factor and other cytokines in immune-mediated vascular pathology. *Int Arch An*
776 *allergy Appl Immunol.*; **88**:34-39.
- 777 Greenwell, P. (1997). Blood group antigens: molecules seeking a function? *Glycoconj J.*;
778 **14**:159-173
- 779 Greenwood, B.M., Fidock, D.A., Kyle, D.E., and et al. (2008) Malaria: progress, perils, and
780 prospects for eradication. *J Clin Invest.*; **118**:1266-1276.
- 781 Guerra, C.A., Gikandi, P.W., Tatem, A.J., and et al. (2008) .The limits and intensity of
782 *Plasmodium falciparum* transmission: implications for malaria control and elimination
783 worldwide.*PLoS Med.*; 5:e38
- 784 Guo, R.F, Ward PA. (2005). Role of C5a in inflammatory responses. *Annu Rev Immunol.*;
785 **23**:821-852.
- 786 Gupta, M. and Chowdhuri, A.N. (1980). Relationship between ABO blood groups and malaria.
787 *Bull World Health Organ.* **58**:913-915.
- 788 Haest, C.W. (1982) Interactions between membrane skeleton proteins and the intrinsic domain of
789 The erythrocyte membrane. *Biochim Biophys Acta.*; **694**:331-352.
- 790 Harris, J.R. (1990). Erythroid cells. New York: Plenum Press.
- 791 Harrison, J. and Ridley, D.S. (1975). Heterologous reactions involving parasites, blood group
792 Antibodies and tissue components. *Trans R Soc Trop Med Hyg*; **69**:312-317.
- 793 Hay, S.I., Guerra, C.A., Tatem, A.J., Noor, A.M. and Snow, R.W. (2004). The global distribution
794 and Population at risk of malaria: past, present, and future. *Lancet Infect Dis.*; **4**:327-336.
- 795 Hebbel, R.P., Morgan, W.T., Eaton, J.W., and Hedland, B.E. (1988) Accelerated autoxidation
796 And heme loss due to instability of sickle haemoglobin. *Proc Natl Acad Sci U S A.*;
797 **85**:237-241.
- 798 Horrocks, P., Pinches, R.A., Chakravorty, S.J. and et al. (2005). PfEMP1 expression is reduced
799 on the surface of knobless *Plasmodium falciparum* infected erythrocytes. *J Cell Sci*;
800 **118**:2507- 2518
- 801 Horuk, R., Chitnis, C.E., Darbonne, W.C. and et al. (1993). A receptor for the malaria parasite
802 *Plasmodium vivax*: the erythrocyte chemokine receptor. *Science.* **261**:1182-1184Hebbel,
803 R.P. (2003). Sickle haemoglobin instability: a mechanism for malarial protection. *Redox*
804 *Rep.* **8**:238-240.
- 805 Hosoi, E. (2008). Biological and clinical aspects of ABO blood group system. *J Med*
806 *Invest.***55**:174-182.
- 807 Hume, J.C., Lyons, E.J. and Day, K.P. (2003) Human migration, mosquitoes and the evolution of
808 *Plasmodium falciparum*. *Trends Parasitol*; **19**:144-149.
- 809 Ibhanebor, S.E., Otokob, E.S., and Ladipo, O.A. (1996). Prevalence of malaria parasitaemia in
810 Transfused donor blood in Benin City, Nigeria. *Ann Trop Paediatr.* **16**:93-95

- 811 Idro, R., Jenkins, N.E., and Newton, C.R. (2005). Pathogenesis, clinical features, and
812 neurological Outcome of cerebral malaria. *Lancet Neurol.*; **4**:827-840.
- 813 Igbineghu, C., Odaibo, G.N., Olaleye, D, O. and Odiabo, A.(2012).Malaria infection and ABO
814 blood Grouping in Iwo community, Southwestern Nigeria. *Res. Journ.Med. Sci.***6**:247-250
- 815 Ilozumba, P, C. and Uzozie, C, R. (2009). Prevalence of malaria parasitaemia and its association
816 with ABO Blood Group in Odoakpu Area of Onitsha south local government area,
817 Anambra State Nigeria.*Nigeria Annals of Natural Sciences.***8** (2):1-8
- 818 Janeway, C.A, Jr. (2002). Medzhitov R. Innate immune recognition. *Annu Rev Immunol*; **20**:197-
819 216.
- 820 Jarolim, P., Palek, J., Amato, D., and *et al.* (1991). Deletion in erythrocyte band 3 gene in
821 malaria-
- 822 Jeremiah, Z.A. (2006). Abnormal haemoglobin variants, ABO and Rh groups among student of
823 African Descant in Port Harcourt. *Africa. Health Si.***6**:177-181
- 824 Joshi, H., Raghavendra, K., Subbarao, S.K., and Sharma, V.P. (1987). Genetic markers in
825 malaria Patients of Delhi. *Indian J Malariol*; **24**:33-3
- 826 Joysey, V.C. (1959). The relation between animal and human blood groups. *Br Med Bull.*;
827 **15**:158-164.
- 828 Judd, W.J. (1999). Elution--dissociation of antibody from red blood cells: theoretical and
829 Practical considerations. *Transfus Med Rev.*; **13**:297-310.
- 830 Kaminsky, R., Kruger, N., Hempelmann, E. and Bommer, W. (1986). Reduced development of
831 *Plasmodium falciparum* in beta-thalassaemic erythrocytes. *Z Parasitenkd*; **72**:553-556
- 832 Kano, K., McGregor, I.A., Milgrom, F. (1968). Hemagglutinins in sera of Africans of Gambia.
833 *Proc Soc Exp Biol Med.*; **129**:849-853
- 834 Kassim, O.O., and Ejezie, G.C. (1982). ABO blood groups in malaria and schistosomiasis
835 Haematobium. *Acta Trop.*; **39**:179-184.
- 836 Kidson, C., Lamont, G., Saul, A., and Nurse, G.T. (1981) . Ovalocytic erythrocytes from
837 Melanesians are resistant to invasion by malaria parasites in culture. *Proc Natl Acad Sci U*
838 *S A.* **78**:5829-5832.
- 839 Klein, H.G., Mollison, P.L., Anstee, D.J., and Mollison, P.L. (2005). Mollison's blood
840 transfusion In clinical medicine (Ed 11th). Malden, Mass.; Oxford: Blackwell Pub.
- 841 Kutner, S., Breuer, W.V., Ginsburg, H., Aley, S.B. and Cabantchik, Z.I. (1985).
842 Characterization Of permeation pathways in the plasma membrane of human erythrocytes
843 infected with Early stages of *Plasmodium falciparum*: association with parasite
844 development. *J Cell Physiol*; **125**:521-527.
- 845 Kwiatkowski, D., Hill, A.V., Sambou, I. and *et al.* (1990). TNF concentration in fatal cerebral,
846 non-fatal cerebral and uncomplicated *Plasmodium falciparum* malaria *Lancet*; **336**:1201-
847 1204.
- 848 Kwiatkowski, D.P. (2005) How malaria has affected the human genome and what human
849 genetics Can teach us about malaria. *Am J Hum Genet.* **77**:171-192.

850 Kyes, S., Pinches, R. and Newbold, C. (2000). A simple RNA analysis method shows var and
851 Rif Multigene family expression patterns in *Plasmodium falciparum*. *Mol Biochem*
852 *Parasitol.* **105**:311-315.

853 Landsteiner, K. and Miller, C.P. Jr. (1925). Serological Observations on the Relationship of the
854 Bloods of Man and the Anthropoid Apes. *Science*; **61**:492-493

855 Lanzillotti, R., and Coetzer, T.L. (2006). The 10 kDa domain of human erythrocyte protein
856 4.1 Binds the *Plasmodium falciparum* EBA-181 protein. *Malar J.*; **5**:100-150

857 Lee, S.K., Tham, K.T., Cheung, K.P. and Jenkins, W.J. (1982) Rh (D) fraction incompatibility
858 Causing hemolytic disease of the newborn. Report of two cases in a Chinese family. *Am J*
859 *ClinPathol*; **78**:95-96.

860 Lell, B., May, J., Schmidt-Ott, R.J. and *et al.* (1999). The role of red blood cell polymorphisms
861 in Resistance and susceptibility to malaria. *Clin Infect Dis.* **28**:794-799.

862 Lin, Y., and Pavenski, K. (2008). Red Cell Physiology 101. Canadian Blood Services :*Annual*
863 *International Symposium*.

864 Liu, Q.P., Sulzenbacher, G., Yuan, H. and *et al.* (2007). Bacterial glycosidases for the production
865 Of universal red blood cells. *Nat Biotechnol.* **25**:454-464.

866 Livinestone, F.B. (1971). Malaria and human polymorphisms. *Annu Rev Genet.*; **5**:33-64

867 Loscertales, M.P. and Brabin, B.J. (2006). ABO phenotypes and malaria related outcomes in
868 Mothers and babies in The Gambia: a role for histo-blood groups in placental Malaria
869 *Malar J*; **5**:72-80

870 Loscertales, M.P., Owens, S., O'Donnell, J., Bunn, J., Bosch-Capblanch, X., and Brabin,
871 B.J.(2007) ABO blood group phenotypes and *Plasmodium falciparum* malaria:Unlocking
872 a pivotal mechanism. *Adv Parasitol*; **65**:1-50

873 Lukyanenko, L.M., Kozlova, N.M. and Slobozhanina, E.I. (2004). Activity of membrane-bound
874 NADH- methemoglobin reductase and physical state of lipids in erythrocyte Membranes.
875 *Bioelectrochemistry*; **62**:191-193.

876 Lutz, H.U., Bussolino, F., Flepp, R. and *et al.* (1987). Naturally occurring anti-band-3 antibodies
877 And complement together mediate phagocytosis of oxidatively stressed human
878 Erythrocytes. *Proc Natl Acad Sci U S A.* **84**:7368-7372

879 Luzzatto, L., Usanga, F.A. and Reddy, S. (1969). Glucose-6-phosphate dehydrogenase deficient
880 Red cells: resistance to infection by malarial parasites. *Science*; **164**: 839-842.

881 MacKinney, A.A. (2003). Hematology for Students: Taylor and Francis Group;.

882 MacPherson, G.G., Warrell, M.J., White, N.J., Looareesuwan, S. and Warrell, D.A. (1985)
883 Human Cerebral malaria. A quantitative ultra structural analysis of parasitized erythrocyte
884 Sequestration. *Am J Pathol.* **119**:385-401.

885 Madigan, C., and Malik, P. (2006). Pathophysiology and therapy for haemoglobinopathies. Part
886 I: sickle cell disease. *Expert Rev Mol Med.*; **8**:1-23.

887 Mak, T.W. and Saunders, M.E. (2006). The immune response: basic and clinical principles.
888 New York: Elsevier/Academic.

- 889 Makela, O., Ruoslahti, E., and Ehnholm, C. (1969). Subtypes of human ABO blood groups and
890 Subtype- specific antibodies. *J Immunol.* **102**:763-77
- 891 Marion, Reid, C.L-F. (2004).The blood group antigen facts book (ed 2nd ed). New York:
892 Elsevier Academic Press;
- 893 Marionneau, S., Cailleau-Thomas, A., Rocher, J. and *et al.* (2001). ABH and Lewis histo-blood
894 Group antigens, a model for the meaning of oligosaccharide diversity in the face of A
895 changing world. *Biochimie.*; **83**:565-573.
- 896 Marsh, W.L., Nichols, M.E., Oyen, R. and *et al.* (1978). Naturally occurring anti-Kell stimulated
897 By E. coli enterocolitis in a 20-day-old child. *Transfusion*; **18**:149-154.
- 898 Martin, S.K., Miller, L.H., Hicks, C.U., David-West, A., Ugbo, C. and Deane, M.
899 (1979).Frequency of blood group antigens in Nigerian children with falciparum malaria.
900 *Trans R Soc Trop Med Hyg*; **73**:216-218.
- 901 Mayer DC, Kaneko O, Hudson-Taylor DE, Reid ME, Miller LH. (2001). Characterization of a
902 *Plasmodium falciparum* erythrocyte-binding protein paralogous to EBA-175. *Proc Natl*
903 *Acad Sci U S A.*; **98**:5222-5227
- 904 Mbanugo, J. I. and Emenalo, S.E.(2004).Prevalence of malaria parasitaemia among blood
905 donors in Owerri, imo state, Nigeria.*Nigeria Journal of Parasitology.* **25**:75-80
- 906 McGilvray, I.D., Serghides, L., Kapus, A., Rotstein, O.D. and Kain, K.C. (2000). Nonopsonic
907 Monocyte/macrophage phagocytosis of *Plasmodium falciparum*-parasitized Erythrocytes:
908 a role for CD36 in malarial clearance. *Blood*; **96**:3231-3240.
- 909 Menendez C, Fleming AF, Alonso PL. Malaria-related anaemia. *Parasitol Today.* (2000).**16**:469-
910 476.
- 911 Mercereau-Puijalon, O., Guillotte, M., and Vigan-Womas, I. (2008) Rosetting in *Plasmodium*
912 *falciparum*: a cytoadherence phenotype with multiple actors. *Transfus Clin Biol.*; **15**:62-
913 71.
- 914 Migot-Nabias, F, Mombo, L, E., Luty, A.J., Dubois, B., Nabias,R., Bisseye , C., Millet, P., Lu,
915 C,Y. and Delovon, P. (2000).Human genetic factors related to susceptibility to mild
916 malaria in Gabon. *Genes Immun.***1**:435-44
- 917 Miller, J., Golenser, J., Spira, D.T., and Kosower, N.S. (1997) *Plasmodium falciparum*: thiol
918 status And growth in normal and glucose-6-phosphate dehydrogenase deficient human
919 Erythrocytes. *Exp*
- 920 Miller, L.H., Baruch, DI, Marsh, K. and Doumbo, O.K. (2002). The pathogenic basis of
921 malaria.*Nature*; **415**:673-679.
- 922 Miller, L.H., Mason, S.J., Dvorak, J.A., McGinniss, M.H. and Rothman, I.K. (1975. Erythrocyte
923 Receptors for (*Plasmodium knowlesi*) malaria: Duffy blood group determinants. *Science.*
924 **189**:561-563.
- 925 Mohan, KaS, M.M. (1998). Acquired immunity to asexual blood stages. In: Sherman IW,
926 ed.Malaria: Parasite Biology, Pathogenesis and protection.
- 927 Mohandas, N. Gallagher, P.G. (2008). Red cell membrane: past, present, and future.*Blood*;
928 **112**:3939-3948.

- 929 Mollison, P.L., Contreras, M., and Engelfriet, C.P. (1997). Blood transfusion in clinical
930 medicine (ed 10th). Oxford [England]; Malden, MA, USA: Blackwell Science.
- 931 Montoya, F., Restrepo, M., Montoya, A.E. and Rojas, W. (1994). Blood groups and malaria. *Rev*
932 *Inst Med Trop Sao Paulo*. **36**:33-38
- 933 Morgan, W.T. and Watkins, W.M. (1948) . The detection of a product of the blood group O
934 Gene and the relationship of the so-called O-substance to the agglutinates A and B. *BrJ*
935 *Exp Pathol*. 29:159-173.
- 936 Morgan, W.T. and Watkins, W.M. (2000). Unravelling the biochemical basis of blood group
937 ABO And Lewis antigenic specificity. *Glycoconj J.*; **17**:501-530
- 938 Moulds, J.M., Nowicki, S., Moulds, J.J. and Nowicki, B.J. (1996). Human blood groups:
939 incidental Receptors for viruses and bacteria. *Transfusion*; **36**:362-374
- 940 Mourant, A.E. (1983). Blood Relations: Blood groups and anthropology. New York: Oxford
941 University Press.
- 942 Mourant, A.E., Kopec, A.C., Domaniewska-Sobczak, K. (1971) Blood-groups and blood-
943 clotting. *Lancet*; **1**:223-227.
- 944 Nagel, R.L. and Fleming, A.F. (1992). Genetic epidemiology of the beta s gene. Baillieres *Clin*
945 *Haematol*; **5**:331-365.
- 946 Naik, R.S., Branch, O.H., Woods, A.S. and et al. (2000) Glycosylphosphatidylinositol anchors of
947 *Plasmodium falciparum*: molecular characterization and naturally elicited antibody
948 Response that may provide immunity to malaria pathogenesis. *J Exp Med.*; **192**:1563-
949 1576.
- 950 Nash, G.B., Cooke, B.M., Carlson, J. and Wahlgren, M. (1992). Rheological properties of
951 rosettes formed by red blood cells parasitized by *Plasmodium falciparum*. *Br J Haematol*;
952 **82**:757-763.
- 953 Nasr, A.(2012). Blood group O protects against complicated *Plasmodium falciparum* malaria by
954 the Mechanism of inducing high levels of antimalarial IgG antibodies. *Saudi Arabia journ.*
955 *Med* .**1**:16-22
- 956 Natta, C.L., Chen, L.C. and Chow, C.K. (1990). Selenium and glutathione peroxidase levels in
957 sickle Cell anaemia. *Acta Haematol.* ; **83**:130-132.
- 958 Ndambaa, Y., Gonoa, E., Nyazemab, N., Makazaa, N. and Kaonarea, K.C.(1997).
959 Schistosomiasis Infection in relation to the ABO blood groups among school children in
960 Zimbabwe. *Acta. Trop.* **65**:181-190.
- 961 Nebe, O, J., Adeoye, G, O. and Agomo, P.U. (2002).Prevalence and clinical profile of malaria
962 among the Coastal dwellers of Lagos state, Nigeria. *Nigeria Journal of*
963 *Parasitology*.**23**:61-68
- 964 Nigg, E.A. and Cherry, R.J. (1980) Anchorage of a band 3 population at the erythrocyte
965 Cytoplasmic membrane surface: protein rotational diffusion measurements. *Proc Natl*
966 *Acad Sci U S A.*; **77**:4702-4706.

- 967 Nkwo-Akenji, Paul, Wepngong. And Jane, F.A.(2004).Effects of ABO Rh blood groups G-6-P-
968 D enzyme Acuity and haemoglobin genotypes on malaria parasitaemia and parasite
969 density.*Afr J Health Science*. **11**:93-97.
- 970 Ockenhouse, C.F, Schulman, S., and Shear, H.L. (1984) Oxidative killing of malaria parasites by
971 Mononuclear phagocytes. *Prog Clin Biol Res.*; **155**:93-108
- 972 Odokuma, E, I., A, C., Okolo. And P.C. Aloamaka. (2007).Distribution of ABO and Rhesus
973 Blood groups in Abraka. Frequency of blood groups ABO and rhesus D in the Guinea
974 population.*Nig. J. Physiol.Sci.***22**:89-91
- 975 Oliver-Gonzalez, JaTM. (1944). A substance in animal parasites related to the human
976 Isoagglutinogens. *J infect Dis.*; **74**:173-200
- 977 Olsson, M.L. Chester, M.A. (2001). Polymorphism and recombination events at the ABO locus:a
978 major challenge for genomic ABO blood grouping strategies. *Transfus Med*; **11**:295-313.
- 979 Omodeo-Sale, F., Motti, A., Basilico, N., Parapini, S., Olliaro, P. and Taramelli, D.
980 (2003).Accelerated senescence of human erythrocytes cultured with *Plasmodium*
981 *falciparum*. *Blood*; **102**:705-711.
- 982 Omotade, O, O., Adeyemo, A, A., Kayode, E, Falade, S,L. and Ikpeme, S.(1999).Gene
983 frequencies of ABO And RH (D) blood group alleles in a healthy infant population in
984 Ibadan, Nigeria .*West African Journal of Medicine*. **18**:294-297
- 985 O'Neil, D. (2009). Distribution of Blood Types. Vol. San Marcos.
- 986 Opera, K.N. (2007). Onchocerciasis and ABO blood group status; a field based study. *Int J Trop*
987 *Med*.**2**:123-125.
- 988 Orlandi, P.A., Sim, B.K., Chulay, J.D. and Haynes, J.D. (1990). Characterization of the 175-Kilo
989 dalton erythrocyte binding antigen of *Plasmodium falciparum*. *Mol Biochem Parasitol*;
990 **40**:285-294.
- 991 Osisanya, J.O. (1983). ABO blood groups and infections with human malarial parasites in vivo
992 And in vitro. *East Afr Med J*. **60**:616-621.
- 993 Otajevwo, F, D. (1997).ABO Blood group Association with malaria parastaemia among
994 residents in Warri Delta State.*Warri Journal of Science band Technology*.**4**:32-36
- 995 Pant C.S., Gupta, D.K., Bhatt, R.M., Gautam, A.S. and Sharma, R.C. (1992).An epidemiological
996 Study of G- 6-PD deficiency, sickle cell haemoglobin, and ABO blood groups in relation
997 to malaria incidence in Muslim and Christian communities of Kheda, Gujarat, (India). *J*
998 *Commun Dis.*; **24**:199-205.
- 999 Pant, C.S, Gupta, D.K., Bhatt, R.M., Gautam, A.S. and Sharma, R.C.(1993). Three genetic
1000 markers And malaria in upper caste Hindus of Kheda District of Gujarat State. *Indian J*
1001 *Malariol*;**30**:229- 233.
- 1002 Pant, C.S., and Srivastava, H.C. (1997) Distribution of three genetic markers and malaria in
1003 Other backward castes of Kheda district, Gujarat. *Indian J Malariol*; **34**:42-46.
- 1004 Pant, C.S., Gupta, D.K., Sharma, R.C., Gautam, A.S., and Bhatt, R.M. (1992) Frequency of ABO
1005 Blood groups, sickle-cell haemoglobin, G-6-PD deficiency and their relation with malaria

- 1006 in Scheduled castes and scheduled tribes of Kheda District, Gujarat. *Indian J Malariol*;
1007 **29**:235-239.
- 1008 Pantaleo, A., Giribaldi, G., Mannu, F., Arese, P. and Turrini, F. (2008). Naturally occurring anti-
1009 Band 3 antibodies and red blood cell removal under physiological and pathological
1010 Conditions. *Autoimmun Rev.*; **7**: b457-462
- 1011 Pasvol, G., Wainscoat, J.S. Weatherall, D.J. (1982) Erythrocytes deficiency in glycophorin resist
1012 Invasion by the malaria parasite *Plasmodium falciparum*. *Nature.* ; **297**:64-66
- 1013 Pasvol, G., Weatherall, D.J. Wilson, R.J. (1978). Cellular mechanism for the protective effect of
1014 haemoglobin S against *P. falciparum* malaria. *Nature.* **274**:701-703.
- 1015 Patel, S.N., Serghides, L., Smith, T.G. and *et al.* (2004). CD36 mediates the phagocytosis of
1016 *Plasmodium falciparum*-infected erythrocytes by rodent macrophages. *J Infect*
1017 *Dis.***189**:204-213.
- 1018 Pathirana, S.L., Alles, H.K., Bandara, S. and *et al.* (2005). ABO-blood-group types and
1019 protection Against severe, *Plasmodium falciparum* malaria. *Ann Trop Med Parasitol.* ;
1020 **99**:119-124.
- 1021 Pattanapanyasat, K., Yongvanitchit, K., Tongtawe, P., and *et al.* (1999). Impairment of
1022 *Plasmodium falciparum* growth in thalassemic red blood cells: further Evidence by using
1023 biotin labeling and flow cytometry. *Blood.* **93**:3116-3119.
- 1024 Pepple, D.J. (1992). On the protective mechanism of the sickle cell trait (HbAS). *Med*
1025 *Hypotheses.* **39**:284-285.
- 1026 Pettenkofer, H.J. and Bickerich, R.(1960). [On common antigen properties between the human
1027 Blood group ABO and the pathogen of diseases dangerous for the community.Preliminary
1028 report.]. *Zentralbl Bakteriol.* ; **179**:433-436.
- 1029 Pettenkofer, H.J., Stoess, B., Helmbold, W. and Vogel, F. (1962). Alleged causes of the present-
1030 Day world distribution of the human ABO blood groups. *Nature.* ; **193**:445-446.
- 1031 Pipitaporn, B., Sueblinvong, T., Dharmkrong-at, A. and Udomsangpetch, R. (2000). Rosetting of
1032 *Plasmodium falciparum* required multiple components of the Uninfected erythrocytes.
1033 *Asian Pac J Allergy Immunol*; **18**:29-35.
- 1034 Pongponratn, E., Riganti, M., Punpoowong, B. and Aikawa, M. (1991). Microvascular of
1035 parasitized erythrocytes in human falciparum malaria: a pathological study. *Am J Trop*
1036 *Med Hyg.* **44**:168-175.
- 1037 Portilo, W.R., and Sullivan, C.C. (1979).Genetic factors that influence immune regulatory
1038 potentials in females against some diseases. *Journ Biotech.* **2**:111-115
- 1039 Pruss, A., Salama, A., Ahrens, N. and *et al.* (2003). Immune hemolysis-serological and clinical
1040 aspects. *Clin Exp Med.*; **3**:55-64.
- 1041 Ramon, chin, Beng-Ong, Patricia, Kim, Chooi Lim, Joon Wah, Mak. (2011). Detection of
1042 glycoprotein from human erythrocytes of different ABO blood groups infected with
1043 *Plasmodium falciparum*. **5**(2):18-28
- 1044 Rank, B.H., Carlsson, J. and Hebbel, R.P. (1985). Abnormal redox status of membrane-protein
1045 thiols in sickle erythrocytes. *J Clin Invest.* ; **75**:1531-1537.

- 1046 Raper, A.B. (1968). ABO blood groups and malaria. *Trans R Soc Trop Med Hyg*; **62**:158- 159.
- 1047 Reid, M.E., and Lomas-Francis, C. (2003). The blood group antigen factsbook (ed
- 1048 2nd).Amsterdam; Boston: Elsevier/Academic Press;
- 1049 Rifkind, J.M., Nagababu, E., Ramasamy, S., and Ravi, L.B., (2003). Hemoglobin redox reactions
- 1050 and Oxidative stress. *Redox Rep.*; **8**:234-237.
- 1051 Riganti, M., Pongponratn, E., Tegoshi, T., Looareesuwan, S., Punpoowong, B., Aikawa, M.
- 1052 (1990). Human cerebral malaria in Thailand: a clinico-pathological correlation *Immunol*
- 1053 *Lett.*; **25**:199-205.
- 1054 Rivadeneira, E.M., Wasserman, M. and Espinal, C.T. (1983). Separation and concentration of
- 1055 Schizonts of *Plasmodium falciparum* by Percoll gradients. *J Protozool*; **30**:367-370.
- 1056 Roberts, D.J, Craig, A.G., Berendt, A.R. and *et al.* (1992). Rapid switching to multiple antigenic
- 1057 And adhesive phenotypes in malaria. *Nature.* ; **357**:689-692
- 1058 Roberts, D.J. and Williams, T.N. (2003). Haemoglobinopathies and resistance to malaria.*Redox*
- 1059 *Rep.*; **8**:304-310.
- 1060 Rosenfield, R.G. (1979). Blood typing and associated immunohematological test procedures.In:
- 1061 manual of clinical immunology, American society for Microbiology, Washington,
- 1062 DC.**526**:111-230
- 1063 Rosenthal, P.J., McKerrow, J.H., Aikawa, M., Nagasawa, H. and Leech, J.H. (1988). A malarial
- 1064 Cysteine proteinase is necessary for hemoglobin degradation by *Plasmodium falciparum*. *J*
- 1065 *Clin Invest.* ;**82**:1560-1566.
- 1066 Ross, M.H., Kaye, G.I. and Pawlina, W. (2003). Histology: a text and atlas (ed 4th).
- 1067 Philadelphia:Lippincott Williams & Wilkins.
- 1068 Roth, E.F., Jr., Raventos-Suarez, C., Rinaldi, A. and Nagel, R.L. (1983) .Glucose-6-phosphate
- 1069 Dehydrogenase deficiency inhibits in vitro growth of *Plasmodium falciparum*. *Proc Natl*
- 1070 *Acad Sci U S A.*; **80**:298-299.
- 1071 Rowe, A., Obeiro, J., Newbold, C.I. and Marsh, K.(1995). *Plasmodium falciparum* rosetting is
- 1072 Associated with malaria severity in Kenya. *Infect Immun.*; **63**:2323-2326.
- 1073 Rowe, J.A., Handel, I.G., Thera, M.A. and *et al.* (2007). Blood group O protects against severe
- 1074 *Plasmodium falciparum* malaria through the mechanism of reduced rosetting. *Proc Natl*
- 1075 *Acad Sci U S A.* 2007; **104**:17471-17476.
- 1076 Rubaschkin, W.J., Leisermann, L.I. (1927). Blutgruppen und Krankheiten. Ukrainisches Zentrabl
- 1077 Blutgruppenforsch; **3**:301-310
- 1078 Ruwende, C. and Hill, A. (1998). Glucose-6-phosphate dehydrogenase deficiency and malaria. *J*
- 1079 *Mol Med.*; **76**:581-588.
- 1080 Ruwende, C., Khoo, S.C., Snow, R.W. and *et al.* (1995). Natural selection of hemi- and
- 1081 Heterozygotes for G6PD deficiency in Africa by resistance to severe malaria.
- 1082 *Nature*;**376**:246-249.
- 1083 Saitou, N. and Yamamoto, F. (1997). Evolution of primate ABO blood group genes and their
- 1084 Homologous genes. *Mol Biol Evol.* ; **14**: 399-411.

- 1085 Schacter, L., Warth, J.A., Gordon, E.M., Prasad, A. and Klein, B.L. (1988). Altered amount and
1086 Activity of superoxide dismutase in sickle cell anaemia. *Faseb J.*; **2**:237-243.
- 1087 Schellenberg, D., Menendez, C., Kahigwa, E. and *et al.* (1999). African children with malaria in
1088 An area of intense *Plasmodium falciparum* transmission: features on admission toThe
1089 hospital and risk factors for death. *Am J Trop Med Hyg?* ; **61**:431-438.
- 1090 Schenkel-Brunner H. (2000). Human blood groups: chemical and biochemical basis of antigen
1091 Specificity (ed 2nd, completely revised). Wien; New York: Springer.
- 1092 Schenkel-Brunner, H. (1981). Studies on Blood-Groups A1 and A2. *EurJBiochem.* ; **122**:511-
1093 514.
- 1094 Schofield, A.E., Reardon, D.M. and Tanner, M.J. (1992). Defective anion transport activity of
1095 The abnormal band 3 in hereditary ovalocytic red blood cells. *Nature*.**355**:836-838.
- 1096 Schofield, L., and Grau, G.E. (2005). Immuno N 0logical processes in malaria pathogenesis. *Nat*
1097 *Rev Immunol.* **5**:722-735.
- 1098 Schrier, S.L. (1994). Thalassemia: pathophysiology of red cell changes. *Annu Rev Med.* **45**:211-
1099 218.
- 1100 Schwarzer, E., Turrini, F., and Arese, P. (1994). A luminescence method for the quantitative
1101 Determination of phagocytosis of erythrocytes, of malaria-parasitized erythrocytes And of
1102 malarial pigment. *Br J Haematol.* ; **88**:740-745.
- 1103 Schwarzer, E., Turrini, F., Ulliers, D., Giribaldi, G., Ginsburg, H. and Arese, P.
1104 (1992).Impairment of macrophage functions after ingestion of *Plasmodium falciparum*-
1105 infected Erythrocytes or isolated malarial pigment. *J Exp Med.*; **176**:1033-1041.
- 1106 Seltsam, A., Hallensleben, M., Kollmann, A. and Blasczyk, R. (2003). The nature of diversity
1107 and Diversification at the ABO locus. *Blood.* **102**:3035-3042.
- 1108 Senga, E., Loscertales, M.P., Makwakwa, K.E., and *et al.* (2007). ABO blood group phenotypes
1109 Influence parity specific immunity to *Plasmodium falciparum* malaria in Malawian
1110 Women. *Malar J.*; **6**:102-112
- 1111 Senok, A.C., Li, K., Nelson, E.A., Yu, L.M., Tian, L.P. and Oppenheimer, S.J.(1997). Invasion
1112 and Growth of *Plasmodium falciparum* is inhibited in fractionated thalassaemic
1113 erythrocytes.*Trans R Soc Trop Med Hyg.***91**:138-143.
- 1114 Serghides, L., and Kain, K.C. (2001). Peroxisome proliferator-activated receptor gamma-retinoid
1115 X Receptor agonists increase CD36-dependent phagocytosis of *Plasmodium falciparum*-
1116 Parasitized erythrocytes and decrease malaria-induced TNF-alpha secretion by
1117 Monocytes/macrophages. *J Immunol.* ; **166**: 6742-6748.
- 1118 Serghides, L., Patel, S.N., Ayi, K. and Kain, K.C. (2006). Placental chondroitin sulfate A-
1119 binding Malarial isolates evade innate phagocytic clearance. *J Infect Dis.*; **194**:133-139.
- 1120 Serghides, L., Smith, T.G., Patel, S.N., and Kain, K.C. (2003). CD36 and malaria: friends or
1121 foes?*Trends Parasitol.* ; **19**:461-469.
- 1122 Severe *falciparum* malaria. (2000). World Health Organization, Communicable Diseases Cluster.
1123 *Trans R Soc Trop Med Hyg.* ; **94** Suppl 1:S1-90.

- 1124 Shear, H.L., Srinivasan, R., Nolan, T. and Ng, C. (1989) Role of IFN-gamma in lethal and
1125 nonlethal Malaria in susceptible and resistant murine hosts. *J Immunol.* ; **143**:2038-2044.
- 1126 Sheffield, W.P., Tinmouth, A. and Branch, D.R.(2005). Blood group biochemistry: a Canadian
1127 Blood services research and development symposium. *Transfus Med Rev.*; **19**:295-307.
- 1128 Sherman, I.W. (2005). Molecular approaches to malaria. Washington, D.C.: ASM Press.
- 1129 Sherman, I.W., Eda, S., and Winograd, E. (2004). Erythrocyte aging and malaria. *Cell Mol Biol*
1130 (Noisy- le-grand). ; **50**:159-169.
- 1131 Silamut, K., Phu, N.H., Whitty, C., and *et al.* (1999). A quantitative analysis of the
1132 microvascular Sequestration of malaria parasites in the human brain. *Am J Pathol.* ;
1133 **155**:395-410.
- 1134 Sim, B.K., Chitnis, C.E., Wasniowska, K., Hadley, T.J. and Miller LH. (1994) Receptor and
1135 ligand Domains for invasion of erythrocytes by *Plasmodium falciparum*. *Science.* ;
1136 **264**:1941-1944.
- 1137 Singh, N., Shukla, M.M., Uniyal, V.P. and Sharma, V.P. (1995). ABO blood groups among
1138 malaria Cases from district Mandla, Madhya Pradesh. *Indian J Malariol.* ; **32**:59-63.
- 1139 Snow, R.W., Guerra, C.A., Noor, A.M., Myint, H.Y., and Hay, S.I. (2005). The global
1140 distribution of Clinical episodes of *Plasmodium falciparum* malaria. *Nature.* ; **434**:214-
1141 217.
- 1142 Spector, W.G. (1968). The origin of mononuclear cells in inflammatory exudates. *Res Publ*
1143 *Assoc Res Nerv Ment Dis.*; **44**:1-13.
- 1144 Springer, G.F. and Horton, R.E. (1969). Blood group isoantibody stimulation in man by feeding
1145 Blood group-active bacteria. *J Clin Invest.* **48**:1280-1291.
- 1146 Springer, G.F. and Wiener, A.S. (1962). Alleged causes of the present-day world distribution of
1147 The human ABO blood groups. *Nature.* **193**:444-445.
- 1148 Stockelberg, D., Hou, M., Rydberg, L., Kutti, J., and Wadenvik, H.(1996) Evidence for an
1149 Expression of blood group a antigen on platelet glycoproteins IV and V. *Transfus*
1150 *Med.***6**:243-248
- 1151 Stormoken, H.E.J. (1977). Plasma antithrombin III and factor VIII antigen in relation to
1152 Angiographic findings, angina and blood groups in middle-aged men. *Thrombosis And*
1153 *Haemostasis.* ; **38**:874-880.
- 1154 Stubbs, J., Simpson, K.M., Triglia, T., and *et al.* (2005). Molecular mechanism for switching of
1155 *P. falciparum* invasion pathways into human erythrocytes. *Science.* ; **309**: 1384-1387.
- 1156 Tanabe, K. (1990). Ion metabolism in malaria-infected erythrocytes. *Blood Cells.* ; **16**:437- 449.
- 1157 Tang, T.K. (1997S). Free radical theory of erythrocyte aging. *J Formos Med Assoc.*; **96**:779-
1158 783.
- 1159 Thakur, A., and Verma, I.C. (1992). Malaria and ABO blood groups. *Indian J Malariol.* ;
1160 **29**:241- 244.
- 1161 Trager, W., and Jensen, J.B. (1976). Human malaria parasites in continuous culture. *Science*;
1162 **193**:673-675.

- 1163 Trampuz, A., Jereb, M., Muzlovic, I., and Prabhu, R.M. (2003). Clinical review: Severe malaria.
1164 Crit Care. ; **7**:315-323.
- 1165 Trentin, J.J. (1967). National Research Council. Committee on Tissue Transplantation. Cross-
1166 Reacting antigens and neoantigens; with implications for autoimmunity and cancer
1167 Immunity. Baltimore: Williams & Wilkins.**123**:233-288
- 1168 Tsaras, G., Owusu-Ansah, A., Boateng, F.O. and Amoateng-Adjepong, Y. (2009) .Complication
1169 Associated with sickle cell trait: a brief narrative review. *Am J Med.*; **122**:507-512.
- 1170 Turner, G.D., Morrison, H., Jones, M. and *et al.* (1994). An immunohistochemical study of the
1171 Pathology of fatal malaria. Evidence for widespread endothelial activation and a Potential
1172 role for intercellular adhesion molecule-1 in cerebral sequestration. *Am J Pathol.* ;
1173 **145**:1057-1069.
- 1174 Turrini, F., Ginsburg, H., Bussolino, F., Pescarmona, G.P., Serra, M.V. and Arese,
1175 P.(1992Phagocytosis of *Plasmodium falciparum*-infected human red blood cells by human
1176 Monocytes: involvement of immune and nonimmune determinants and dependence on
1177 Parasite developmental stage. *Blood.* ; **80**:801-808.
- 1178 Turrini, F., Giribaldi, G., Carta, F., Mannu, F. and Arese, P. (2003). Mechanisms of band 3
1179 oxidation And clustering in the phagocytosis of *Plasmodium falciparum*-infected
1180 erythrocytes. *Redox Rep.* **8**:300-303.
- 1181 Tursen,U.,Tiftik,En., Unal,S., Gunduz,O.,Kaya,T.I.and Camdevinen,H.(2005).Relationship
1182 between ABO blood groups and skin cancers. *Dermatol.Journ.***11**:44-47
- 1183 Tuteja, R. (2007). Malaria - an overview. *Febs J.***274**:4670-4679.
- 1184 Udomsangpetch, R., Sueblinvong, T., Pattanapanyasat, K., Dharmkrong-at, A., Kittikalayawong,
1185 A. and Webster, H.K. (1993). Alteration in cytoadherence and rosetting of *Plasmodium*
1186 *falciparum*-infected thalassemic red blood cells. *Blood.* ; **82**:3752- 3759.
- 1187 Udomsangpetch, R., Todd, J., Carlson, J., and Greenwood, B.M. (1993). The effects of
1188 Hemoglobin genotype and ABO blood group on the formation of rosettes by *Plasmodium*
1189 *falciparum*-infected red blood cells. *Am J Trop Med Hyg?*;**48**:149-153
- 1190 Uneke, C, J., Ogbu, O. and Nwojiji, V.(2006). Potential risk of induced malaria by blood
1191 transfusion in South-eastern Nigeria. *Meagul J Med.***9**:8-13
- 1192 Uneke, C.J. (2007). *Plasmodium falciparum* malaria and ABO blood group: is there any
1193 Relationship? *Parasitol Res.*; 100:759-765.
- 1194 Uzoegwu, P, N. and A.E.Onuwurah.(2003).Prevalence of heamoglobinopathy and malaria
1195 diseases in the population of old Aguata Division, Anambra state, Nigeria. *Biokemistri.*
1196 **15**:57-66
- 1197 Wambua, S., Mwangi, T.W., Kortok, M., and et al. (2006).The effect of alpha+-thalassaemia on
1198 the incidence of malaria and other diseases in children living on the coast of Kenya. *PLoS*
1199 *Med.*; 3:e158.
- 1200 Weatherall, D.J. (1998). Pathophysiology of thalassaemia. *Baillieres Clin Haematol.* ; **11**:127-
1201 146.

- 1202 Weatherall, D.J. (2008). Genetic variation and susceptibility to infection: the red cell and
1203 Malaria. *Br J Haematol.* ; **141**:276-286.
- 1204 Westhoff, C.M., Ferreri-Jacobia, M., Mak, D.O., and Foskett, J.K. (2002) . Identification of the
1205 Erythrocyte Rh blood group glycoprotein as a mammalian ammonium transporter. *J Biol*
1206 *Chem.*; 277:12499-12502.
- 1207 WHO, (1991).Basic malaria microscopy learner's guide. Geneva
1208 WHO. Vol. (2009).
- 1209 Widmaier, E.P., Raff, H., and Strang, K.T. (2008). Vander's human physiology: the mechanisms
1210 of body function (ed 11th). Boston: McGraw-Hill Higher Education; .
- 1211 Wiener, A.S. (1951). Origin of naturally occurring hemagglutinins and haemolysins; a review. *J*
1212 *Immunol.* ; **66**:287-295.
- 1213 Williams, T.N. (2006). Human red blood cell polymorphisms and malaria. *Curr Opin Microbiol.*
1214 **9**:388-394.
- 1215 Williams, T.N., Mwangi, T.W., Wambua, S. and *et al.* (2005) .Sickle cell trait and the risk of
1216 *Plasmodium falciparum* malaria and other childhood diseases. *J Infect Dis.*; **192**:178-186.
- 1217 Winograd, E., and Sherman, I.W.(1989). Characterization of a modified red cell membrane
1218 Protein expressed on erythrocytes infected with the human malaria parasite *Plasmodium*
1219 *falciparum*: possible role as a cytoadherent mediating protein. *J Cell Biol.*; 108:23-30.
- 1220 Winograd, E., Prudhomme, J.G. and Sherman, I.W.(2005). Band 3 clustering promotes the
1221 Exposure of neoantigens in *Plasmodium falciparum*-infected erythrocytes. *Mol Biochem*
1222 *Parasitol.* ; **142**:98-105.
- 1223 Wood, C.S. (1974). Preferential feeding of *Anopheles gambiae* mosquitoes on human subjects of
1224 blood group O: A relationship between the ABO polymorphism and malaria vectors.*Hum*
1225 *Biol.*; 46:385-404.
- 1226 Wood, C.S., Harrison, G.A., Dore, C. and Weiner, J.S. (1972).Selective feeding of *Anopheles*
1227 *Gambiae* according to ABO blood group status. *Nature.* ; **239**:165-178
- 1228 Worlledge, S., Ogiemudia, S.E., Thomas, C.O., Ikoku, B.N. and Luzzatto, L. (1974). Blood
1229 group antigens and antibodies in Nigeria. *Ann Trop Med Parasitol*; **68**:249-264.
- 1230 Yamamoto, F-i. (2009). ABO Blood Types. Vol. 200. California: The Owen Wilson Foundation
- 1231 Yawata, Y. (2003).Cell membrane: the red blood cell as a model. Weinheim: Wiley-VCH.
- 1232 Yayon, A., Vande, Waa JA., Yayon, M., Geary, T.G., and Jensen, J.B. (1983). Stage-dependent
1233 Effects of chloroquine on *Plasmodium falciparum* in vitro. *J Protozool.* ; **30**:642-647.
- 1234 Yenchitsomanus, P., Summers, K.M., Board, P.G., and *et al.* (1986). Alpha-thalassemia in Papua
1235 New Guinea. *Hum Genet.* ; **74**:432-437
- 1236 Yulia, Yin, k.P. (2008). Red Cell Physiology 101. Canadian Blood Services, 6th Annual
1237 International Symposium.
- 1238 Yuthavong, Y., Bunyaratvej, A. and Kamchonwongpaisan, S. (1990). Increased susceptibility of
1239 Malaria-infected variant erythrocytes to the mononuclear phagocyte system. *Blood Cells.* ;
1240 **16**:591-597.

- 1241 Yuthavong, Y., Butthep, P., Bunyaratvej, A. and Fucharoen, S. (1987). Inhibitory effect of beta
1242 Zero- thalassaemia/haemoglobin E erythrocytes on *Plasmodium falciparum* growth In
1243 vitro. *Trans R Soc Trop Med Hyg.* ; **81**:903-906.
- 1244 Zerihum, Abraham, Degarege. and Berhanu Erko. (2011).Association of ABO blood group and
1245 *Plasmodium falciparum* malaria in done Bafeno area. Southern Ethiopia. *Asia Journal of*
1246 *Tropical cell Biomedicine*.**6**:289-294.
- 1247 Zouali, M., Druilhe, P., Gentilini, M. and Eyquem. A. (1982). High titres of anti-T antibodies
1248 and other haemagglutinins in human malaria. *Clin Exp Immunol.* ; **50**:83-91.
- 1249
- 1250
- 1251
- 1252
- 1253
- 1254