

Review Article

Therapeutic arsenal against leishmaniasis: a review

ABSTRACT

Leishmaniasis are included in the group of reemerging neglected tropical diseases caused by flagellated parasites of the *Leishmania* genus. WHO recommends the use of chemotherapy as the main control measure for all the types of the disease. Antimonials are the first drug of choice, but they can be replaced by amphotericin B, miltefosine, paromomycin and pentamidine isethionate, or combinations. Generally, all the medicines that contain these drugs have several drawbacks. In view of this, chemotherapy for leishmaniasis treatment is increasingly the object of study, searching for safer or more effective drugs or new drug delivery vehicles, where nanotechnology is crucial. Thus, the purpose of this review is to present the therapeutic arsenal available in the market and that has been investigated for this application.

Keywords: neglected tropical diseases (NTD); leishmaniasis; chemotherapy; leishmanicidal; nanotechnology.

INTRODUCTION

Leishmaniasis is a parasite disease transmitted by the hematophagy of a Phlebotominae (*Phlebotomus* or *Lutzomyia* genus) whose major reservoir are the humans [1, 2, 3]. It is caused by more than 20 different species of a protozoa of the *Leishmania* genus, which is subdivided into two different subgenera, *L. (Leishmania)* spp. and *L. (Viannia)* spp., of the Kinetoplastida order, Tripanosomatidae family. Depending on the lifecycle stage, these protozoa take one of the two forms: with flagellum, named promastigote and without flagellum, amastigote. Based on the species of the infectious agent and the spectrum of their clinical manifestation, leishmaniasis are classified by World Health Organization (WHO) [4] into cutaneous leishmaniasis (CL) and visceral leishmaniasis (VL).

According to the Drugs for Neglected Diseases initiative (DNDi), leishmaniasis is one of the most important neglected diseases, as it occurs in 98 countries with 350 million people at risk. VL is endemic in 88 countries (across Asia, East Africa, South America and the Mediterranean region), of which 72 are low and middle income countries, but its impact on global public health is underestimated, since the notification of the disease is mandatory in only 32 of the affected countries. The most affected countries are Bangladesh, Brazil, India, Ethiopia, Kenya, Nepal and Sudan, which represent more than 90% of new cases; whereas the countries most affected by CL are Afghanistan, Algeria, Colombia, Brazil, Iran, Syria, Ethiopia, Sudan, Costa Rica and Peru [5].

This disease appears as reemerging in several regions, already described in countries such as Brazil and India [6-8]. This panorama is attributed to several factors such as: the range of zoonotic *Leishmania* species of unknown etiology that can affect humans; the occurrence of infection in other secondary hosts; human invasion of zoonotic foci; changes in susceptibility to infection of the human host [9], as for example, the co-infection *Leishmania*-HIV [10].

However, regardless of the form of manifestation of leishmaniasis, WHO recommends chemotherapy as the primary form of control.

LEISHMANICIDAL DRUGS IN THE MARKET

Antimonials

According to the WHO, antimonials are the first drug of choice [4].

Antimony is classified as a metalloid, and is found in nature as the isotopes ¹²¹Sb (57.25%) and ¹²³Sb (42.75%). Its most common oxidation states are Sb(III) and Sb(V) [11].

Comment [d1]: Everywhere Leishmaniasis need to be with capital L

Comment [d2]: Chemotherapy for Leishmaniasis

Comment [d3]: Leishmanicidal Drugs

The first global report on the use of antimonials in leishmaniasis treatment was by the Brazilian doctor Gaspar Vianna in 1912, who described the use of tartar emetic or antimony III and potassium tartrate [12]. However, in view of its toxicity (it normally accumulates in vascularized organs and tissues, especially liver and kidney, besides having high affinity for the spleen and blood) other antimony compounds have been tested, arriving to pentavalent antimonials [13]. In both cases, the compounds with therapeutic action are the result of the formation of water soluble complexes between organic acids and derived carbohydrates (tartaric acids or gluconic acid) and antimony [14].

Pentavalent antimonials

The antimonials used in leishmaniasis treatment are listed in figure 1 [13].




Figure 1. Proposed chemical structure of antimonials [13, 14]

N-methylglucamine antimoniate (Glucantime®, Sanofi-Aventis, France) is the antimonial compound used in Latin countries; and sodium stibogluconate (Pentostam®, GlaxoSmithKline, United Kingdom) is used in English-speaking countries. The difference in the choice of these drugs is not only due to the different *Leishmania spp.* species that cause the disease, but also to the inherent differences in sensitivity of the species to the available drugs and the site of the infection (CL, ML and VL forms), thus imposing different requirements on drug pharmacokinetics [15].

Dosage

Glucantime® is marketed in 5mL vials, containing 1.5g crude antimony and 405 mg/SbV. Then, each mL contains 81mg of SbV. According to the WHO, the dosage must be calculated in mg of SbV/kg/day aiming at standardizing the treatment. In 1982, WHO suggested optimal dose of 20mg/kg/day, for both, CL and VL treatments, limited to a maximum dosage of 850 mg/Sb/day for up to 28 days (usually 20 days for CL and 28 days for VL, with possible dose and/or therapy duration adjustments in individual cases – [5]). This dosage is well tolerated by children and malnourished people. The drug should be preferably intravenously administered by slow injection over 5 minutes, but gluteal intramuscular administration can also be used [16].

Marsden reports that conventional pentavalent antimonials cannot be orally administered due to poor absorption and/or inactivation in the stomach [17].

Mechanism of action

From the discovery of pentavalent antimonial drugs in 1940 until nowadays, they have been the first-line treatment in almost all geographic areas, through intramuscular, intravenous or intralesional administration for any infection by *Leishmania* [5; 18]. However, the mechanism

Comment [d4]: Is this proposed or is it the scientifically proven structure?

Comment [d5]: Sentence requires editing. What does Indians use? What to the Kenyans use? Only Latin and English speaking countries are mentioned here

of action and the reason of the selective toxicity of these leishmanicidal agents remain little known [19].

According to Sereno et al. [20], there are indications that trivalent antimony is substantially more potent than its pentavalent oxidation state against both parasite forms (amastigotes and promastigotes) of, at least three *Leishmania* species. Some authors have reported that trivalent antimony is the toxic element to intracellular *Leishmania* [20; 21]. Such results strengthen some hypotheses, as that from Miekeley et al. [22], who verified that after intramuscular administration of N-methylglucamine antimoniate in leishmaniasis patients, the stibiate (organic compound) is converted *in vivo* to the ionic forms SbIII and SbV. This result indicates that SbV is bio-reduced to its trivalent form, in accordance with other studies that show that the *in vivo* formation of SbIII is responsible for both the drug toxicity and its therapeutic activity. This theory was first proposed by Goodwin and Page, who believed that SbV could act as a pro-drug, thus needing to be converted to its active form, SbIII [23]. Frézard et al. [24] pointed out the importance of the thiols as a reducing agent in this conversion, emphasizing four different ones: glutathione (GSH), which is the main thiol in the cytosol of mammalian cells; cysteine (Cys) and cysteinyl-glycine (Cys-Gly), which are the predominant thiols within lysosomes, and the glutathione-spermine conjugate, trypanothione (T(SH)₂), which is the predominant thiol within the parasite.

The *Leishmania* species present an organelle characteristic of the Trypanosomatidae family, the glycosome, which contains nine enzymes involved in reactions of the glycolytic pathway, including glyceraldehyde 3-phosphate dehydrogenase (GAPDH). It acts both, in human hosts and in trypanosome, catalyzing the conversion of glyceraldehyde 3-phosphate to 1,3-diphosphoglycerate, in the presence of NAD⁺ and inorganic phosphate. This biochemical path is considered the main source of energy for promastigotes but it is also important for amastigotes. As a result, the depletion of intracellular ATP levels due to the interference of glycolysis and beta-oxidation of fatty acids in amastigotes is regarded as the main mechanism of action of antimonials against different *Leishmania* species [25].

Another possible mechanism of action of antimonials is the inactivation of a zinc-dependent metalloprotease in the amastigote form, through the substitution of antimony by zinc in that enzyme, thus inhibiting the development of the parasite [26].

Adverse effects

It is known that the pentavalent form does not bind to erythrocytes. Therefore, they reach higher concentration levels in plasma than the trivalent compounds. After consecutive administrations, there is an increase of the retention levels, transforming the trivalent form and accumulating in tissues such as spleen and liver, which causes the toxic effects related to antimony administration. The trivalent compounds bind in greater amount to the tissues and red blood cells, justifying their higher toxicity [27].

According to the systematic review by Oliveira et al., the most frequently reported adverse effects of antimonials are musculoskeletal pain, nausea, vomiting, diarrhea, abdominal pain, headache, anorexia, asthenia, fatigue, fever, pancreatitis, rash, erythema and hives; besides frequently reported reactions, such as phlebitis and edema associated with intravenous administration [28]. Side effects such as cardiotoxicity, pancreatitis and nephrotoxicity [29-31], can lead to patient hospitalization and monitoring and interruption of antimonial treatment.

Clinical tests show elevation of serum levels of pancreatic enzymes associated with abdominal pain. In addition, increased creatine phosphokinase and alkaline phosphatase, renal failure, eosinophilia, leucopenia and thrombocytopenia have also been reported; leading to treatment discontinuation in some cases [32].

Factors that lead to antimonial treatment substitution

The parenteral administration of these drugs requires a multiple dose regimen due to rapid renal clearance. When multiple consecutive doses are not administered, the dosage is not fulfilled, resulting in drug resistance [33]. Resistance can also occur because of the intrinsic differences among the different species that cause the disease in diverse geographic regions. Thus, reassessment of each type of treatment depending on VL and/or CL endemic areas is necessary [18].

When there is no satisfactory response and in some limited cases such as co-infection with HIV and treatment of pregnant women, treatment with pentavalent antimonials is not recommended and the adoption of second choice drugs is required [4].

Amphotericin B

In Bihar, an Indian state where VL is more endemic, the parasites are increasingly less susceptible to treatment with antimonials. This led to failure of the general treatment therewith, resulting in changing leishmaniasis treatment to conventional amphotericin B [18].

Amphotericin B is a polyene antibiotic (Figure 2) naturally produced by the actinomycete *Streptomyces nodosus* widely used in the treatment of most systemic mycoses affecting immunocompromised patients [35]. It was introduced in the pharmaceutical market in 1950 [36] and the reports on its use for leishmaniasis treatment in northern Bihar and districts adjacent to Nepal are from the 80s when the response to antimonials became less than 50% [37]. Nowadays, it is the first drug of choice for treating pregnant women and the second-line drug when there is no response to treatment with antimonials or when antimonials are not recommended such as cases of immunocompromised and HIV-VL co-infected patients [38].




Figure 2. Amphotericin B chemical structure

Four formulations are currently marketed: amphotericin B deoxycholate (Fungizone®, Bristol Meyers Squibb, United States), liposomal amphotericin B (AmBisome®, Gilead Sciences, United States), amphotericin B lipid complex (ABLC; Abelcet®, Enzon Pharmaceuticals, United States) and amphotericin B colloidal dispersion (ABCD; Amphotec™, InterMune Corp, Canada) [39]; however AmBisome® is the drug recommended by the WHO for leishmaniasis treatment [4].

Amphotericin B liposomal formulation was introduced in the European pharmaceutical market in 1989 [39] and the combination of the drug with the nanocarrier (small unilamellar vesicles – SUVs) allowed a reduction of the adverse effects, due to a specific release to macrophages of the liver, spleen and bone marrow, affected by leishmaniasis [39].

Dosage

For VL treatment it is also recommended the intravenous administration of 0.75–1 mg/kg of amphotericin B for 15–20 doses (daily or alternate days) and for CL treatment it is also recommended the administration of 0.7 mg/kg/day (intravenous infusion – 25/30 doses) or 0.7–1 mg/kg/day (intravenous infusion – 25/40 doses) [40].

In case of liposomal amphotericin B, WHO recommends a daily administration of 0.75 to 1mg/kg of, which comes in vials containing 50mg of freeze dried amphotericin B sodium deoxycholate through intravenous infusion during 14 to 20 days, with a 50mg minimum daily dose for VL treatment. For CL treatment it is also recommended the administration of 0.75 to 1mg/kg daily or on alternate days, limited to 50mg/day [5; 41]. Treatment duration should be determined based on clinical evolution, considering the speed of the response and the presence of co-morbidities [42].

Immunocompromised patients present higher tolerance and, therefore require higher amphotericin B total doses. Currently, 3mg/kg liposomal amphotericin B are intravenously administered on days 1 to 5, 14 and 21; or 3 mg/kg/day during 7 to 10 days; or 10 mg/kg/day for 2 days VL infected patients. Immunocompromised patients, HIV-VL co-infected patients receive 0.5 to 1.0 mg/kg/day until reaching a total dose of 1.5 to 2.0g amphotericin B and 2 to 4 mg/kg/day liposomal amphotericin B in consecutive days or 4mg/kg on alternate days, accumulating a total dose of 20 to 60mg/kg [43].

Mechanism of action

According to Chattopadhyay and Jafurulla, *Leishmania* binding and internalization into host cells involves its interaction with the cell membrane, thus requiring the presence of cholesterol to occur the pathogen-host interaction [44]. Then, cholesterol would play a key role

Comment [d6]: Kindly make a table to show different Ampho regimens currently in use and their recommendations for clarity and ease of reading for the reader

in protein organization and function of membranes and receptors. Amphotericin B binds to the cell membrane causing a specific change to sterols, resulting in pore formation, which would allow the leakage of intracellular substances and the parasite's cell death [35]. Liposomal amphotericin B specifically interacts with membrane sterols of the parasite (in this case, macrophages) membrane, sequestering them, and consequently, effectively reducing the ability to interact with cholesterol [36-42].

Adverse effects

The use of amphotericin B in the clinic is limited due to the development of nephrotoxicity, the most common adverse effect that may manifest as an acute kidney injury and tubular injury, including renal failure, hypokalemia and hypomagnesemia [46, 47]. However, side effects such as anemia, heart toxicity effect, nausea, vomiting, phlebitis, chills and fever are reported sometimes requiring treatment discontinuation [48]. Liposomal amphotericin B is preferred to conventional because it presents a milder toxicity profile [18], in addition to high potential and action spectrum, resulting in greater reliability of use [35]. The main advantage of the liposomal formulation is that it reduces renal and cardiac toxicity [35, 49].

Pentamidine

Pentamidine (Figure 3) is also used for CL systemic treatment [18]. It was introduced in the pharmaceutical market in 1937 but it was used as leishmanicide only in 1973. It is the drug of choice in cases of resistance to antimonials, when they are contraindicated or when the patient shows signs of liver or cardiac toxicity as a consequence of its use [50].




Figure 3. Pentamidine chemical structure

Dosage

Usually, pentamidine isethionate (Pentcarinat[®], Sanofi-Aventis, France) is administered at a dose of 4mg/kg on alternate days, with 20 intramuscular applications for CL treatment [51].

Mechanism of action

Its mechanism of action remains unknown. According to Basselin et al., amastigotes and promastigotes do not metabolize, but accumulate pentamidine against a concentration gradient through the cell membrane [52]. This accumulation results in polyamine transport inhibition, causing an imbalance between intra and extracellular polyamines, with consequent parasite growth inhibition.

Adverse effects

Pentamidine was the first drug to be used in patients resistant to antimony and this treatment was effective, although associated with adverse effects including pain at the injection site, nausea, fever, bitter taste, development of hypoglycemia, hypotension and type 1 diabetes [23]. Cardiac toxicity and nephrotoxicity have also been reported [53], besides effectiveness decreasing after consecutive administrations [39].

Paromomycin

Paromomycin (PM) or aminosidine is a broad spectrum antibiotic (Figure 4), active against *Leishmania* (promastigotes and amastigotes), and some enteric protozoa and bacteria [33, 34]. Its leishmanicidal action was discovered in 1980 and has been successfully administered for VL treatment by parenteral route and as topical agent for CL treatment as will be discussed in the relevant chapter of this manuscript ("Topic Treatment of American Tegumentary Leishmaniasis Lesions") [18]. It is indicated when the use of antimonials and amphotericin B is contraindicated [35], whether because of resistance to antimonials or because of adverse effects caused by them or amphotericin B. Paromomycin has fewer side effects than amphotericin B and is not sensitive to light [36].




Figure 4. Paromomycin chemical structure

Dosage

It is marketed as Humatin® (Pfizer, United States) in vials or capsules (parenteral or oral treatment). The former is indicated for leishmaniasis treatment, with a dose of 14-16 mg/kg/day for up to 3 weeks [37]. However, some studies report that a smaller dose of 11mg/kg/day intramuscular administered is also effective [38-39, 39].

Mechanism of action

According to Davidson et al., paromomycin binds to the 30S ribosomal subunit, impairing protein synthesis of the parasite causing its death through inhibition of its metabolism and mitochondrial respiration [34].

Adverse effects

Adverse effects are not common, but include elevation of liver function enzymes, reversible ototoxicity and renal failure [33].

Resistance to the drug and its *in vitro* induction for *Leishmania* have been reported. However, unlike resistant bacteria, *Leishmania* species do not present mutations in the 30S ribosomal subunit (main mechanism leading to resistance), but a lower paromomycin absorption [34].

Miltefosine

Miltefosine, trade name Impavido® (Paladin Labs, Canada), is an alkylphosphocholine analog (hexadecylphosphocholine) (Figure 5), which was originally developed as an antitumor agent, but that has proven to be clinically effective. It was recently introduced in the global pharmaceutical market for VL treatment [39].




Figure 5. Miltefosine chemical structure

Although the leishmanicidal biochemical mechanism of miltefosine remains uncertain, it shows good *in vitro* and *in vivo* efficiency against *Leishmania*, when orally administered to animals. In a study that investigated oral miltefosine tolerance for Indian VL, 29 out of 30 patients (97%) were healed with 100mg miltefosine/day (an average of 2.5mg miltefosine/kg/day) during 28 days [39].

Taking into consideration, that the other drugs used in leishmaniasis treatment are currently parentally administered, miltefosine is the only one with the possibility of oral administration [13; 31].

Dosage

Miltefosine is formulated in capsules. A daily dose from 1.5 to 2.5mg/kg is indicated for VL treatment of children above 3 years of age, adolescents, adults and the elderly; whereas for CL treatment, the daily dose indicated for children above 12 years of age weighing at least 30kg, adolescents and adults weighing less than 45kg is 100mg miltefosine, and 150mg/day for patients weighing more than 45kg [3].

Mechanism of action

Miltefosine is a phospholipid that, despite not knowing exactly how it acts as leishmanicidal in the human body, interacts with the protozoan's cell membrane, causing its death. This is attributed to the close similarity between protozoan and human plasma membranes [32]. Miltefosine is not considered toxic to mammalian cells because it does not induce the activation of natural killer cells, of cytotoxic spleen cells, the phagocytic activity of macrophages, or a humoral response per se [33].

Adverse effects

The limitations of this drug are: high cost, long half-life thus becoming vulnerable to rapid resistance development, and the need for monitoring gastrointestinal and liver side effects besides nephrotoxicity [34]. Increased serum levels of aminotransferases and creatine phosphokinase were also reported [35]. In addition, it is potentially teratogenic [5; 34].

COMBINATION THERAPY

According to Van Griensven and Diro, combination therapy has been increasingly exploited, especially in highly endemic areas, aiming at identifying a short, inexpensive and well-tolerated combination regimen [36]. The combination of drugs with different mechanisms of action can also help to delay resistance emergence and increase the therapeutic life of drugs, bringing new therapies to current administration possibilities [37].

Several therapeutic regimens using this strategy have been reported [34-35; 37]. Some examples are described as highly efficient and have been adopted and indicated by WHO: combination of liposomal amphotericin B (5mg/kg per infusion, single dose) and miltefosine (2.5mg/kg/day during 7 days) and the combination of liposomal amphotericin B (5mg/kg per infusion, single dose) and paromomycin (11mg/kg during 10 days) used for the treatment of VL caused by *L. donovani* in Bangladesh, Bhutan, India and Nepal. The combination between pentavalent antimonials (20mg/kg/day intramuscular or intravenously administered) and paromomycin (15mg/kg/day intramuscular administered) during 17 days is also indicated by WHO for the treatment of VL caused by *L. donovani* in East Africa (Ethiopia, Eritrea, Kenya, Somalia, Sudan and Uganda) and Yemen [4].

UNDER STUDY

The long list of adverse effects described for all the drugs currently marketed, together with parenteral administration, except for miltefosine, leading to frequent treatment abandonment by the patients, thus favoring the development of resistant strains [38], explains that leishmaniasis chemotherapy is increasingly being investigated. There are researches in literature on new chemical substances and plant extracts, as well as new formulations of known drugs, where nanotechnology is highlighted. These, then, are the object of the next topics of this paper.

Sitamaquine

Sitamaquine is an 8-aminoquinoline analog (Figure 6) whose formulation is under development by GlaxoSmithKline (GSK) and is currently undergoing phase III clinical trials in India and Kenya. It is known that sitamaquine interacts with anionic phospholipids, such as phosphatidylinositol (PI) or phosphatidylglycerol (PG), while no interaction has been described with zwitterions or sterols. After sitamaquine initial interaction with the lipid monolayer, the hydrophobic interactions between the aromatic rings of the former and the acyl groups of the

phospholipids of the latter, allow the complete insertion of the drug, with subsequent accumulation inside the parasite cells causing its death [68, 70].




Figure 6. Sitamaquine chemical structure

Sitamaquine, similar to miltefosine, is active by the oral route; however it has not shown to be efficient when topically administered for cutaneous leishmaniasis treatment [64, 71].

The side effects common to sitamaquine administration are: vomiting, abdominal pain, headache, methemoglobinemia, cyanosis, nephrotic syndrome, glomerulonephritis and nephrotoxicity [72].

Sitamaquine has shown *in vitro* activity against a range of species causing CL, with ED_{50} values against amastigotes varying between 2.9-19.0 μ M [73]. Another study by Jha et al., where 4 different sitamaquine doses (1.5; 1.75; 2.0 and 2.5mg/kg/day) were used during 28 days, for a total of 30 patients, concluded that the oral administration of sitamaquine is efficient for VL treatment under test conditions and that it was, in general, well tolerated [74]. Similarly, the oral administration of 2mg/kg/day sitamaquine during 28 days has been efficient and well tolerated in the treatment of 61 Kenyan patients. However, further studies are necessary to define dosage and duration of the ideal administration of this drug to better understand its adverse effects on the kidneys [75, 76].

Natural products

Plants are described as a potential source of new antiprotozoal drugs [77]. Many groups of natural products, including alkaloids, flavonoids, terpenoids and essential oils, have leishmanicidal activity already reported in the literature.

Alkaloids

Alkaloids are secondary metabolites, nitrogenated and generally alkaline (except colchicine, piperine and oximes), and, among other activities, their high leishmanicidal activity has been highlighted in scientific circles [78]. The literature reports were listed in chronological order in table 1.

Table 1. Investigation of alkaloid leishmanicidal activity

Alkaloid	Type	Evaluated species	Type of test	Result	Reference
N-demethylconodurine	indole	<i>L. amazonensis</i> amastigotes	<i>in vitro</i>	IC ₅₀ = 25µg/mL	[76]
dictylomide-A and dictylomide-B	quinoline	<i>L. amazonensis</i> promastigotes	<i>in vitro</i>	total lysis of promastigotes at 100 µg/mL	[77]
N-demethylholacurtine, 15α-hydroxyholaminee holacurtine	steroid	<i>L. donovani</i> promastigotes	<i>in vitro</i>	6.25 > IC ₅₀ > 1.56µg/mL	[78]
holamine				1.56 > IC ₅₀ > 0.39µg/mL	
dihydrocorynantheine	indole	<i>L. major</i> promastigotes	<i>in vitro</i>	IC ₅₀ = 1.65µM	[79]
corynantheine				IC ₅₀ = 1.12µM	
corynantheidine				IC ₅₀ = 2.81µM	
E-3-quinol-2-yl-prop-2-en-1-ol	quinoline	<i>L. amazonensis</i> amastigotes	<i>in vivo</i> administered by oral route in BALB/c mice	25mg/kg, during 15 days, reduced the parasite load in 54% and showed a significant reduction of the lesion weight of 22% Meglumine antimoniate (100mg/kg) reduced 98.4% and 87.3%, respectively	[80]
		<i>L. infantum</i> amastigotes		25 mg/kg, during 10 days, were able to reduce the parasite load by 83% Meglumine antimoniate (100mg/kg) reduced by 65%	
		<i>L. donovani</i> amastigotes		25 or 12.5mg/kg, during 10 days, were able to reduce the parasite load in 61% and 59%, respectively Mitelfosine (7.5mg/kg) reduced by 72%	
15,22-O-diacetyl-19-oxo-dihydroatisine	diterpenoid	<i>L. infantum</i> promastigotes	<i>in vitro</i>	IC ₅₀ = 12.8mg/L (after 72-hour culture), low toxicity (74.28mg/L)	[81]

azitine	isoquinoline	<i>L. braziliensis</i> promastigotes	<i>in vitro</i>	IC ₅₀ = 10.12mg/L (after 72-hour culture), high toxicity (> 200mg/L)	[82]
isoazitine				IC ₅₀ = 7.39mg/L (after 72-hour culture), high toxicity (> 300mg/L)	
O-methylmoschatolinee				IC ₅₀ = 320.8µg/mL	
liriodenine				IC ₅₀ = 58.5µg/mL	
neolitsine	isoquinoline	<i>L. mexicana</i> promastigotes	<i>in vitro</i>	IC ₅₀ = 15µM	[83]
cryptodorine				IC ₅₀ = 3µM	
duguetine β-N-oxide	isoquinoline	<i>L. braziliensis</i> promastigotes	<i>in vitro</i>	IC ₅₀ = 0.11µM	[84]
dicentrinone				IC ₅₀ = 0.01µM	

Flavonoids

Flavonoids are a group of polyphenolic compounds widely found throughout the plant kingdom and with known antioxidant activity [33].

Tasdemir et al. conducted a comprehensive study to assess the efficiency of several flavonoids and their analogs against *L. donovani* amastigotes [34]. The study showed that the most potent metabolites were fisetin ($IC_{50} = 0.6\mu g/mL$), 3-hydroxyflavone ($IC_{50} = 0.7\mu g/mL$), and luteolin ($IC_{50} = 0.8\mu g/mL$). The drug used as control, miltefosine, had IC_{50} of $0.34\mu g/mL$.

Vila-Nova et al. isolated rutin from *Dimorphandra gardneriana* grains, a native plant of northeastern Brazil caatinga biome and tested it against *L. infantum chagasi* parasites [35]. The compound presented results that were not significantly different from the results of the control drug – EC_{50} of $30.3\mu g/mL$ and $43.3\mu g/mL$ for promastigotes and amastigotes, respectively, whereas for amphotericin B the results were: no effect and $19.75\mu g/mL$, respectively.

Muzitano et al. [36] isolated quercitrin from *Kalanchoe pinnata* (Crassulaceae) and proved their leishmanicidal action by the reduction of parasitemia (57%) in BALB/c mice infected with *L. amazonensis* (murine model of CL), compared to Pentostam (8mg/kg twice a week). Some years before, Muzitano et al. had suggested the importance of quercetin aglycone-type structure and the rhamnosyl unit linked at C-3, in an unusual flavonoid (quercetin 3-O- α -L-arabinopyranosyl (1 $\rightarrow$ 2) α -L-rhamnopyranoside) isolated from the same plant, for the leishmanicidal activity. A $IC_{50} \cong 45mg/mL$ (78mM) against *L. amazonensis* amastigotes was obtained [39].

Quercetin (3,3',4',5'-7-penta-hidroxy-flavone) stands out among the flavonoids, because its leishmanicidal activity was confirmed in several studies, as shown in table 2 (in chronological order).

Table 2. Leishmanicidal activity of the flavonoid quercetin

Evaluated species	Type of test	Result	Reference
<i>L. donovani</i> amastigotes	<i>in vitro</i>	$IC_{50} = 45.5\mu M$	[30]
	<i>in vivo</i> orally administered to golden hamster	14mg/kg of the compound was able to reduce the spleen parasite load in 90%	
<i>L. donovani</i> amastigotes	<i>in vitro</i>	$IC_{50} = 1.0\mu g/L$	[36]
	<i>in vivo</i> intraperitoneal administered to BALB/c mice	30mg/kg during 5 days, was able to reduce infection in 15.3%	
<i>L. amazonensis</i> promastigotes	<i>in vitro</i>	$IC_{50} = 1.0\mu g/mL$	[31]
<i>L. infantum chagasi</i> amastigotes	<i>in vitro</i>	$EC_{50} = 10.6\mu g/mL$ (Amphotericin B: $ED_{50} = 19.75\mu g/mL$)	[32]
<i>L. infantum chagasi</i> promastigotes		$EC_{50} = 26\mu g/mL$ (Amphotericin B: $ED_{50} =$ no effect)	
<i>L. amazonensis</i> amastigotes	<i>in vitro</i>	$IC_{50} = 4.30\mu M$	[32]
<i>L. amazonensis</i> promastigotes	<i>in vitro</i>	$IC_{50} = 31.4\mu M$	[33]

Several authors tried to elucidate quercetin's leishmanicidal activity. Fonseca-Silva et al. attributed it to a pro-oxidant action generating reactive oxygen species (ROSs), which would lead to mitochondria dysfunction and ultimate death of the parasite [34]. Quercetin can be seen as a non-cytotoxic drug. Da Silva et al. elucidated, through docking analyses, the interaction of quercetin, quercitrin and isoquercitrin with Asp129 amino acid, which is involved in the formation of the bridge between the metal and Mn^{2+} A and B cofactors in arginase active site [34]. The

three compounds interacted with Asp129, which may indicate a correlation between leishmanicidal action and arginase-flavones interaction. But, from the three compounds, only quercetin interacted with L-arginine substrate and the Mn^{2+} cofactor of *L. amazonensis*, at pH 9.6 and was thus considered a mixed inhibitor.

Terpenoids

Terpenoids represent one of the largest and most diverse classes of secondary metabolites among the natural compounds. Although terpenoids do not derive from isoprene, but from mevalonic acid (MVA) and/or methylerythritol phosphate (MEP), this class is usually classified according to the number of isoprene units [32]. Table 3 lists the studies found in the literature for this application in chronological order.

Table3. Leishmanicidal activities of terpenoids

Terpenoid	Type of terpenoid	Evaluated species	Type of test	Result	Reference	
dihydrobetulinic acid	triterpenoid	<i>L. donovani</i> amastigotes	<i>in vitro</i>	IC ₅₀ = 4.1μM	[95]	
		<i>L. donovani</i> promastigotes		IC ₅₀ = 2.6μM		
		<i>L. donovani</i> promastigotes	<i>in vivo</i> orally administered to golden hamsters	10mg/kg during 6 weeks allowed to reduce the parasite load by 95.5% and 92.5% in spleen and liver, respectively; the compound did not present toxicity		
			<i>in vivo</i> intramuscular administered to golden hamsters	10mg/kg during 6 weeks allowed to reduce the parasite load by 97% and 99% in spleen and liver, respectively; the compound did not present toxicity		
maesabalide III	triterpene saponins	<i>L. donovani</i> amastigotes	<i>in vivo</i> subcutaneously administered to golden hamsters	0.8mg/kg allowed to reduce 94.2% of the amastigotes in liver (7-day treatment) AmBisome® (5 mg/kg) reduced 99.4%	[96]	
nerolidol	sesquiterpene	<i>L. amazonensis</i> amastigotes	<i>in vitro</i>	IC ₅₀ = 67μM	[97]	
		<i>L. amazonensis</i> promastigotes		IC ₅₀ = 85μM		
		<i>L. braziliensis</i> promastigotes		IC ₅₀ = 74μM		
		<i>L. chagasi</i> promastigotes		IC ₅₀ = 75μM		
		<i>L. amazonensis</i> amastigotes	<i>in vivo</i> intraperitoneally administered to BALB/c mice	100mg/kg during 12 days were able to reduce (P < 0.05) the size of the lesion, but, in the long term the disease was not healed. No toxic effects were observed		
		promastigotas de <i>L. amazonensis</i>				
ursolic acid	triterpenoid	<i>L. donovani</i> promastigotes	<i>in vitro</i>	IC ₅₀ = 3.7μg/mL (the drugs pentamidine and amphotericin B, were used as control and presented IC ₅₀ values of 1.9μg/mL and 0.7μg/mL, respectively)	[98]	

hautriwaic acid lactone				IC ₅₀ = 7.0µg/mL (idem)	
8,13-diacetyl- piptocarphol	lactone sesquiterpene	<i>L. amazonensis</i>		IC ₅₀ = 0.2µM	93
ursolic acid				IC ₅₀ = 0.99µM	
8-acetyl-13-O-ethyl- piptocarphol				IC ₅₀ = 0.37µM	
cumanin	sesquiterpene	<i>L. amazonensis</i> promastigotes	<i>in vitro</i>	growth inhibition higher than 80% at 5µg/mL (19µM) for both evaluated species	100
		<i>L. braziliensis</i> promastigotes			

441
442

Essential oils

Essential oils are derived from aromatic plants and include highly concentrated lipophilic compounds. As they are widely used in medical applications and have reported virucidal and antimicrobial activity, their leishmanicidal properties have been investigated [101].

The activity of *Chenopodium ambrosioides* essential oil on *L. amazonensis* has been investigated. BALB/c mice were infected and were treated with *Chenopodium ambrosioides* essential oil administered by intraperitoneal, oral and intralesional routes. Its activity against the parasite amastigotes and promastigotes was confirmed except for the intralesional administration (3%), which had no effect – the size of the lesion did not decrease ($P > 0.05$). The treatments by intraperitoneal and oral routes with 30mg/kg of essential oil resulted in a leishmanicidal activity greater than the control, 1mg/kg amphotericin B ($P > 0.05$). However, lesion size increase was observed during oral treatment ($P < 0.01$), and the authors concluded that the intraperitoneal route was the most efficient, although *Chenopodium ambrosioides* essential oil showed toxicity at that concentration causing a small abscess in the peritoneal cavity and death of 2 animals, after 25 injections. Healing was not observed in none of the animals treated [102].

In a recent study, which assessed the susceptibility of *L. infantum*, *L. tropica* and *L. major* to the essential oil *Cymbopogon citratus* and its main compounds mrycene and citral, *C. citratus* and citral exhibited a potential inhibition of *L. infantum*, *L. tropica* and *L. major*, with IC_{50} between 25, 52 and 38, and 42, 34 and 36 μ g/mL, respectively. Mrycene proved to be the less active compound, with an IC_{50} value of 164 μ g/mL for *L. infantum* and no effect on the other species evaluated [103].

According to Monzote et al., the essential oil linalol-rich, from *Croton cujacara* leaves has been successfully used against leishmaniasis parasites [104]. The oil proved to be efficient against *L. amazonensis* promastigotes and amastigotes with IC_{50} of 8.3 and 22ng/mL, respectively. No toxic effect on mammalian cells was observed.

Extracts

Plant extracts or compounds derived from plants are likely to provide a valuable source of new medicinal agents [105, 106], including those with potential leishmanicidal activity [107, 108]. The activity of these extracts largely depends on the solvent used for extraction, the part of the plant used and the presence of one or more natural products [109].

Moreira et al. demonstrated the leishmanicidal activity of *Stachytarpheta cayennensis* (Verbenaceae) hydroalcoholic extract [110]. In folk medicine, its crushed leaves and roots are already applied to the ulcerated lesion of tegumentary leishmaniasis. *L. braziliensis* and *L. amazonensis* amastigotes and promastigotes were used in the study. The leaf extract proved to be more efficient against *L. braziliensis* (IC_{50} of 3.7 μ g/mL) than against *L. amazonensis* (IC_{50} of 382.5 μ g/mL).

The leishmanicidal activity of the methanol extract of *Tridax procumbens* (Asteraceae) whole plant has already been described [111]. The research was conducted on the amastigotes of *L. mexicana*, causative agent of CL. *In vitro* studies with this extract showed growth inhibition of that parasite, with IC_{50} of 3 μ g/mL, whereas soxylipin (3S)-16,17-didehydrofalcarninol, a compound obtained from a *T. procumbens* exhibited greater inhibition, with IC_{50} of 0.478 μ g/mL. Both, the extract and the compound were obtained through chromatographic methods.

García et al. assessed the leishmanicidal activity of 48 extracts of 46 Cuban plants against *L. amazonensis* by *in vitro* bioassay [112]. The best result against amastigotes of that parasite were obtained with ethanol extracts of *Hura crepitans* leaves, with IC_{50} (27.7 $\pm$ 0.6 μ g/mL), followed by the extract from *Bambusa vulgaris* leaves and roots and the extract from *Simarouba glauca* leaves (IC_{50} (41.5 $\pm$ 0.6 μ g/mL and 45.5 $\pm$ 0.3 μ g/mL, respectively).

Pontin et al. assessed *in vitro* the leishmanicidal effect of different concentrations of Brazilian green propolis hydroalcoholic extract against *L. braziliensis* promastigotes (1, 10, 30, 50, 100, 250, 500 and 750 μ g/mL) and amastigotes (10, 100 and 250 μ g/mL) [113]. The results showed that the extract was inactive against amastigotes and it caused the lysis of 79.3% amastigotes at 500 μ g/mL, whereas amphotericin B used as control caused the lysis of 68.3% at the same concentration. *In vivo* effects were also evaluated using Brazilian green propolis extract in BALB/c mice infected with *L. braziliensis* promastigotes. For systemic evaluation, the infected animals were treated by the oral route. The extract concentration used was 1.5mg/kg/day and the lesion diameter was measured monthly. The intraperitoneal administration of 20mg/kg/day of Glucantime[®] was used as a control. The oral treatment with

propolis extract caused a significant reduction of 78.6% of the lesion diameter over 90-day treatment. **Glucantime®** caused a 57.7% reduction of the size lesion.

Ferreira et al. investigated the water extract of same kind of propolis and concluded that oral doses of this were able to reduce parasite load of in the liver of mice infected with *Leishmania infantum* [137].

Lantana ukambensis (Verbenaceae) methanol extract showed leishmanicidal activity with IC_{50} of 6.9 µg/mL against *L. donovani* promastigotes [138].

Nanotechnology

The research and development in nanotechnology involve structures with one (lamellae), two (nanotubes and nanofilaments) or three (nanoparticles) dimensions in the nanometric scale [139]. The last group stands out in Pharmaceutical Technology, where liposomes, polymeric nanoparticles and cyclodextrins are the most investigated nanocarriers for leishmaniasis chemotherapy, although metallic nanoparticles and lamellar clays have also been reported.

Liposomal amphotericin B is the sole example of drug based on nanotechnology, which has already been used in leishmaniasis treatment, as already mentioned in the present paper (AmBisome®, Gilead Sciences, Inc, Foster City, USA). It is basically composed by phospholipid molecules (hydrogenated soy phosphatidylcholine and distearoyl phosphatidylglycerol) arranged in a lipid bilayer [140,141], where the drug is included. Although it has been the standard treatment for VL, there are several limitations in the use, including high cost and nephrotoxicity [142]. According to Filippin and Souza, the formation of liposomal complexes has been investigated as resource to reduce this nephrotoxicity characteristic of amphotericin B administration [143].

Frézard et al. encapsulated meglumine antimoniate in liposomes by dehydration-rehydration procedure and obtained a high degree of encapsulation efficiency (28-58%), besides low indices of lipid weight [144]. Opposite to the formulations obtained through the conventional methods, this one can be stored lyophilized and reconstituted just prior to administration.

In other study, liposomal meglumine antimoniate reduced parasite load in liver and spleen of mice infected with *Leishmania infantum* [145]. Furthermore, the authors evaluated its association (single dose by intraperitoneal route) with water extract of green propolis (oral doses). Unexpectedly, there was no synergism in reducing the parasite load.

In an attempt to neutralize the damaging effects of the oxidative stress caused by the use of SbIII, Castro et al. [146] prepared liposomal nanoparticles of trivalent antimony associated with ascorbic acid. For evaluating their leishmanicidal action, BALB/c mice were infected with promastigotes of *L. infantum* by intravenous route. The results showed the preservation of leishmanicidal activity and confirmed the reduction of toxic effects on the liver and kidneys.

Amphotericin B was nanoencapsulated in poly (lactic-co-glycolic acid) (PLGA)/dimercaptosuccinic acid (DMSA) and associated (Nano-D-AMB-MG) or not (Nano-D-AMB) to magnetic nanoparticles, and evaluated in C57BL mice infected. The Nano-D-AMB achieved a greater reduction in leishmaniasis parasites if compared to the group treated with D-AMB (Fungizone®). Furthermore, a fewer frequency of dosages was required to achieve the same therapeutic levels. Regarding the Nano-D-AMB-MG, the formation of aggregates difficulted the delivery of the drug at the targeted site, so that the authors suggest further investigation [147].

Demicheli et al. assessed the complexation of meglumine antimoniate with β -cyclodextrin, which resulted in an increase of antimony absorption by the oral route and turned it orally active in a murine model infected by *L. amazonensis* [148]. The efficiency of the oral formulation was equivalent to that of meglumine antimoniate formulation intraperitoneally administered with a two times higher antimony dose. And in a more recent study, the authors also observed that lyophilization increased the oral absorption of Sb and contributed to a better controlled drug release [149].

Das et al. developed quercetin conjugates with gold nanoparticles [150]. Although quercetin shows leishmanicidal activity, it has a serious problem of systemic transportation due to its low water solubility. The nanostructured system proved to be efficient, with a complete absorption by macrophages with IC_{50} of 15, 40 and 30 µM against *L. donovani*, wild type, and strains resistant to sodium stibogluconate and paromomycin, respectively. These results were compared with quercetin (34; 150 and 75 µM, resp.), amphotericin B (0.2; 0.4 and 0.35 µM, resp.), sodium stibogluconate (3.6; 130 and 115 µM, resp.) and paromomycin (10; 380 and

330µM, resp.). The authors attributed the effect to synergism between quercetin and the nanocarrier, which can cause deficiencies in the oxygen metabolism of the parasite. The toxicity of those nanoparticles was considered low.

The effects of several metal and metal oxide nanoparticles were investigated against *L. major* in the presence of ultra-violet light (UV), infrared (IR) and in the dark. These nanomaterials have a known antibacterial effect, which mechanism is based on the production of reactive oxygen species (presence of UV), or on the generation of heat (IR), causing damage to the parasites. The results confirmed the leishmanicidal activity of all tested nanoparticles, especially the nanoparticles of silver and of titanium dioxide, which demonstrated the best results [124].

Menezes et al. produced a nanocomposite containing SbIII and lamellar clay (Mg, Al HDL) as nanocarrier, which showed ability to be internalized by macrophages, although its leishmanicidal activity is still under investigation [125]. Different from other nanocarriers, this one, which is inorganic, degrades at acidic pH and presents biocompatibility and thus, some authors considered it a potential nanocarrier in the treatment of intracellular diseases, such as leishmaniasis.

TOPIC TREATMENT OF AMERICAN TEGUMENTARY LEISHMANIASIS LESIONS

American tegumentary leishmaniasis is caused by species of the *Leishmania* or *Viannia* sub-genera [40]. The high incidence and severity of this disease make an early diagnosis essential in order to immediately begin treatment to prevent further tissue damages or, when not treated, the death of the patient [126].

Following diagnosis, the choice between topic or systemic treatment is done according to the evidence degree, geographic distribution, clinical manifestation and *Leishmania* species involved [127]. There is an increasingly number of studies on the efficiency of the topic treatment for one of the clinical manifestations of this type of leishmaniasis, cutaneous leishmaniasis.

WHO recommends paromomycin (ointment composed by 15% paromomycin and 12% methylbenzethonium chloride) twice a day as the first drug of choice for CL (*L. major*, *L. tropica*, *L. aethiopica* and *L. infantum*) treatment. However, the benefits obtained with topic treatment, were considered modest when compared with the placebo and the antimonials [128]. The intralesional administration of 1 to 5mL of antimony is considered as the treatment of second choice [129].

Imiquimod (Aldara®, Bayer, Germany), an imidazoquinoline amine used for the treatment of skin lesions caused by the human papillomavirus (HPV), genital warts and pre-malignant conditions [128] has shown high leishmanicidal potential [129]. It acts inducing the release of nitric oxide, which has a potent cytotoxic effect against *Leishmania* species, and subsequent macrophage activation [128].

Arevalo et al. proved its efficiency when associated to intramuscular administered meglumine antimoniate in Peruvian regions where the infection by *L. Peruvians* was endemic [130]. Twelve patients with active lesions which had not responded to previous intravenous or intramuscular treatment with antimonials (20mg/kg/day) were selected for imiquimod evaluation. 250mg of ointment (5% of drug) were spread on the lesions for 20 days, on alternate days, in conjunction with 20mg SbV/kg/day intramuscular administered during 20 consecutive days. The response to treatment was daily assessed by the same physician. The results suggest that the combined therapy of imiquimod and meglumine antimoniate should be considered for CL patients who did not respond to the initial treatment with only meglumine antimoniate.

Another study observed that one patient who did not present a good response to liposomal amphotericin B was successfully treated with imiquimod. However, the authors are not sure whether this improvement was due to previous treatments received by the patient, including amphotericin or even to spontaneous healing of the disease [131].

CONCLUSIONS

Leishmaniasis is a neglected disease which can trigger mild, moderate or severe reactions in infected patients. A small number of cases can be resolved by the patient's immune system, but the vast majority of cases require a pharmacological treatment. However, the drugs currently used and available in the pharmaceutical market are potentially toxic which may result on treatment interruption and consequent resistance of the parasites that cause the disease or even on the death of the patient, depending on the species. Connected to this, the need for

Comment [d7]: Topical treatment

Comment [d8]: Can omit

Comment [d9]: Sentence needs correction. Leishmaniasis does not cause "reactions"

Comment [d10]: Needs revision

622 affordable treatments is today the most serious drawbacks of existing therapies. In addition, this
623 review showed that most of the researches evaluate the effects of the compounds under test on
624 the promastigotes. This suggests the need for further studies on the human infective form
625 (amastigote), and a better knowledge of the vector and its behavior, the dynamics of the
626 disease and different hosts and reservoirs involved in the biological cycle for the development
627 and choice of the most adequate treatment (one or more chemotherapy agents, and its
628 administration route) as well as conducting a better control of the disease. So, the development
629 of non-profit organizations has helped to increase R&D, but until the lack of commercial and
630 political is overcome, the situation of millions of people at risk of infection unfortunately has no
631 perspectives to be improved.

Comment [d11]: Use abbreviations once full form has been given at least once

REFERENCES

1. Ngure PK, Kimutai A, Ng'ang'a ZW, Rukunga G, Tonui WK. A review of Leishmaniasis in Eastern Africa. *Journal of Nanjing Medical University*. 2009;23:79-86.
2. Pavli A, Maltezos HC. Leishmaniasis, an emerging infection in travelers. *International Journal of Infectious Diseases*. 2010;14:1032–1039.
3. Control of the leishmaniasis: report of a meeting of the WHO Expert Committee on the Control of Leishmaniases, Geneva, *World Health Organ Tech Rep Ser*. 1990;793:1-158.
4. CDC – Centers for Disease Control and Prevention (2013) Parasites –Leishmaniasis. Accessed 23 June 2014. Available: <http://www.cdc.gov/parasites/leishmaniasis/>.
5. DNDi - Drugs for Neglected Diseases initiative. Accessed 14 April 2014. Available: <http://www.dndi.org/diseases-projects/diseases/vl/current-treatment/current-treatment-cl.html>.
6. Arias JR, Monteiro PS, Zicker F. The reemergence of visceral leishmaniasis in Brazil. *Emerging Infectious Diseases*. 1996;2:145-146.
7. Sharma U, Redhu NS, Mathur P, Singh S. Re-emergence of visceral leishmaniasis in Gujarat, India. *Journal of Vector Borne Diseases*. 2007;44:230-232.
8. Arce A, Estirado A, Ordobas M, Sevilla S, García N, Moratilla L et al. Re-emergence of leishmaniasis in Spain: community outbreak in Madrid, Spain, 2009 to 2012. *Eurosurveillance*. 2013;18:1-9.
9. Ashford RW. The leishmaniases as emerging and reemerging zoonoses. *International Journal for Parasitology*. 2000;30:1269-1281.
10. Ostfeld RS, Glass, GE, Keesing F. Spatial epidemiology: an emerging (or reemerging) discipline. *Trends in Ecology & Evolution*. 2005;20:328-336.
11. Yan S, Jin L, Sun H. Antimony in Medicine. In: Gielen M, Tiekink ER, editors. *Metallotherapeutic Drugs and Metal-Based Diagnostic Agents: The Use of Metals in Medicine*. John Wiley & Sons; 2005:441-461.
12. Vianna G. Tratamento da Leishmaniose Tegumentar com Injeções Intravenosas de Tártaro Emético. In: *Anais do 7º Congresso Brasileiro de Medicina e Cirurgia*. 1912;4:426-428.
13. Rath S, Trivelin LA, Imbrunio TR, Tomazela DM, Jesús MN, Marzal PC. Antimoniais empregados no tratamento da leishmaniose: estado da arte. *Química Nova*. 2003;26:550-555.
14. Roberts WL, McMurray WJ, Rainey PM. Characterization of the antimonial antileishmanial agent meglumine antimonate (glucantime). *Antimicrobial Agents and Chemotherapy*. 1998;42:1076-1082.
15. Croft SL, Coombs GH. Leishmaniasis – current chemotherapy and recent advances in the search for novel drugs. *Trends in Parasitology*. 2003;19:502-508.
16. Davies CR, Kaye P, Croft SL, Sundar S. Leishmaniasis: new approaches to disease control. *BMJ*. 2003;326:377-382.
17. Marsden PD. Pentavalent antimonials: old drugs for new diseases. *Revista da Sociedade Brasileira de Medicina Tropical*. 1985;18:187-198.

18. Dorlo TPC, Balasegaram M, Beijnen JH, Vries PJ. Miltefosine: a review of its pharmacology and therapeutic efficacy in the treatment of leishmaniasis. *Journal of Antimicrobial Chemotherapy Advance*. 2012;67:2576-2597.
19. Soares-Bezerra RJ, Leon L, Genestra M. Recentes avanços da quimioterapia das leishmanioses: moléculas intracelulares como alvo de fármacos. *Brazilian Journal of Pharmaceutical Sciences*. 2004;40:139-149.
20. Sereno D, Cavaleira M, Zemzoumi K, Maquaire S, Ouaisi A, Lemesre JL. Axenically grown amastigotes of *Leishmania infantum* used as an in vitro model to investigate the pentavalent antimony mode of action. *Antimicrobial agents and chemotherapy*. 1998;42:3097-3102.
21. Gil ES, Cunha LC, Paula JR, Bezerra JCB, Aguiar FA. Leishmaníase: arsenal terapêutico e alvos moleculares. *Vita et Sanitas*. 2007;1:90-115.
22. Miekeley N, Mortari SR, Schubach AO. Monitoring of total antimony and its species by ICP-MS and on-line ion chromatography in biological samples from patients treated for leishmaniasis. *Analytical and Bioanalytical Chemistry*. 2002;372:495-502.
23. Goodwin LG, Page JE. A study of the excretion of organic antimonials using a polarographic procedure. *The Biochemical Journal*. 1943;37:198-209.
24. Frézard F, Demicheli C, Ribeiro RR. Pentavalent Antimonials: New Perspectives for Old Drugs. *Molecules*. 2009;14:2317-2336.
25. Balaña-Fouce R, Reguera RM, Cubria JC, Ordonez D. The pharmacology of leishmaniasis. *General Pharmacology*. 1998;30:435-443.
26. Bangs JD, Ransom DA, Nimick M, Christie G, Hooper NM. Cytocidal effects on *Trypanosoma brucei* and inhibition of *Leishmania major* GP63 by peptidomimetic metalloprotease inhibitors. *Molecular and Biochemical Parasitology*. 2011;114:111-117.
27. Brasil. Ministério da Saúde (2011) Manual de recomendações para diagnóstico, tratamento e acompanhamento de pacientes com a coinfeção Leishmania-HIV. Secretaria de Vigilância em Saúde. Brasília. Editora do Ministério da Saúde, 2011. 106 p. – ISBN 978-85-334-1771-7. Accessed 6 December 2013. Available: http://portal.saude.gov.br/portal/arquivos/pdf/leishmania_hiv_web_25_01_11.pdf.
28. Oliveira LF, Schubach AO, Martins MM, Passos SL, Oliveira RV, Marzochi MC et al. Systematic review of the adverse effects of cutaneous leishmaniasis treatment in the New World. *Acta Tropica*. 2011;118:87-96.
29. Gasser JRRA, Magill AJ, Oster CN, Franke ED, Grogl M, Berman JD. Pancreatitis induced by pentavalent antimonial agents during treatment of leishmaniasis. *Clinical Infectious Diseases*. 1994;18:83-90.
30. Zaghoul IY, Al-Jasser M. Effect of renal impairment on the pharmacokinetics of antimony in hamsters. *Annals of Tropical Medicine and Parasitology*. 2004;98:793-800.
31. Matoussi N, Ameer HB, Amor SB, Fitouri Z, Becher SB. Cardiotoxicity of N-methyl-glucamine antimoniate (Glucantime). A case report. *Médecine et Maladies Infectieuses*. 2007;37(suppl. 3):257-259.
32. Shahian M, Alborzi A. Effect of meglumine antimoniate on the pancreas during treatment of visceral leishmaniasis in children. *Medical Science Monitor*. 2009;15:290-293.
33. Freitas-Junior LH, Chatelain E, Kim HA, Siqueira-Neto JL. Visceral leishmaniasis treatment: What do we have, what do we need and how to deliver it? *International Journal for Parasitology: Drugs and Drug Resistance*. 2012;2:11-19.

34. Frézard F, Michalick MSM, Soares CF, Demicheli C. Novel methods for the encapsulation of meglumine antimoniate into liposomes. *Brazilian Journal of Medical and Biological Research*. 2000;33:841-846.
35. Filippin FB, Souza LC. Eficiência terapêutica das formulações lipídicas de anfotericina B. *Revista Brasileira de Ciências Farmacêuticas*. 2006;42:167-94.
36. Gallis HA. Amphotericin B: a commentary on its role as an antifungal agent and as a comparative agent in clinical trials. *Clinical Infectious Diseases*. 1996;22:145-147.
37. Sinha PK, Bhattacharya S. Single-dose liposomal amphotericin B: an effective treatment for visceral leishmaniasis. *The Lancet Global Health*. 2013;2:7-8.
38. Sundar S, Chakravarty J. Liposomal amphotericin B and leishmaniasis: dose and response. *Journal of Global Infectious Diseases*. 2010;2:159–160.
39. Brajtburg J, Bolard J. Carrier effects on biological activity of amphotericin B. *Clinical Microbiology Reviews*. 1996;9:512–531.
40. Goto H, Lindoso JAL. Cutaneous and Mucocutaneous Leishmaniasis. *Infectious Disease Clinics of North America*. 2012; 26:293–307.
41. WHO – World Health Organization (2014b) Box 1. Recommended treatment regimens for visceral leishmaniasis, ranked by preference. Accessed 14 March 2014. Available: http://who.int/leishmaniasis/research/978924129496_pp67_71.pdf.
42. Murray HW, Berman JD, Davies CR, Saravia NG. Advances in leishmaniasis. *The Lancet*. 2005;366:1561-1570.
43. Bern, C. Treatment of visceral leishmaniasis. UpToDate (2013). Accessed 23 April 2014. Available: <http://www.uptodate.com/contents/treatment-of-visceral-leishmaniasis>.
44. Chattopadhyay A, Jafurulla MD. A novel mechanism for an old drug: amphotericin B in the treatment of visceral leishmaniasis. *Biochemical and Biophysical Research Communications*. 2011;416:7-2.
45. Foglia F, Lawrence MJ, Demè B, Fragneto G, Barlow D. Neutron diffraction studies of the interaction between amphotericin B and lipid-sterol model membranes. *Scientific reports*. 2012;2(778):1–6.
46. Laniado-Laborín R, Cabrales-Vargas MN. Amphotericin B: side effects and toxicity. *Revista Iberoamericana de Micología*. 2009;26:223-227.
47. Ulozas E. Amphotericin B-Induced Nephrotoxicity. *Comprehensive Toxicology*. 2010;7:347-357.
48. Pimentel MIF, Baptista C, Rubin EF, Vasconcellos ECF, Lyra MR, Salgueiro MM. American cutaneous leishmaniasis caused by *Leishmania (Viannia) braziliensis* resistant to meglumine antimoniate, but with good response to pentamidine: a case report. *Revista da Sociedade Brasileira de Medicina Tropical*. 2011;44:254-256.
49. Cohen BE. Amphotericin B toxicity and lethality: a tale of two channels. *International Journal of Pharmaceutics*. 1998;162:95-96.
50. Lai A Fat EJ, Vrede MA, Soetosenojo RM, Lai A Fat RF. Pentamidine, the drug of choice for the treatment of cutaneous leishmaniasis in Surinam. *Tropical Medicine Rounds*. 2002;41:796–800.
51. Liegl U, Bogner JR, Goebel FD. Insulin-dependent diabetes mellitus following pentamidine therapy in a patient with AIDS. *Clinical Investigation*. 1994;72:1027-1029.

52. Costa JML, Moraes MSC, Saldanha ACR, Barral A, Burattini MN. Diabetes melitus associated with pentamidine isethionate in diffuse cutaneous leishmaniasis. *Revista da Sociedade Brasileira de Medicina Tropical*. 1995;28:405-407.
53. Basselin M, Lawrence F, Robert-Gero M. Pentamidine uptake in *Leishmania donovani* and *Leishmania amazonensis* promastigotes and axenic amastigotes. *Biochemical Journal*. 1996;315:631-644.
54. Davidson RN, Den Boer M, Ritmeijer K. Paromomycin. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 2009;103:653-660.
55. Santos A, Noronha E, Ferreira L, Carranza-Tamayo C, Cupolillo E, Romero G. Efeito de uma formulação hidrofílica de paromomicina tópica na leishmaniose cutânea em pacientes com contra-indicações de tratamento com antimonial pentavalente. *Revista da Sociedade Brasileira de Medicina Tropical*. 2008;41:444-448.
56. Wiwanitkit V. Interest in paromomycin for the treatment of visceral leishmaniasis (kala-azar). *Therapeutics and Clinical Risk Management*. 2012;8:323-328.
57. Singh S, Sivakumar R. Challenges and new discoveries in the treatment of leishmaniasis. *Journal of Infection and Chemotherapy*. 2004;10:307-315.
58. Seaman J, Boer C, Wilkinson R, de Jong J, de Wilde E, Sondorp E et al. Liposomal amphotericin B (AmBisome) in the treatment of complicated kala-azar under field conditions. *Clinical Infectious Diseases*. 1995;21:188-193.
59. UKMi – UK Medicines Information (2013) New Drugs Online. Accessed 18 February 2014. Available: http://www.ukmi.nhs.uk/applications/ndo/record_view_open.asp?newDrugID=2877.
60. Sundar S, Jha TK, Thakur CP, Engel J, Sindermann H, Fischer C. Oral miltefosine for Indian visceral leishmaniasis. *The New England Journal of Medicine*. 2002;347:1739-1746.
61. Paladin Labs Inc. Application for Inclusion of Miltefosine on WHO Model List of Essential Medicines (2010). Accessed 17 April 2014. Available: http://www.who.int/selection_medicines/committees/expert/18/applications/Miltefosine_application.pdf.
62. Croft SL, Engel J. Miltefosine – discovery of the antileishmanial activity of phospholipid derivatives. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 2006;100:S4-S8.
63. Perez-Victoria FJ, Sánchez-Cañete MP, Seifert K, Croft SL, Sundar S, Castanys S, Gamarro F. Mechanisms of experimental resistance of *Leishmania* to miltefosine: Implications for clinical use. 2006;9:26-39.
64. Jha TK, Sundar S, Thakur CP, Felton JM, Sabin AJ, Horton J. A phase II dose-ranging study of sitamaquine for the treatment of visceral leishmaniasis in India. *The American Journal of Tropical Medicine and Hygiene*. 2005;73:1005-1011.
65. Van Griensven J, Diro E. Visceral leishmaniasis. *Infectious Disease Clinics of North America*. 2012;26:309-22.
66. Sundar S, Sinha PK, Verma DK, Kumar N, Alam S, Pandey K. Ambisome plus miltefosine for Indian patients with kala-azar. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 2011;105:115-117.
67. Thakur CP, Oliaro P, Gothoskar S, Bhowmick S, Choudhury BK, Prasad S et al. Treatment of visceral leishmaniasis (kala-azar) with aminosidine (=paromomycin)-antimonial combinations,

a pilot study in Bihar, India. Transactions of the Royal Society of Tropical Medicine and Hygiene. 1992;86:615-616.

68. Laguna F, Videla S, Jiménez-Mejías ME, Sirera G, Torre-Cisneros J, Ribera E et al. Amphotericin B lipid complex versus meglumine antimoniate in the treatment of visceral leishmaniasis in patients infected with HIV: a randomized pilot study. Journal of Antimicrobial Chemotherapy. 2003;52:464–468.

69. Dueñas-Romero AM, Loiseau PM, Saint-Pierre-Chazalet M. Interaction of sitamaquine with membrane lipids of *Leishmania donovani* promastigotes. Biochimica et Biophysica Acta. 2007;1768:246–252.

70. Coimbra ES, Libong D, Cojean S, Saint-Pierre-Chazalet M, Salgadi A, Le Moyec L. Mechanism of interaction of sitamaquine with *Leishmania donovani*. The Journal of Antimicrobial Chemotherapy. 2010;65:2548–2555.

71. Loiseau PM, Cojean S, Schrével J. Sitamaquine as a putative antileishmanial drug candidate: from the mechanism of action to the risk of drug resistance. Parasite. 2011;18:115-119.

72. Garnier T, Brown MB, Lawrence MJ, Croft SL. *In vitro* and *in vivo* studies on a topical formulation of sitamaquine dihydrochloride for cutaneous leishmaniasis. Journal of Pharmacy and Pharmacology. 2006;58:1043–1054.

73. Wasunna MK, Rashid JR, Mbui J, Kirigi G, Kinoti D, Lodenyo H et al. A phase II dose-increasing study of sitamaquine for the treatment of visceral leishmaniasis in Kenya. The American Journal of Tropical Medicine and Hygiene. 2005;73:871-876.

74. Tiuman TS, Santos AO, Ueda-Nakamura T, Dias Filho BP, Nakamura CV. Recent advances in leishmaniasis treatment. International Journal of Infectious Diseases. 2011;15:525-532.

75. Mishra BB, Kale RR, Singh RK, Tiwari VK. Alkaloids: Future prospective to combat leishmaniasis. Fitoterapia. 2009;80:81-90.

76. Munoz V, Morretti C, Sauvain M, Caron C, Porzel A, Massiot G et al. Isolation of bis-indole alkaloids with antileishmanial and antibacterial activities from *Peschiera van heurkii* (syn. *Tabernaemontana van heurkii*). Planta Medica. 1994;60:455-459.

77. Lavaud C, Massiot G, Vasquez C, Moretti C, Sauvain M, Balderrama L. 4-Quinolinone alkaloids from *Dictyoloma peruviana*. Phytochemistry. 1995;40:317-320.

78. Kam T, Sim K, Koyano T, Toyoshima M, Hayashi M, Komiyama K. Cytotoxic and leishmanicidal aminoglycosteroids and aminosteroids from *Holarrhena curtisii*. Journal of Natural Products. 1998;61:1332-1336.

79. Staerk D, Lemmich E, Christensen J, Kharazmi A, Olsen CE, Jaroszewski JW. Leishmanicidal, antiplasmodial and cytotoxic activity of indole alkaloids from *Corynanthe pachyceras*. Planta Medica. 2000;66:531-536.

80. Nakayama H, Loiseau PM, Bories C, Torres de Ortiz T, Schinini A, Serna E et al. Efficacy of orally administered 2-substituted quinolines in experimental murine cutaneous and visceral leishmaniasis. Antimicrobial Agents and Chemother. 2005;49:4950-4956.

81. Gonzalez P, Marin C, Rodriguez-Gonzalez I, Hitos AB, Rosales MJ, Reina M et al. In vitro activity of C20-diterpenoid alkaloid derivatives in promastigotes and intracellular amastigotes of *Leishmania infantum*. International Journal of Antimicrobial Agents. 2005;25:136-141.

82. Costa EV, Pinheiro MLB, Xavier CM, Silva JRA, Amaral ACF, Souza ADL et al. A pyrimidine- β -carboline and other alkaloids from *Annona foetida* with antileishmanial activity. Journal of Natural Products. 2006;69:292-294.

929
930 83. Correa JE, Ríos CH, Castillo ADR, Romero LI, Ortega-Barría E, Coley PD et al. Minor
931 alkaloids from *Guatteria dumetorum* with antileishmanial activity. *Planta Medica*. 2006;72:270-
932 272.
933
934 84. Da Silva DB, Tulli EC, Militão GC, Costa-Lotufo LV, Pessoa C, de-Moraes MO et al. The
935 antitumoral, trypanocidal and antileishmanial activities of extract and alkaloids isolated from
936 *Duguetia furfuracea*. *Phytomedicine*. 2009;16:1059.
937
938 85. Lewin G, Cojean S, Gupta S, Verma A, Puri SK, Loiseau PM. In vitro antileishmanial
939 properties of new flavonoids against *Leishmania donovani*. *Biomedicine and Preventive*
940 *Nutrition*. 2011;1:168-171.
941
942 86. Tasdemir D, Kaiser M, Brun R, Yard V, Schmidt TJ, Tosunc F et al. Antitrypanosomal and
943 antileishmanial activities of flavonoids and their analogues: in vitro, in vivo, structure-activity
944 relationship, and quantitative structure-activity relationship studies. *Antimicrobial Agents and*
945 *Chemotherapy*. 2006;4:1352-1364.
946
947 87. Vila-Nova NS, Morais SM, Falcão MJC, Bevilacqua CML, Rondon FCM, Wilson ME et al.
948 Leishmanicidal and cholinesterase inhibiting activities of phenolic compounds of *Dimorphandra*
949 *gardneriana* and *Platymiscium floribundum*, native plants from Caatinga biome. *Pesquisa*
950 *Veterinária Brasileira*. 2012;32:1164-1168.
951
952 88. Muzitano MF, Falcão CAB, Cruz EA, Bergonzi MC, Bilia AR, Vincieri FF et al. Oral
953 Metabolism and Efficacy of *Kalanchoe pinnata* Flavonoids in a Murine Model of Cutaneous
954 Leishmaniasis. *Planta Medica*. 2009;75:307-311.
955
956 89. Muzitano MF, Tinoco LW, Guette C, Kaiser CR, Rossi-Bergmann B, Costa SS. The
957 antileishmanial activity assessment of unusual flavonoids from *Kalanchoe pinnata*.
958 *Phytochemistry*. 2006;67:2071-2077.
959
960 90. Mitra B, Saha A, Chowdhury AR, Pal C, Mandal S, Mukhopadhyay S et al. Luteolin, an
961 abundant dietary component is a potent anti-leishmanial agent that acts by inducing
962 topoisomerase II-mediated kinetoplast DNA cleavage leading to apoptosis. *Molecular Medicine*.
963 2000;6:527-541.
964
965 91. Muzitano MF, Cruz EA, de Almeida AP, Da Silva SAG, Kaiser CR, Guette C et al. Quercitrin:
966 an antileishmanial flavonoid glycoside from *Kalanchoe pinnata*. *Planta Medica*. 2006;72:81-83.
967
968 92. Da Silva ER, Maquiaveli Cdo C, Magalhães PP. The leishmanicidal flavonols quercetin and
969 quercitrin target *Leishmania (Leishmania) amazonensis* arginase. *Experimental Parasitology*.
970 2012;130:183-188.
971
972 93. Fonseca-Silva F, Inacio JD, Canto-Cavaleiro MM, Almeida-Amaral EE. Reactive oxygen
973 species production and mitochondrial dysfunction contribute to quercetin induced death in
974 *Leishmania amazonensis*. *Plos One*. 2011;6(2:e14666):1-7.
975
976 94. Phillips MA, Leon P, Boronat A, Rodriguez-Concepcion M. The plastidal MEP pathway:
977 unified nomenclature and resources. *Trends in Plant Science*. 2008;13:619-623.
978
979 95. Chowdhury AR, Mandal S, Goswami A, Ghosh M, Mandal L, Chakraborty D et al.
980 Dihydrobetulinic acid induces apoptosis in *Leishmania donovani* by targeting DNA
981 topoisomerase I and II: implications in antileishmanial therapy. *Molecular Medicine*. 2003;9:26-
982 36.
983
984 96. Maes L, Germonprez N, Quirijnen L, Van Puyvelde L, Cos P, Vanden Berghe D.
985 Comparative activities of the triterpene saponin maesabalide III and liposomal amphotericin B
986 (AmBisome) against *Leishmania donovani* in hamsters. *Antimicrobial Agents and*
987 *Chemotherapy*. 2004;48:2056-2060.
988

989 97. Arruda DA, D'Alexandri FL, Katzin AM, Uliana SRB. Antileishmanial activity of the terpene
990 nerolidol. *Antimicrobial Agents and Chemotherapy*. 2005;49:1679-1687.
991
992 98. Da Silva Filho AA, Resende DO, Fukui MJ, Santos FF, Pauletti PM, Cunha WR et al. In vitro
993 antileishmanial, antiplasmodial and cytotoxic activities of phenolics and triterpenoids from
994 *Baccharis dracunculifolia* D. C. (Asteraceae). *Fitoterapia*. 2009;80:478-482.
995
996 99. Odonne G, Herbette G, Eparvier V, Bourdye G, Rojas R, Sauvaine M et al. Antileishmanial
997 sesquiterpene lactones from *Pseudelephantopus spicatus*, a traditional remedy from the
998 Chayahuita Amerindians (Peru). Part III. *Journal of Ethnopharmacology*. 2011;137:875-879.
999
1000 100. Sülsen VP, Cazorla SI, Frank FM, Laurella LC, Muschietti LV, Catalán CA et al. Natural
1001 terpenoids from ambrosia species are active in vitro and in vivo against human pathogenic
1002 trypanosomatids. *PLoS Neglected Tropical Diseases*. 2013;7:1-10.
1003
1004 101. Rohloff J, Hymete A, Tariku Y. Studies in natural products chemistry. In: Atta-ur-Rahman,
1005 editor. *Plant-derived natural products for the treatment of leishmaniasis*. Elsevier. 2013;381-
1006 429.
1007
1008 102. Monzote L, Montalvo AM, Scull R, Miranda M, Abreu J. Activity, toxicity and analysis of
1009 resistance of essential oil from *Chenopodium ambrosioides* after intraperitoneal, oral and
1010 intralesional administration in BALB/c mice infected with *Leishmania amazonensis*: A
1011 preliminary study. *Biomedicine & Pharmacotherapy*. 2007;61:48-153.
1012
1013 103. Machado M, Pires P, Dinis AM, Santos-Rosa M, Alves V, Salgueiro L et al. Aldheides
1014 monoterpenes as potential anti-Leishmania agents: Activity of *Cymbopogon citratus* and citral
1015 on *L. infantum*, *L. tropica* and *L. major*. *Experimental Parasitology*. 2012;130:223-231.
1016
1017 104. Monzote L, Alarcón O, Setzer WN. Antiprotozoal Activity of Essential Oil. *Agriculturae*
1018 *conspectus scientificus*. 2012;4:167-175.
1019
1020 105. Carvalho PB, Ferreira EI. Leishmaniasis phytotherapy. Nature's leadership against an
1021 ancient disease-review. *Fitoterapia*. 2001;72:599-618.
1022
1023 106. Kayser O, Kiderlen AF. In vitro leishmanicidal activity of naturally occurring chalcones.
1024 *Phytotherapy Research*. 2001;15:148-152.
1025
1026 107. Chan-Bacab MJ, Peña-Rodríguez LM. Plant natural products with leishmanicidal activity.
1027 *Natural Product Reports*. 2001;18:674-688.
1028
1029 108. Rocha LG, Almeida JRGS, Macedo RO, Barbosa-Filho JM. Review of natural products with
1030 antileishmanial activity. *Phytomedicine*. 2005;12:514-535.
1031
1032 109. Adebayo OL, Suleman D, Samson AA. Natural products in antileishmanial drug discovery:
1033 a review. *Journal of Asian Scientific Research*. 2013;3:157-173.
1034
1035 110. Moreira RCR, Costa GC, Lopes TC, Bezerra JL, Guerra RNM, Rebêlo JMM et al. Efeito
1036 leishmanicida in vitro de *Stachytarpheta cayennensis* (Rich.) Vahl (Verbenaceae). *Revista*
1037 *Brasileira de Farmacognosia*. 2007;17:59-63.
1038
1039 111. Martín-Quintal Z, Moo-Puc R, González-Salazar F, Chan-Bacab MJ, Torres-Tapia LW,
1040 Peraza-Sánchez SR. In vitro activity of *Tridax procumbens* against promastigotes of *Leishmania*
1041 *mexicana*. *Journal of Ethnopharmacology*. 2009;122:463-477.
1042
1043 112. García M, Monzote L, Scull R, Herrera P. Activity of cuban plants extracts against
1044 *Leishmania amazonensis*. *ISRN Pharmacology*. 2012;2012(104540):1-7.
1045
1046 113. Pontin K, Da Silva Filho AA, Santos FF, Andrade e Silva ML, Cunha WR, Nanayakkara
1047 NPD et al. In vitro and in vivo antileishmanial activities of a Brazilian green propolis extract.
1048 *Parasitology Research*. 2008;103:487-492.

1049
1050 **114** Ferreira FM, Castro RAO, Batista MA, Rossi FMO, Silveira-Lemos D, Frézard F et al.
1051 Association of water extract of green propolis and liposomal meglumine antimoniate in the
1052 treatment of experimental visceral leishmaniasis. *Parasitology Research*. 2014;113:533-543.
1053
1054 **115** Sawadogo WR, Le Douaron G, Maciuk A, Bories C, Loiseau PM, Figadère B et al. In vitro
1055 antileishmanial and antitrypanosomal activities of five medicinal plants from Burkina Faso.
1056 *Parasitology Research*. 2012;110:1779–1783.
1057
1058 **116** ISO/TS 27687:2008 Nanotechnologies – Terminology and definitions for nano-objects –
1059 Nanoparticle, nanofibre and nanoplate.
1060
1061 **117** Patel PA, Patravale VB. AmbiOnp: solid lipid nanoparticles of amphotericin B for oral
1062 administration. *Journal of Biomedical Nanotechnology*. 2011;7:632–639.
1063
1064 **118** Carvalho RF, Ribeiro LF, Miranda-Vilela AL, Filho JS, Martins OP, Silva DOC et al.
1065 Leishmanicidal activity of amphotericin B encapsulated in PLGA–DMSA nanoparticles to treat
1066 cutaneous leishmaniasis in C57BL/6 mice. *Experimental Parasitology*. 2013;135:217–222.
1067
1068 **119** Lima SAC, Silvestre R, Barros D, Cunha J, Baltazar MT, Dinis-Oliveira RJ et al. Crucial
1069 CD8+ T-lymphocyte cytotoxic role in amphotericin B nanospheres efficacy against experimental
1070 visceral leishmaniasis. *Nanomedicine: Nanotechnology, Biology and Medicine*. 2014;10:1021-
1071 1030.
1072
1073 **120** Castro RAO, Silva-Barcellos NM, Licio CSA, Souza JB, Souza-Testasicca MC, Ferreira FM
1074 et al. Association of Liposome-Encapsulated Trivalent Antimonial with Ascorbic Acid: An
1075 Effective and Safe Strategy in the Treatment of Experimental Visceral Leishmaniasis. *Plos One*.
1076 2014;9:e104055.
1077
1078 **121** Demicheli C, Ochoa R, Silva JBB, De Melo AL, Falcão CAM, Rossi-Bergmann B et al. Oral
1079 delivery of meglumine antimoniate-beta-cyclodextrin complex for treatment of leishmaniasis.
1080 *Antimicrobial Agents and Chemotherapy*. 2004;48:100-103.
1081
1082 **122** Frézard F, Martins PS, Bahia APCO, Moyec LL, Melo AL, Pimenta AMC et al. Enhanced
1083 oral delivery of antimony from meglumine antimoniate/ β -cyclodextrin nanoassemblies.
1084 *International Journal of Pharmaceutics*. 2008;347:102–108.
1085
1086 **123** Das S, Roy P, Mondal S, Bera T, Mukherjee A. One pot synthesis of gold nanoparticles and
1087 application in chemotherapy of wild and resistant type visceral leishmaniasis. *Colloids and*
1088 *surfaces. B, Biointerfaces*. 2013;107:27-34.
1089
1090 **124** Jebalia A, Kazemib B. Nano-based antileishmanial agents: A toxicological study on
1091 nanoparticles for future treatment of cutaneous leishmaniasis. *Toxicology in Vitro*.
1092 2013;27:1896-1904.
1093
1094 **125** Menezes JB, da Silva T, dos Santos J, Catari E, Meneghetti M, da Matta C et al. Layered
1095 double hydroxides (LDHs) as carrier of antimony aimed for improving leishmaniasis
1096 chemotherapy. *Applied Clay Science*. 2014;91-92:127-134.
1097
1098 **126** Garcia AL, Parrado R, De Doncker S, Bermudez H, Dujardin JC. American tegumentary
1099 leishmaniasis: direct species identification of *Leishmania* in non-invasive clinical samples.
1100 *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 2007;101:368-371.
1101
1102 **127** WHO – World Health Organization. Control of leishmaniasis. Report of a meeting of the
1103 WHO Expert Committee WHO, Geneve. WHO Technical Report Series. 2010;949:1–186.
1104
1105 **128** Buates S, Matlashewski G. Treatment of experimental leishmaniasis with the
1106 immunomodulators imiquimod and S28463: efficacy and mode of action. *Journal of Infectious*
1107 *Diseases*. 1999;179:1485-1494.
1108

- 1109 129. Aguado M, Espinosa P, Romero-Maté A, Tardío JC, Córdoba S, Borbujo J. Outbreak of
1110 cutaneous leishmaniasis in Fuenlabrada, Madrid. *Actas Dermo-Sifiliográficas (English Edition)*.
1111 2013;104:334-342.
1112
1113 130. Arevalo I, Ward B, Miller R, Meng TC, Najjar E, Alvarez E et al. Successful treatment of
1114 drug-resistant cutaneous leishmaniasis in humans by use of imiquimod, an immunomodulator.
1115 *Clinical Infectious Diseases*. 2001;33:1847-1851.
1116
1117 131. Hervás JA, Martín-Santiago A, Hervás D, Rojo E, Mena A, Rocamora V et al. Old world
1118 *Leishmania infantum* cutaneous leishmaniasis unresponsive to liposomal amphotericin B
1119 treated with topical imiquimod. *The Pediatric Infectious Disease*. 2012;31:97-100.