

Biocidal activities of Solvent extracts of *Azadirachta indica* against Some Endemic Tropical Vector-borne diseases.

ABSTRACT

The biocidal efficacy of Chloroform (CEX), n-Hexane (HEX), Acetone (AEX) and Ethanol (EEX) extracts, of the bark and root of *Azadirachta indica* (Neem), were investigated against *Anopheles gambiae* and *Bulinus globosus* in a two-phased rapid and final screening test. Results of the phytochemical screening, indicated phytochemical constituents like; Flavonoids, Tannins, phenols, saponin, phytates, glycosides and triterpenes. All extracts were active (≥ 500 ppm), during the rapid screening phase; hence final screening phase was also carried out. Results of the final screening shows that, the snail (*B. globosus*), were more susceptible to all solvent extracts compared to the mosquito larvae (*An. gambiae*). The ethanol extract of the bark against the snail (EEXbS) induced the highest mortality with LC_{50} value of 0.35ppm compared to the least, chloroform extract of the root against the larvae (CEXrL, $LC_{50}=46.0$ ppm). The positive control induced mortality to both snails and larvae at 1ppm, while there was no mortality induced by the negative control. The results confirm solvent extracts of neem bark and root as potential biocidal agent against disease vectors.

Keywords: Vector-borne diseases, *Azadirachta indica*, *Anopheles gambiae*, *Bulinus globosus*, Solvent extracts.

1.0 INTRODUCTION

Over the past decades, malaria and schistosomiasis are becoming hyper-endemic vector-borne diseases in Africa. Malaria is transmitted by female mosquito belonging to the genus *Anopheles* (Ohimain et al., 2014; Aina et al., 2009), while schistosomiasis is transmitted by aquatic snails belonging to the genera *Biomphalaria*, *Bulinus* and *Oncomelania* (Angaye, 2013). There are about 3,500 species of mosquitoes worldwide, which are grouped into about 41 genera (Ohimain et al., 2014). Furthermore, only about 30-40% of the *Anopheles* mosquitoes transmit malaria (NCID, 2004; Ohimain et al., 2014). Diseases caused by Mosquito are prevalent in more than 100 countries across the world, with a morbidity burden affecting over 700million people globally (Ghosh et al., 2012). A global Statistic by the World Health Organization, shows that more than 3 billion people in endemic area are prone to malaria (WHO/UNICEF, 2005), with about 59, 38 and 3% of clinical cases in Africa, Asia, and America, respectively (Dibua et al., 2013).

In terms of morbidity burden, data show that schistosomiasis is next to malaria in ranking, amongst vector-borne diseases (Bassey et al., 2013; Angaye et al., 2013), schistosomiasis affects about 4-5% of the world population (Clerk et al., 1997; Angaye, 2013). The morbidity burden of schistosomiasis may vary depending on the intermediate host responsible for the disease (Angaye, 2013). For instance, Angaye (2013), reported that the genus *Bulinus* host parasites is mostly responsible for urinary schistosomiasis, while the genus *Biomphalaria* is the intermediate host for the parasite responsible for intestinal schistosomiasis. These vectors are endemic in regions such as Africa, Asia, South America and Middle East (Angaye, 2013).

Control measures like, chemotherapy and environmental manipulation had been put in place to check vector-borne diseases (Angaye, 2013). Chemotherapy has witnessed several challenges including insect resistance, inadequate resources, vector adaptation (Okumu et al., 2007; Ohimain et al., 2014). Furthermore, chemotherapy only abates the morbidity burden as the risk of re-infection still abounds (Agboola et al., 2011; Angaye, 2013; Ohimain et al., 2014), and even the use of synthetic pesticides affects non-targeted species which might infringe on the health of the ecosystem (Angaye, 2013; Angaye et al., 2014; Ohimain et al., 2014).

Multifaceted and eco-friendly vector control measures are recommended for an integrated vector management (Angaye, 2013; Angaye et al., 2014) especially the environmental manipulation of vectors prior to maturity (Ohimain et al., 2014). Contemporary, research focuses on eco-friendly vector control measures, especially the use of botanicals with biocidal efficacy (Okumu et al., 2007; Ohimain et al., 2014). Neem products contains over a hundred comprehensive bioactive metabolites (Drabu et al., 2012), that are potentially eco-friendly tools used for vector control (Adewole et al., 2012; Okumu et al., 2007; Mordue et al., 2000). Notwithstanding, it is established in literature that the efficacy of a plant-derived pesticide varies, depending on the solvent used

during extraction (Angaye, 2013), as such it became necessary to investigate the biocidal activities of some Solvent extracts of *Azadirachta indica* against malaria and schistosomiasis vectors.

2.0. MATERIALS AND METHODS

2.1. Collection of Plant Materials

The fresh bark and root of *A. indica* (Neem plant) were purchased from an alternate therapy center (Amolab Centre for Alternative Therapy), in Yenagoa Metropolis, Bayelsa State, Nigeria in March 2014. The plants were identified at the center by Dr. Adesekan (a Phyto-pathologist), and transported to the Federal Medical Center Yenagoa for bioassay.

2.2. Phytochemical Screening

Phytochemical screening of the bark and root extracts was carried out following the standard procedures (AOAC, 1995; Khan, 2012; Harbone), with slight modification. Prior to the bioassay, all the prepared plant extracts were subjected to preliminary phytochemical screening for the presence of alkaloids, flavanoids, tannins, saponin, phytates, glucosides, triterpenes and phenolic compounds. The degree of turbidity of the different extracts were ranked using the following scales; abundance (++), present (+) and absent (-).

2.3. Plant Extraction process

The fresh bark and root of the plant were pulverized with mortar and pestle. Two hundred grams of the pulverized plant were macerated in 500ml chloroform, hexane, acetone and ethanol for 72 hours. The filtrates were concentrated to dryness in a rotary evaporator at 65°C, and preserved at -4°C until it was ready for use.

2.3. Vector Collection/culture of Mosquito

Mosquito Larvae belonging to the genus *Anopheles* (*An. gambiae*), as well as aquatic snails belonging to the genus *Bulinus* (*B. globosus*), were used for the study. The larvae were cultured *in-vivo* in the wild using baits positioned around breeding conspicuous sites (4.94886N 6.34046E; 4.93816N 6.34006E). The baits used were slightly modified as per method described by Ohimain et al., (2014), using plastic container half-filled with water and stuffed with cotton wool as substrate for egg laying. The baits were constantly monitored for the conspicuous emergence of larvae. Prior to their transportation for the *in-vitro* bioassay, the emerging larvae (wrigglers) were identified as 3rd instar larvae by Dr. Bassey (a Parasitologist at the Niger Delta University). Prior to the bioassay, the larvae were placed on an enamel tray with de-chlorinated water (pH 7.4), and acclimatized to laboratory condition, using methods as described by Dibua et al., (2013). The Snails were collected from the wild and bred in a laboratory-designed aquarium. The aquaria were designed to suit the snail's natural habitat with stick, sand, some stones, and aerators to improve oxygen in the water (Angaye et al., 2013; Bassey et al., 2013). They were fed with lettuce (salad leaves), and acclimatize to laboratory conditions.

2.5. Experimental Set Up

Triplicate samples of 20 healthy larvae and snails (8±1mm), were distinctly placed in a 500ml solution of the various extracts, in a 24 hours exposure static non-renewal test. It was performed in accordance with the World Health Organization guidelines (WHO, 1965), with slight modification. The average mortality rates of organisms were observed and recorded from the replicates. Copper sulphate (1ppm) was used as the positive control (Bassey et al., 2013; Agboola et al., 2011), while 500ml of distilled water adjusted with 2.5ml of 10% dimethyl sulfoxide (DMSO) at pH 7.3, was used as the negative control (Dibua et al., 2013). The screening was carried out in two phases; the rapid screening test and the final screening test. These screening methods, with slight modifications are as described by Agboola et al., (2011).

2.6. Rapid Screening and Final Screening Test

Triplicate concentrations of 1000 ppm and 500 ppm were used to screen the larva and snails for total (i.e. 100%) mortality within 24 hours in order to detect the range of activity. The replicates of the extracts showed average mortality (i.e. 100% mortality) on larva or snails at 500ppm during the rapid screening. For the final screening, the extracts which were active at 500ppm and below (≤500ppm), were further screened (at different concentrations in descending order), in order to determine the average minimal lethal concentrations (LC₁₀₀).

2.8. Statistical Analysis

The median Lethal doses (LC₅₀) of all tested extracts of the bark and root, were estimated from the average minimal lethal concentrations by probit analysis using SPSS Version 20 (IBM SPSS Inc.). Furthermore, the dose-time dependency of the larvae and snails to each of the 4 extracts of the bark and root, were determined using Microsoft 2013 excel package with 5% error (Bassey et al., 2013).

3.0 RESULTS AND DISCUSSIONS

The results of the biocidal efficacy of the four solvent (i.e. chloroform, hexane, acetone and ethanol) extracts of the bark and root of neem plant (*A. indica*), against *An. Gambiae* and *B. globosus* are presented in Tables 1-3 and Figures 1 and 2. This was carried out according to the standards methods (WHO 1965; WHO/UNICEF, 2005), of assessing the toxicity of the different solvent extracts against the vectors.

Results of the phytochemical screening (Table 1) indicated phytochemical constituents like Flavonoids, Tannins, phenols, saponin, phytates, glycosides and triterpenes. Results of the Bark-extracts of *A. indica* showed stronger presence of most phytochemicals compared to the root-extracts. Notwithstanding, triterpenes was the predominant phytochemicals. Furthermore, it was also worthy of note that the ethanol extract yielded more phytochemicals compared to other solvents (Chloroform, n-hexane, acetone). Results of the rapid screening shows that all extracts tested were active as a result of the average total mortalities of the replicates (ALC₁₀₀), within range of 500-1000ppm (i.e. ALC₁₀₀=100±0.00%), as such all extracts tested proceeded to the final screening phase.

Table 1: Phytochemical screening of solvent extracts of the bark and root of *A. indica*

S/N	Phytochemicals	Chloroform Extracts		Hexane Extracts		Acetone Extracts		Ethanol Extracts	
	Plant organ	Bark	Root	Bark	Root	Bark	Root	Bark	Root
1.	Alkaloids	-	-	-	-	-	-	-	-
2.	Flavonoids	+	+	+	+	+	+	++	+
3.	Tannins	+	-	++	+	-	-	-	-
4.	Phenols	-	-	+	-	-	-	++	-
5.	Saponins	-	-	+	+	-	-	++	+
6.	Phytates	-	-	+	-	-	-	++	-
7.	Glycosides	+	+	++	-	+	+	++	+
8.	Triterpenes	++	-	++	+	++	+	++	++

++: Present in abundance; +: Present; -: Absent

Results of the final screening phase for all solvent extracts of the bark (SOLEx^B) against the larvae and snails (Table 2) shows that the highest LC₅₀ was induced by the ethanol extract (EEXb) against the snails (0.35ppm), followed by Hexane Extract of the bark (HEXb), whose LC₅₀ value was 0.45ppm. The acetone (AEXb) and Chloroform (CEXb) extracts had LC₅₀ values of 0.60ppm and 0.65ppm respectively. On the other hand, the larvae (which were less susceptible compared to the snails), had LC₅₀ values of 6.50ppm for the ethanol, 11.02ppm for Hexane, 12.14ppm for Acetone and 13.50ppm for Chloroform extracts. Meanwhile, the snails and larvae survived in the negative control, but the positive control was lethal to both snails and larvae at 1ppm.

Table 2: Results of final Screening for Solvent extracts of the Bark of *A. indica* against the Larvae and snail

Concentrations (ppm)	% MORTALITY RATES ± SD								CONTROLS	
	Chloroform Extract		Hexane Extract		Acetone Extract		Ethanol Extract		+ve	-ve
	Larvae	Snail	Larvae	Snail	Larvae	Snail	Larvae	Snails	1ppm Copper Sulphate	Distilled Water adjusted with DMSO
90-100	100±0.00	100±0.00	100±0.00	100±0.00	100±0.00	100±0.00	100±0.00	100±0.00		
80-90	100±0.00	100±0.00	100±0.00	100±0.00	100±0.00	100±0.00	100±0.00	100±0.00		
70-80	100±0.00	100±0.00	100±0.00	100±0.00	100±0.00	100±0.00	100±0.00	100±0.00		
60-70	100±0.00	100±0.00	100±0.00	100±0.00	100±0.00	100±0.00	100±0.00	100±0.00		
50-60	****	100±0.00	100±0.00	100±0.00	****	100±0.00	100±0.00	100±0.00		
40-50	****	100±0.00	****	100±0.00	****	100±0.00	100±0.00	100±0.00		
30-40	****	100±0.00	****	100±0.00	****	100±0.00	100±0.00	100±0.00		
20-30	***	100±0.00	****	100±0.00	***	100±0.00	****	100±0.00		
10-20	****	100±0.00	***	100±0.00	****	100±0.00	****	100±0.00		
0-10	****	100±0.00	****	100±0.00	****	100±0.00	****	100±0.00		
ALC ₁₀₀	67.50ppm	3.25ppm	55.10ppm	2.25ppm	60.07ppm	3.00ppm	32.50ppm	1.75ppm	1ppm	0ppm
LC ₅₀	13.50ppm	0.65ppm	11.02ppm	0.45ppm	12.14ppm	0.60ppm	6.50ppm	0.35ppm	0.2ppm	0ppm

*ALC100= Average Minimal Lethal dose; **** = Average Mortality less than 100%.

Result of the final screening for solvent extracts of the root ($SOLEx^R$) is presented in Table 3. $SOLEx^R$ were less active compared to $SOLEx^B$ (Table 2). Notwithstanding, for the $SOLEx^R$, the larvae were likewise less susceptible to all extracts compared to the snails, as similarly observed amongst $SOLEx^B$. Amongst the $SOLEx^R$ tested against the snails, Ethanol extract of the root ($EEXr$), was the most active ($LC_{50}=9.35\text{ppm}$), while Chloroform extract of the root ($CEXr$) was least active with LC_{50} value of 15.50ppm. In addition, Hexane extract of the root ($HEXr$), and Acetone extract of the root ($AEXr$) induced mortality with LC_{50} values of 10.31 and 13.00ppm respectively. On the other hand, $SOLEx^R$ against the larvae induced mortalities with LC_{50} values of 46.00ppm for $CEXr$, 28.03ppm for $HEXr$, 39.82ppm for $AEXr$ and 21.50ppm for $EEXr$ (Table 3).

Table 3: Results of final Screening for Solvent extracts of the Root of *A. indica* against the Larvae and snail

Concentrations (ppm)	% MORTALITY RATES $\pm$ SD								CONTROLS	
	Chloroform Extract		Hexane Extract		Acetone Extract		Ethanol Extract		+ve	-ve
	Larvae	Snail	Larvae	Snail	Larvae	Snail	Larvae	Snails	1ppm Copper Sulphate	Distilled Water adjusted with DMSO
225-250	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00		
200-225	****	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00		
175-200	****	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00		
150-175	****	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00	****	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00		
125-150	***	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00	****	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00		
100-125	****	100 $\pm$ 0.00	****	100 $\pm$ 0.00	****	100 $\pm$ 0.00	100 $\pm$ 0.00	100 $\pm$ 0.00		
75-100	****	100 $\pm$ 0.00	****	100 $\pm$ 0.00	****	100 $\pm$ 0.00	****	100 $\pm$ 0.00		
50-75	****	****	****	100 $\pm$ 0.00	****	****	****	100 $\pm$ 0.00		
25-50	***	****	***	****	****	****	****	100 $\pm$ 0.00		
LC ₁₀₀	230ppm	77.50ppm	140.14ppm	51.54ppm	199.1ppm	65.00ppm	***	46.75ppm	1ppm	0ppm
LC ₅₀	46.00ppm	15.50ppm	28.03ppm	10.31ppm	39.82ppm	13.00ppm	21.50ppm	9.35ppm	0.2ppm	0ppm

Notwithstanding, the biocidal activities of all extracts of neem plant in this studies is comparable to previous studies, wherein the biocidal activities of *A. indica* induced mortality with LC_{50} values of 0.53ppm against *Culex quinquefasciatus* for aqueous extract of the seeds (Tandon and Sirohi 2010), 6ppm for methanolic extract against *An. gambiae* (Adewole et al., 2012), 62.5ppm against *Culex pipiens* after 24 hours (Mustafa and Al Khazraji, 2008). Field trials of neem oil formulation showed 95.5% fecundity suppression of *Cx. quinquefasciatus* under 24hours and subsequently 80% reduction in about 3weeks (Dua et al., 2009).

Neem oil formulated diet induced an LC_{50} value of 11 ppm against *An. gambiae* after 8 days (Okumu et al., 2007). Tiwari (2012) reported the molluscicidal activities of neem bark formulated feed pellet against *Lymnaea acuminata* in a 24-144 hour exposure investigation with LC_{50} values of 1.35ppm in 24hours, 0.92ppm in 48hours, 0.32ppm in 72hours, 0.20ppm in 96hours, 0.12ppm in 120hours and 0.09ppm in 144hours. Generally, the variation in susceptibility of the snails (to all extracts tested, i.e. $SOLEx^B$ and $SOLEx^R$), compared to the larvae may have been largely attributed to their genetic makeup (Angaye, 2013).

Furthermore, the reason for higher activity of the bark compared to the root might be attributed to the distribution/ quantity of phytochemical in various organs of the plant. The phytochemicals identified in our studies were similar to phytochemicals reported in an earlier studies by other authors (Adewole et al., 2012; Sofowora 1993; Hamid, 1993;Kabar and Gichia, 2001;Okumu et al., 2007). Adewole et al., (2013), screened and quantified some phytochemicals in neem, which were reported, Alkaloids (3.57%), flavonoids (4.07%) and saponins (1.76%), steroids, tannins, trepenoids and glycosides. The metabolites in neem were attributed to their biocidal efficacy (Adewole et al., 2013; Okumu, 2007).

Time-dose dependency assessment are presented in Figures 1 and 2. Results for the $SOLEx^B$ against the snails induce mortality in less than 12 hours of exposure as compared to the larvae whose mortality occurred above 18 hours (Figure 1). Meanwhile, amongst $SOLEx^R$, the larvae induced mortality above 18 hours as well; the snails were lethal in 18 hours (Figure 2). These findings are comparable to Arunpandiyana (2011), whose investigation showed 30% and 70% mortalities with the leaves of neem plant against *culex* mosquitoes within 6 and 12 hour of exposure respectively.




Figure 1: Result of final screening for SOLEx^B based on exposure.

Result of the time-dose mortality assessment of the SOLEx^B (Figure 1), shows that none of the extracts induced total mortality within 06-Hours, however the highest mortality was induced by Ethanol extract of the bark with mortality rate of 78.50% (i.e. Ethanol ext. vs. snail). As the exposure increased (12-Hours), the ethanol extract further against the snail induced 90.33% mortality. Furthermore, as the exposure approached the 18-Hour mark; Ethanol ext. vs. snail, Acetone ext. vs. snail, Hexane ext. vs. snail and chloroform ext. vs. snail had all induced total mortality. At the end of (24-Hours), all extracts had induced mortality.




Figure 2: Result of final screening for SOLEx^R based on exposure.

Compared to the SOLEx^B, results of the SOLEx^R was less toxic to the snails and larvae. In 06-Hour experiment, ethanol extract of the bark induced the highest mortality against the snail with a value of 88.50%. Furthermore, at the 12-Hour, total mortality were observed in the Ethanol ext. vs. snail, Acetone ext. vs. snail, Hexane ext. vs. snail and Chloroform ext. vs. snail. As the exposure increased to the 18-Hour, mortality rates remain constant, but at 24-Hour all extracts against the snail and larvae induced total mortality.

Results of this study showed that SOLEx^B and SOLEx^R produced secondary metabolites (Table 1), which were capable of inducing larvicidal and molluscicidal properties as reported by other authors (Sofowora 1993; Hamid, 1993; Kabaru and Gichia, 2001; Okumu et al., 2007; Adewole et al., 2013). Whereas we used the solvent extracts of the bark and root in this study; the pressed cake, bark, leaf and oil have been reported for their potent molluscicidal activities against *Lymnaea acuminata* and *Indoplanorbis exustus* (Singh et al., 1996, 2013).

In other studies, the chemical constituents of Chloroform, Hexane, ethyl acetate and methanol extracts of neem leaves were reported for their anti-oxidant activities with the chloroform extract yielding more chemicals compounds (order of activities: chloroform >ethyl acetate> hexane> methanol extracts); compared to our study. It has been established in literature that the activities of neem plant are solely reliant on their bioactive chemicals (Hossain et al., 2013;Kabaru and Gichia, 2001; Su and Mulla 1999; Sofowora 1993; Okumu et al., 2007).

Furthermore, Sofowora (1993) reported that neem phytochemicals possessed larvacidal and insecticidal properties. In addition, the crude extract of neem plant as well as the oil were reported to inhibit insect metamorphosis (Kabaru and Gichia, 2001; Adewole et al., 2012), and pupation (Kabaru and Gichia, 2001; Su and Mulla 1999; Okumu et al., 2007). In another study, some chemicals also found in neem (Nimbinin, nimbandiol nimbidin, nimbin nimbidolas and Azadirachtin), have been shown to possess insecticidal, nematocidal, and molluscicidal properties (Bhowmik et al., 2012). Others include quercetin which has anti-protozoal, antioxidant properties, as well as Salanin with repellent and anti-feedant (Drabu et al., 2012).

Neem plant, in an earlier study showed anti-fecundity properties against mosquitoes (Su and Mulla, 1998; Okumu et al., 2007; Kabaru and Gichia, 2001), as well as anti-feedant properties to snails (Tiwari, 2012). It is capable of disrupting the prothoracic gland from the synthesis and release of ecdysteroids, which is a molting hormones (Mordue et al., 1993).Coincidentally, the vectors responsible for malaria and schistosomiasis begins their metamorphosis in water which is favorable medium for larviciding and mollusciciding (Ohimain et al., 2014). Besides, neem plant is less toxic to most aquatic non-targeted species; hence their application as biocidal agents.

Conclusion

The morbidity and mortality burden of vector-borne diseases in developing nation has become a global challenge. The ecosystem has suffered inimical damages induced by the application of synthetic pesticides, as such contemporary research now focuses on cost-effective and eco-friendly plant-derived pesticides. Consequent upon our findings, the bark and root of neem possess moderate toxicity against the malaria and schistosomiasis vectors. We therefore recommend further investigation of the field application especially with regards to their toxicities against non-targeted species.

References

1. Adewole A, Oderinde A, Bankole O, Faparusi F and Oyede RT. Larvacidal activities of three plant extracts of common wire weed (*Sida acuta*), Catnip (*Nepeta cataria*) and Neem (*Azadirachta indica*) against the larva of mosquito (*Anopheles gambiae*). Academia Journal of Medicinal Plants.2012; 039.
2. AOAC (1995). Official methods of analysis (16th edn) Association of Official Analytical Chemists. Arlington, V. A. USA. 1995.
3. Agboola IO, Ajayi GO, Adesegun SA, Adesanya SA. Comparative Molluscicidal activity of fruit pericarp, leaves, seed and stem Bark of *Blighia unijugata* Baker. Pharmacology Journal. 2011; 3:63-66.
4. Aina SA, Banjo AD, Lawal OA, Jonathan K. Efficacy of Some Plant Extracts on *Anopheles gambiae* Mosquito Larvae. Academic Journal of Entomology. 2009; 2(1): 31-35.
5. Angaye TCN, Zige DV, Didi B, Biobelemoye N, Gbodo EA. Comparative Molluscicidal Activities of Methanolic and Crude Extracts of *Jatropha curcas* Leaves against *Biomphalaria pfeifferi*. Greener Journal of Epidemiology and Public Health. 2014;2 (1): 016-022.
6. Angaye, TCN. In-vitro Comparative Molluscicidal Activities of Aqueous and Methanolic Extracts of *Jatropha curcas* Leaves against *Bulinus globosus* and *Bulinu srholfsi*; vectors of urinary schistosomiasis. 2013; M.Sc. thesis, Niger Delta University, Wilberforce Island, Bayelsa State, Nigeria.
7. Arunpandiyam G. Toxicity of aqueous crude neem leaf extract against *culex* mosquitoes. Int J Pharm Biomed Sci. 2011; 2(1): 1-3.
8. Bassey SE, Ohimain EI, Angaye TCN. The Molluscicidal Activities of Methanolic and Aqueous Extracts of *Jatropha curcas* leaves against *Bulinus globosus* and *Bulinus rholfi*, Vectors of Urinary Schistosomiasis. Journal of Parasitology. 2013; 115-122.

9. Bhowmik D, Chiranjib1, Yadav J, Tripathi KK, Kumar KPS. Herbal Remedies of *Azadirachta indica* and its Medicinal Application. *J. Chem. Pharm. Res.* 2010; 2(1): 62-72.
10. Dibua, Uju Marie-Esther; Odo, Greg Ejikeme; Nwabor, Ozioma Forstinus; Ngwu, Goddy Ikechukwu (2013). Larvicidal activity of *Picralimanitida*, an environmental approach in malaria vector control. *American Journal of Research Communication*, 1(12): 451-469.
11. Drabu S, Khatri S, Babu S. *Neem: Healer of All Ailments*. Maharaja Surajmal Institute of Pharmacy C-4, Janak Puri, India. 2012; 3:120-127.
12. Dua V.K, Pandey A.C, Raghavendra K, Gupta A, Sharma T, Dash A.P. (2009). Larvicidal activity of neem oil (*Azadirachta indica* formulation against mosquitos. *Malaria Journal*. 8, 124.
13. Ghosh A, Chowdhury N and Chandra G. (2012) Review on Plant extracts as potential mosquito larvicides *Indian J Med Res* 135, pp. 581-598.
14. Hamid, A.A. Molluscicidal activity of neem tree (*Azadirachta indica* A- Jusa). M.Sc. Thesis, Khartoum University; Sudan. 1993.
15. Harbone, J.B. (1998). *Phytochemical Methods- A Guide to Modern Techniques of Plant Analysis*, Chapman and Hall London.
16. Hossain MA, Al-Toubi WAS, Weli AM, Al-Riyami QA, Al-Sabahi JN. Identification and Characterization of Chemical Compounds in Different Crude Extracts from Leaves of Omani Neem. *Journal of Taibah University for Science*. 2013; 7:181–188.
17. Kabaru J.M, Gichia L. (2001). Insecticidal activity of extracts derived from different parts of the mangrove tree *Rhizophora mucronata* (rhizophoraceae) Lam. against three anthropoids: *Afr. J. Sci. Tech (AJST)*. *Sci. Eng. Series* 2(2):44-49.
18. Khan I., Srikakolupu S.R., Darsipudi S., Gotteti S.D and Amaranadh H (2012). Phytochemical studies and screening of leaf extracts of *Azadirachta indica* for its anti-microbial activity against dental pathogens. *Archives of Applied Science Research*, 2 (2):246-250.
19. Mordue (Luntz) AJ, Nisbet A.J (2000). *Azadirachtin* from the neem tree *Azadirachta indica*: its actions against insects. *Ann Entomol Soc Brasil*, 29:615-632.
20. Mordue, A.J., Blackwell, A. (1993) *Azadirachtin*: an update. *J. Insect Phys.*, 39, 903–924.
21. Mustafa M.A, Al Khazaraji A. (2008). Effect of some plant extracts on the *Culex pipiens molestus* Forskal larvae. *Iraqi J. of Vet. Sci.* 22, 9-12.
22. National Centre for Infectious Diseases (NCID), 2004. Mosquitoes. Division of parasitic diseases, Atlanta, pp: 6.
23. Ohimain E.I., Angaye T. C.N and Bassey S.E. (2014). Comparative Larvicidal activities of the Leaves, Bark, Stem and Root of *Jatropha curcas* (Euphorbiaceae) against malaria vector *Anopheles gambiae*. *Sky Journal of Biochemistry*. 3 (4): 024-027.
24. Okumu F.O, Knols B.G.J. and Fillinger U. (2007) Larvicidal effects of a neem (*Azadirachta indica*) oil formulation on the malaria vector *Anopheles gambiae*.
25. Singh A, Singh DK, Mishra TN, Agarwal RA. Molluscicidal of plant origin and their environmental Impact on non-targeted organisms. *Biol. Agric Hort.* Ph.D thesis, Gorakhpur University, Gorakhpur. 1996; 13: 205-252.
26. Singh SK, Singh SK, Singh A. Molluscicidal and piscicidal properties of three medicinal plants of family Apocynaceae - a review. 2013. *J Biol Earth Sci*. 2013; 3(2): B1 94-B205.
27. Sofowora, A., (1993). *Medicinal Plants and Traditional Medicine in Africa*. Spectrum Books Ltd., Ibadan, Nigeria, 191-289.
28. Su T, Mulla MS (1998). Antifeedancy of neem products containing *Azadirachtin* against *Culex tarsalis* and *Culex quinquefasciatus* (Diptera: Culicidae). *J Vector Ecol*, 23:114-122.
29. Tandon, P and Sirohi A (2010). Assessment of Larvicidal Properties of Aqueous Extracts of four plants against *Culex quinquefasciatus* larvae. *Jordan Journal of Biological Sciences*. Vol. 3, 1.
30. Tiwari, F. (2012). Bait Formulation Toxicity of Plant Derived Molluscicides in Attractant Food Pellets against Vector Snail, *Lymnaea acuminata*. *World Journal of Zoology* 7 (1): 55-59.
31. WHO/UNICEF. World Malaria Report. Roll Back Malaria partnership, WHO/UNICEF 2005. <http://rbm.who.int/wmr2005/>.