

Original Research Article

Antifungal Activity of Phytochemicals Against *Penicillium* Strains

ABSTRACT

Aims: The fungi have become, over the last two decades, a major cause of illness in humans. Fungi of the genus *Penicillium* can be found in various substrates and affect immunocompromised people, hospitalized patients, many animals and plants, as well as compromise the quality of air indoors. The indiscriminate use of antibiotics and the resulting framework of resistance of microorganisms to conventional antimicrobial therapy has been stimulating researchers to seek alternative sources of antimicrobial compounds, among them the medicinal plants. A tendency to getting phytochemicals from extracts, fractions, fixed or essential oils obtained from plant species is currently observed. In this context, the present study aims to evaluate the in vitro antifungal activity of seven phytochemicals (geraniol, carvacrol, thymol, linalool, p-cymene, terpinolene and citral) against twelve strains of *Penicillium*.

Place of Study: Laboratory tests were carried out at the Mycology Laboratory Department of Pharmaceutical Sciences, located in the Health Sciences Center (CCS) of the Federal University of Paraíba (UFPB).

Methodology: Firstly, screening was carried out to find the phytochemical with the best activity by determining the minimum inhibitory concentration (MIC) by the broth microdilution technique. Following, the tests were proceeded with thymol to determine of minimum fungicidal concentration (MFC).

Results: Through analysis of results, it is observed that carvacrol, thymol and citral showed the best activities of the strains of *Penicillium* studied. The MIC values were 256 mg / mL, for carvacrol and citral, and 128 mg / mL, for thymol (compound showed better results). The thymol had its MIC 90 established at 128 mg / mL, inhibiting, at this concentration, 92% of the tested strains. Analyzing compared to MIC and MFC, it was observed that thymol presented CFM values similar to CIM values for 1/3 of the strains, showing also values greater than 1024 mg / mL for only two strains. The CFM value ranged from 128µg/ml to 1024µg/ml.

Conclusion: The thymol is a promising new possibility among the products with antifungal activity against *Penicillium*, however if the performance is needed further studies, deeper, about their mechanism of action and toxicity, as well as in vivo tests, aiming a possible application therapy.

Keywords: Antifungal activity; Phytochemicals; *Penicillium*; Thymol.

1. INTRODUCTION

The incidence of fungal infections has increased over the last ten years [1]. Airborne fungi use the atmospheric air as a mean of dispersal [2]. Inhalation of airborne fungi spores by man can lead to the development of complications in the respiratory tract, such as asthma and rhinitis [3]. In addition, airborne fungi can cause sometimes fatal infections in immunocompromised individuals, which are called opportunistic infections [4,5].

Among airborne fungi, the genus *Penicillium* deserves mention, it is one of that has the largest number of species and it can be found in various substrates [6]. Fungal infections by

21 *Penicillium marneffe* ranks third in Thailand in relation to opportunistic infections in patients
22 with HIV [7].

23 The indiscriminate use of antibiotics by the general population has led to a complex
24 framework of resistance of microorganisms to conventional antimicrobial therapy, what may
25 be a relevant factor in the emergence of difficult control infectious diseases and has
26 stimulated researchers to seek alternative sources of antimicrobial compounds, among them
27 the medicinal plants [8].

28 The plants provide an important source of new biologically active compounds [9,10,11]
29 containing a number of substances that can be used to treat different infectious diseases
30 [12], for which reason they have been intensively studied to find compounds more effective
31 and less toxic.

32 The term phytochemical relates to chemical compounds, non-nutritive, which naturally occur
33 in plants and exhibit biological activity [13]. Studies involving phytochemicals are of great
34 importance, because they facilitate the utilization of individual components, instead of a
35 mixture like in essential oils, giving more predictability and probably less collateral effects.

36 Several studies point to the various activities of phytochemicals, which are: antimicrobial,
37 antioxidant, anti-inflammatory, analgesic, cardioprotective, anti-hemorrhagic,
38 hepatoprotective, antitussive, antitumor, immunostimulating, anticancer, antiviral, among
39 other [14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32. Among these, some
40 studies attribute considerable antimicrobial activity to phytochemicals commonly found in
41 plants belonging to the family Lamiaceae.

42 These findings prompted the investigation of the antifungal profile of seven phytochemicals
43 commonly found in plants of the Lamiaceae family (geraniol, carvacrol, thymol, linalool, P-
44 cymene, terpinolene and citral) against different strains of *Penicillium* by microbiological
45 screening. The aim of the present work was to investigate the antifungal activity of
46 phytochemicals, choosing among them that which showed the best antifungal profile for an
47 after in-depth study of its fungicidal and/or fungistatic effects against *Penicillium* strains.

48 **2. MATERIAL AND METHODS**

49

50 **2.1 Phytochemicals and synthetic antifungal**

51

52 The phytochemicals (geraniol, carvacrol, thymol, linalool, P-cymene, terpinolene and
53 citral) and amphotericin B (standard drug) were acquired from Sigma-Aldrich®. All them were
54 dissolved in 2% Tween 80 (INLAB®) and up to 10% dimethyl sulfoxide – DMSO (MERCK®)
55 in sterile distilled water to obtain 1,024 µg/mL solutions [33].

56

57 **2.2 *Penicillium* strains**

58

59 For testing of antifungal activity were selected and used twelve strains of *Penicillium*
60 previously isolated from the ambient air of a food dry industry and identified by the Mycology
61 Laboratory of the Department of Pharmaceutical Sciences, Health Sciences Center, Federal
62 University of Paraíba. The strains were maintained on Sabouraud Dextrose Agar - SDA
63 (DIFCO®) at room temperature (28 ° to 30 °C) and under refrigeration (4 °C).

64 **2.3 Inoculum**

Stock inoculums of *Penicillium* strains were prepared from 10-day cultures in SDA at 28 °C to induce sporulation. Fungal colonies were covered with 5 mL of sterile saline solution (NaCl 0.85 % w/v), the surface gently scraped with a sterile loop and the resultant mixture of fungal units was then transferred to a sterile tube. After heavy particles were allowed to settle for 3 to 5 min, the upper homogeneous suspension was collected and vortexed for 15 s. The turbidity of the final inoculum was standardized according to a McFarland scale 0.5 tube and adjusted to a fungal population of 10⁶ colony former units (CFU). The confirmation of inoculum quantification was done by plating 0.01 mL of inoculum suspension in SDA. The dishes were incubated at 28 °C and examined daily for the presence of fungal colonies which were counted as soon as growth became visible [34,35].

2.4 Antifungal activity screening

Initially, microbiological screening was performed with the phytochemicals (geraniol, carvacrol, thymol, linalool, P-cymene, terpinolene and citral) at a concentration of 1,024 µg/mL. Microbiological screening was performed based on the broth microdilution technique [33,36,37,38].

Sterile 96-U-shaped-well microplates were used and each well of the plates contained 100 µL of Sabouraud dextrose broth - SDB (DIFCO®). Then, 100 µL of the products (1,024 µg/mL) were individually added to each line of wells, so that each line of wells corresponded to a phytochemical tested. Finally, 10 µL of fungal inoculum of each strain of *Penicillium* were added to wells, so that each column corresponded to a strain. The microplates were incubated at 28 °C being selected those phytochemicals who showed better inhibition profile visual growth of microorganisms after seven days incubation. Negative control (without drugs) was performed to confirm the viability of the sporangiospores. Sensitivity control for Tween 80 and DMSO was also performed.

Phytochemicals that showed better activity were subsequently selected, which were evaluated for their potential antifungal activity from a concentration of 1,024 to 4 µg/mL, in order to determine their MIC [39].

2.5 Determination of MIC and MFC

Broth microdilution bioassay was used to determine MIC of thymol, carvacrol, citral and amphotericin B against *Penicillium* strains [40,41]. For this, sterile 96-U-shaped-well microplates and caps were used. The 96-well dishes were prepared by dispensing 100 µL of double concentrated Sabouraud dextrose broth- SDB (Difco®) into each well. A 100 µL volume of double concentrated products was placed into the first wells, to obtain a concentration of 1,024 µg/mL in them. Next, serial twofold dilutions in culture medium were prepared to obtain concentrations ranging from 1,024 to 4 µg/mL. The last well contained 100 µL of broth inoculated with fungal inoculums to confirm cell viability (viability control). Finally, 10 µL of fungal inoculum of each strain were added to wells, so that each column corresponded to one of the twelve strains. All the dishes were aseptically sealed then mixed on a plate shaker (300 rpm) for 30 seconds, incubated at 28 °C and read after 5 days of incubation. The MIC values were determined by visual inspection of the growth inhibition of each well compared with that of the control (without drugs) well. Sensitivity control for Tween 80 and DMSO was also performed. The MIC was determined from three independent experiments and was defined as the lowest drug concentration that showed absence of growth or complete fungal growth inhibition (100% inhibition) compared with that of the control (without drugs) well.

Under the same conditions, it was also used the product chlorine sanitizer Tecsa Clor®- TC (SERQUIMICO LTDA., São Paulo, SP, Brazil), composed of stabilized chlorine dioxide 5%. The TC was prepared in the same concentrations used to the phytochemicals (1,024-4 µg/mL) and furthermore in the concentration recommended by the manufacturer for obtaining the fungicide effect (50 ppm).

The MFC was determined for the phytochemical that showed stronger antifungal activity and for amphotericin B. After determining the MIC, 10 µL were subcultured from each well that showed complete inhibition (100% or an optically clear well) on SDA plates. The plates were incubated at 28 °C for five days, and the MFC was the lowest concentration that showed either no growth or fewer than three colonies [42,43,44,45]. The MFC was determined from two independent experiments on different occasions.

2.6 Statistical analysis

The results are expressed as mean ± S.E.M. Statistical analysis for the study of the effects of the phytochemical on mycelial growth and spore germination was performed to determine statistically significant differences (P <0.05) employing analysis of variance (one-way ANOVA), followed by the Bonferroni post-test. For this, the implementation of statistical analysis was performed using GraphPad Prism version 5.0 for Windows, San Diego, California, USA.

3. RESULTS AND DISCUSSION

At 1,024 µg/mL, the concentration of phytochemicals used in the screening, it was found that carvacrol, citral and thymol showed the best activity on the investigated strains of *Penicillium* (Table 1), having been observed, for all three phytochemicals, growth of only one strain among those tested (Pen 9).

Table 1. Antifungal activity of phytochemicals against strains of *Penicillium*-Microdilution technique.

Microorganisms	Phytochemicals (1,024 µg/mL)						
	geraniol	carvacrol	thymol	linalool	p-cymene	citral	terpinolene
<i>Penicillium</i> Pen 2	-	-	-	+	+	-	+
<i>Penicillium</i> Pen 3	-	-	-	+	+	-	+
<i>Penicillium</i> Pen 7	-	-	-	+	+	-	+
<i>Penicillium</i> Pen 8	+	-	-	+	+	-	+
<i>Penicillium</i> Pen 9	+	+	+	+	+	+	+
<i>Penicillium</i> Pen 10	-	-	-	+	+	-	+
<i>Penicillium</i> Pen 11	-	-	-	+	+	-	+
<i>Penicillium</i> Pen 12	-	-	-	+	+	-	+
<i>Penicillium</i> Pen 13	-	-	-	+	+	-	+
<i>Penicillium</i> LM 28	-	-	-	+	+	-	+
<i>Penicillium</i> LM 63	-	-	-	+	+	-	+
<i>Penicillium</i> LM 120	-	-	-	+	+	-	+

146 (+): Microbial growth in culture medium (-): Absence of microbial growth.
147

148 After initial screening, in which carvacrol, thymol and citral showed better results in inhibiting
149 the growth of strains of *Penicillium*, we proceeded to a new screening, this time with the MIC
150 determination of selected phytochemicals. To this end, carvacrol, thymol and citral were
151 tested at concentrations ranging from 1024 to 4 µg/mL, with concomitant control of viability
152 of strains, sterility of broth and interference of solvents DMSO and Tween 80 in the results.
153

154 The MIC values found were 256 µg/mL, for carvacrol and citral, and 128 µg/mL, for thymol,
155 what triggered interest in further study of the antifungal action of thymol. These results were
156 compatible with the findings of other authors.
157

158 Studies with microencapsulation of thymol and carvacrol to evaluate the potential use of
159 these agents as antimicrobials in food packaging showed the action of thymol and carvacrol
160 was evaluated against *Escherichia coli*, *Staphylococcus aureus*, *Listeria innocua*,
161 *Saccharomyces cerevisiae* and *Aspergillus niger*, yielding MICs ranging from 125 to 250
162 µg/mL for thymol and 75-375 µg/mL for carvacrol [46]. Trials with various essential oils
163 against *Botrytis cinerea*, *Fusarium sp.* and *Clavibacter michiganensis subsp. Michiganensis*
164 showed that the growth of these microorganisms was completely inhibited by oils of
165 oregano, thyme and marjoram in relatively low concentrations, 85-300 µg/mL, thymol being
166 pointed out as the main component of oregano oil and carvacrol as a major component of
167 oils of thyme and marjoram [47].
168

169 In a study using the disk diffusion method, thymol showed antimicrobial activity against 23
170 different genera of bacteria with inhibition zones ranging from 25.8 to 53.1 mm [48]. In other
171 study designed to evaluate the inhibitory activity of thymol on the formation and maturation
172 of biofilms of *Candida albicans*, thymol significantly interfered by reducing the formation and
173 maturation of the biofilm [18]. In partial agreement with this study, using the methods of broth
174 dilution, disk diffusion and microatmosfera, found in his work that thymol showed the best
175 antifungal activity against *Aspergillus niger*, among the phytochemicals tested (thymol,
176 carvacrol and eugenol), based on the study of MICs [16].
177

178 Given that thymol showed the best result in microbiological screening, it is justified the
179 choice to continue the investigation of its antifungal activity. After determination of the MIC,
180 the fungicidal effect of thymol was investigated. The MFC values of thymol varied from 128
181 to 1,024 µg/mL and the MFCs of this phytochemical corresponded to MIC or 2 × MIC for a
182 half of the *Penicillium* strains (Table 2). The Amphotericin B, standard antifungal used as
183 control, showed MIC of 128 µg/mL, concentration in which 100% of strains were inhibited.
184 Whereas the sanitizing control Tecsa clor®, showed MIC greater than 1024 µg/mL for 75% of
185 strains of *Penicillium*, that were resistant to the product at tested concentrations.
186

187 **Table 2. Minimal inhibitory concentration (MIC) of thymol, amphotericin B and**
188 **Tecsa clor® and minimal fungicidal concentration (MFC) of thymol and amphotericin**
189 **B in *Penicillium spp* strains.**
190

Strain <i>Penicillium</i>	Thymol (µg/mL)		Amphotericin B (µg/mL)		Tecsa clor® (µg/mL)	Control of strain	Sterility control
	MIC	MFC	MIC	MFC	MIC		
Pen 2	128	1024	128	1024	NF*	+	-

Pen 3	128	1024	128	NF*	NF*	+	-
Pen 7	128	128	128	128	512	+	-
Pen 8	128	128	128	128	NF*	+	-
Pen 9	256	NF*	128	128	NF*	+	-
Pen 10	128	128	128	128	1024	+	-
Pen 11	128	512	128	128	1024	+	-
Pen 12	128	1024	128	128	NF*	+	-
Pen 13	128	128	128	256	NF*	+	-
LM28	128	NF*	128	NF*	NF*	+	-
LM63	128	256	128	512	NF*	+	-
LM120	128	256	128	512	NF*	+	-

Note: *NF: not found- the value is greater than the highest concentration tested, 1024µg/mL; +, fungal growth.

Research indicate a compound with MIC ranging from 50 to 500 µg/mL as having optimal antimicrobial activity, and attribute moderate activity to compounds which have a variation of 500 to 1500 µg/mL in MIC. Thus, according to these parameters, it can be stated that thymol showed an optimal antimicrobial activity [49].

Other authors have tested the antimicrobial activity of thymol, finding satisfactory results, consistent with the findings of this study. Thymol was tested against strains of *E. coli*, *S. aureus*, *B. cereus* and *L. monocytogenes*, finding MIC values ranging between 300 and 500 µg/mL [50]. Previous studies reported MIC values of 62-250 µg/mL for thymol on strains of *Streptococcus mutans*, *Staphylococcus aureus*, *Bacillus subtilis*, *Staphylococcus epidermidis* and *Escherichia coli* [51]. Another study found thymol MICs values of 15-250 µg/mL on strains of *Staphylococcus aureus*, *Bacillus cereus*, *Listeria monocytogenes*, *Proteus mirabilis*, *Escherichia coli*, *Klebsiella pneumoniae* and *Pseudomonas aeruginosa* [52].

In another study that evaluated the activity of thymol nanospheres against *Staphylococcus aureus*, *Escherichia coli* and *Pseudomonas aeruginosa* strains and observed that free thymol showed a MIC ranging from 10 to 156 µg/mL and MBC ranging 156 - 313 µg/mL, whereas the encapsulated thymol was less effective with MIC values of 10-313 µg/mL and MBC values of 156-1250 µg/mL [53]. Researchers Studies evaluated the activity of thymol on seven species of fish pathogenic bacteria (*Aeromonas salmonicida subsp. masoucida*, *A. salmonicida subsp. salmonicida*, *A. hydrophila*, *Edwardsiella tarda*, *Vibrio vulnificus*, *V. parahaemolyticus* and *V. anguillarum*) and found that thymol showed MIC and MBC values ranging from 10 to 320 µg/mL [54]. While other investigation used the essential oil (EO) of *Thymus vulgaris* and its constituents thymol and p-cymene, in order to determine their antifungal activity against *Rhizopus oryzae*, this study revealed that the MIC of thymol and EO ranged 128-512 µg/mL, since the MFC ranged from 512-1024 µg/mL for the EO and 128-1024 µg/mL for thymol [55]. Studied the activity of thymol on various bacteria (*Escherichia coli*, *Pseudomonas aeruginosa*, *Salmonella typhimurium*, *Proteus mirabilis*, *Listeria monocytogenes*, *Bacillus cereus*, *Micrococcus flavus* and *Staphylococcus aureus*) and fungi (*Aspergillus flavus*, *Aspergillus fumigates*, *Aspergillus niger*, *Aspergillus ochraceus*, *Penicillium funiculosum*, *Penicillium ochrochloron*, *Trichoderma viride* and *Candida albicans*), getting through microdilution broth and MIC, MBC/MFC values and found

227 that for bacteria MIC values ranged from 10-100 µg/mL and MBC ranged from 50 to 150
228 µg/mL, as for fungi MIC and MFC values ranged from 10 to 50 µg/mL [56].

229
230 Macromolecules whose functionality is related to growth, survival, virulence or cellular
231 morphogenesis are pointed out as promising targets for new antifungal agents [57]. Thus,
232 thymol is considerate as having a promising antifungal activity.

233 234 **4. CONCLUSION**

235
236 Among the tested phytochemicals, thymol showed the best results. The thymol is a
237 promising new possibility among the products with antifungal activity against *Penicillium*,
238 however if the performance is needed further studies, deeper, about their mechanism of
239 action and toxicity, as well as in vivo tests, aiming a possible application therapy.

240 241 242 **CONSENT (WHERE EVER APPLICABLE)**

243
244 All the authors declare that no consent was obtained for this study.

245 246 **REFERENCES**

- 247
248 1. Tortora GJ, Funke BR, Case CL. Microbiology, 10^a Ed. Porto Alegre: Artmed; 2012.
- 249 2. Trabulsi, L. B.; Alterthum, F. Microbiology. 5^a ed. São Paulo: Atheneu; 2008.
- 250 3. Mezzari A, Perin C, Júnior SAS, Bern LAG, Gesu GD. Airborne fungi and
251 sensitization in atopic individuals in Porto Alegre, RS, Brazil. Rev. Assoc. Med. Bras.
252 São Paulo. 2003;49(3).
- 253 4. Sidrim JJC, Moreira JLB. Clinical and laboratory Fundamentals of medical
254 mycology. Rio de Janeiro: Guanabara Koogan; 1999.
- 255 5. Wanke B, Lazéra MS, Nucci M. Fungal infections in the immunocompromised host.
256 Mem. Inst. Oswaldo Cruz, Rio de Janeiro. 2000;95(Supl. I):153-158.
- 257 6. Pitt JI, Hocking AD. Fungi and Food Spoilage. 2.ed. Cambridge: Chapman & Hall;
258 1997.
- 259 7. Ranjana KH, Priyokumar K, Singh ThJ, Gupta ChC, Sharmila L, Singh PN,
260 Charkrabarti A. Disseminated *Penicillium marneffei* Infection among HIV-infected
261 patients in Manipur State, India. J. Infect. 2002;45(4):268-271.
- 262 8. Akinpelu DA. Antimicrobial activity of *Anacardium occidentale* bark. Fitoterapia
263 2001;72: 286-287.
- 264 9. Nakamura CV, Ishida K, Faccin LC, Filho BPD, Cortez DAG, Rozental S. In vitro
265 Microbiol, 2004;155 (7): 579–586.
- 266 10. Oliveira DR, Leitão GG, Santos SS, Bisso HR, Lopes D, Alviano CS, Alviano DS,
267 Leitão SG. Ethnopharmacological study of two *Lippia* species from Oriximina, Brazil.
268 J Ethnopharmacol, 2006;108(1):103-108.

- 269 11. Prabuseenivasan S, Jayakumar M, Ignacimuthu S. In vitro antibacterial activity of
270 some plant essential oils. BMC Complement Altern Med, 2006; 6(39):1-8.
- 271 12. Saad A, Fadli M, Bouaziz M, Benharref A, Mezrioui NE, Hassani L. Anticandidal
272 activity of the essential oils of *Thymus maroccanus* and *Thymus broussonetii* and
273 their synergism with amphotericin B and fluconazol. Phytomedicine,
274 2010;17(13):1057–1060.
- 275 13. Anjorin T, S, Salako EA, Makun HA. Control of Toxigenic Fungi and Mycotoxins with
276 Phytochemicals: Potentials and Challenges Mycotoxin and Food Safety in
277 Developing Countries. InTech. 2013. Accessed 5 July 2013. Available:
278 [http://www.intechopen.com/books/mycotoxin-and-food-safety-in-](http://www.intechopen.com/books/mycotoxin-and-food-safety-in-developingcountries/control-of-toxigenic-fungi-and-mycotoxins-with-phytochemicals-potentialsand-challenges)
279 [developingcountries/control-of-toxigenic-fungi-and-mycotoxins-with-phytochemicals-](http://www.intechopen.com/books/mycotoxin-and-food-safety-in-developingcountries/control-of-toxigenic-fungi-and-mycotoxins-with-phytochemicals-potentialsand-challenges)
280 [potentialsand- challenges" title="Control of Toxigenic Fungi and Mycotoxins with](http://www.intechopen.com/books/mycotoxin-and-food-safety-in-developingcountries/control-of-toxigenic-fungi-and-mycotoxins-with-phytochemicals-potentialsand-challenges)
281 [Phytochemicals: Potentials and Challenges">Control of Toxigenic Fungi and](http://www.intechopen.com/books/mycotoxin-and-food-safety-in-developingcountries/control-of-toxigenic-fungi-and-mycotoxins-with-phytochemicals-potentialsand-challenges)
282 [Mycotoxins with Phytochemicals: Potentials and Challenges.](http://www.intechopen.com/books/mycotoxin-and-food-safety-in-developingcountries/control-of-toxigenic-fungi-and-mycotoxins-with-phytochemicals-potentialsand-challenges)
- 283 14. Ayoola GA, Coker HAB, Adesegun SA. Phytochemical Screening and Antioxidant
284 Activities of Some Selected Medicinal Plants Used for Malaria Therapy in
285 Southwestern Nigeria. Trop J Pharm Res, 2008;7(3):1019-1024.
- 286 15. Araújo-Júnior JX, Oliveira MSG, Aquino PGV. A Phytochemical and
287 Ethnopharmacological Review of the Genus *Erythrina*. In: RAO, V. (Ed.).
288 Phytochemicals - A Global Perspective of Their Role in Nutrition and Health. New
289 York: InTech. 2012;538. Accessed 20 June 2013. Available:
290 [http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-theirrole-](http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-theirrole-in-nutrition-and-health)
291 [in-nutrition-and-health" title="Phytochemicals - A Global Perspective of Their Role in](http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-theirrole-in-nutrition-and-health)
292 [Nutrition and Health">Phytochemicals - A Global Perspective of Their Role in](http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-theirrole-in-nutrition-and-health)
293 [Nutrition and Health.](http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-theirrole-in-nutrition-and-health)
- 294 16. Bouddine L, Louaste B, Achahbar S. Comparative study of the antifungal activity of
295 some essential oils and their major phenolic components against *Aspergillus niger*
296 using three different methods. Afr J Biotechnol, 2012;11(76):14083-14087.
- 297 17. Braga PC, Culici M, Alfieri M, Dal Sasso M. Inhibitory activity of thymol against the
298 formation and viability of *Candida albicans* hyphae. Mycoses, 2007;50(6):502–506.
- 299 18. Braga PC, Culici M, Alfieri M, Dal Sasso M. Thymol inhibits *Candida albicans* biofilm
300 formation and mature biofilm. Int J Antimicro Ag, 2008;31:472–477.
- 301 19. Canuto KM, Silveira ER, Bezerra AM. Phytochemistry, Pharmacology and
302 Agronomy of Medicinal Plants: *Amburana cearensis*, an Interdisciplinary Study. In:
303 RAO, V. (Ed.). Phytochemicals - A Global Perspective of Their Role in Nutrition and
304 Health. New York: InTech. 2012;538. Accessed 20 June 2013. Available:
305 [http://www.intechopen.com/books/phytochemicals-a-global-perspective-oftheir-role-](http://www.intechopen.com/books/phytochemicals-a-global-perspective-oftheir-role-in-nutrition-and-health)
306 [in-nutrition-and-health" title="Phytochemicals - A Global Perspective of Their Role in](http://www.intechopen.com/books/phytochemicals-a-global-perspective-oftheir-role-in-nutrition-and-health)
307 [Nutrition and Health">Phytochemicals - A Global Perspective of Their Role in](http://www.intechopen.com/books/phytochemicals-a-global-perspective-oftheir-role-in-nutrition-and-health)
308 [Nutrition and Health.](http://www.intechopen.com/books/phytochemicals-a-global-perspective-oftheir-role-in-nutrition-and-health)
- 309 20. Combrincka S, Regniera T, Kamatoub GPP. In vitro activity of eighteen essential oils
310 and some major components against common postharvest fungal pathogens of fruit.
311 Ind Crop Prod, 2011;33(2):344–349.

- 312 21. Irkin R, Korukluoglu M. Effectiveness of *Cymbopogon citratus* L. Essential Oil to
313 Inhibit the Growth of Some Filamentous Fungi and Yeasts. *J Med Food*,
314 2009;12(1):193–197.
- 315 22. Krishnaiah D, Nithyanandam R, Sarbatly R. Phytochemical Constituents and
316 Activities of *Morinda citrifolia* L. In: RAO, V. (Ed.). *Phytochemicals- A Global*
317 *Perspective of Their Role in Nutrition and Health*. New York: InTech 2012:538p.
318 Accessed 20 June 2013. Available:
319 [http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-theirrole-](http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-theirrole-in-nutrition-and-health)
320 [in-nutrition-and-health" title="Phytochemicals - A Global Perspective of Their Role in](http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-theirrole-in-nutrition-and-health)
321 [Nutrition and Health">Phytochemicals - A Global Perspective of Their Role in](http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-theirrole-in-nutrition-and-health)
322 [Nutrition and Health.](http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-theirrole-in-nutrition-and-health)
- 323 23. Kumar JK, Prasad AGD, Richard SA. In vitro Antioxidant activity and preliminary
324 phytochemical analysis of medicinal Legumes. *J Pharmaceut Res*, 2012;5(6):3059-
325 3062.
- 326 24. Maisuthisakul, P. Phenolic Constituents and Antioxidant Properties of some Thai
327 Plants. In: RAO, V. (Ed.). *Phytochemicals - A Global Perspective of Their Role in*
328 *Nutrition and Health*. New York: InTech. 2012:538p. Accessed 20 June 2013.
329 Available: [http://www.intechopen.com/books/phytochemicals-a-global-perspective-](http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-theirrole-in-nutrition-and-health)
330 [of-theirrole-in-nutrition-and-health" title="Phytochemicals - A Global Perspective of](http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-theirrole-in-nutrition-and-health)
331 [Their Role in Nutrition and Health">Phytochemicals - A Global Perspective of Their](http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-theirrole-in-nutrition-and-health)
332 [Role in Nutrition and Health.](http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-theirrole-in-nutrition-and-health)
- 333 25. Marei GIK, Rasoul MAA, Abdelgaleil SAM. Comparative antifungal activities and
334 biochemical effects of monoterpenes on plant pathogenic fungi. *Pest Biochem*
335 *Physiol*, 2012;103:56–61.
- 336 26. Naithani R, Huma LC, Holland LE. Antiviral activity of phytochemicals: a
337 comprehensive review. *Mini Rev Med Chem*, 2008;8(11):1106-33.
- 338 27. Efouet N, Pépin A. Phytochemicals from *Beilschmiedia anacardioides* and Their
339 Biological Significance. In: RAO, V. (Ed.). *Phytochemicals - A Global Perspective of*
340 *Their Role in Nutrition and Health*. New York: InTech. 2012:538p. Accessed 20 June
341 2013. Available: [http://www.intechopen.com/books/phytochemicals-a-global-](http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-theirrole-in-nutrition-and-health)
342 [perspectiveof-theirrole-in-nutrition-and-health" title="Phytochemicals - A Global](http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-theirrole-in-nutrition-and-health)
343 [Perspective of Their Role in Nutrition and Health">Phytochemicals - A Global](http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-theirrole-in-nutrition-and-health)
344 [Perspective of Their Role in Nutrition and Health.](http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-theirrole-in-nutrition-and-health)
- 345 28. Riju A, Sithara K, Nair SS. In Silico Screening Major Spice Phytochemicals for their
346 Novel Biological Activity and Pharmacological Fitness. *J Bioequiv Availab*,
347 2009;1:063-073.
- 348 29. Thapa D, Losa R, Zweifel B, Wallace RJ. Sensitivity of pathogenic and commensal
349 bacteria from the human colon to essential oils. *Microbiology*, 2012;158:2870–2877.
- 350 30. Wang L, Chen J, Xie H. Phytochemical Profiles and Antioxidant Activity of Adlay
351 Varieties. *J Agric Food Chem*, 2013;61(21):5103–5113.
- 352 31. Yang JL, Liu LL, Shi YP. Phytochemicals and biological activities of *Gentiana*
353 species. *Nat Prod Commun*, 2010;5(4):649-64.

- 354 32. Yousuf M, Aslam K, Wani BA. In vitro antibacterial activity and phytochemical
355 studies of methanolic extract of leaves of *Hypericum perforatum* L. growing wild in
356 Kashmir Himalaya. *Asian J Plant Sci Res*, 2012;2(4):414-420.
- 357 33. Cleeland R, Squires E. Evaluation of new antimicrobials in vitro and experimental
358 animal infection In: Lorian V. *Antibiotics in laboratory medicine*. 3th ed. Baltimore:
359 Williams and Wilkiam; 1991.
- 360 34. Hadacek F, Greger H. Testing of antifungal natural products: methodologies,
361 comparability of results and assay choice. *Phytochem Analysis*, 2000;11(3):137-147.
- 362 35. Santos DA, Hamdan JS. Evaluation of broth microdilution antifungal susceptibility
363 testing conditions for *Trichophyton rubrum*. *J Clin Microbiol*, 2005;43(4):1917-1920.
- 364 36. Koneman EW, Allen SD, Dowel JR V. R, Sammers HM. *Microbiological Diagnosis*
365 *Text and Atlas*. 2th ed. Medical Editorial Panamericana. São Paulo: Panamericana;
366 1993.
- 367 37. Eloff JN. A sensitive and quick microplate method to determine the minimal
368 inhibitory concentration of plant extracts for bacteria. *Planta Méd*, 1998;64(8):711-
369 713.
- 370 38. Hadacek F, Greger H. Testing of antifungal natural products: methodologies,
371 comparability of results and assay choice. *Phytochem. Analysis*, 2000;11(3):137-
372 147.
- 373 39. Allegrini J, Bouchberg MS, Maillols H. Emulsions d'huiles essentielles, fabrication et
374 applications in microbiologia. *Society Pharmacy Montpellier*, 1973;33(1):73-86.
- 375 40. Sahin F, Gulluce M, Daferera D, Sokmen A, Polissiou M, Sokmen M, Agar G, Ozer
376 H. Biological activities of the essential oils and methanol extract of *Origanum vulgare*
377 ssp. *Vulgare* in the Eastern Anatolia region of Turkey. *Food Control*, 2004;15(7):549-
378 557.
- 379 41. Moreira ACP, Lima EO, Souza EL, Van Dingenen MA, Trajano VN. Inhibitory effect
380 of *Cinnamomum zeylanicum* Blume (Lauraceae) essential Oil and beta-pinene on
381 the growth of dematiaceous moulds. *Braz J Microbiol*, 2007;38(1):33-38.
- 382 42. Espinel-Ingroff A, Fothergill A, Peter J, Rinaldi MG, Walsh TJ. Testing conditions for
383 determination of, minimum fungicidal concentrations of new and established
384 antifungal agents for *Aspergillus* spp: NCCLS collaborative study. *J Clin Microbiol*,
385 2002;40(9):3204-3208.
- 386 43. Pereira FO, Wanderley PA, Viana FAC, Lima RB, Sousa FB, Lima EO. Growth
387 inhibition and morphological alterations of *Trichophyton rubrum* induced by essential
388 oil from *Cymbopogon winterianus* Jowitt ex bor. *Braz J Microbiol*. 2011a;42(1):233-
389 242.
- 390 44. Pereira, F. O.; Wanderley, P. A.; Viana, F. A. C.; Lima, R. B.; Sousa, F. B.; Santos,
391 S. G.; Lima, E. O. Effects of *Cymbopogon winterianis* Jowitt ex Bor essential oil on
392 the growth and morphogenesis of *Trichophyton mentagrophytes*. *Braz J Pharm Sci*,
393 2011b;47(1).

- 394 45. Trajano VN, Lima EO, Souza EL, Travassos AER. Inhibitory effect of the essential
395 oil from *Eugenia caryophyllata* Thumb leaves on coalho chesse contaminating
396 microorganisms. *Ciênc Tecnol Aliment*, 2010;30(4):1001-1006.
- 397 46. Guarda A, Rubilar JF, Miltz J. The antimicrobial activity of microencapsulated thymol
398 and carvacrol. *Int J Food Microbiol*, 2011;146:144–150.
- 399 47. Daferera DJ, Ziogas BN, Polissiou MG. The effectiveness of plant essential oils on
400 the growth of *Botrytis cinerea*, *Fusarium* sp. and *Clavibacter michiganensis* subsp.
401 *Michiganensis*. *Crop Protection*, 2003;22(1):39–44.
- 402 48. Dorman HJD, Deans SG. Antimicrobial agents from plants. *J Appl Microbiol*,
403 2000;88:308–316.
- 404 49. Sartoratto A, Machado ALM, Delarmelina C, Figueira GM, Duarte MCT, Rehder
405 VLG. Composition and antimicrobial activity of essential oils from aromatic plants
406 used in Brazil. *Braz J Microbiol*, 2004;35:275-280.
- 407 50. Tippayatum P, Chonhenchob V. Antibacterial Activities of Thymol, Eugenol and
408 Nisin Against Some Food Spoilage Bacteria. *Nat Sci*, 2007;41:319-323.
- 409 51. Mathela CS, Singh KK, Gupta VK. Synthesis and in vitro antibacterial activity of
410 thymol and carvacrol derivatives. *Acta Pol Pharm*, 2010;67(4):375-380.
- 411 52. Mohammed MJ, Al-Bayati FA. Isolation and identification of antibacterial compounds
412 from *Thymus kotschyanus* aerial parts and *Dianthus caryophyllus* flower buds.
413 *Phytomedicine*, 2009;16:632–637.
- 414 53. Wattanasatcha A, Rengpipat S, Wanichwecharungruang S. Thymol nanospheres as
415 an effective anti-bacterial agent *Int J Pharm*, 2012;434(1–2):360–365.
- 416 54. Heo GJ, Kim CH, Park SC. et al. Antimicrobial Activity of Thymol against Pathogenic
417 Gram-negative Bacteria of Fishes. *Philipp. J. Vet. Med.* 2012;49(2):103-106.
- 418 55. Mota KSL, Pereira FO, Oliveira WA. et al. Antifungal Activity of *Thymus vulgaris* L.
419 Essential Oil and Its Constituent Phytochemicals against *Rhizopus oryzae*:
420 Interaction with Ergosterol. *Molecules*, 2012;17:14418-14433.
- 421 56. Giweli A, Džamic AM, Sokovic M. et al. Antimicrobial and Antioxidant Activities of
422 Essential Oils of *Satureja thymbra* Growing Wild in Libya. *Molecules*, 2012;17:4836-
423 4850.
- 424 57. Odds FC, Brown AJP, Gow NAR. Antifungal agents: mechanisms of action. *Trends*
425 in Microbiology, 2003;11:272–279.