

Original Research Article

Cryptosporidial diarrhoea in children at a paediatric hospital in Accra, Ghana.

ABSTRACT

Background: Diarrhoeal diseases are common among children in developing countries, and is caused by several aetiological agents including *Cryptosporidium* sp. Several species of this parasite exist which may belong to either anthroponotic or zoonotic forms. With recent application of molecular tools, species involved in human transmission in any locality and sources of infection can now be determined.

Aim: We screened children with acute diarrhoea at a paediatric hospital in Accra, Ghana for enteric parasites to determine frequency of cryptosporidial diarrhoea. *Cryptosporidium* isolates were then characterized by molecular methods to determine the genetic species in transmission.

Methodology: A total of 365 diarrhoeic children of age ≤ 5 years were used in this cross-sectional study. Stool samples were collected and tested for enteric parasites by microscopy and ELISA. *Cryptosporidium* isolates were subsequently genotyped by PCR-RFLP and confirmed by sequencing of the *18S rRNA* gene. Demographic and clinical data were obtained by a structured questionnaire and data analysed for possible association with cryptosporidial diarrhoea.

Results: Enteric parasites detected were *Cryptosporidium* sp. (22.2%), *G. lamblia* (5.8%) and *E. histolytica* (0.8%). Neither gender nor breastfeeding habits, presence of domestic animals, source of children's food, seasons (dry or rainy) appeared to be associated with infection of *Cryptosporidium* sp. However, age of children, source of drinking water, and education level of mother seems to have association with infection of the parasite.

Genotyping results show that *C. parvum* is the main species involved in transmission.

Conclusion: *Cryptosporidium parvum* is the commonest enteric parasite causing diarrhoea among children with acute diarrhoea. Children ≤ 3 years and those who drank sachet water were most affected. A carefully planned health education among illiterate mothers and improved sanitary conditions could reduce rate of infections. Further sub-genotyping of *C. parvum* is needed to determine whether source of infection is zoonotic or anthroponotic.

Keywords: *Cryptosporidium parvum*, cryptosporidial diarrhoea, *18SrRNA* gene, anthroponotic, zoonotic.

1. INTRODUCTION

Diarrhoeal diseases are common among people in developing countries, and they cause considerable amount of morbidity and mortality, especially among children (1, 2). According to a United Nations Children's Fund (UNICEF)/ World Health Organization (WHO) joint report (3), diarrhoea remains the most common cause of death among children under five years globally. The report indicated that each year, about 1.5 million children in this age group die as a result of diarrhoea, and more than 80 per cent of these deaths occur in Africa and South Asia. Cryptosporidiosis caused by *Cryptosporidium* sp. is among the most common gastroenteritis in humans worldwide. Symptomatic infection is characterized by diarrhoea, epigastric pain, nausea, vomiting, and weight loss, though many infections are asymptomatic. The diarrhoea can develop into a persistent life threatening type especially in immune-deficient individuals and malnourished children. *Cryptosporidium* is widespread in the developing world, with 10–30% of individuals being asymptomatic oocyst excretors (4).

The frequency of cryptosporidiosis worldwide is often dependent on HIV status. Besides humans, cryptosporidiosis is also a common cause of diarrhoeal disease in a number of other mammals (5, 6, 7), which suggests the zoonotic potential of the parasite. The oocyst, which is apparently not host specific, has been found in cattle, sheep, rats, mice, cats, dogs, rabbits and guinea pigs (8, 9).

Cryptosporidium is a protozoan parasite which belongs to the Phylum Apicomplexa, (Class Sporozoasida, Subclass Coccodiasina, Order Eucoccidiorida, Suborder Eimeriorina, Family Cryptosporidiidae). The genus *Cryptosporidium* now comprises 14 species, namely *C. hominis* in humans and monkeys, *C. parvum* in cattle, other mammals, and humans, *C. andersoni* in cattle, *C. muris* in rodents, *C. suis* in pigs, *C. felis* in cats, *C. canis* in dogs, *C. wrairi* in guinea pigs, *C. bailey* in poultry, *C. meleagridis* in turkeys and humans, *C. galli* in finches and chicken, *C. serpentis* in reptiles, *C. saurophilum* in lizard, and *C. molnari* in fish (10). Humans are most frequently infected with *C. hominis* and *C. parvum* (11, 12, 13, 14). Molecular characterization of the 60-kDa glycoprotein (GP60) gene of *C. hominis* and *C. parvum* has enabled further division into subtype families and subtypes (14). Molecular analyses indicate that *C. parvum* which is the major cause of cryptosporidiosis in humans comprises of at least two different genotypes (11). These are genotypes 1 (or human type), and genotype 2 (or calf type). Whilst genotype 1 is restricted to humans, genotype 2 is found in livestock as well as humans. Anthropophilic and zoonotic species such as *C. meleagridis* in turkeys, *C. muris* in mice, and *C. felis* in cats have also all been implicated in human illness (15). Identification of *C. parvum* genotypes has also been achieved through the amplification of the *Cryptosporidium* oocyst wall protein (COWP) (5). Other *Cryptosporidium* gene targets for amplification include the 18SrDNA, Hsp70, Actin, β -Tubulin, GP60, Microsatellites, Minisatellites, and Extrachromosomal double-stranded RNA (16).

Currently in many African countries, through application of genotyping and sub-genotyping techniques, the knowledge and understanding of the transmission dynamics of this parasite has improved. For instance, in Tunisia, *C. hominis*, *C. parvum*, and *C. meleagridis* were all identified, with *C. hominis* and *C. parvum* being higher in diarrhoeal stools than in formed ones (17). Also, in that study, *C. hominis* was more prevalent in children from urban areas than those from rural areas. Among children at the Mulago hospital, Kampala, Uganda (18), it was revealed through application of the PCR-RFLP tool that the main species involved in transmission in the area was *C. parvum*. Similarly, the tool has been used to determine the extent of genetic diversity and transmission pathways of the parasite in the Kaduna state of Nigeria (19). According to that study *C. hominis* (Ia), *C. parvum* (Ile) were the commonest in the population. Importantly, the authors recognized that mode of transmission in this population was likely to be anthroponotic, as revealed by their genotyping results. In an earlier study at Osun State, Nigeria, similar techniques were employed to determine the species, genotypes, and subgenotypes of *Cryptosporidium* sp. in the area (20). The authors reported a high diversity of the parasite in the area, which include three species, namely *C. hominis*, *C. parvum*, and *C. meleagridis*, as well as *Cryptosporidium* rabbit genotype, the cervine genotype and *C. canis*. Observations from this study showed that source of infection could be anthroponotic, zoonotic and/ or environmental.

A recent study of paediatric diarrhoea in southern Ghana (21) revealed that *Cryptosporidium* was one of the aetiological agents. Although a prevalence rate of 8.7% was reported by the authors, like other previous studies in the country (22) the particular species involved in transmission was not determined. In many Ghanaian communities companion animals including cats and dogs, usually kept as pets as well as farm animals reared in close proximity to human habitations is a common practice. Our observations of frequent contamination of homes by faecal material of these animals and poor sanitary conditions

prompted us of need to investigate possible association of environmental risk factors with cryptosporidiosis. In our study, we determined rate of cryptosporidial diarrhoea in children hospitalized for acute diarrhoea and the species involved in transmission by polymerase chain reaction (PCR) and nucleotide sequence analysis of the (18S) subunit of rRNA gene. Our results show the clinical significance of the parasite in paediatric diarrhoea in Accra, Ghana and provide preliminary information on the extent of genetic diversity. We also report on aspects of socioeconomic, behavioural and environmental risk factors needed to be considered in the planning and future implementation of intervention methods against transmission of this parasite in Ghanaian children.

2. MATERIAL AND METHODS

Study site and study participants

The study which was hospital-based, prospective and cross-sectional was conducted between the periods of March, 2010 and June, 2011 at the Princess Marie Louise Children's hospital (PML) at Accra, Ghana. The hospital is the main paediatric health facility located within Accra and accessed by people from all parts of the city. These include people from different socio-cultural, economic, political or educational background. The hospital has both out-patients and in-patients departments which are fully and well patronized by patients. According to the hospital's recent report, the daily average attendance of patients to the hospital is 143. This comprises of an average of 85 new attendants, and 58 old attendants. It is estimated that about 90 percent of these patients are children below 5 years, of which about 11 percent report with diarrhoea and other related problems (PML records).

The study participants were children (of age ≤ 5 years) who had been hospitalized at PML primarily due to acute diarrhoea and dehydration. Diarrhoea was defined as passage of loose or watery stools in the previous 24 hours and still present when the faecal specimen was collected at the hospital. Patients who developed diarrhoea after admission at the hospital, as well as patients with diarrhoea resulting from food intolerance were all excluded from our study. In all, 485 patients were referred to the study by paediatricians and nurses at the ward, 365 with diarrhoea and 120 without diarrhoea (control).

Socio-demographic and clinical data

Socio-demographic and clinical data were obtained by a study nurse at the hospital ward by use of a structured questionnaire. These included sex, age, parents educational background, source of drinking water, breastfeeding habits and source of food for children, and presence of domestic animals at home. Clinical data included vomiting, malaise, fever, abdominal pain, frequency of passage and consistency of stools, drugs used for treatment or management, as well as duration, if admitted, in the hospital.

Sample collection

A single stool sample was collected from each patient by parents or guardians and placed into a clean disposable plastic tube with tight fittings. The consistency of the stool was directly observed, classified and recorded by the study nurse as loose, semi-formed, formed, mucoid, slimy, or watery. Each sample was then divided into two portions (one portion was preserved in 10% formalin whilst the other was unpreserved with any reagent) and stored frozen at a temperature of -20°C .

Stool Microscopy and Enzyme immunoassay test

Each stool sample preserved in 10% formalin was processed by direct smear after formal-ether concentration procedures as previously described (23). Slides were examined under microscopy for enteric parasites including *Giardia* and *Entamoeba* cysts. The modified Ziehl-

Neelsen method (23) was used to detect *Cryptosporidium* sp oocysts. The enzyme immunoassay test was performed on the -20°C stored faecal samples using the Wampole™ *Giardia/Cryptosporidium/E. histolytica* Check® ELISA kit (TechLab Inc. Blacksburg VA) following the manufacturer's instructions. Results were read both visually by assessing the colour formed in each well of a microtitre plate, and quantified by measuring the absorbance at 450nm on a microplate ELISA reader (Labsystems Multiskan MS, Finland, serial RS-232C). The recommended values (cut-off points) were <0.150 OD₄₅₀, for the negative control, and ≥0.500 for the positive control. In the test wells, any value greater than, or equal to 0.150 (≥ 0.150) was considered positive, and recorded.

DNA extraction, RFLP- PCR and Sequencing of samples

Cryptosporidium DNA was directly extracted from the stool samples by using the MO BIO UltraClean® Fecal DNA Isolation kit. The procedure was performed by following manufacturer's instructions and the purified DNA was stored frozen at -40°C until further use. *Cryptosporidium* 18S rRNA gene was amplified by a semi-nested PCR following previously described methods (24). Amplifications were carried out in an Applied Biosystem thermocycler (USA), model 2720. Restriction digests were carried out directly on the second-round PCR products in a final volume of 20µL containing 10µL of PCR product, 10U Vsp1 or SSp1 (Thermoscientific, EU, Lithuania), 1X reaction buffer and 0.2µg/µl BSA. Reaction was incubated at 37°C for 30 minutes. All reactions were controlled with negative and positive samples. PCR (Fig. 1) and RFLP products were run, alongside 100bp ladder, on 2% agarose gel stained with ethidium bromide. Purification and sequencing of PCR products were performed by Macrogen Inc, Amsterdam, Netherlands. BLAST searches and multi sequence alignments were performed in MEGA V5 (25).

Statistical analysis

All data were entered into Microsoft Excel 2010 and analyzed using SPSS V17. The results were expressed as means for the continuous variables and as numbers and percentages for the categorical data. Univariate analysis was performed using independent-sample t-test, Pearson X² test or Fisher's exact tests, as appropriate. A P-value of < 0.05 was considered statistically significant.

3. RESULTS

Prevalence of *Cryptosporidium parvum* and other intestinal parasites

The parasitological results of stool from both diarrhoeal and non-diarrhoeal cases are represented in Tables 1 and 2. Among children with diarrhoea, 81/365 (22.2%) were positive *Cryptosporidium* sp. cases, 21/365 (5.8%) *G. lamblia*, and 3/365 (0.6%) were *E. histolytica* cases. None of the non diarrhoeal cases had *E. histolytica* infection but had both *Cryptosporidium* sp and *G. lamblia* present in significantly lower rates compared with the diarrhoeal samples (P < 0.0001). We also identified 3/365 (0.8%) diarrhoeal cases as dual infections (*Cryptosporidium* sp. and *G. lamblia*).

The risk factors suspected to be associated with *Cryptosporidium* sp infection were investigated among children with diarrhoea only. These factors included sex and age of child, educational background of mother, source of drinking water for the child, breastfeeding habits of mother, presence of animals at home, source of food for the child, and seasonality (Table 3).

Sex and age of child

We observed that 21.8% (N=211) males had *Cryptosporidium* sp. infection whilst 22.2% (N=154) females were infected with the parasite. The difference in infection rates was statistically insignificant (P >0.05) (Table 3), suggesting that the infection is not associated

183 with sex (OR=0.948). Although no infection was recorded among children of age 3 years and
 184 above, our results show that there was relatively high risk of *Cryptosporidium* infection
 185 associated with ages less than 3 years. Children of age 25-36 months had the highest risk
 186 (OR=3.432), followed by 6-12 months old (3.111), and then less than 6 months (OR= 2.856).
 187

188 Maternal education, source of drinking water, and breastfeeding habits

189 The highest *Cryptosporidium* infection (33.3%) occurred among children whose mothers had
 190 no formal education. Their odds ratio (OR= 3.629) also suggests that they could be
 191 associated with risk of infection more than any other group (Table 4). Although children
 192 belonging to mothers of all levels of education were infected at significantly different levels
 193 (P= 0.011), our study shows that children whose mothers have tertiary education could be
 194 protected (OR=0.926).
 195

196 The children depended on two main sources of water, namely Sachet (bagged) water and
 197 Pipe borne water. We recorded 23.1% infections for children who drank sachet water
 198 (N=337) and 11.5% for those who drank pipe-borne water (N= 26). Although there was no
 199 significant difference (P =0.293) between the two sources, pipe-borne water appears to be
 200 protective (OR= 0.725), whilst drinking sachet water could be associated with some risk
 201 (OR= 1.935).
 202

203 In Ghana, mothers are encouraged to practice 6-months exclusive breastfeeding, and we
 204 observed from our study that 21.6% of children who were exclusively breastfed (N= 218) had
 205 infections. Infection among children who were not exclusively breastfed was 23.7% (N=
 206 131). We did not find any significant difference (P= 0.850) between the two groups but rather
 207 observed that there could be a higher risk associated with exclusive breastfeeding (OR=
 208 3.118).
 209

210 Presence of domestic animals and seasonal variations

211 From this study, 25.6% of children who had domestic animals at home (N=156) had
 212 *Cryptosporidium* infection whilst 19.6% of children who did not have any domestic animal at
 213 home (N= 209) had infection of the parasite. The difference however was not significant (P=
 214 0.171).
 215

216 *Cryptosporidium* infections in the dry season (December – March) was 37/189 (19.6%), and
 217 44/176 (25%) during the rainy season (April- November). The observations made suggest

218 **Table 1. *Cryptosporidium* sp., *G. lamblia* and *E. histolytica* infections among 365**
 219 **diarrhoeic and 120 non-diarrhoeic children at the PML hospital, Accra, Ghana**

Parasite	Number and Percentage (%) infected			P- value
	Symptomatic n (%)	Asymptomatic n (%)	Total, n (%)	
<i>G. lamblia</i>	21 (5.8)	6 (5.0)	27 (5.6)	P< 0.0001 P<0.0001 P=0.0143
<i>Cryptosporidium</i> sp.	81 (22.2)	4 (3.3)	85 (17.5)	
<i>E. histolytica</i>	3 (0.8)	0 (0.0)	3 (0.6)	

220
 221

Table 2: Detection of *Cryptosporidium sp.* infection in diarrhoeic and non-diarrhoeic stool by Microscopy and Enzyme immunoassay

<i>Diarrhoea stool</i>					
	No. tested	No. Positive	%	No. Negative	%
Microscopy	365	32	8.77	333	91.23
ELISA	365	81	22.19	344	94.25
<i>Non-diarrhoea stool</i>					
	No. tested	No. Positive	%	No. Negative	%
Microscopy	120	2	1.67	118	98.33
ELISA	120	4	3.33	116	96.67

225
226
227
228

Table 3. Clinical characteristics of *Cryptosporidium sp.* infections in 81 positive cases

		Number (N)	Percentage (%)
Consistency of stool	Mucoid	1	1.2
	Slimy	5	6.2
	Loose	75	92.6
Symptoms	Diarrhoea alone	0	0
	Diarrhoea & vomiting	69	85.2
	Diarrhoea, vomiting & fever	12	14.8
Days of hospitalization	1 day	3	3.7
	3 days	32	39.5
	4 days	46	56.8

229
230
231
232

Table 4. Odds ratios and 95% confidence intervals for *Cryptosporidium* infection in children and associated risk factors (N = 365)

Character/Risk factor		No	No infected	% infected	X ²	P-value	OR	95% CI
1. Gender	Male	211	46	21.8	0.044	0.883	0.948	0.576-1.561
	Female	154	35	22.7				
2. Age (months)	< 6	74	13	17.6	13.043	0.011	2.856	1.672-3.042
	6 - 12	165	49	29.7			3.111	1.973-3.201
	13 - 24	107	16	15			2.050	0.432-2.854
	25 - 36	9	3	33.3			3.432	0.325-3.654
	37 - 48	10	0	0			-	-
3. Education level of mother	Tertiary	28	13	17.6	13.043	0.011	0.926	0.654-1.453
	Secondary	250	49	29.7			2.437	1.768-2.763
	Primary	40	16	15			1.232	1.108-2.319
	None	47	3	33.3			3.629	2.456-3.896
4. Source of drinking water	Sachet	337	78	23.1	2.457	0.293	1.935	0.763-2.176
	Pipe-borne	26	3	11.5			0.725	0.564-1.342
	Both	2	0	0			1.000	0.895-1.241
5. Breast-feeding habits	Exclusive	218	47	21.6	0.325	0.850	3.118	0.567-3.344
	Not exclusive	131	31	23.7			2.827	0.679-3.129

	N/R	16	3	18.8			1.724	0.768-1.881
6.Presence of domestic animals at home	Yes	156	40	25.6	1.877	0.171	1.433	0.861-2.320
	No	209	41	19.6				
7.Seasonal variations	Dry	189	37	19.6	1.553	0.213	0.730	0.443-1.199
	Rainy	176	44	25.0				

that the parasite has no specific preference for any of the seasons, as difference between seasonal occurrence was not significant ($P=0.213$, $OR=0.73$, $95\%CI=0.443-1.199$).

Microscopy and enzyme immunoassay tests

Of the 365 diarrhoea samples tested, 81 and 32 were positive for ELISA and microscopy, respectively (Table 2). Also, of the 120 non-diarrhoea samples tested, 2 and 4 samples were positive by microscopy and ELISA, respectively. From a total of 85 positive samples by ELISA 72 tested positive by PCR, which amplified a 450bp fragment of the 18S rRNA gene.

Clinical data

The observed clinical characteristics associated with *Cryptosporidium* sp. infection have been reported in Table 3. Majority (92.6%) of samples tested were loose diarrhoeic samples. Diarrhoea and vomiting constituted the predominant clinical symptom (85.2%). Days of hospitalization varied between 1-4 days, and most patients spent 3-4 days on admission at the hospital.

PCR-RFLP analysis and sequencing of *Cryptosporidium* sp.

PCR amplification was successful for 72 ELISA-positive samples (Fig. 1). The RFLP with *SspI* and *VspI* enzymes gave single patterns of fragments for all PCR products which when compared with previously reported work (24) (Table 5) were indicative of *C. parvum*. Sequencing was carried out for 6 representative PCR products, and all the samples, EMO312, EMO303, EMO289, EMO311, EMO325 and EMO346, had good sequence reads.




Fig 1 A Nested PCR at the 18S ribosomal RNA gene locus for identification of *Cryptosporidium* sp. (Lane 1 is 100bp molecular ladder, Lanes 3 to 16 are patient samples. Lanes 17 and 2 are positive & negative controls respectively).


263 **Table 5. *Cryptosporidium* sp. and genotypes determined by RFLP of the amplicon**
 264 **defined by CPB-DIAGR/F primers after digestion with the enzymes VspI, DraI and DdeI**
 265 **(Nichols et al., 2003)**

<i>Cryptosporidium</i> spp. and genotypes(total amplicon length in bp)					GenBank accession number
	Amplicon length(s) (bp) as defined with				
	VspI	DraI	DdeI		
<i>C. parvum</i> 1 (438)	222,104,112	None	204,68,166	L16997	
<i>C. parvum</i> 2 (435)	219,104,112	None	201,68,166	L16996,AF161856	
<i>C. muris</i> (432)	320,112	None	42,224,166	AF093498,AF093498,AF093497	
<i>C. andersoni</i> (431)	319,112	None	265,166	AF093496,49069	
<i>C. felis</i> (455)	239,104,112	50,405	221,68,166	AF087577	
<i>C. baileyi</i> (428)	212,104,112	84,344	262,166	L19068,AF093495	
<i>C. maleagris</i> (434)	47,171,104,112	None	200,68,166	AF112574	
<i>C. serpentis</i> (430)	312,112	None	264,166	AF093502	
<i>C. wrairi</i> (435)	219,104,112	None	201,68,166	AF115378	

266
 267 BLAST searches of the NCBI database however confirmed only one of the sequences,
 268 EMO346, with >95% sequence identity, as *C. parvum*. In a separate phylogenetic analysis
 269 (Fig. 2), sample EMO346 formed a clade with *C. parvum* KKU and *C. parvum* M31, together
 270 with other 10 strains, also previously reported as *C. parvum*. Samples EMO312, EMO303,
 271 EMO289, EMO311, and EMO325 formed a separate cluster with unspecific identity (Fig. 2).

272 273 DISCUSSION

274 In many developing countries including Ghana, diarrhoeal diseases remain a major cause of
 275 morbidity and mortality among children. *Cryptosporidium* sp. is an important enteric parasite
 276 associated with diarrhoea particularly in communities without proper sanitation and potable
 277 water.
 278



279
280
281
282
283
284
285
286

Fig 2. Phylogram (maximum likelihood) of all the *Cryptosporidium* 18S ribosomal RNA gene partial sequences and selected members of the *Cryptosporidium* sp. species inferred from 18S rDNA sequence comparisons. Bootstrap values (within branches at relevant nodes) are reported when equal or greater than 40%. Emo 346, Emo 312, Emo 303, Emo 289, Emo 325, Emo 311 are isolates from the present study at PML, Accra, Ghana. at Mulago hospital in Uganda (18), Cuban children (31, 12) and Ethiopian children (28). In many places worldwide, children of both sexes in same location are likely to be equally

287 In our hospital based study among children hospitalized for acute diarrhoea at PML, Accra,
288 we recorded a 22.2% prevalence of *Cryptosporidium* sp., and observed cryptosporidial
289 diarrhoea to be more common than diarrhoea caused by other enteric parasites. The
290 prevalence appears to be reasonably high and an increase over previous observations of
291 8.7% in the year 2010 at the same hospital (21). Our findings suggest that the parasite has
292 increasingly become more important in paediatric diarrhoea and now needs urgent attention
293 by health authorities and researchers in the country than ever. The association of
294 *Cryptosporidium* with acute to very severe forms of diarrhoea in children leading to
295 hospitalization and deaths has been reported worldwide (19, 26, 27, 28).
296

297 Knowing the importance of accurate reporting, we employed a combination of two separate
298 laboratory methods for diagnosis of parasites in the stool samples. Although microscopy has
299 a high specificity, it could be less sensitive in its detection of parasites. Immunoassays
300 however have high sensitivity and can be used as effective tool for epidemiological
301 investigations (29, 30). We are convinced to a large extent that an application of the two
302 methods for each sample has enabled us to report the magnitude of the disease correctly
303 and accurately. Our findings underscore a critical look at the laboratory methods currently
304 used for diagnosis of *Cryptosporidium* sp. in our hospitals. The use of microscopy alone, as
305 being done now in many hospitals may not be enough to diagnose the infection.
306

307 An effective prevention strategy requires a good understanding of factors associated with
308 occurrence of the disease. We investigated risk factors that could be associated with
309 cryptosporidial diarrhoea among the children and found no association with gender. Our
310 observation was similar to those made among Ugandan children hospitalized for diarrhoea
311

312 exposed to any environmental contamination. Based on this assumption, observations made
313 in the present study were expected.
314

315 *Cryptosporidium* infection was relatively common among age group 3 (13-24 months) and
316 below, with no infection among children above 3 years. In Ghana, a previous study reported
317 a peak of infection in children 12-24 months, and a decrease with age thereafter (22).
318 Reports from investigators worldwide indicate that most infections occur among younger
319 children of ages 2 years and below (32, 33, 22). The observations from the present study
320 therefore agree with many others worldwide. An exception however was in Lagos, Nigeria,
321 where the parasite was found to be most common among the age group 4- 5 years (34), in
322 contrast to findings in our study.
323

324 High prevalence of the parasite among children whose mothers had no formal education
325 appears to conform to the general notion that illiterate mothers practice less personal
326 hygiene. In a semi-urban slum in India, children whose mothers had no formal education
327 suffered multiple symptomatic *Cryptosporidium* infections significantly (35). We presume that
328 in Ghana majority of illiterate mothers lack knowledge of transmission of such infections
329 which could have resulted in the high prevalence of cryptosporidiosis observed in this study.
330 It will be a laudable idea to expand health education among prospecting mothers to cover
331 knowledge on common infectious diseases at antenatal clinics.
332

333 Breast milk has been highly recommended for use in feeding infants worldwide because of
334 its immunological and nutritional protection value. In Ghana, there have been so much
335 campaign by the Ghana Health Service (GHS), an outfit of the Ministry of Health (MOH) for
336 exclusive breastfeeding by mothers as it is more protective (12, 36). In Israel (37), children
337 who were exclusively breastfed received at least, some protection from *Cryptosporidium* sp.
338 infections within the first 3 months after birth. According to our study, majority of mothers
339 practiced exclusive breastfeeding but unfortunately it appears not to have provided
340

expected protection against *Cryptosporidium* parasites. We are of the view that the subject of exclusive breastfeeding in relation to enteric parasitic infections in Accra needs further investigation.

Consumption of untreated water is one of the major routes by which the parasite is transmitted into humans. In our study, most infections occurred in children whose drinking water was from sachet (bagged) water. It is not surprising, as the purity and safety of sachet water in Accra has been previously challenged by a study (38). According to that study 77% of the samples screened contained parasitic agents including oocysts of *Cryptosporidium* sp. To ensure public health safety of the consumption of sachet water in Accra, regular screening and monitoring of brands of sachet water sold in the city of Accra is largely recommended.

Cryptosporidiosis being a zoonotic infection, we suspected that there could be some association between infection of the parasite and pet ownership at home. The keeping of pets and rearing of domestic animals at home is a common practice in the Ghanaian society. We observed that a majority of children had animals at home, and these were either dogs or cats, or both. Most of these animals contaminate the environment which also serves as playing grounds for children by their faeces. We found that having pets or domestic animals at home was not associated with *Cryptosporidium* infection. Perhaps good hygiene and sanitation in homes with such animals played a role to prevent infection.

It was noticed in our study that the parasite was present all year round and seasonal variations were not significant. The pattern seems to be in line with erratic supply of potable water in Accra for domestic use which occurs throughout the year. Large population of the city depends on untreated water which is also likely to increase the risk of infection all year round. Elsewhere, *Cryptosporidium* infections vary significantly from one season to the other (39, 40, 41).

Our genotyping results indicate that *Cryptosporidium parvum* is a major cause of paediatric cryptosporidial diarrhoea at the hospital. *C. parvum* comprises of both anthroponotic and zoonotic species (subtypes 1 and 2). Subtype 1 of *C. parvum* is anthroponotic whilst subtype 2 is zoonotic. Being unable to perform sub-genotyping analysis to determine specific subtypes, our study is limited in reporting whether all infections were human to human, or animals involved in transmission. That notwithstanding, we consider our study to be significant for being able to determine the particular species of *Cryptosporidium* in paediatric diarrhoea at the hospital for the first time.

Further investigations are needed to confirm the identity of other genotypes with unclear identity found in our study. They could be new genetically modified species being introduced into the population of Accra.

4. CONCLUSION

Our study suggests that *Cryptosporidium parvum* is the commonest enteric parasite which causes diarrhoea among children with acute diarrhoea at the Princess Marie Louise children's hospital at Accra, Ghana. Infection was highest among children 3 years and below, and children who drank sachet water had infections most. The findings of our study shows that laboratory methods presently employed in routine diagnosis of the parasite in many hospitals need a critical consideration. A carefully planned health education among illiterate mothers and improved personal hygiene could reduce rate of infections among children. Further studies on sub-genotyping of *C. parvum* isolates are needed to determine whether source of infection is zoonotic or anthroponotic.

393 **CONSENT**

394

395 Selection of participants (patients) in the study was strictly voluntary and based on informed
396 consent (verbal) of their guardians or parents, after being informed of the objectives and
397 goals of the study.

398

399

400 **ETHICAL APPROVAL**

401 An ethical approval of the study (SAHS-Et/SAHS/PSM-ML/1/26A/2010-2011) was obtained
402 from the Research and Ethical Review Committee of the University of Ghana School of
403 Allied Health Sciences, College of Health Sciences, Korle Bu, Accra.

404

405

406 **REFERENCES**

407

- 408 1. Kosek M, Bern C, Guerrant RL. The global burden of diarrheal disease, as
409 estimated from studies published between 1992 and 2000. Bull World Health Organ.
410 2003; 81, 197–204.
- 411 2. Boschi-Pinto C, Velebit L, Shibuya K. Estimating child mortality due to diarrhoea in
412 developing countries. Bull World Health Organ. 2008; 86:710–717.
- 413 3. United Nations Children's Fund and World Health Organization 'WHO/UNICEF Joint
414 Report: Diarrhoea: Why children are still dying and what can be done'. UNICEF,
415 New York/ WHO Library Cataloging-in-Publication Data; 2009.
- 416 4. Current WL, Garcia LS. Cryptosporidiosis. Clin Microbiol Rev. 1991; 4: 325-358.
- 417 5. McLauchlin J, Amar C, Pedraza-Di'az S, Nichols G L. Molecular Epidemiological
418 Analysis of *Cryptosporidium* spp. in the United Kingdom: Results of Genotyping
419 *Cryptosporidium* spp. in 1,705 Fecal Samples from Humans and 105 Fecal Samples
420 from Livestock Animals. J Clin Microbiol. 2000; 38(11), 3984–3990.
- 421 6. Winkworth CL, Learmonth JL, Matthaci CD, Townsend CR. Molecular
422 Characterisation of *Giardia* Isolates from calves and humans in a region in which
423 dairy farming has recently intensified. Appl Environ Microbiol. 2008; 74 (19): 5100-
424 5105.
- 425 7. Epe C, Rehker G, Schnieder T, Lorentzen L, Kreienbrock L. *Giardia* in symptomatic
426 dogs and cats in Europe- Results of a European study. Vet Parasitol. 2010; 173, 32-
427 38.
- 428 8. Hamnes IS, Gjerde BK., Robertson L J. A longitudinal study on the occurrence of
429 *Cryptosporidium* and *Giardia* in dogs during their first year of life. Acta veterinaria
430 scandinavica. 2007; 49: 22.
- 431 9. Kutz SJ, Thompson RCA, Polley L, Kandola K, Nagy J, Wielenga C M et al. *Giardia*
432 assemblage A: human genotype in muskoxen in the Canadian Artic. Parasit Vectors.
433 2008; 1:32.
- 434 10. Xiao L, Fayer R, Ryan U, Upton SJ. Cryptosporidium taxonomy: recent advances
435 and implications for public health," Clin Microbiol Rev. 2004; 17(1): 72– 97.
- 436
- 437
- 438
- 439
- 440
- 441
- 442
- 443
- 444

- 445 11. Cama VA., Bern C, Roberts J, Cabrera L., Sterling C R., Ortega Y et al.
446 *Cryptosporidium* Species and Subtypes and Clinical Manifestations in Children,
447 Peru. *Emerg Infect Dis* 2008; 14 (10), 1567-1573.
448
- 449 12. Pelayo L, Nuñez FA, Rojas L, Wilke H, Furuseth Hansen E, et al. Molecular and
450 epidemiological investigations of cryptosporidiosis in Cuban children. *Ann Trop Med*
451 *Parasitol.* 2008; 102 (8), 659-669.
452
- 453 13. Yakoob J, Abbas Z, Beg MA, Naz S, Khan R, Islam M et al. Prevalences of *Giardia*
454 *lamblia* and *Cryptosporidium parvum* infection in adults presenting with chronic
455 diarrhoea. *Ann Trop Med Parasitol.* 2010; 104(6): 505-510.
456
- 457 14. Sulaiman IM, Hira PR, Zhou L, Al-Ali FM, Al-Shelahi FA, Shweiki HM et al. Unique
458 endemicity of cryptosporidiosis in children in Kuwait. *J Clin Microbiol.* 2005; 43: 6,
459 2805-2809
460
- 461 15. Morgan UM, Pallant L, Dwyer BW, Forbes DA, Rich G, Thompson RC. Comparison
462 of *Cryptosporidium parvum* in human faecal specimens: clinical trial. *J Clin Microbiol.*
463 1998; 36:995–998.
- 464 16. Cacciò SM, Thompson RC, McLauchlin J, Smith HV. Unravelling *Cryptosporidium*
465 and *Giardia* epidemiology, *Trends Parasitol.* 2005; (21) 9: 430-438.
466
- 467 17. Essid R, Mousli M, Aoun K, Abdelmalek R, Mellouli F, Kanoun F et al. Identification
468 of *Cryptosporidium* Species Infecting Humans in Tunisia. *Am J Trop Med Hyg.* 2008;
469 79(5):702–705.
470
- 471 18. Tumwine JK, Kekitiinwa A, Nabukeera N, Akiyoshi DE, Rich SM, Widmer G et al.
472 *Cryptosporidium parvum* in children with diarrhea in Mulago Hospital, Kampala,
473 Uganda. *Am J Trop Med Hyg.* 2003; 68(6), 710-715.
474
- 475 19. Maikai BV, Umoh JU, Lawal IA, Kudi AC, Ejembi CL, Xiao L. (2012) Molecular
476 Characterizations of *Cryptosporidium*, *Giardia* and *Enterocytozoon* in humans in
477 Kaduna State, Nigeria. *Exp Parasitol.* 2012; 131: 452-456.
478
- 479 20. Molloy SF, Smith HV, Kirwan P, Nichols RA, Asaolu SO, Connelly L et al.
480 Identification of a High Diversity of *Cryptosporidium* Species Genotypes and
481 Subtypes in a Pediatric Population in Nigeria. *Am J Trop Med Hyg.* 2010; 82(4):
482 608–613
- 483 21. Opintan JA, Newman MJ, Ayeh-Kumi PF, Afrim R, Gepi-Attee R, Sevilleja JE.
484 Pediatric Diarrhea in Southern Ghana: Etiology and Association with Intestinal
485 Inflammation and Malnutrition. *Am J Trop Med Hyg.* 2010; 83(4): 936–943.
- 486
- 487 22. Adjei A, Lartey M, Adiku TK, Rodrigues O, Renner L, Sifah E et al. (2003)
488 *Cryptosporidium* oocysts in Ghanaian AIDS patients with diarrhoea. *East African*
489 *Med J.* 2003; 80 (7); 369-372.
490
- 491 23. World Health Organization. Basic Laboratory Manuals In Medical Parasitology.
492 WHO Library Cataloguing in Publication Data (NLM Classification: WC 25); 1999.
493

- 494 24. Nichols RA, Campbell BM, Smith HV. (2003) Identification of *Cryptosporidium* spp.
495 Oocysts in United Kingdom Noncarbonated Natural Mineral Waters and Drinking
496 Waters by Using a Modified Nested PCR-Restriction Fragment Length
497 Polymorphism Assay. Appl Environ Microbiol. 2003; 69 (7): 4183-4189.
498
499 25. (<http://www.technelysium.com.au/chromas>. html
500
501 26. Al-Braiken FA, Amin A, Beeching NJ, Hommel M, Hart CA. (2003). Detection of
502 *Cryptosporidium* amongst diarrhoeic and asymptomatic children in Jeddah, Saudi
503 Arabia. Ann Trop Med Parasitol. 2003; 97 (5): 505-510.
504
505 27. Velasquez DE, Arvelo W, Cama VA, López B, Reyes L, Roellig DM et al. Molecular
506 Insights for *Giardia* , *Cryptosporidium* , and Soil Transmitted Helminths from a
507 Facility-Based Surveillance System in Guatemala. Am J Trop Med Hyg. 2011; 85
508 (6): 1141–1143.
509
510 28. Eyasu T, Beyene P, Tekola E. Prevalence of Giardiasis and Cryptosporidiosis
511 among children in relation to water sources in Selected Village of Pawi Special
512 District in Benishangul-Gumuz Region, Northwestern Ethiopia. Ethiop J Health Dev.
513 2010; 10;24 (3): 205-213.
514
515 29. den Hartog J, Rosenbaum L, Wood Z, Burt D, Petri Jr WA. Diagnosis of Multiple
516 Enteric Protozoan Infections by Enzyme-Linked Immunosorbent Assay in the
517 Guatemalan Highlands. Am J Trop Med Hyg. 2013; 88 (1): 167–171.
518
519 30. Goldin AJ, Apt W, Aguilera X, Zulantay I, Warhurst DC. Efficient Diagnosis of
520 Giardiasis Among Nursery and Primary School Children in Santiago, Chile by
521 Capture Elisa for the Detection of Fecal Giardia Antigens. Am J Trop Med Hyg.
522 1990; 42 (6): 538-545.

523 31. Bello J, Nunez FA, Gonzalez OM, Fernandez R, Almirall P, Escobedo AA. Risk
524 factors for Giardia infection among hospitalized children in Cuba. Annals of Tropical
525 Medicine & Parasitology. 2011; 105 (10): 57–64
526
527 32. Alexander KA, Herbein J, Zajac A. The Occurrence of *Cryptosporidium* and *Giardia*
528 Infections among Patients Reporting Diarrheal Disease in Chobe District, Botswana.
529 Advances in Infectious Diseases. 2012; 2: 143-147.

530 33. Wangeci G, Wamae C, Mbae C, Waruru A, Mulenge E, Waithera T, et al.
531 *Cryptosporidiosis* prevalence, genotype analysis, and symptoms associated with
532 infections in children in Kenya. Am J Trop Med Hyg. 2006; 75 (1): 78–82.

533 34. Wellington O, Chika O, Teslim O, Oladipo O, Adetayo F, Godswill I. *Cryptosporidium*
534 and other Intestinal Protozoans in Children with Diarrhoea in Lagos, Nigeria. The
535 internet Journal of Trop Med. 2009; 5: 2.

536 35. Ajjampur SSR, Sarkar R, Sankaran P, Kannan A, Menon VK, Muliyl J, et al.
537 Symptomatic and Asymptomatic *Cryptosporidium* Infections in Children in a Semi-
538 Urban Slum Community in Southern India. Am J Trop Med Hyg. 2010; 83 (5):
539 1110–1115.
540

- 541 36. Mahmud MA, Chappell CL, Hossain MM, Huang DB, Habib M, DuPont HL. Impact of
542 breast-feeding on *Giardia lamblia* infections in Bilbeis, Egypt. Am J Trop Med Hyg.
543 2001; 65 (3): 257–260.
544
- 545 37. Bilenko N, Ghosh R, Levy A, Deckelbaum RJ, Fraser D. Partial breastfeeding
546 protects Bedouin infants from infection and morbidity: prospective cohort study.
547 Asia Pac J Clin Nutr. 2008; 17 (2): 243-249.
548
- 549 38. Kwakye-Nuako G, Borketey PB, Mensah-Attipoe I, Asmah RH, Ayeh-Kumi PF.
550 Sachet drinking water in Accra: The potential threats of transmission of enteric
551 pathogenic protozoan organisms. Ghana Medical Journal. 2007; 41 (2): 62-67.
552
- 553 39. Wongstitwilairoong B, Srijan A, Serichantalergs O, Fukuda CD, McDaniel P,
554 Bodhidatta L, et al. Intestinal Parasitic Infections among Pre-School Children in
555 Sangkhlaburi, Thailand. Am J Trop Med Hyg. 2007; 76 (2): 345–350.
556
- 557 40. Hussein TK. Prevalence and related risk factors for *Giardia lamblia* infection among
558 Children with Acute Diarrhea in Thi-Qar, Southern Iraq. Thi-Qar Medical Journal.
559 2010; 4 (4): 68- 74.
560
- 561 41. Siwila J, Phiri IGK, Enemark HL, Nchito M, Olsen A. Seasonal Prevalence and
562 Incidence of *Cryptosporidium* spp. and *Giardia duodenalis* and associated
563 diarrhoea in Children attending Pre-School in Kafue, Zambia. Trans Roy Soc Trop
564 Med Hyg. 2011; 105 (2): 102-108.
565