

# A COMPARISON OF THE ACCURACY OF THE USE OF LAST MENSTRUAL PERIOD AND SYMPHYSIO-FUNDAL HEIGHT FOR GESTATIONAL AGE DETERMINATION AMONG NIGERIAN WOMEN

Ogbe AE<sup>1\*</sup>, Ekwempu CC<sup>2</sup>, Musa J<sup>2</sup>, Anzaku AS<sup>3</sup>

<sup>1</sup>Department of Obstetrics and Gynaecology, Dalhatu Araf Specialist Hospital, Lafia, Nigeria

<sup>2</sup>Department of Obstetrics and Gynaecology, Jos university Teaching Hospital, Jos, Nigeria

<sup>3</sup>Department of Obstetrics and Gynaecology, College of Health Sciences, Bingham

University, Karu, Jos Campus, Jos, Nigeria

## ABSTRACT

**Aims:** The aim of the study was to evaluate the accuracy of the symphysiofundal height (SFH) in comparison to the last menstrual period (LMP) for gestational age assessment

**Study design:** Hospital-based prospective cross sectional study.

**Place and Duration of Study:** Maternity unit of the Jos University Teaching Hospital, Jos, Nigeria, between December 2012 and April 2013.

**Methodology:** A total of 289 consecutive consenting women with singleton uncomplicated pregnancies at gestational ages of less than or equal to 20 weeks were recruited at the maternity unit of Jos University Teaching Hospital between December 2012 and April 2013. Ultrasound scan (USS) was used to confirm eligibility after which other information including the LMP were documented on a questionnaire. The women returned after 22 weeks' gestation based on ultrasound recorded GA for SFH assessment and some weeks thereafter for a second SFH assessment.

**Results:** Mean age of the women was  $28.9 \pm 4.8$  years with a range of 16-42 years. Most of them were of parity 1 – 4 (58.1%). The mean GA at booking was  $15.3 \pm 3.1$  weeks based on LMP and  $14.9 \pm 3.1$  from early ultrasound scan. The mean percentage accuracy for SFH method compared to USS dating was 95.8% while that of LMP was 91.0%. This difference was found to be statistically significant ( $P = .02$ ).

**Conclusion:** The study showed a significant difference between the LMP and early ultrasound scan dating but not between SFH and early ultrasound scan. Also, the mean percentage accuracy was statistically higher for SFH, suggesting that SFH was a more accurate tool for gestational age assessment among these women.

*Keywords: Comparison, Gestational age, Menstrual date, Symphysiofundal height, Nigeria*

## 1. INTRODUCTION

The accurate determination of gestational age is the cornerstone of optimal obstetric care. The diagnosis and interventions given to a pregnant woman including antepartum haemorrhage, intrauterine growth restriction, pre-term and post-term deliveries, the categorisation of pre-labour rupture of membranes (whether at term or pre-term) and the need for steroid administration, tocolysis or induction of labour, all rest on the correct assessment of the gestational age at the time of evaluation. In addition, it is important to the pregnant woman and her family who wants to know when to expect the baby<sup>1-4</sup>.

There are different ways of assessing the gestational age. Traditionally, the first day of the last menstrual period has been used to calculate the day of delivery (which is supposed to occur 280 days later) and date the pregnancy<sup>5,6</sup>. But this Naegele's rule is based on certain assumptions hence if a woman has irregular menstrual cycles, has been on hormonal contraception, had lactational amenorrhoea prior to conception, or has poor recollection of her last menstrual period, then the application of the rule becomes difficult<sup>2,3,7,8,9</sup>.

The symphysiofundal height (SFH) measurement in centimetres using a non-elastic tape, can be used after the 20 weeks to 34 weeks of gestation but it has its own limitations in dating pregnancy in multiple pregnancy, co-existing fibroid and pregnancy, polyhydramnios, oligohydramnios, fetal transverse lie<sup>2,10,11</sup>.

The gold standard for assessing gestational age has been the use of early ultrasound scan done at first half of the pregnancy<sup>2,4,12</sup>. In resource endowed countries, ultrasound scan has become an essential part of obstetric practice but the same cannot be said of low resource countries like Nigeria especially in the rural areas<sup>13</sup>. In this setting, when available, the ultrasound machine is often of poor quality and operated by undertrained technicians. Furthermore, late presentation of our women for antenatal care in our environment, is a factor that limits the usefulness of ultrasound in assessing gestational age<sup>3,7,14,15</sup>.

Previous studies conducted in Nigeria suggest that accurate measurement of symphysiofundal height is a reliable method of gestational age assessment in the second half of pregnancy<sup>13,16</sup>. Similarly, a study done in Pakistan suggested that symphysio-fundal height measurement was more accurate than the reported last menstrual period as a tool for assessing gestational age when the ultrasound scan was used as a reference<sup>2</sup>.

In developing countries where illiteracy compounds the general poor recollection of the LMP among pregnant women and where ultrasound facility is still not so widely available<sup>2-4,7,13,17</sup>, an alternative method that is independent of patient's recollection but based on the accurate measurement on the spot by the physician, will go a long way to solve the dilemma of inaccurate assessment of gestational age. Hence, this study was carried out in our center to compare the accuracy of the LMP with the symphysio-fundal height in assessing gestational age, using the ultrasound as a reference. This method will be particularly useful in our rural areas where ultrasound equipment and the needed expertise are not readily available.

## 2. MATERIAL AND METHODS

This hospital-based cross-sectional prospective research was carried out in the maternity unit of the Department of Obstetrics and Gynaecology, Jos University Teaching Hospital (JUTH), a tertiary health institution situated in Jos, Plateau state. The hospital is one of the three federal teaching hospitals in the North-Central zone of Nigeria. It serves as a referral center for the neighbouring states of Bauchi, Gombe, Benue, Kogi, Nasarawa, Taraba, Adamawa, part of Kaduna States. Jos is the capital city of Plateau State.

The study population comprised of consecutive pregnant women with singleton pregnancy in the antenatal clinic with estimated gestational age less than or equal to 20 weeks' gestation who consented to participate in the study. Women with co-existing pelvic mass, polyhydramnios or oligohydramnios, a fetus in transverse lie or fetal intrauterine growth restriction were excluded from the study.

## 2.1 Study Procedure

On presentation for booking at the maternity unit of JUTH, eligible women were directed to the investigator who explained the purpose of the study to them. Those who consented to participate signed a written consent form. The investigator, who has received training in obstetric ultrasonography, then proceeded to have their gestational age determined by ultrasound to confirm eligibility after which other information were collected on a questionnaire including the first day of their last menstrual period (LMP) as they recalled. For those who could not recall the first day of their LMP, the 15<sup>th</sup> day of the recalled month was used. The number of weeks between the LMP and the day of enrolment provided the gestational age estimates.

The ultrasound scan was performed by means of a real-time Toshiba model (OTPS- 320A ultrasound machine fitted with a 3.75MHz **transabdominal** sector probe (Otawara, Japan, 2001). The fetal gestational sac, crown-rump (CRL) and biparietal diameter (BPD) were used depending on the gestational age (GA), using reliable landmarks and planes<sup>19</sup>.

The women were allowed to go home to return for the study after 22 weeks of the ultrasound recorded gestational age for SFH assessment by a senior registrar who was blinded to the GA estimated previously. The subjects had their bladder emptied and after lying down in supine position, each fundus of the uterus was outlined by gentle palpation. A non-elastic tape was then used with the graduation in centimetres facing the abdomen (to reduce bias) to take the measurement in the midline from the fundus uteri to the upper part of the pubic symphysis. Two measurements were taken for each subject and each measurement was rounded up to the nearest centimetre. The average of the two measurements was taken and the number of centimetres considered as corresponding to GA in weeks<sup>20,21</sup>. A second measurement of the symphysiofundal height was done in a similar manner **by the same senior registrar (who was the one that did it for all the women)** two to six weeks later. The women were then seen by their respective managing units in subsequent antenatal clinics until delivery.

The data was collected using the study questionnaires including results of the early ultrasound findings and the subsequent SFH measurements. The information obtained was entered into a pre-design program in the Epi-info software version 3.5.4(CDC, Atlanta Georgia, USA) and analyzed. The mean gestational ages were calculated for the Ultrasound scan estimation, LMP and SFH readings for the first and second visits after the 22weeks of gestation and the mean differences determined. The test of hypothesis was done using the student t-test and a *P*-value of less than .05 was considered statistically significant. **Approval for this study was granted by the research and ethical committee of the Jos University Teaching Hospital, Jos.**

## 3. RESULT

A total of 301 women were screened between December 2012 and April 2013 for the study at booking. Nine of them were excluded as a result of the diagnosis of twin pregnancy and another 3 for pregnancy coexisting with uterine fibroids.

Table 1 shows the socio-demographics features of the 289 study population analyzed. The mean age of the women was  $28.9 \pm 4.8$  years with a range of 16-42 years. Most of them were within the age group of 20-34 years (82.0%) and 9 (3.1%) were teenagers. Most of the women were of tribes from Plateau state (Plateau tribes) (50.9%), Christians (80.6%), had at least secondary school education (88.2%), Married (99.3%) and Civil servants (45.0%). One hundred and twenty one of them were nulliparous (41.9%), while 168 (58.1%) have had at least one prior delivery, out of which 21 (7.2%) were grandmultiparous women.

The average gestational age at booking from the LMP was  $15.3 \pm 3.1$  weeks while that from the early ultrasound scan done was  $14.9 \pm 3.1$  **with a range of 8-20 weeks**. The total number of women who were unsure of the first day of their LMP was 24, giving a prevalence of 8.3% of the total population. Table 2 presents the mean differences between the gestational ages obtained by the ultrasound scan and those of the last menstrual period (LMP) and symphysio-fundal height (SFH) for both the first and the second visits after 22 weeks of gestation. The mean differences were higher for the LMP values and these were found to be statistically significant ( $P = .001$ ). The mean differences for the SFH values were not statistically significant.

Table 3 shows the mean percentage accuracy which is higher for the SFH compared to the LMP (95.8% Vs 91%) and this is statistically significant ( $P = .02$ ).

**Table 1: Baseline Characteristics of the Women**

| Variable                       | Total (N=289) | Percent (%) |
|--------------------------------|---------------|-------------|
| <b>Age Group (Mother)</b>      |               |             |
| ≤19 years                      | 9             | 3.1         |
| 20-34 years                    | 237           | 82.0        |
| ≥ 35 years                     | 43            | 14.9        |
| <b>Ethnicity</b>               |               |             |
| Hausa                          | 40            | 13.8        |
| Igbo                           | 20            | 6.9         |
| Plateau tribes                 | 147           | 50.9        |
| Yoruba                         | 22            | 7.6         |
| Others                         | 60            | 20.8        |
| <b>Religion</b>                |               |             |
| Christian                      | 233           | 80.6        |
| Muslim                         | 56            | 19.4        |
| <b>Education of the Mother</b> |               |             |
| None                           | 28            | 9.7         |
| Primary                        | 6             | 2.0         |
| Some Secondary                 | 182           | 63.0        |
| Completed Secondary            | 17            | 5.9         |
| Tertiary                       | 56            | 19.4        |
| <b>Education of Spouse</b>     |               |             |
| None                           | 4             | 1.4         |
| Primary                        | 8             | 2.8         |
| Some Secondary                 | 9             | 3.1         |
| Completed Secondary            | 180           | 62.3        |
| Tertiary                       | 88            | 30.4        |
| <b>Marital Status</b>          |               |             |
| Married                        | 287           | 99.4        |
| Never Married                  | 1             | 0.3         |
| Widowed                        | 1             | 0.3         |
| <b>Occupation of Mother</b>    |               |             |
| Civil Servant                  | 130           | 45.0        |
| Trader                         | 41            | 14.2        |
| Seamstress                     | 85            | 29.4        |
| Housewife                      | 21            | 7.3         |
| Others                         | 12            | 4.1         |
| <b>Parity</b>                  |               |             |
| 0                              | 121           | 41.9        |
| 1-4                            | 147           | 50.9        |
| ≥5                             | 21            | 7.2         |

144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158

159 **Table 2: Comparing the accuracy of the LMP and SFH using Ultrasound as the Standard**

| Variables            | Mean difference | Standard deviation | 95% Confidence Interval | t-test | P-Value |
|----------------------|-----------------|--------------------|-------------------------|--------|---------|
| EGAUSS-EGALMP        | -0.391          | 1.501              | (-0.565 to -0.217)      | 4.429  | .001    |
| EGA2USS-EGA2LMP      | -0.391          | 1.501              | (-0.565 to -0.217)      | 4.429  | .001    |
| EGAUSS-EGASFH1       | -0.028          | 1.504              | (-0.202 to -0.146)      | 0.313  | .755    |
| EGA2USS-<br>EGA2SFH2 | -0.031          | 1.449              | (-0.199 to -0.137)      | 0.365  | .715    |

160

161 EGAUSS= Ultrasound estimated gestational age first visit after 22 weeks

162 EGALMP= Last menstrual period estimated gestational age first visit after 22 weeks

163 EGA2USS= Ultrasound estimated gestational age second visit after 22 weeks

164 EGA2LMP= Last menstrual period estimated gestational age second visit after 22 weeks

165 EGASFH1= SFH estimated gestational age first visit after 22 weeks

166 EGA2SFH2= SFH estimated gestational age second visit after 22 weeks

167

168 **Table 3: Comparison of Mean Percentage accuracy of LMP and SFH**

| Method | No of Subjects | Gestational age Assessment |                 | P-value |
|--------|----------------|----------------------------|-----------------|---------|
|        |                | Mean Accuracy* n(%)        | Inaccurate n(%) |         |
| SFH    | 289            | 277 (95.8%)                | 12 (4.2%)       | .02     |
| LMP    | 289            | 263 (91%)                  | 26 (9%)         |         |

169

170 \*Accuracy was defined by the number of Subjects within 2 weeks of USS estimated gestational age.

171

#### 172 4. DISCUSSION

173

174 Our study showed that symphysiofundal height is a better method for gestational age assessment compared to the last  
 175 menstrual period. This was reflected in the fact that the paired mean differences of the LMP with the early ultrasound scan  
 176 were higher and found to be statistically significant ( $P = .001$ ) compared to the mean differences of the SFH with the early  
 177 ultrasound scan which were not as high and statistically insignificant ( $P = .755$  and  $.715$ ). This implied that the SFH  
 178 assessment from the study was not statistically different from the early ultrasound scan as a tool for gestational age  
 179 assessment and may be more useful than the LMP in the absence of an early ultrasound scan. Furthermore, the mean  
 180 percentage accuracy of the last menstrual period compared to early ultrasound scan was 91% as against the mean

percentage accuracy of 95.8% for the symphysis-fundal height compared to the early ultrasound scan. And this higher accuracy with the symphysis-fundal height method was found to be statistically significant ( $P = .02$ ). Although traditionally the LMP is used to date pregnancy, it is subjective compared to the SFH which may be considered more objective with identifiable landmarks. The subjective problem with the LMP could be because a woman may fail to remember for sure the first day of her last menstrual period and just assigned one of the days around it. This is generally compounded by the level of literacy. However, even a literate woman who conceives after a period of lactational amenorrhoea or oral contraceptive pill will obviously have a problem knowing her LMP. Up to 24 (8.3%) of the women from our study could not recall the first day of their LMP. It has been estimated that 10-45% of pregnant women cannot provide useful information about their LMP and that 18% of women with certain menstrual dates have significant differences between their menstrual and ultrasound dating<sup>17,22</sup>.

Our finding is similar to the finding from the study by Adewale and Ijaiya that the SFH is a reliable method of gestational age assessment in the second half of pregnancy<sup>13</sup>. Although in that study, the researchers did not compare the SFH with the LMP to ascertain the more accurate tool. In a prospective study conducted in Cape Town, South Africa to compare different dating methods commonly used in their setting, the authors concluded that ultrasound dating was most accurate of all the other methods, but their study also revealed a better agreement between the ultrasound dating method and the SFH compared with the LMP<sup>23</sup>.

In a longitudinal study conducted in rural Guatemala between 1996 and 1999 for 171 women-infant pairs, comparing LMP, neonatal examination and SFH using ultrasound as the reference, the authors concluded that the recalled LMP provided the best estimate of gestational age<sup>3</sup>. They observed in addition, that the SFH measure during the second trimester may provide a reasonable alternative when the LMP is unavailable<sup>3</sup>. In their study however, the researchers used SFH measured between 15 and 24 weeks of gestation on the assumption that this earlier measure would be less influenced by growth retardation. But as a tool for determining gestational age (GA), it is generally reported that SFH in centimeters equals GA in weeks between 20 and 34 weeks<sup>10,11</sup>. Hence, the early measure could have provided a suboptimal assessment of the SFH in that population.

In Nigeria and other resource-constraint developing countries where ultrasound machines are not readily available especially in the rural areas, and where even when available the expertise may be lacking, the knowledge that SFH which is simple, cheap, easy to learn method is more accurate than the LMP will make estimation of gestational age easier and less doubtful than previously perceived.

The strength of the study is that, it was a prospective study and early ultrasound scan was done for all the women. In addition a good proportion of the women (88.2%) had at least secondary school education (lower educational attainment is one of the sociodemographic characteristics associated with uncertain LMP<sup>24</sup>).

The limitation of this study is that although ultrasound was used as the gold standard in this study, it has its own error margin even with the best biometric measurement which is the CRL ( $\pm 5$  to 7 days). Also, Parity and body mass index which are covariates for SFH measurement in this study were not considered during data collection. Another limitation of the study is the fact that the few women (8.3%) who could not recall their LMP, the 15<sup>th</sup> day of the month was used to calculate their EGA. The convenience sampling technique is subject to sampling bias, hence, the sample might not be representative of the entire population. This limits the ability to generalize the findings of the study to the entire population.

## 5. CONCLUSION

The findings from this study showed a significant difference between the last menstrual period and the early ultrasound scan as tools for gestational age assessment, but not between symphysis-fundal height and early ultrasound scan. In addition, the mean percentage accuracy of the symphysis-fundal height was higher and statistically more significant compared to that of the last menstrual period. This suggests that barring factors which may erroneously increase the symphysiofundal height estimation like multiple pregnancy, polyhydramnios, co-existing pelvic mass or other factors which may abnormally decrease the symphysiofundal height estimation like fetal transverse lie, oligohydramnios or intrauterine growth restriction, the SFH estimation is a more accurate tool for gestational age assessment after the 22<sup>nd</sup> week of gestation compared to the LMP in this Nigerian obstetric population.

It is our recommendation that, there should be training and re-training of Doctors, Midwives and Nurses at different levels of health care in the proper technique for symphysis-fundal height measurement for GA assessment as described in this study especially in low resource settings where ultrasound is a luxury and where women have poor recollection of their LMP.

## ACKNOWLEDGEMENTS

We acknowledge all the nursing and record staff of the maternity unit of the Jos University Teaching Hospital for their assistance and cooperation.

## COMPETING INTERESTS

We declare that no competing interest exists.

## AUTHORS' CONTRIBUTIONS

Dr. Ogbe AE designed the study, wrote the protocol, performed the statistical analysis and wrote the first draft. Dr Musa J and Dr Ekwempu CC designed the study objectives and managed the analyses in addition to contributing to the second draft. Dr Anzaku AS managed the literature search and contributed to the final draft.

## ETHICAL APPROVAL AND PATIENT CONSENT

The proposal for the study was presented to the institutional research and ethical committee of the Jos University Teaching Hospital and approval was duly obtained for it. Consent was obtained from each subject before enlistment.

## REFERENCES

1. Peek MJ, Devonald KJ, Beiby R, Ellwood D. The value of routine early pregnancy ultrasound in the antenatal booking clinic. *Aust NZJ Obstet Gynaecol*, 1994; 34(2): 140-3.
2. Jehan I, ZaidiS, Rizvi S, et al. Dating gestational age by last menstrual period, symphysis-fundal height, and ultrasound in Urban Pakistan. *International journal of Obstetrics and Gynaecology*, 2010;110(3): 231-234.
3. Neufeld LM, Haas JD, Grajeda R, Martorell R. Last menstrual period provides the best estimates of gestation length for women in rural Guatemala. *PaediatrperinatEpidemiol*, 2006; 20 (4): 290-298.
4. Mongelli M, Gardosi J. Evaluation of gestation. e-medicine Obstetrics and Gynaecology. Updated 2010. [Online]. Available from: <http://www.emedicinemedscape.com/article/259269-overview>. (Accessed 30/03/11).
5. Naegele FC. *Lehrburch der Beburtshtilfe fur Hebammen*. 3<sup>rd</sup> ed. 1836.
6. Saunders N, Paterson C. Can we abandon Naegele's rule? *Lancet*, 1991; 337: 600-601.
7. KalishRB, Chervenak FA. Sonographic determination of gestational age. *Ultrasound Rev ObstetGynaecol*, 2005; 5 (4): 254-258.
8. Mongelli M, Wilcox M, Gardosi J. Estimating the date of confinement: Ultrasonographic biometry versus certain menstrual dates. *Am J Obstet Gynaecol*1996 ; 174 (1): 278-281.
9. Savitz DA, Terry JW, Dole N, Thorp JM, Siegal-Riz AM, Herring AH. Comparison of pregnancy dating by last menstrual period, ultrasound scanning and their combination. *Am J ObstetGynaecol*, 2002; 187 (6): 1660-1666.
10. Calvert JP, Crean EE, NewcombeRG, PearseJF. Antenatal screening by measurement of symphysis-fundal height. *Br Med J*, 1982; 285: 846-849.
11. Quaranta P, Currell R, Redman CWG, RoinsonJS. Prediction of small-for-date infants by measurements of symphysis-fundal height. *Br J Obstet Gynaecol*, 1981; 88: 115-119.
12. Taipale P, Hiilesmaa V. Predicting delivery date by ultrasound and last menstrual period in early gestation. *Obstet Gynecol*, 2001; 97(2): 189-194.
13. Adewale FB, Ijaiya MA. Symphysio-fundal Height as a means of Gestational age Assessment in the second half of pregnancy at the University of Ilorin Teaching Hospital, Nigeria. *NJOG*, 2011; 6(2): 27-32.
14. Ibrahim SA, Galadanci HS, Omale AE. Gestational age at first antenatal attendance in Kano, northern Nigeria. *Highland Medical Research journal*, 2007; 5 (1): 75-78.
15. Mutihir JT, Maduka WE, Ujah IAO. Gestational age at booking of primigravidae in Jos, north-central Nigeria. *Journal of med in the tropics*, 2004; 6 (1): 13-18.
16. Kuti O, Osinusi BO, Ojengbede OA. Assessment of Gestational Age in the third trimester of pregnancy: Comparison of Symphysio-fundal Height Measurements with Ultrasonic Methods. *Trop J ObstetGynaecol* 2003; 20: 123-126.
17. GeirrsonRT, Busby- Earle RM. Certain dates may not provide a reliable estimate of gestational age. *Br J ObstetGynaecol*, 1991; 98(1): 108-109.

18. Plateau state people and culture. Plateau state Gov.org 2006. [Online]. Available from: <http://www.plateaustategov.org/history/people-cul.html>. (Accessed 30/03/11).
19. Romero R, Quintero R, and Brekus C. Assessment of gestational age. In: Divon M (ed). Abnormal fetal growth. Elsevier science: New York, NY, 1991.
20. Westin B. Gravidogram and fetal growth (Comparison with biochemical supervision). Acta obstet Gynecol scand, 1977; 56:273-282.
21. Lindhard A, Nielsen PV, Mouritsen LA, Zachariassen A, Sorensen HU, Roseno H. The implications of introducing the symphyseal-fundal height-measurement. A prospective randomized controlled trial. BJOG, 1990; 97:675-680.
22. Buekens P, Delvoye P, Wollast E, Robyn C. Epidemiology of pregnancies with unknown last menstrual period. *Journal of Epidemiology and Community Health* 1984; 38: 79-80.
23. Geerts I, Poggenpoel E, Theron G. A comparison of pregnancy dating methods commonly used in South Africa: A prospective study. *S Afr Med J*, 2013; 103 (8): 552-556.
24. Hall MH, Carr-Hill RA, Fraser C, Campbell D, Samphier ML. The extent and antecedents of uncertain gestation. *British journal of Obstetrics and Gynaecology* 1985; 92: 445-451.

## ABBREVIATIONS

EGAUSS= Ultrasound estimated gestational age first visit after 22 weeks  
EGALMP= Last menstrual period estimated gestational age first visit after 22 weeks  
EGA2USS= Ultrasound estimated gestational age second visit after 22 weeks  
EGA2LMP= Last menstrual period estimated gestational age second visit after 22 weeks  
EGASFH1= SFH estimated gestational age first visit after 22 weeks  
EGA2SFH2= SFH estimated gestational age second visit after 22 weeks