

1
2
3
4
5
6
7
8

9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

Original Research Article

Incidence of nontreponemal reactive test in women at childbirth in a maternity facility in the central region of Rondonia state, Brazil

ABSTRACT

Background: Syphilis is an infectious sexually transmitted disease caused by the spirochaeta *Treponema pallidum*. This disease is still a problem in public health, particularly in developing countries.

Aims: This study aimed to analyze the incidence of syphilis in women at childbirth in a maternity facility in the city of Ji-Paraná, state of Rondônia in Brazil.

Study design: An analytical, cross-sectional, prospective study.

Place and Duration of Study: Department of Pharmaceutical Sciences (CEULJI/ULBRA) and the public maternity facility of the Ji-Paraná city, between August 2 to October 2, 2014.

Methodology: We investigated a total of 300 pregnant women at childbirth. When VDRL test was positive a semi-structured interview was administered.

Results: We found 5 women (1.66%) with VDRL positive tests, however only one newborn, whose mother did not perform prenatal or therapy care, had a significant serology titration according to Brazilian diagnostic guidelines. Diagnosed women in this research were all adults with an average of 21.8±2.8 years of age, with basic education, married, and low income.

Conclusion: The estimated prevalence of VDRL positive tests in women at childbirth was similar to that reported in the latest studies in Brazil. However, the results found in this study show that Brazilian Unified Health System is far from the established goal for 2015 and strongly suggests that it may not be reached.

Keywords: Venereal Disease Research Laboratory (VDRL); Unified Health System (SUS); Childbirth; Brazil.

1. INTRODUCTION

Syphilis is a Sexually Transmitted Disease (STD), caused by the spirochaeta *Treponema pallidum* in a systemic way [1]. When untreated or treated improperly, the disease evolves compromising the skin, internal organs, and central nervous system. Syphilis is still a problem in public health, especially in developing countries. The World Health Organization (WHO) estimated that in 2008, 12 million people, including about 2 million pregnant women were infected by *T. pallidum*. In addition, there is an increasing trend in the incidence of syphilis and congenital syphilis (CS) worldwide [2,3]. The WHO [21] estimative for developing countries is between 10 and 15% pregnant women infected by *T. pallidum*.

24 Congenital syphilis occurs with the transmission and hematogenous dissemination of *T.*
25 *pallidum* in pregnant women untreated or treated inadequately to the fetus through the
26 placenta. When infection happens in the first 6 months of pregnancy, vertical transmission in
27 untreated women occurs in 70 to 100% of the cases, and in approximately 30% of the cases,
28 for infection after the 6th month of pregnancy [3-8].

29
30 In places funded by the Brazilian Unified Health System (SUS), pregnant women during
31 prenatal and birth must perform the screening for syphilis through the nontreponemal test
32 (VDRL - Venereal Disease Research Laboratory). The screening must be performed in the
33 first prenatal medical appointment, in the third trimester of pregnancy, and at childbirth or
34 curettage [6-8].

35
36 Until the year of 2013, the nontreponemal test VDRL was used for CS screening, therapeutic
37 follow-up and control. After this date, the rapid treponemal test which is target-specific was
38 introduced, but it is still not available in all country regions. In the facilities where only VDRL
39 is available, laboratory confirmation with treponemal tests for reactive VDRL, even though
40 being decisive is not mandatory according to SUS policy [4,6,9-11]. In cases when the
41 mother is reactive in the VDRL test, the newborn must be tested, if the result is of the child is
42 also reactive, with clinical evidence or not, the treatment proceeds according to WHO
43 guidelines [2,4,6].

44
45 Since 2005, the Brazilian Ministry of Health considers syphilis a notifiable disease. Previous
46 works have reported flaws in the notification system, and it is known that the disease
47 frequency actually is higher than shown by official data [6,8]. In this context, in which this
48 disease is still a challenge in Brazil, this study aimed to analyze the prevalence of syphilis in
49 women at childbirth in a maternity facility in the city of Ji-Paraná.

50 51 2. METHODS AND MATERIALS

52 53 2.1 Study area

54
55 This study was conducted in the city of Ji-Paraná, Rondônia, Brazil, located in the North
56 region of the country (Geographic coordinates: 10°53'07" S - 61°57'06" W). The estimated
57 population for the year 2014 is 129,242 inhabitants and the Human Development Index
58 (HDI) is 0.714, according to the census of the Brazilian Institute of Geography and Statistics
59 (IBGE) in 2010. The maternity facility of the public Hospital was chosen because it serves
60 the majority of women in the central area of the State.

61 62 2.2 Study Design

63
64 An analytical, cross-sectional, prospective study with qualitative and quantitative approach
65 was carried out in a two-month period from August 2 to October 2 in 2014.

66 67 2.3 Data Collection

68 69 2.3.1 Sample method

70
71 We investigated 300 pregnant women attending the public maternity facility for childbirth,
72 covering the entire sampling space. Data collection was performed every day of the week
73 during the study period, regardless of childbirth type (natural or C-section).

2.3.2 VDRL Test

Blood samples of all participants were obtained by peripheral venous puncture, collecting 3 mL of mother's blood. Newborn blood samples were collected for VDRL test only when the mother's sample was reactive. All investigations were carried out according to manufacturers' instructions. VDRL test (Record N°.10009010223; Labtest Diagnostica SA, Lagoa Santa, MG, Brazil) in newborn was considered significant when the titration was at least four times mother titration [6].

2.3.3 Interviews

A semi-structured interview was administered for collection of demographic and clinical information in women with reactive VDRL test. The studied variables in the interview are presented in Table 1.

3. RESULTS AND DISCUSSION

We analyzed 300 pregnant women in this study, and in 1.66% of the cases VDRL tests were reactive. Diagnosed women in this research were all adults with an average of 21.8 ± 2.8 years of age, with basic education, married, and low income. Table 1 presents the risk factors for syphilis diagnostic.


Table 1. Profile of patients positive for VDRL

Variables	Woman 1	Woman 2	Woman 3	Woman 4	Woman 5
Age	20	22	25	18	24
Color	White	Brown	White	Brown	White
Marital status	Married	Married	Married	Married	Married
Education	Basic	High School	Basic	Basic	Basic
Income ¹	234.50	469.00	> 234.50	234.50	469.00
Number of pregnancy	2	2	2	1	1
Age of 1 st sexual relationship	17	18	17	15	18
Used condom?	No	No	No	No	No
Number of sexual partners	2	3	1	2	2
If condom is always used	No	Sometimes	No	Sometimes	No
Knowledge about STD	Yes	Yes	No	No	Yes
Communication method	Lecture	Radio	-	-	Radio
When syphilis was diagnosed?	Prenatal care	Prenatal care	Pre childbirth	Prenatal care	Prenatal care
If during pregnancy, which trimester?	1 st	2 nd	3 rd	2 nd	1 st
Medical treatment	Yes	Yes	No	Yes	Yes

¹ US\$ per month

102 The WHO [21] estimated that developing countries had between 10 and 15% of pregnant
 103 women infected by *T. pallidum*. Research work on syphilis in Brazil reported that each year
 104 about three million women give birth and syphilis prevalence rate is estimated in 1.6% [3, 12-
 105 15]. The Brazilian Ministry of Health had joined efforts to eradicate syphilis in Brazil. The
 106 official goal was set to reach eradication (0.5 cases per 1000 live births) in 2015. In order to
 107 eradicate the disease, the government had intensified awareness campaigns and treatment
 108 programs, as well as improving the notification system. Syphilis has been reported more
 109 commonly in single women with active sex life which suggests that number of sexual
 110 partners is more significant than frequency of sexual intercourse for this disease [3].

111
 112 The results of serological tests were reported in Figure 1. We found that only the newborn
 113 from Woman 3, who did not receive prenatal care or therapy, had significant serology
 114 titration (1:16) associated with syphilitic characteristics.



115
 116 **Fig. 1. Patients' titration. *Significant value according Brazilian Ministry of Health.**
 117

118 According to previous works, titrations 1:02 should be considered suspect and the
 119 treponemal test should be performed. This is because neonates may be in immunologic
 120 window period in which low titrations occur, even false negative or false positive due to the
 121 presence of maternal antibodies and/or cross-reactivity [22-25]. Our results confirm that, the
 122 timing of antenatal care interventions makes a significant difference in the risk of having an
 123 adverse outcome due to syphilis. Women who sought care in the first two trimesters of their
 124 pregnancy, and received the appropriate intervention, were more likely to have a healthy
 125 infant, compared to women screened and treated in the third trimester [24,26].

126 127 4. CONCLUSION

128
 129 The estimated prevalence of VDRL positive tests in women at childbirth was similar to that
 130 reported in the latest studies in Brazil. However, the results found in this study show that
 131 Brazilian unified health system is far from the established goal for 2015 and strongly
 132 suggests that it may not be reached.

CONSENT

All authors declare that “written informed consent” was obtained from the patients for publication of this research. The original copies are archived in the Department of Biomedicine, CEULJI/ULBRA. The questionnaires were kept separated from the written consent, making it impossible to know which data belongs to a particular participant.

ETHICAL APPROVAL

The authors have obtained ethical approval from the Research Ethic Committee (CEP-CEULJI/ULBRA: 79.807/2014) and Legal authorization from the Hospital manager.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

5 REFERENCES

1. Hook EW, Peeling RW. Syphilis control - a continuing challenge. The New England Journal of Medicine. 2004; 351:122-124.
2. Saraceni V, Madeira RMSD, Vellozo V, Lauria LM, Dias MAB, Ratto KMN, Durovni B. Vigilância da sífilis na gravidez. Epidemiologia e Serviços de Saúde. 2007;16(2):103-111. Portuguese.
3. Araújo CL, Shimizu HE, Sousa AIA, Hamann EM. Incidência da sífilis congênita no Brasil e sua relação com a Estratégia Saúde da Família. Revista de Saúde Pública. 2012;46(3):479-486. Portuguese.
4. Chaudhary M, Kashyap B, Bhalla P. Congenital syphilis, still a reality in 21st century: a case report. Journal of Medical Case Reports, 2004; 90(1):1-13.
5. Sidana R, Mangala HC, Murugesh SB, Ravindra K. Prozone phenomenon in secondary syphilis. Indian journal of sexually transmitted diseases. 2011;32(1):47-49.
6. Departamento de DST, AIDS e Hepatites Virais Secretaria de Vigilância em Saúde Ministério da Saúde. Available: www.aids.gov.br. Accessed on Setembro/2014. Portuguese.
7. Lumbiganon P, Piaggio G, Villar J, Pinol A, Bakketeig L, Bergsjö P, Berendes H. The epidemiology of syphilis in pregnancy. International journal of STD & AIDS. 2002;13(7):486-494.
8. Solanke VN, Kawane UP, Karmarkar MG, Mehta PR. A Relevance of Serological Tests in Diagnosis of Early Congenital Syphilis: A Case Report. International Journal of Prevention and Treatment 2014, 3(1): 8-10.
9. Qin JB, Feng TJ, Yang TB, Hong FC, Lan LN, Zhang CL, Dong W. Risk factors for congenital syphilis and adverse pregnancy outcomes in offspring of women with syphilis in Shenzhen, China: a prospective nested case-control study. Sexually transmitted diseases. 2014;41(1):13-23.
10. French P, Gomberg M, Janier M, Schmidt B, Van Voorst Vader P, Young H. IUSTI: 2008 European guidelines on the management of syphilis. International Journal of STD & AIDS. 2009;20(5):300-309.
11. Domingues RMSM, Saraceni V, Hartz ZMDA, Leal MDC. Sífilis congênita: evento sentinela da qualidade da assistência pré-natal. Revista de Saúde Pública. 2013;47(1):147-157.

- 184 12. da Costa CC, Freitas LV, do Nascimento Sousa DM, Lopes O, de Castro
185 Damasceno AK. Sífilis congênita no Ceará: análise epidemiológica de uma década.
186 Revista da Escola de Enfermagem da USP, 2013;47(1):152-159. Portuguese.
187 13. Carvalho IS, Brito RS. Sífilis congênita no Rio Grande do Norte: estudo descritivo do
188 período 2007-2010. Epidemiologia e Serviço de saúde. 2014;23(2):287-294.
189 Portuguese.
190 14. Magalhães DMS, Kawaguchi IAL, Dias A, Calderon IMP. Sífilis materna e congênita:
191 ainda um desafio. Caderno de Saúde Pública, 2013;29(6):1109-1120. Portuguese.
192 15. Amaral E. Sífilis na gravidez e óbito fetal: de volta para o futuro. Revista Brasileira
193 de Ginecologia e Obstetrícia. 2012;34(2):970.
194 16. Komka MR, Lago EG. Sífilis congênita: notificação e realidade. Scientia Medica.
195 2007;17(4):205-1. Portuguese.
196 17. Lima MG, Santos RFR, Barbosa GJA, RGS. Incidência e fatores de risco para sífilis
197 congênita em Belo Horizonte, Minas Gerais, 2001-2008. Ciência e Saúde Coletiva.
198 2013;18(2):499-506.
199 18. Mascola L, Blount JH, Alexander CE, Cates W Jr. Congenital syphilis. Why is it still
200 occurring? JAMA 1984;252:1719-22.
201 19. Genç M, Ledger WJ. Syphilis in pregnancy. Sexually transmitted infections,
202 2000;76(2):73-79.
203 20. Peeling RW, Ye H. Diagnostic tools for preventing and managing maternal and
204 congenital syphilis: an overview. Bulletin of WHO. 2004;82:439-446.
205 21. World Health Organization. Investment case for eliminating mother-to-child
206 transmission of syphilis: promoting better maternal and child health and stronger
207 health systems. Geneva: World Health Organization; 2012.
208 22. Sá RAM, Bornia RBG, Cunha A A, Oliveira C A, Rocha GPG, Giordano EB. Sífilis e
209 gravidez: avaliação da prevalência e fatores de risco nas gestantes atendidas na
210 Maternidade Escola – UFRJ. Revista Brasileira de Doenças Sexualmente
211 Transmissíveis. 2001;13:6-8.
212 23. Brasil. Portaria nº 3242 do Ministério da Saúde. Diário Oficial da União, Brasília, p.
213 50, 2 jan. 2012. Seção 1. Portuguese.
214 24. Duarte SJH. Motivos que levam as gestantes a fazerem o pré-natal: um estudo das
215 representações sociais. Ciencia y Enfermeria. 2012;18(2):75-82. Portuguese.
216 25. Finelli L, Berman SMK. Congenital syphilis. Bull World Organ. 1998; 76(2):126.
217 26. Hawkes, S. J., Gomez, G. B., & Broutet, N. (2013). Early antenatal care: does it
218 make a difference to outcomes of pregnancy associated with syphilis? A systematic
219 review and meta-analysis. *PloS one*, 8(2), e56713.
220