

Promising therapeutics against Ebola virus disease

Abstract

Ebola virus disease (EVD) has become a major threat to global peace and security. Since 1976, when Ebola virus was discovered, over 20 outbreaks have been reported. Most of these outbreaks occurred in rural areas of East and Central Africa. But the current 2014 outbreak, which started in Guinea in March 2014, spread to Liberia, Sierra Leone, Nigeria, Senegal and Mali, and traveled beyond Africa into Europe and US. Thus far, there have been no approved therapeutics and preventive vaccines and hence response is limited to supportive care, barrier nursing, and management of patient complications. Spurred by the global threat, research has identified promising drug candidates against the disease. This review presents the current status of promising drug candidates against EVD. The current development status of the experimental drugs ZMapp, TKM-Ebola, Favipiravir (T-705 or Avigan), AVI 6002, BCX 4430 and Brincidofovir (CMX-001) is given. In the absence of licensed drugs, these first generation anti-Ebola virus experimental drugs, which are currently in phase 1 clinical trial, were administered to a limited number of healthcare workers during the current EVD outbreak.

Key Words: Antiviral, Anti-coagulant, Filovirus, Hemorrhagic fever

1.0 Introduction

One of the disease that threatened the economy and people of West Africa in 2014 is Ebola virus disease (EVD) formerly called Ebola hemorrhagic fever (EHF) caused by Ebola virus. Ebola virus was discovered in 1976 with simultaneous outbreaks in Zaire (now Democratic Republic of Congo, DRC) and Sudan (World Health Organization (WHO), 2014a). Thereafter, several sporadic outbreaks have occurred in different years and different parts of Africa including DRC (1977, 1995, 2001 – 2003, 2007, 2008 – 2009, 2012), Sudan (1979, 2004), Gabon (1994, 1996 – 1997, 2001 – 2002), Uganda (2000 – 2001, 2007 – 2008 and 2012) and Cote d'Ivoire (1994). The current 2014 outbreak of Ebola virus disease (EVD) in West Africa was alleged to have first occurred in a 2 year baby who died on 6 December 2013 in a remote Gueckedou prefecture of Guinea (Gatherer, 2014; Onwuakor, 2014), a forested area near the border with Sierra Leone and Liberia (The Economist, 2015). Thereafter, nothing was heard of the disease, until a major outbreak was reported in West Africa 3 months after. EVD outbreak was reported in Guinea on 22 March 2014, Liberia on 31 March 2014 and Sierra Leone on 26 May 2014. The disease subsequently spread to Nigeria on 20 July 2014 (Fasina et al., 2014; Shuaib et al., 2014), DRC on 24 August 2014, Senegal on 26 August 2014, and Mali on 25 October 2014 (Ohimain, 2015). In the current outbreak, the first imported case of EVD from Africa was recorded in the US on 30 September 2014 and a second case was reported in Spain on 6 October 2014, when a Spanish nurse developed EVD after treating an infected priest that contacted EVD while working on voluntary/humanitarian basis in West Africa. Other cases of EVD were also reported in other parts of the world including United Kingdom (UK). EVD was spread by the rapidity of modern travel (Olinger et al., 2012; Sarwar et al., 2014), religion, culture, customs, as well as virus characteristics.

The 2014 EVD outbreak which was caused by Zaire Ebola virus (Beeching et al., 2014; Geisbert, 2014; WHO Ebola Response Team, 2014), is considered the largest outbreak of the disease since

1976, when it was first discovered (Geisbert, 2014; Hampton, 2014; Qiu et al., 2014). Bishop (2014) and WHO Ebola Response Team (2014) reported that the current epidemic is larger than all the previous epidemics of EVD combined. Sarwar et al. (2014) described the current epidemic as the worst outbreak to date and the first to be localized primarily in urban areas. On 8 August 2014, the WHO declared EVD outbreak to be an epidemic of public health emergency of international concern (WHO Ebola Response Team, 2014). Statistics abound on the devastating effects of EVD in West Africa. In July 2014, the total cases of Ebola was 1,407 with 743 death (WHO, 2014a). As of 15 August 2014, there were 2,127 total cases of EVD with 1,145 deaths in Guinea, Sierra Leone, Liberia and Nigeria (Qiu et al., 2014) and by 14 September 2014, the total number of cases had increased to 4,507 with 2,296 deaths (WHO Ebola Response Team, 2014). By 28 December, a total of 20,206 cases and 7,905 deaths from EVD have been reported worldwide, with the largest incidence and fatalities in the 3 West African countries of Guinea (2,707 cases with 1,709 deaths), Liberia (8,018 cases with 3,423 deaths) and Sierra Leone (9,446 cases with 2,753 deaths) (The Economist, 2015). While, the spread of Ebola virus have been contained in Nigeria and Senegal, the disease is still ravaging Guinea, Sierra Leone and Liberia.

Ebola virus disease is a severe and often fatal disease in humans and other primates caused by the virus in the genus Ebola and belonging to the Filoviridae family. Five species of Ebola virus have been documented including Zaire Ebola virus, Sudan Ebola virus, Bundibugyo Ebola virus, Tai Forest Ebola virus and Reston Ebola virus (WHO, 2014a, b). Of these, the Sudan and Zaire Ebola virus are the most virulent (Bausch, 2014). The current outbreak in West Africa is caused by the Zaire Ebola virus. According to WHO (2014a), the commonest symptoms of EVD are sudden, onset of fever, intense weakness, muscles pain and sore throat, which is followed by vomiting, diarrhea, bleeding (internal and external) and multi-organ damage particularly the kidney, spleen and liver especially in advanced stage. In the 2014 EVD outbreak, the commonest symptoms were fever (87.1%), fatigue (76.4%), loss of appetite (64.5%), vomiting (67.6%), diarrhea (65.5%), head ache (53.4%), abdominal pain (44.3%), and unexplained bleeding (18%) (Beeching et al., 2014). The virulence of Ebola virus appears to differ among these virus species, with case fatality being highest in Zaire species (60 – 90%) followed by Sudan species (40- 60%) (Feldmann and Geisbert, 2011). Many other authors have reported the case fatality of Zaire Ebola virus to be up to 90% (Basler and Amarasinghe, 2009; Geisbert et al., 2010; Johansen et al., 2013; Marzi et al., 2012; Oestveich et al., 2014; Qiu et al., 2013a, b; Sarwar et al., 2014).

Recently, studies show that host genetic diversity affects Ebola virus pathogenesis and resistance (Rasmussen et al., 2014). But the fatality rate of the 2014 EVD outbreak is less being 55 – 60% globally (Hampton, 2014) and 40% in Nigeria (Fasina et al., 2014; Ohimain, 2015; Shuaib et al., 2014) as against the well documented 90% fatality rate of Zaire Ebola virus.

The Filoviruses, particularly the Ebola virus have been listed among emerging and re-emerging viral disease of global threat (Marston et al., 2014; Peter et al., 1994). Ebola viruses are evasive, with the capacity of inhibiting type I interferon activity, and therefore reducing or destroying the host (Basler and Amaransinghe, 2009; Geisbert et al., 2010; Watanabe et al., 2007), and these viruses then reduce or destroy the host capacity to combat the infection. Though, research has revealed promising therapeutics and vaccines for EVD, none have been approved by US Food and Drug Agency (FDA) or WHO. Hence, Ebola virus has been described as a possible target for

bioterrorism/bio-weapon (Johansen et al., 2013; Sarwar et al., 2014; Swenson et al., 2009; Warfield et al., 2006). Ebola virus have been classified as biosafety level 4 (BSL4) pathogen (Bente et al., 2009; Enterlein et al., 2006) and “Category A” agents of bioterrorism (Bente et al., 2009; Heald et al., 2014; Pettitt et al., 2013), which therefore requires urgent attention. Because of the antecedents of Ebola virus including its virulence, high fatality rates, evasiveness and the potential for global spread through air travel, it has become urgent to develop therapeutics for treatment and vaccines for prevention of EVD. Hence, this review is focused on potential anti-EVD drugs including antibodies, plasma/serum and antiviral and non-anti-viral drugs with reported activity against EVD. The study methodology was based on critical review of literature including published journals, newspaper reports and broadcast media. The study used the following recent background literature among others: Beeching et al. (2014), Bishop (2014), Butler (2014), Feldmann and Geisbert (2011), IDCRP (2014), Norwegian Institute of Public Health (2014), WHO (2014c-e). Prior to presenting the list of possible drugs and their current development status, Ebola genomes, which are linked to the mode of action of the drugs, shall be first summarized.

2.0 Ebola virus Genome

The genome of Ebola virus, which is about 19kb long, consists of genes that encode for 7 proteins (Enterlein et al., 2006; Feldmann and Geisbert, 2011; Giesbert et al., 2010; Peters et al., 1994; Sobarzo et al., 2012). Four of the genes encode for structural proteins including virion envelope glycoprotein (GP), nucleoprotein (NP) and 2 viral (matrix) proteins VP24 and VP40, while the nonstructural proteins include VP30 and VP35, and the RNA dependent viral polymerase (L) (Sullivan et al., 2003). The order of occurrence of the genes is as follows; 3' Leader, NP, VP35, VP40, GP, VP30, VP24, L and 5' trailer (Geisbert et al., 2010). A simplified version of the genome is presented in Figure 1.

3' Leader	NP	VP35	VP40	GP	VP30	VP24	L	5' Trailer
-----------	----	------	------	----	------	------	---	------------

Key

NP= Nucleoprotein

VP= Virus protein

GP= Glycoprotein

L= RNA dependent RNA polymerase

VP35=Phosphoprotein IFN antagonist.

VP40= Membrane-Associated matrix protein.

VP30=Ribonucleoprotein.

VP24= Membrane-Associated protein.

Figure 1: Simplified structure of Ebola virus genome

Most of these genes have various functions (Table 1), some are potential targets for novel therapeutic agents. For instance, the glycoprotein is responsible for binding and viral entry (Enterlein et al., 2006; Geisbert et al., 2010; Sobarzo et al., 2012; Watanabe et al., 2007), NP, VP35, VP30 and L are responsible for replication and transcription of viral RNA (Enterlein et al., 2006; Sobarzo et al., 2012; Watanabe et al., 2007), while VP 40 and VP 24 are responsible

for assembly, budding and release of **virion particles** (Enterlein et al., 2006; Sobarzo et al., 2012).

The NP encapsulates the genome and forms a complex with **VP30, VP35** and L, which are required for both genomic replication and transcription of viral genes (Kondratowicz and Maury, 2012). The 3 other proteins, GP, **VP40**, and **VP24** are membrane associated proteins (Watanabe et al., 2007), which may be important in the expression of antibodies against Ebola virus antigens. Watanabe et al. (2007) **suggested that VP24** is very important for Ebola virus to evade the antiviral activities of interferons. Also, **VP24**, NP and **VP35** are involved in the formation of nucleocapsid (Watanabe et al., 2007). Geisbert et al. (2010) reported that **VP24** and **VP35** are potential targets for antiviral interventions since both genes have inhibitory effects against host type I interferon response. Also, according to Geisbert et al. (2010), the L proteins which provides the RNA dependent RNA polymerase is a potential target for antiviral drugs because of two reasons, 1) suppression could result in the **inhibition** of viral replication and, 2) such protein is not found in mammalian systems.

Table 1: Functions of Ebola virus genes

Gene	Functions	References
NP	Transcription and replication	Enterlein et al., 2006; Geisbert et al., 2010; Sobarzo et al., 2012; Watanabe et al., 2007
VP35	Transcription and replication	Enterlein et al., 2006; Sobarzo et al., 2012
	RNA synthesis, type 1 interferon antagonist	Enterlein et al., 2006; Watanabe et al., 2007; Warren et al., 2015
VP40	Virus assembly and budding	Enterlein et al., 2006; Kondratowicz and Maury, 2012
GP	Mediate viral entry into cell	Enterlein et al., 2006; Martinez et al., 2011; Sobarzo et al., 2012
VP30	Transcription and replication	Enterlein et al., 2006; Sobarzo et al., 2012; Watanabe et al., 2007
VP24	Virus assembly and budding	Enterlein et al., 2006; Watanabe et al., 2007
	Type 1 interferon antagonist	Geisbert et al., 2010; Warren et al., 2015
L	Transcription and replication	Enterlein et al., 2006; Sobarzo et al., 2012; Watanabe et al., 2007

3.0 Potential EVD therapies

Promising therapies against EVD based on **Research and Development** can be broadly classified into antibody therapies, **specific** antiviral therapies for Ebola virus, anti-inflammatory therapy and anticoagulants. Others include mannose binding lectin and sepsis-related therapies (Table 2). Promising therapeutics were tested in small animal and non-human primates before being considered in human clinical trials. Table 3 presents the leading/ first generation experimental drugs with proven efficacy against EVD that have started preclinical or clinical trials. Though not yet licensed, the WHO and /or FDA have endorsed their limited use on compassionate grounds during an outbreak. Table 4 listed some FDA approved drugs for other treatments that have exhibited antiviral activities, which could also be tried against EVD during outbreaks because of the unavailability of approved therapeutics and vaccines.

168 Table 2: Research towards the development of therapies for Ebola virus disease

Therapy		Humans	Non-Human Primates (NHP) model	Small animal (rodent) model	Remarks/ references
Antibodies	Convalescent plasma	Emond et al., 1977; Mupapa et al., 1999; Sadeket al., 1999; WHO2014d, e	Jahrling et al., 2007		
	Immunoglobulin	Kudoyarova-Zubavichene et al., 1999	Dyeetal., 2012;Jahrlinget al.,1996		
	Monoclonal Antibodies	Maruyama et al., 1999	Geisbert 2014; Qiu et al., 2012a, Qiu et al 2013a, Qiu et al., 2013b; Qiu et al., 2014; Olinger et al., 2012; Oswald et al., 2007	Parrenet al., 2002; Qiu et al., 2012b	
	Polyclonal antibodies			Gupta et al 2001	Feldman and Geisbert 2011; WHO 2014c, d, e, f
	Hyperimmune serum				WHO 2014c, d, e, f
Anti-Viral	LNP/siRNA		Geisbertet al., 2010; Thiet al., 2014	Geisbert et al.,2007	
	PMOs	Healdet al., 2014	Warfield et al., 2006; Warrenet al., 2010	Iversenet al., 2012	
	Polymerase inhibitors		Warrenet al.,2014	Oestereichet al., 2014; Smither et al., 2014	
	Garcinai kola (bitter kola)	Iwu et al., 1999			The kola has antiviral activity, but has not been tested with Ebola virus
	Nanosilver				Speshock and Hussain, 2009
Anti-Inflammatory modulators	interferons (Type 1 (α , β))		Qiu et al., 2013b; Smith et al.,2013	Qiu et al., 2013b	Modulation of immune system. Feldman and Geisbert, 2011
	S-adenosyl homocysteine hydrolase inhibitors				Modulation of immune system. Feldman and Geisbert, 2011
Anti-Coagulants	rhAPC		Geisbert et al., 2003; Richardson et al., 2010		Limited success with NHP. 33% efficacy in the treatment of Zaire Ebola virus infected macaque
	rNAPc2, Tissue factor pathway inhibitors		Geisbert et al., 2003; Hensley et al 2007; Richardson et al 2010; Zhong et al., 2014		Not tested in rodents but confer partial protection in NHPs. Feldman and Geisbert, 2011
	Heparin sulphate				Feldman and Geisbert, 2011
Others	Sepsis-Related Therapies		Hensley et al., 2007		
	Mannose-binding lectin				Ji et al., 2005; Michelow et al., 2011

Abbreviations:
PMOs=Anti-sense phosphoro-diamidatemorpholino oligomers
LNP/siRNA= Lipid nanoparticle/ smallinterfering RNA
rhAPC =Recombinant human activated protein C
rNAPc2 = Recombinant nematode anti-coagulant protein c2

Antibodies

Treatments based on antibodies will be the first to be tried and approved for EVD despite the limited validity and scientific conflicts of the therapy. Research results abounds on the use of blood from convalescent patients for the treatment of EVD (Emond et al., 1977; Gupta et al., 2001; Kudoyarova-Zubavichene et al., 1999; Mupapa et al., 1999). The use of blood transfusion from convalescent patients confers passive immunity on the recipients (Mupapa et al., 1999).

During the 1995 EVD outbreak in DRC, 8 patients who were severely ill of EVD were transfused with blood from convalescent patients. Most of these patients fully recovered (87.5%) i.e. 12.5% fatality in contrast with the overall case fatality rate of 80% in 1995 (Mupapa et al., 1999).

An occasion when serum from convalescent patient given with human interferon was used to successfully treat an EVD patient, viremia decreased rapidly and the patients recovered (Emond et al., 1977; Gupta et al., 2001; Mupapa et al., 1999). The success of these cases and many other unreported cases in West Africa, have fuelled the black market trading of convalescent blood. Despite the successes in humans, few authors have reported failures of blood from convalescent patients to protect humans, mice and guinea pigs against EVD. For instance, Jahrling et al. (1999) reported fatality of all the monkeys transfused with whole blood from convalescent monkeys that were challenged with EVD. Sadek et al. (1999) reported an instance where over 80% of the 250 patients given blood from convalescent patients died of EVD in DRC in 1995. Other authors presented conflicting results on the use of hyperimmune serum for prophylaxis and treatment of EVD (Jahrling et al., 1999; Kudpyarova-Zubavichene et al., 1999). Jahrling et al. (1999) reported that guinea pig were completely protected by injection of hyper immune equine IgG when treatment was initiated early but not after viremia had developed. The authors also reported that the immune IgG could not protect mice and monkeys. Kudpyarova-Zubavichene et al. (1999) used multiple immunization of sheep and goats to obtained hyper immune serum, which demonstrated effective prophylaxis during challenge experiments in guinea pigs and baboon. This therapy was demonstrated to be safe in phase I clinical trials. Convalescent serum was successfully used to treat EVD patients in Spain during the current 2014 EVD outbreak (Almansa et al., 2014). Notwithstanding the conflicting scientific findings, WHO have approved the use of hyper-immune serum and blood from convalescent patients for the treatment of EVD (Gulland, 2014; WHO, 2014c-e). The WHO Blood Regulators Network (BRN) subsequently released the guidelines for the collection and use of convalescent plasma or serum for response against EVD (BRN, 2014). A study testing the use of convalescent serum started in Guinea in February 2015 (Kupferschmidt and Cohen, 2015).

Antibodies produced by convalescent patients, against Ebola virus primarily targeted viral proteins, NP, VP 40, and GP (Sobarzo et al., 2012). For instance, Sobarzo et al. (2013) reported that anti-Ebola IgG was primarily targeted to viral proteins NP, VP 40 and GP. In NHP (non-human primates) and other smaller animal models, it was demonstrated that the adaptive immunity, which contributes to the protection against EVD, is associated with both cellular and humoral immunity (Sobarzo et al., 2012, 2013). Neutralizing monoclonal antibodies can provide

protection against viruses in animal models (Parren et al., 2002; Marzi et al., 2012). Hence, the survival of patients has been linked to the early and robust production of antibodies (Sobarzo et al., 2012); alternately, the antibodies produced may have been consumed due to viral replication (Marzi et al., 2012). Parren et al. (2002) reported an instance when a neutralizing human monoclonal antibody, KZ52, protected guinea pigs from lethal Ebola Zaire virus infection. Hence, KZ52 was able to confer pre- and post-exposure prophylaxis against Ebola virus in a small animal model by passive transfer of a neutralizing human antibody. But Oswald (2007) demonstrated that transfer of neutralizing human monoclonal antibodies failed to protect macaque monkeys against a clinical challenge with Ebola virus and also failed to halt the replication of the virus. Qui et al. (2013b) demonstrated that monoclonal antibodies combined with Adenovirus-vectored interferon significantly extended the treatment window in Ebola virus infected guinea pigs. Dye et al. (2012) obtained polyclonal IgG from NHP that survived EVD and used it as post exposure treatment (delayed for 48 hours) for NHP challenged with Ebola virus. Results show that the IgG protected the macaques. Cocktails of monoclonal antibodies have been successful used for the treatment of NHP infected with Ebola virus. Qui et al. (2012) reported the successful treatment of cynomolgus macaques using a combination of 3 neutralizing antibodies. A cocktail of humanized monoclonal antibodies protected NHP from EVD (Pettit et al., 2013; Qui et al., 2013a, b; Olinger et al., 2012). Based on these success stories, ZMapp was developed. ZMapp is composed of three monoclonal antibodies that have been humanized by genetic engineering and contains monoclonal antibody C13C6 from MB003 and C2G4 and C4G7 from Zmab (ADI, 2014; Geisbert, 2014). ZMapp components are produced in tobacco plant *Nicotiana benthamiana* (ADI, 2014; Qiu et al., 2012a; Qui et al., 2014). Qiu et al. (2014) demonstrated that administration of ZMapp to Ebola virus infected rhesus macaques exhibiting advanced disease symptoms with high viremia and liver enzymes was rescued 100%. ZMapp was able to rescue the monkeys even after 5 days of infection. Though, no approval has been given, but the FDA have permitted its limited use on compassionate ground (Butler, 2014; Hampton, 2014). In the current outbreak, 7 patients were given ZMapp and 2 of these patients died (Butler, 2014). ZMapp contains neutralizing antibodies conferring passive immunity by specifically targeting and reacting with Ebola virus glycoprotein (ADI, 2014; MaCarthy, 2014). The US has signed a contract with Mapp Biopharmaceuticals to accelerate the development of the drug (MaCarthy, 2014). Testing of ZMapp on EVD patients is scheduled to begin in Liberia in February 2015, while Guinea and Sierra Leone could follow after resolving some issues pertaining to research design (Kupferschmidt and Cohen, 2015).

Antiviral Agents

A number of promising antiviral substances with proven efficacy against EVD are being developed (Table 2). Some of the leading antiviral substances are polymerase inhibitors, anti-sense oligomers, lipid nanoparticles/small interfering RNA and selective estrogen receptors. RNA interference (RNAi) based therapies are fast becoming a viable tool for the treatment of many diseases. RNAi is the process of sequence specific, post-transcriptional gene silencing, which suppress the expression of pathologically or physiologically important genes by using small interfering RNA (si RNA) (Uchino et al., 2013). RNAi represents a powerful natural occurring biological strategy for the inhibition of gene expression and has the ability to inhibit viral replication (Geisbert et al., 2006). Ursic- Bedoya et al. (2014) reported protection of guinea pigs against lethal Marburg virus mediated by lipid encapsulated siRNA. Similarly, Geisbert et al. (2006) reported that post exposure protection of guinea pigs against lethal Ebola virus

challenge using siRNAs targeting the polymerase (L) gene of Zaire Ebola virus. The siRNA therapy completely protected guinea pigs against viremia and death. After being successful in small animal models, the use of RNAi was tried in NHP.

In a proof of concept study, Geisbert et al. (2010) demonstrated the post-exposure protection of NHP against a lethal Zaire Ebola virus challenge using siRNA targeted at L, VP 24 and VP 35 genes. The drug TKM-Ebola manufactured by Tekmira Pharmaceutical Corporation, Vancouver, Canada was developed based on RNAi technology. The drug was produced targeting 3 Zaire Ebola virus proteins L, VP 35 and VP 24 (ADI, 2014). These 3 molecules were formulated as stable nucleic acid-lipid particles (SNALPs) in a novel lipid nanoparticle delivery technology. In 2010, Tekmira Pharmaceutical Co signed a contract of \$140 million with US government to fast track the development of the drug. The company started phase 1 clinical trials in January 2014 and as of May 2014 has successfully completed the ascending single dose portion of the phase 1 clinical trials in healthy patients. TKM-Ebola was administered to a limited number of patients during the 2014 EVD outbreak.

Another method for the development of antiviral drug is antisense technology. The potential of oligo-deoxynucleotides to act as antisense agents that inhibit viral replication (Kurreck, 2003) has great promise against EVD. Phosphoro-diamidate morpholino oligomers (PMOs) are synthetic molecules used to inhibit viral replication by specifically binding to RNA, which suppress the expression of selected genes by steric blockage (Warren et al., 2015). The use of antisense PMOs as gene-specific counter measures against EVD in have been well demonstrated in rodents (Enterlein et al., 2006; Swenson et al., 2009; Warfield et al., 2006). Warfield et al. (2006) used a combination of Ebola virus specific PMO, targeted at viral mRNAs for VP 24, VP 35 and L protected rodents in both pre- and post-exposure therapeutics regimes. In a proof of prophylaxis, the authors showed that PMOs also protected 75% of rhesus macaques from lethal Ebola virus infection. Warren et al. (2010) had demonstrated that AVI -6002 containing two PMOs, one targeting VP24 (AVI-7537) and the other targeting VP35 (AVI-7539) protected over 60% of rhesus monkeys against lethal Ebola virus challenge. Warren et al. (2015) further demonstrated that a single PMO targeting VP24 protected rhesus monkeys against lethal Ebola virus infection. Iversen et al. (2012) reported the discovery and early development of PMO AVI-6002 comprising of AVI-7357 and AVI-7539 against Ebola virus. Sarepta Therapeutics (USA) is developing and commercializing AVI-6002 containing AVI-7357 and AVI-7539. Results of phase 1 clinical studies indicated that AVI-6002 is effective against Ebola virus and is safe and well tolerated.

Drugs with broad spectrum antiviral activity are also being considered as possible therapy for EVD. For instance, favipiravir , 6-fluoro-3-hydroxy-2-pyrazine carboxamide(T-705 or Avigan) have broad spectrum antiviral properties against many RNA viruses including influenza virus, Arenaviruses, Bunyaviruses, flaviviruses, Alphaviruses, Picornaviruses and Noroviruses (Furuta et al., 2013). The authors also described favipiravir as a novel compound that selectively and potently inhibit the RNA-dependent RNA polymerase of influenza and many other RNA viruses. Caroline et al. (2014) demonstrated the efficacy of favipiravir for protection from highly lethal inhalation Rift Valley fever in rats. Oestereich et al. (2014) demonstrated the efficacy of favipiravir against Zaire Ebola virus in small animal model both in-vitro and in-vivo. in the small animal model experiments, this author also reported initiation of favipiravir administration at 6

day post infection induced rapid virus clearance, reduced biochemical parameters of disease severity and prevented fatality in 100% of the tested animals. In a mouse model for norovirus infection, Arias et al. (2014) showed that the broad-range antiviral nucleoside favipiravir mode of action involves the reduction of viral load in vivo by exerting antiviral mutagenesis. In mice infected with Ebola virus, results have shown that the initiation of 150 mg/kg favipiravir twice daily within 6 days of infection induced rapid virus clearance, reduced biochemical parameters of disease severity, which led to 100% survival (Mentré et al., 2015; Oestereich et al., 2014; Smither et al., 2014).

Favipiravir, which is produced by Toyama chemicals, Japan is currently under phase III clinical evaluation for influenza in Japan and has completed phase II in US (Furuta et al., 2013) and has commenced phase 1 clinical trial for Ebola virus. During the 2014 EVD outbreak, a French nurse who contracted the disease recovered after taking favipiravir. Favipiravir was recently reported to have halved mortality in one group of Ebola virus patients in Guinea (Kupferschmidt and Cohen, 2015).

Another promising therapy for EVD is BCX.4430 produced by Biocryst Pharmaceuticals, USA. BCX.4430 is an analog of adenosine (a nucleotide) that has a broad spectrum antiviral properties. BCX.4430 acts as a broad spectrum RNA polymerase inhibitor with activity against over 20 RNA viruses in various families (Yelle, 2014). According to Bishop (2014), BCX-4430 indirectly inhibit the RNA polymerase of viruses through non-obligate RNA chain termination resulting in the termination of transcription and replication of the virus. BCX-4430 blocks viral RNA synthesis. Warren et al. (2014) demonstrated that BCX-4430 could confer protection of mouse against lethal doses of Ebola virus challenge up to 96 hours post exposure. BCX-4430 also protected monkeys infected with lethal doses of Marburg virus after 48 hours post infection.

Brincidofovir (CMX 001) manufactured by Chimerix Inc. is also listed among the leading therapeutics against EVD (Table 3). Brincidofovir has antiviral activity against Ebola virus, which is an RNA virus, because Brincidofovir is a phosphonate derivatives of Cidofovir, under development for Cytomegalovirus, Adenovirus and other DNA viruses. Though, there are no published data on the safety and efficacy of Brincidofovir in the treatment of EVD in human, NHPs or small animal models, it was recommended for use on compassionate grounds. In the 2014 EVD outbreak, it was used to treat 5 (Bishop, 2014) Americans infected with Ebola virus. Other drugs that have been recently discovered showing broad-spectrum antiviral, but still under preclinical development include FGI-106 (USAMRIID), FGI-104 (functional Genetics/USAMRIID), FGI-103 (UCLA) and LJ-001 (UCLA) (Norwegian Institute of Public Health, 2014).

Table 3: Leading first generation experimental therapies for Ebola virus disease

Therapy	Manufacturer	Mode of action	Phase of development	Administration route	Remarks
ZMapp	Mapp Biopharmaceuticals Inc, USA	Cocktail of 3 monoclonal antibodies (c13c6, h-13F6, c6D8) binds and inactivates virus	Phase 1	Intravascular	Administered to 7 people of which 2 died. Completely protected monkeys from EVD
TKM Ebola	Tekmira Pharmaceuticals, Canada	si RNA which interferes with L, VP24, VP35. Gene silencing	Phase 1	Intravascular	Protect monkeys infected with Ebola virus. Used for

					humans under emergency in the 2014 EVD outbreak
Favipiravir (T-705 or Avigan)	Toyama Chemical/Fuji Film, Japan and Medi vector USA	Nucleotide analog that inhibits L. Broad spectrum antiviral agent. RNA chain terminator or lethal mutagenesis	Phase 1 (Ebola virus); Phase 3 (influenza)	Oral	Protect mice infected with Ebola virus
AVI 6002	Sarepta Therapeutics, USA	Antisense PMO which inhibits VP24. Gene silencing	Phase 1	Oral or intramuscular	
BCX 4430	Biocryst Pharmaceuticals, USA	Nucleoside which inhibits L. Broad spectrum antiviral agent. RNA chain terminator	Phase 1	Oral or intramuscular	Protect monkeys infected with Ebola virus
Brincidofovir (CMX-001)	ChimerixInc, USA	Ebola: unknown; CMV inhibits DNA synthesis	Phase 1 for Ebola virus, Phase 3 for CMV & ADV	Oral	Used for humans under emergency in the 2014 EVD outbreak

Sources: Bishop (2014); Butler (2014); Goeijenbier et al. (2014); Norwegian Institute of Public Health (2014); WHO (2014c, d, e, f); Yelle (2014)

Anti-inflammatory modulators

Anti-inflammatory modulators have also been listed as effective against EVD including Type I (α , β) interferons, and S-adenosyl homocysteine hydrolase inhibitor (Table 2). Interferons are protein synthesized by the body to fight infections and have been used to successfully treat some diseases such as hepatitis and certain cancers. However, filovirus can evade the protective effects of interferons. Ying et al. (2014) reported that when Ebola virus attacks reticuloendothelia cells, it over-produces cytokines that cause exaggerated inflammatory response, which is not protective. Smith et al. (2012) reported that EVD is associated with robust production interferon- α with plasma concentrations that greatly exceed those observed in other viral infections (60 – 100 fold), but little IFN- β production. Kondratowicz and Maury (2012) reported that filovirus encodes 2 different proteins that block induction of cytokine. Ebola virus VP 24 directly blocks IFN- α / β and IFN- γ signaling, while VP 35 inhibits Type I IFN induction. Sullivan et al. (2003) reported VP 35 acts as an IFN- α / β antagonist. Similarly, Reid et al. (2006) demonstrated that VP 24 inhibits interferon signaling. Furthermore, Kondratowicz and Maury (2012) reported that Ebola virus directly inhibits the antiviral activities of RNAi pathway through the actions of VP 30, VP 35 and VP 40. Hence, the complete absence or reduction of type I interferons in infected macrophages and dendritic cells, which is associated with the dysregulation of the immune system, and the pronounced and aberrant cytokine profile is thought to be responsible for the inappropriately stimulated and quickly depleted immune response characteristics of fatal cases. Smith et al. (2012) reported that the imbalance interferon production (robust production of IFN- α and absence of IFN- β) during Ebola virus infection leads to inefficient antiviral activity and dysregulated lymphocyte apoptosis. Smith et al. (2012) demonstrated that early post-exposure treatment with IFN- β significantly increased survival time (i.e., treatment window) of rhesus macaques infected with a lethal dose of Ebola virus, but failed to prevent fatality/mortality. Similarly, Jahrling et al. (1999) demonstrated that recombinant interferon - α 2b delayed viremia and death several days in monkeys challenged with Ebola virus. Leung et al. (2010) reported that EBOV VP 35 contributes to virulence through the evasion of host immune system and viral RNA synthesis. Similarly, Watanebe et al. (2007) reported that V 24 is important for EBOV to

evade the antiviral effects of IFNs. Hence, interferon therapies is often considered in combination with other therapies. For instance, Qiu et al. (2013b) demonstrated that monoclonal antibodies combined with Adenovirus-vectored interferon significantly extend the treatment window in Ebola virus infected guinea pigs. Administration of the combination therapy at 3 days post infection (dpi) provided 100% survival, which was a significant improvement over survival with either treatment alone. In a follow up study, Qiu et al. (2013b) demonstrated that administration of monoclonal antibodies (ZMAS) combined with Adenovirus vectored IFN- α therapy rescued Ebola virus infected cynomolgus and rhesus macaques and 100% rhesus macaques survived when treatment was administered after detection of viremia at 3 dpi.

Huggins et al. (1999) reported that S-Adenosyl homo-cysteine hydrolase inhibitors inhibit Ebola virus in vitro in a lethal mouse model. They demonstrated 100% survival when treatment was initiated at 0 or 1 dpi, 90% survival at 2 dpi and 40% survival at 3dpi. Based on these results, they claimed that carbocyclic 3- deazadenosine is the first compound demonstrated to cure animals with a lethal Ebola virus infections.

Due to the dysfunctional effect of Ebola virus on the blood system (coagulopathy), anti-coagulants are also being considered as possible cure for EVD. Some of the commonest anticoagulants with demonstrated efficacy include recombinant human activated protein C (rhAPC) and recombinant nematode protein C2 (rNAPC2) and to a lesser extent heparin sulphate (Table 2). Hensley et al. (2007) reported that infection of primate with ZEBOV leads to hypertension, coagulation disorders and impaired immune system similar to Sepsis, which is characterized by rapid decreases in plasma levels of protein C. The authors demonstrated that human activated protein C (Xigris, licensed for treating patients with severe sepsis) delayed death of monkeys to about 12.6 days while only 2 out of 11 treated monkeys i.e. 18% survived. Geisbert et al. (2003) reported that Ebola virus infection induces over expression of pro-coagulant tissue factor in primate monocytes and macrophages, suggesting an inhibition of the tissue-factor pathway. The authors further demonstrated that treatment with rNAPC2 (a potent inhibitor of tissue factor initiated blood coagulation) results in prolong survival of rhesus monkeys challenged with Ebola virus with 33% survival rates. These modest results obtained could be enhanced by the use of combination therapies. Other anti-coagulants such as melatonin have been tested in rats and humans (Tunali et al., 2005; Wirtz et al., 2008a, b). According to Tan et al. (2014), melatonin is a potent free radical scavenger and an anti-inflammatory agent. Masters-Israilov et al. (2015) reported the potential usefulness of melatonin in the treatment of EVD. The authors reported that melatonin has certain features useful against complications arising from EVD such as antioxidant, anti-inflammatory and anti-coagulopathic properties.

Other purpose drugs

The FDA has approved drugs for other purposes, and some of these have potential usage against Ebola virus (Table 4). These drugs include anti-malarial drugs such as chloroquine and amodiaquine (Ekins et al., 2014; Kouznetsova et al., 2014), selective estrogen receptor modulators; clomiphene and toremifene (Johansen et al., 2013) and cationic amphiphiles, which induce a Niemann-Pick C phenotype that inhibit Ebola virus entry and infection (Shoemaker et al., 2013). Kouznetsova et al. (2014) screened several FDA approved drugs for other purposes and found about 53 candidates that can block Ebola virus entry. Of these, about 11 have been

demonstrated for anti-Ebola virus activity (Table 4). Other purpose drugs that have been considered against Ebola virus disease include antivirals (Lamivudine), vitamins (vitamin P/Rutin, Vitamin C, retinazole/vitamin A and statins (Table 4). Kesel et al. (2014) reported that retinazole inhibit certain blood-borne human virus including ZEBOV. Mannose-binding lectin have been reported to be effective against Ebola and Marburg viruses (Ji et al., 2005; Michelow et al., 2007).

Finally, three potential anti-EVD therapies that have been strongly criticized are: bitter kola (*Garcina kola*), Ribavirin and nanosilver. Ribavirin is a broad spectrum antiviral agent that has been used to treat many viral infections including other hemorrhagic fevers, but some authors have reported that the drug is ineffective against Ebola virus (Mupapa et al., 1999; Huggins et al., 1999). Ribavirin when incorporated into the viral genome causes lethal mutagenesis and a subsequent decrease in the specific infectivity of RNA viruses (Vignuzzi et al., 2015). It is therefore surprising that ribavirin, which is an antiviral nucleoside analog that is mutagenic to several RNA viruses is not effective against Ebola virus. More research is therefore required to demonstrate the infectiveness of Ribavirin against Ebola virus.

Iwu et al. (1999) reported the antiviral properties of bitter kola, but the drugs has never been demonstrated as effects against EVD in small animals, NHPs or human models. No published report on the use of nanosilver against EVD apart from un-published 2009 US Department of Defense funded study (Speshock and Hussain, 2009). It has been well demonstrated that nanosilver is effective against monkey pox virus (Rogers et al., 2008), HIV (Elechiguerra et al., 2005), Tacaribe virus (Speshock et al., 2010) and some bacteria (Tyagi et al., 2013; Zhang et al., 2013). Since there are still no approved therapies or prevention vaccines against EVD, and the death toll continues to increase, the research community and drug manufacturing companies should not be in haste to discard any potential lead, but all options should remain on the table for consideration.

Table 4: FDA approved other purpose drugs with potential use against Ebola virus disease

Treatment types	Drugs	Usual uses	Mechanism against Ebola virus	References
Cationic amphiphilic	Clomiphene ^	Ovulation reduction	Induce a Niemann-Pick C-like phenotypes and block the entry of Ebola virus through late endosomes	Bishop, 2014; Johansen et al., 2013; Kaniklidis, 2014; Lai et al., 2014; Shoemaker et al., 2013
	Toremifene*^	Breast cancer		
	Amiodarone	Anti-arrhythmics		
	Dronedarone	Anti-arrhythmics		
	Verapamil	Anti- hypertension	Inhibits virus entry	Bishop, 2014; Johansen et al., 2013; Kouznetsova et al., 2014
	Clomiphene*	Ovulation reduction		
Anti-malaria	Toremifene*	Breast cancer		
	Chloroquine*	Anti-malaria	It prevents acid dependent cleavage of Ebola virus GP ₁ by endosomal proteases. Attacks Ebola virus VP35	Bishop, 2014; Ekins et al., 2015; Kouznetsova et al., 2014; Lai et al., 2014`
	Amodiaquine*	Anti-malaria	Attacks Ebola virus VP35	Ekins et al., 2015; Kouznetsova et al., 2014; Lai et al., 2014

Antiviral	Lamivudine	Antiviral drug approved for the treatment of HIV and hepatitis B virus		Bishop, 2014
	Ribavirin	Broad spectrum antiviral agent. It is a synthesized guanosine analog that could inhibit RNA dependent RNA polymerase used for the treatment of hepatitis C and several other viral hemorrhagic diseases		Feldman and Geisbert, 2011; Gupta et al., 2001; Huggins et al., 1999; Jahrling et al., 1999; Mupapa et al 1999; Ying et al., 2014; Zhong et al., 2014
Statins		A class of drugs used to lower cholesterol levels	Anti-inflammatory and anti immuno-modulatory agent; positive effect on coagulation	Bishop, 2014; Kaniklidis, 2014 Fedson, 2015
Vitamins	Vitamin P /Rutin	Used to treat Crimean-Congo hemorrhagic fever	Protect barrier integrity via its ability to inhibit adhesion of viral particles	Kaniklidis, 2014;Zhong et al., 2014
	Vitamin C	Used to treat Crimean-Congo hemorrhagic fever	Prevent endothelial dysfunction	Kaniklidis, 2014; Zhong et al., 2014
	Retinazone*	Antiviral agent		Kezel et al., 2013
Na ⁺ /K ⁺ -ATPase pump inhibitors	Ouabain, digoxin, digitoxin,	Cardiac glycosides	Inhibit viral replication	Lai et al., 2014
Na ⁺ /K ⁺ exchangers	Amiloride& its derivatives	Used as potassium sparing diuretics to treat hypertension and congestive heart failure	It inhibits entry and infection of Ebola virus. It inhibit RNA synthesis	Lai et al., 2014
Antioxidants	N-acetyl-cysteine	Pregnancy drug	Possibly manage Ebola virus induced cytokine dysregulation	Lai et al., 2014
Other approved drugs	Melatonin*	An hormone used to cure sleeping disorders	Anti-inflammatory and anti-coagulant , free radical scavenger	Kaniklidis, 2014; Tan et al., 2014; Tunali et al., 2005; Wirtz et al., 2008a, 2008b
	Nocodazole*	Widely-used as a cell cycle synchronizing agent to induce mitotic arrest		Kouznetsova et al., 2014
	Tamoxifen*	used to treat some types of breast cancer		
	Cepharanthine*	used for the treatment of many acute and chronic diseases. Antitumor agent		
	Zoloft (Sertraline)*	used to treat depression, obsessive-compulsive disorder, panic disorder, anxiety/stress disorders		
	Raloxifene*^	Treats and prevents osteoporosis (weak or thin bones) in women who are past menopause.		
	Nilotinib*	used to treat certain types of		

		chronic myeloid leukemia		
	Imipramine*	Treats depression		

* tested for anti-Ebola virus activity (Kouznetsova et al., 2014)

^ also called Selective Estrogen Receptor Modulators (SERMS)

Conclusion

Ebola virus has now become a major threat to global peace and security. Ebola virus was discovered in South Sudan and DRC in 1976 and since then, over 20 outbreaks have occurred in nearly 40 years, which translates to an average of one outbreak every 2 years. The 2014 EVD outbreak is unique in several ways because it is the largest outbreak thus far, and even larger than all the other outbreaks combined. Most of the previous outbreaks occurred mostly in rural areas in East and Central Africa. But the current outbreak occurred in urban areas in West Africa, reaching Lagos, a mega city with population of about 21 million people. So far, there are no licensed drugs or preventive vaccines for the treatment of EVD. Hence, response is limited to the provision of supportive care, contact tracing, barrier nursing and management of attendant complications such as blood pressure and opportunistic infections. Recent research using both small animal and non-human primate models has identified some promising therapeutics against the dreaded Ebola virus. This paper reviewed the progress made thus far in the development of drugs against EVD. Some of the promising therapeutics includes ZMapp, TKM-Ebola, Favipiravir (T-705 or Avigan), AVI 6002, BCX 4430 and Brincidofovir (CMX-001). The current development status of these first generational experimental drugs is presented. These drugs, which have commenced phase 1 clinical trials, were also used on compassionate grounds for the treatment of limited number of healthcare workers that were infected during the 2014 EVD outbreak.

References

1. ADI, 2014. Ebola and Marburg virus vaccine ELISA Kits, recombinant protein and antibodies. pp.1 -8. Alpha Diagnostic Lab Inc, USA.
2. Almansa, R., Eiros, J.M., Fedson, D., Bermejo-Martin, J.F., 2014. Hyper immune serum from healthy vaccinated individuals for Ebola virus disease?. The Lancet. 2,e686.
3. Arias, A., Thorne, L., Goodfellow, I., 2014. Favipiravir elicits antiviral mutagenesis during virus replication in vivo. eLife. 3, e03679.
4. Basler, C.F., Amarasinghe, G.K., 2009. Evasion of interferon responses by Ebola and Marburg Viruses. J. Interferon Cytokine Res. 29(9), 511 – 520.
5. Bausch, D.G., 2014. One step closer to an Ebola virus vaccine. The New England J. Med. Editorial.
6. Beeching, N.J., Fenech, M., Houlihan, C.F., 2014. Ebola virus disease. BMJ. 349,g7348.
7. Bente, D., Gren, J., Strong, J.E., Feldmann, H., 2009. Disease modelling for Ebola and Marburg viruses. Disease Models and Mechanisms, 2, 12 – 17.
8. Bishop, B.M., 2014. Potential and emerging treatment options for Ebola virus disease. Annals of Pharmacotherapy. 1 - 11.
9. Butler, D., 2014. Ebola drug trials set to begin amid crises. Nature. 513, 13 – 14.
10. Caroline, A.L., Powell, D.S., Bethel, L.M., et al., 2014. Broad Spectrum Antiviral Activity of Favipiravir (T-705): Protection from Highly Lethal Inhalational Rift Valley Fever. PLoS Negl. Trop. Dis. 8 (4), e2790.

11. Dye, J.M., Herbert, A.S., Kuehne, A.I., et al., 2012. Postexposure antibody prophylaxis protects nonhuman primates from filovirus disease. *PNAS*. 109(13), 5034 – 5039.
12. Ekins, S., Freundlich, J.S., Coffee, M., 2015. A common feature pharmacophore for FDA – approved drugs inhibiting the Ebola virus. *F1000Research*. 3, 277.
13. Elechiguerra, J.L., Burt, J.L., Morones, J.R., et al., 2005. Interaction of silver nanoparticles with HIV-1. *J. Nanobiotech*, 3, 6.
14. Emond, R.T.D., Evans, B., Bowen, E.T.W., et al., 1977. A case of Ebola virus infection. *Brit. Medical J.* 2, 541 – 544.
15. Enterlein, S., Warfield, K.L., Swenson, D.L., et al., 2006. VP 35 knockdown inhibits Ebola virus amplification and protects against lethal infection in mice. *Antimicrob. Agent Chemother.* 50(3), 984 – 993.
16. Fasina, F.O., Shittu, A., Lazarus, D., et al., 2014. Transmission dynamics and control of Ebola virus disease outbreak in Nigeria, July to September 2014. *Eurosurveillance*: 19 (3), 1-7
17. Feldmann, H., Geisbert, T., 2011. Ebola haemorrhagic fever. *Lancet*, 377, 849 – 862.
18. Fedson, D., 2015. A Practical Treatment for Patients with Ebola Virus Disease. *J Infect Dis.* 211 (4): 661-662
19. Furuta, Y., Gowen, B.B., Takahashi, K., et al., 2013. Favipiravir (T-705), a novel viral RNA polymerase inhibitor. *Antiviral Res.* 100(2), 1 – 23.
20. Gatherer, D., 2014. The 2014 Ebola virus disease outbreak in West Africa. *J. Gen.Virol.* 95 (Pt 8),1619-24. doi: 10.1099/vir.0.067199-0.
21. Geisbert, T.W., 2014. Ebola therapy protects severely ill monkeys. *Nature*. 1 – 2.
22. Geisbert, T.W., Feldmann, H.,2011. Recombinant vesicular stomatitis virus-based vaccines against Ebola and Marburg virus infections. *The J. Infect. Dis.* 204, S1075 – S1081.
23. Geisbert, T.W., Hensley, L.E., Jahrling, P.B., et al., 2003. Treatment of Ebola virus infection with a recombinant inhibitor of factor VIIa/tissue factor: a study in rhesus monkeys. *The Lancet*. 362, 1953
24. Geisbert, T.W., Hensley, L.E., Kagan, E., et al., 2007. Post exposure protection of guinea pigs against a lethal Ebola virus challenge is conferred by RNA interference. *J. Infect. Dis.* 193, 1650 – 1657.
25. Geisbert, T.W., Hensley, L.E., Kagan, E., et al., 2006. Post exposure Protection of Guinea Pigs against a Lethal Ebola Virus Challenge Is Conferred by RNA Interference. *The J. Infect. Dis.* 193 (12), 1650 – 1657.
26. Geisbert, T.W., Lee, A.C.H., Robbins, M., et al., 2010. Post exposure protection of non-human primates against a lethal Ebola virus challenge with RNA interference: a proof – of –concept study. *The Lancet*, 375, 1896 – 1905.
27. Goeijenbier, M., van Kampen, J.J.A., Reusken, C.B.E.M., et al., 2014. Ebola virus disease: a review on epidemiology, symptoms, treatment and pathogenesis. *The J. Med.* 72(9), 442 – 448.
28. Gulland, A., 2014. First Ebola treatment is approved by WHO, *BMJ*, 349, g5539.
29. Gupta, M., Mahanty, S., Bray, M., et al., 2001. Passive transfer of antibodies protects immunocompetent and immunodeficient mice against lethal Ebola virus infection without complete inhibition of viral replication. *J. Virol.* 75(10), 4649 – 4654.
30. Hampton, T., 2014. Largest-Ever outbreak of Ebola virus Disease thrust experimental therapies, vaccines into spotlight. *JAMA*, 312(10): 987 – 989
31. Heald, A.E., Iversen, P.L., Saoud, J.B., et al., 2014. Safety and pharmacokinetic profile of phosphorodiamidatemorpholino oligomers with activity against Ebola virus and Marburg virus: Results and two single-ascending- dose studies. *Antimicrob. Agent Chemother.* 58(11), 6639– 6647.
32. Hensley, L.E., Stevens, E.L., Yan, S.B., 2007. Recombinant human activated protein C for the postexposure treatment of Ebola Hemorrhagic fever. *The J. Infect. Dis.* 196, S390 – S399.

33. Hoenen, T., Feldmann, H., 2014. Ebolavirus in West Africa, and the use of experimental therapies or vaccines. *BMC Biology*, 12, 80.
34. Huggins, J., Zhang, Z-X., Bray, M., 1999. Antiviral drug therapy of filovirus infections: S-Adenosylhomocysteine hydrolase inhibitors inhibit Ebola virus In vitro and in a lethal mouse model. *The J. Infect. Dis.* 179, S240 - S247.
35. IDCRP., 2014. References summary for Ebola therapeutics and vaccines. Ver. 2. Infectious Disease Clinical Research Program.
36. Iversen, P.L., Warren, T.K., Wells, J.B., et al., 2012. Discovery and early development of AVI-7537 and AVI-7288 for the treatment of Ebola virus and Marburg virus infections. *Viruses*, 4, 2806 – 2830.
37. Iwu, M.M., Duncan, A.R., Okunji, C.O., 1999. New antimicrobials of plant origin. Janick, J. (Ed.). *Perspectives of new crops and new uses*. ASHS Press, Alexandria, VA. Pp 457 – 462.
38. Jahrling, P.B., Geisbert, J., Swearingen, J.R., et al. 1996. Passive immunization of Ebola virus infected cynomolgus monkeys with immunoglobulin from hyperimmune horse. *Arch. Virol.* 11, 135 – 140.
39. Jahrling, P.B., Geisbert, T.W., Geisbert, J.B., et al., 1999. Evaluation of immune globuline and recombinant interferon- α 2b for treatment of experimental Ebola virus infections. *The J. Infect. Dis.* 179, S224 – S234.
40. Jahrling, P.B., Geisbert, J.B., Swearingen, R.T., et al., 2007. Ebola hemorrhagic fever: evaluation of passive immunotherapy in nonhuman primates. *J. Infect. Dis.* 196, S400 – S403.
41. Ji, X., Olinger, G.G., Aris, S., et al., 2005. Mannose-binding lectin binds to Ebola and Marburg envelope glycoproteins, resulting in blocking of virus interaction with DC-SIGN and complement-mediated virus neutralization. *J. Gen. Virol.* 86, 2535 – 2542.
42. Johansen, L.M., Brannam, J.M., Delos, S.E., et al., 2013. FDA-approved selective estrogen receptor modulators inhibit Ebola virus infection. *Sci. Trans. Med.* 5 (190), 190ra79.
43. Kaniklidis, C., 2014. Ebola virus disease (EVD): new understandings and new frontiers in therapeutics. [https://www.academia.edu/9160716/Ebola_Virus_Disease_EVD_New_Understandings_and_Ne
w_Frontiers_in_Therapeutics](https://www.academia.edu/9160716/Ebola_Virus_Disease_EVD_New_Understandings_and_New_Frontiers_in_Therapeutics). Accessed 23 Feb 2015
44. Kesel, A.J., Huang, Z., Murray, M.G., et al., 2013. Retinazone inhibits certain blood-borne human viruses including Ebola virus Zaire. *Antiviral Chem. Chemother.* 23, 197 – 215.
45. Kondratowicz, A.S., Maury, W.J., 2012. Ebola virus: a brief review of novel therapeutic targets. *Future Microbiol.* 7(1), 1 – 4.
46. Kouznetsova, J., Sun, E., Martinez-Romero, C., et al., 2014. Identification of 53 compounds that block Ebola virus-like particles entry via a repurposing screen of approved drugs. *Emerging Microbes Infect.* 3, e84.
47. Kudoyarova-Zubavichene, N.M., Sergeyev, N.N., Chepurnov, A.A., et al., 1999. Preparation and use of hyperimmune serum for prophylaxis and therapy of Ebola virus infection. *The J. Infect. Dis.* 179, S218 -223.
48. Kupferschmidt, K., Cohen, J., 2015. 'Positive' results for Ebola drug upsets plans for trials. *Science Insider*. 10 February 2015.
49. Kurreck, J., 2003. Antisense technologies; improvement through novel chemical modifications. *Euro. J. Biochem.* 270, 1628 – 1644.
50. Lai, K.Y., Ng, G.W.Y., Cheng, F., 2014. Human Ebola virus infection in West Africa: a review of available therapeutic agents that target different steps of the life cycle of Ebola virus. *Infect. Dis. Poverty*. 3, 43 – 60.
51. Leung, D.W., Prins, K.C., Baster, C.F., 2010. Ebolavirus VP35 is a multifunctional virulence factor. *Virulence*, 1(6), 526 – 531.

52. Marston, H.D., Folkers, G.K., Morens, D.M., et al., 2014. Emerging viral disease: confronting threats with new technologies. *Sci. Trans. Med.* 6(253), 253ps10.
53. Martinez, O., Tantral, L., Mulherkar, N., et al., 2011. Impact of Ebola mucin-like domain on antiglycoprotein antibody responses induced by Ebola virus-like particles. *J. Infect. Dis.* 204 (Suppl 3), S825-832. doi: 10.1093/infdis/jir295.
54. Maruyama, T., Pareen, P.W.H.I., Sanchez, A., et al., 1999. Recombinant human monoclonal antibodies to Ebola virus. *The J. Infect. Dis.* 179, S235 – S239.
55. Marzi, A., Yoshida, R., Miyamoto, H., et al., 2012. Protective efficacy of neutralizing monoclonal antibodies in a nonhuman primate model of Ebola Hemorrhagic fever. *PLoS One*, 7(4), e36192.
56. Masters-Israilov, A., Pandi-Perumal, S.R., Seixas, A., et al., 2015. Melatonin, the Hormone of Darkness: From Sleep Promotion to Ebola Treatment. *Brain Disorders Therapy*, 4,1-4
57. Mentré, F., Taburet, A-M., Guedj, J., et al., 2015. Dose regimen of favipiravir for Ebola virus disease. *Lancet* 15 (2), 150–151.
58. McCarthy, M., 2014. US signs contract with ZMapp maker to accelerate development of Ebola drug. *BMJ*, 349, g5488.
59. Michelow, I.C., Lear, C., Scully, C., et al., 2011. High-dose mannose-binding lectin therapy for Ebola virus infection. *The J. Infect. Dis.* 203, 175- 179.
60. Mupapa, K., Massamba, M., Kibadi, K., et al., 1999. Treatment of Ebola hemorrhagic fever with blood transfusion from convalescent patients. *The J. Infect. Dis.* 179, S18 – S23.
61. Norwegian Institute of Public Health 2014. Discussion note for the WHO urgent Ebola vaccine access meeting.
62. Oestereich, L., Ludtke, A., Wurr, S., et al., 2014. Successful treatment of advanced Ebola virus infection with T-705 (favipiravir) in a small animal model. *Antiviral Res.* 105, 17 – 21.
63. Ohimain, E.I., 2015. How the spread of Ebola virus was curtailed in Nigeria. *Inter. J. Med. Pharm Case Reports*. In press
64. Olinger, G.G. (Jr.), Pettitt, J., Kim, D., et al., 2012. Delayed treatment of Ebola virus infection with plant-derived monoclonal antibodies provides protection in rhesus macaques. *PNA*, 109(44), 18030 – 18035.
65. Onwuakor, C.E. 2014. Ebola outbreak in Africa: Current issues- A mini review. *Amer. J. Med. Case Reports*, 2(9), 184 – 186.
66. Oswald, W.B., Geisbert, T.W., Davis, K.J., et al., 2007. Neutralizing antibody fails to impact the course of Ebola virus infection in Monkeys. *PLoS Pathogens*. 3(1), 0062 – 0066.
67. Parren, P.W.H.I., Geisbert, T.W., Maruyama, T., et al., 2002. Pre- and post exposure prophylaxis of Ebola virus infection in an animal model by passive transfer of a neutralizing human antibody. *J.Virol.* 76(12), 6408 – 6412.
68. Peters, C.J., Feldmann, S.H., Rollin, P.E., et al., 1994. Filoviruses as emerging pathogens. *Virol.* 5,147 – 154.
69. Pettitt, J., Zeitlin, L., Kim, Do. H., et al., 2013. Therapeutic intervention of Ebola virus infection in rhesus macaques with the MB-003 monoclonal antibody cocktail. *Sci. Transl. Med.* 5,199ra113.
70. Qiu, X., Wong, G., Fernando, L., et al., 2013a. mAbs and Ad-vectored IFN- α therapy rescue Ebola-infected nonhuman primates when administered after the detection of viremia and symptoms. *Sci. Transl. Med.* 5(207), 207ra143.
71. Qiu, X., Audet, J., Wong, G., et al., 2013b. Sustained protection against Ebola virus infection following treatment of infected nonhuman primates with ZMAb. *Scientific Reports*. 3, 3365.
72. Qiu, X., Audet, J., Wong, G., et al., 2012a. Successful treatment of Ebola virus – infected cynomolgus macaques with monoclonal antibodies. *Sci. Transl. Med.* 4 (138), 138ra81.

73. Qiu, K., Fernando, L., Melito, P. L., Audet, J., Feldmann, H., Kobinger, G., Alimonti, J. B. and Jones, S. [M. 2012b](#). Ebola GP-Specific Monoclonal Antibodies Protect Mice and Guinea Pigs from Lethal Ebola Virus Infection. *PLoS Negl Trop Dis*. 6(3): e1575.
74. Qiu, X., Wong, G., Audet, J., et al., 2014. Reversion of advanced Ebola virus disease in nonhuman primates with ZMapp. *Nature*, in press.
75. Rasmussen, A.L., Okumura, A., Ferris, M.T., et al., 2014. Host genetic diversity enables Ebola hemorrhagic fever pathogenesis and resistance. *Science*. 346 (6212), 987 – 991.
76. Reid, S. P., Leung, L. W., Hartman, A. L., Martinez, O., Shaw, M.L., Carbonnelle, C., Volchkov, V. E., Nichol, S. T., Basler, C. F. 2006. Ebola virus VP24 binds karyopherin alpha1 and blocks STAT1 nuclear accumulation. *J Virol*. 80(11):5156-67.
77. Richardson, J.S., Dekker, J.D., Croyle, M.A., et al., 2010. Recent advances in Ebolavirus vaccines development. *Human Vaccines*. 6(6), 439 – 449.
78. Rogers, J.V., Parkinson, C.V., Choi, Y.W., et al., 2008. A preliminary assessment of silver nanoparticles inhibition of [monkey pox](#) virus plaque formation. *Nanoscale Res. Lett*. 3, 129 – 133.
79. Sadek, R.F., Khan, A.S., Stevens, G., et al., 1999. Ebola hemorrhagic fever, Democratic Republic of the Congo, 1995: Determinants of survival. *The J. Infect. Dis*. 179, S24 – S27.
80. Sarwar, W.W., Costner, P., Enama, M.E., et al., (2014). Safety and immunogenicity of DNA vaccines encoding Ebola virus and Marburg wild-type glycoproteins in a phase I clinical trial. *J. Infect. Dis*. 1 – 9.
81. Shoemaker, C.J., Schornberg, K.L., Delos, S.E., et al., 2013. Multiple cationic amphiphiles induce a Niemann-pick C phenotype and inhibit Ebola virus entry and infection. *Plos One*, [8\(2\), e56265](#).
82. Shuaib, F., Gunala, R., Musa, E.O., et al., [2014](#). Ebola virus disease outbreak-Nigeria, July – September 2014. *Morbidity and mortality Weekly Report (MMWR)*, 63 (39), 867-872
83. Smith, L.M., Hensley, L.E., Geisbert, T.W., et al., 2013. Interferon- β therapy prolongs survival in rhesus macaque models of Ebola and Marburg Hemorrhagic fever. *The J. Infect. Dis*. 208: 310 – 218.
84. Smither, S.J., Eastaugh, L.S., Steward, J.A., et al., 2014. Post-exposure efficacy of oral T-705 (favipiravir) against inhalational Ebola virus infection in a mouse model. *Antiviral Res*, 105, 153 – 155.
85. Sobarzo, A., Perelman, E., Groseth, A., et al., 2012. Profiling the native specific human humoral immune response to Sudan Ebola virus strain gulu by chemiluminescence enzyme-linked immunosorbent assay. *Clin. Vaccine Immunol*. 19(11), 1844 – 1852.
86. Sobarzo, A., Groseth, A., Dolnik, O., et al., 2013. Profile and persistence of the virus-specific neutralizing humoral immune response in human survivors of Sudan Ebolavirus (Gulu). *The J. Infect. Dis*. 208, 299 – 309.
87. Speshock, J.L., Murdock, R.C., Braydich-Stolle, L.K., et al., 2010. Interaction of silver nanoparticles with Tacaribe virus. *J. Nanobiotechnol*. 8:19.
88. Speshock, J.L., Hussain, S., 2009. Novel Nanotechnology-Based Antiviral Agents: Silver nanoparticle neutralization of hemorrhagic fever viruses. *Defense Threat Regulatory Agency*. US Department of Defense.
89. Sullivan, N., Yang, Z-Y., Nabel, G.J., [2003](#). Ebola virus pathogenesis: implications for vaccines and therapies. *J. Virol*. 77(18), 9733 – 9737.
90. Swenson, D.L., Warfield, K.L., Warren, T.K., et al., 2009. [Antimicrob. Agents Chemother](#). 53(2), 2089 – 2099.
91. Tan, D-X., Korkmaz, A., Reiter, R.J., et al., 2014. Ebola virus disease: potential use of melatonin as a treatment. *J. Pineal. Res*. 57, 381 – 384.

92. The Economist 2015. Ebola graphics: the toll of tragedy. The Economist 1st January 2015.
93. Thi, E.P., Mire, C.E., Ursic-Bedoya, R., et al., 2014. Marburg virus unfection in nonhuman primates: Therapeutic treatment by lipid-encapsulated siRNA. *Sci. Tranl. Med.* 6(250), 250ra116.
94. Tunali, T., Sener, G., Yarat, A., et al., 2005. Melatonin reduces oxidative damage to skin and normalizes blood coagulation in a rat model of thermal injury. *Life Sci.* 76, 1259 – 1265.
95. Tyagi, P.K., Tyagi, S., Verma, C., et al., 2013. Estimation of toxic effects of chemically and biologically synthesized silver nanoparticles on human gut microflora containing *Bacillus subtilis*. *J. Toxicol. Environ. Health Sci.* 5(9), 172 – 177.
96. Uchino, K., Ochiya, T., Takeshita, F., 2013. RNAi therapeutics and applications of microRNAs in cancer treatment. *Jap. J. Clin. Oncology.* 43(6), 596 – 607.
97. Ursic-Bedoya, R., Mire, C.E., Robbins, M., et al., 2014. Protection against lethal Marburg virus infection mediated by lipid encapsulated small interfering RNA. *The J. Infect. Dis.* 209, 562 – 570.
98. Vignuzzi, M., Stone, J.K., Andino, R., 2005. Ribavirin and lethal mutagenesis of poliovirus: molecular mechanisms, resistance and biological implications. *Virus Res.* 107(2),173-181.
99. Warfield, K.L., Swenson, D.L., Olinger, G.G., et al., 2006. Gene – specific countermeasures against Ebola virus based on antisense phosphorodiamidatemorpholino oligomers. *Plos Pathog.* 2(1), 0005 – 0013.
100. Warren, T.K., Warfield, K.L., Wells, J., et al., 2010. Advanced antisense therapies for postexposure protection against lethal filovirus infections. *Nat. Med.* 16(9), 991-994.
101. Warren, T.K., Wells, J., Panchal, R.G., et al., 2014. Protection against filovirus diseases by a novel broad –spectrum nucleoside analogue BCX4430. *Nature*, 508, 402 – 405.
102. Warren, T.K., Whitehouse, C.A., Wells, J., 2015. A single phosphorodiamidatemorpholino oligomer targeting VP24 protects rhesus monkeys against lethal Ebola virus infection. *mBio* 6(1):e02344-14. doi:10.1128/mBio.02344-14.
103. Watanabe, S., Noda, T., Halfmann, P., et al., 2007. Ebola virus (EBOV) VP24 inhibits transcription and replication of the EBOV genome. *The J. Infect. Dis.* 196, 284 – S290.
104. WHO 2014a. Ebola virus disease (EVD), implications of introduction in the Americas. World Health Organization, Rome, Italy.
105. WHO 2014b. Frequently asked question on Ebola virus disease. World Health Organization, Rome, Italy.
106. WHO 2014c. Background document potential Ebola therapies and vaccines. World Health Organization, Rome, Italy
107. WHO 2014d. Potential Ebola therapies and vaccines. World Health Organization, Rome, Italy.
108. WHO 2014e. New therapeutics and preventive medicines to fight the Ebola epidemic.
109. WHO Blood Regulatory Networks (BRN) 2014. Position paper on collection and use of convalescent plasma or serum as an element in filovirus outbreak response. World Health Organization, Rome, Italy
110. WHO Ebola Response Team, 2014. Ebola virus disease in West Africa-The first 9 months of the epidemic and forward projections. *The New Engl. J. Med.* 1 - 14
111. Wirt, P.H., Bartschi, C., Spillmann, M., et al., 2008. Effect of oral melatonin on the procoagulant response to acute psychosocial stress in healthy in healthy men: a randomized placebo-controlled study. *J. Pineal. Res.* 44, 358 – 365.
112. Wirtz, P.H., Spillman, M., Bartschi, C., et al., 2008. Oral melatonin reduces blood coagulation activity: a placebo-controlled study in healthy young men. *J. Pineal. Res.* 44,127–133.

- 743 113. Yelle, J., 2014. The 2014 Ebola virus outbreak in West Africa: current perspectives for
744 prevention and treatment. *J. Human Virol. Retrovirol.* 1(4), 00024.
- 745 114. Ying, Z., Jun, X.U., TaiSheng, L.I., et al., 2014. Potential clinical treatment for Ebola
746 pandemic. *Life Sci.* 57(10), 982 – 984.
- 747 115. Zhang, Z-Q., Higgins, J., Kim, C.L., et al., 2013. Synthesis and antimicrobial properties
748 of nanosilver, *J. Mater. Environ. Sci.* 4(1), 139 – 142.
- 749 116. Zhong, Y., Xu, J., Li, T., et al. 2014. Potential clinical treatment for Ebola pandemic. *Sci.*
750 *China Life Sci.* 57 (10), 982–984.