

Original Research Article
Asymptomatic *Blastocystis*-infected
as disease carriers in Thailand

ABSTRACT

Aim: *Blastocystis* spp. is currently classified as a gastrointestinal protozoan parasite and has become quite a controversial organism. This study was aimed to obtain the prevalence of *Blastocystis* infection among an otherwise healthy, asymptomatic Thai population.

Study Design: Retrospective study

Place and Duration of study: The study areas were the Border Patrol-policemen (Subdivision 12), Sa Kaeo Province; the Home for Mentally Handicapped Childcare, Ban Rajawadee, in Nonthaburi Province; Silpakorn University, Nakhon Pathom Province and the Intercity Motorway Division, Department of Highways, Bangkok and the study was carried during 2010-2013.

Methodology: Fecal examination by direct simple smear was performed among 3,940 out of 6,537 Thais of different genders, ages, occupations, and residential locations who joined the annual health check-up program serviced by Faculty of Tropical Medicine, Mahidol University.

Results: It was found that 1.0% (40/3,940) presented with *Blastocystis* infection and were asymptomatic of clinical symptoms at a male to female ratio of 2.3:1.0. *Blastocystis* infection was predominantly 47.5% (19/40) in the age group 41-50 years. Border Patrol-policemen (Subdivision 12), who resided and worked in rural areas of Sa Kaeo Province near the border between Thailand and Cambodia, had significantly higher prevalence of *Blastocystis* infection (2.8%) than the staff of the Intercity Motorway Division, Department of Highways, in urban Bangkok (0.1%). In addition, other occupations with work places were in suburban areas, such as workers at the Home for Mentally Handicapped Childcare, Ban Rajawadee, in Nonthaburi Province, who worked closely with mentally handicapped children, and officials and lecturers at Silpakorn University, Nakhon Pathom Province, also had significantly higher prevalent rates than Intercity Motorway Division staff in urban Bangkok, at 4.20% and 2.55%, respectively, while food handlers and cooks who prepared and sold food in Silpakorn University had similar prevalent rates of *Blastocystis* infection to Intercity Motorway Division staff.

Conclusion: This study revealed that *Blastocystis* infected patients probably did not show obviously clinical symptoms in asymptomatic carriers, who can spread infections to others and environments.

Keywords: *Blastocystis* infection, prevalence, asymptomatic, carriers

38 1. INTRODUCTION

39 *B. hominis* is common and distributed worldwide among humans and animals, particularly in
40 poor or developing countries where the prevalence is high due to poor personal hygiene, the
41 consumption of contaminated food/drinking water, exposure to animals, insufficient
42 education, and socioeconomic factors [1, 2]. Currently, *B. hominis* is classified as a
43 eukaryotic, unicellular protozoan of the gastrointestinal tract, particularly the large intestine
44 (colon and cecum), and is commonly found in the feces [3]. *B. hominis* has been changed to
45 *Blastocystis* spp., because the morphology of the organisms in humans and animals is
46 indistinguishable among species. Generally, it is strictly anaerobic, without a cell wall and
47 present in a variety of sizes (ranging 5-40 μm), with a polymorphic morphological shape,
48 from spherical to amoebic in form [4]. Considerable controversy remains as to whether
49 *Blastocystis* spp. is simply a gastrointestinal commensal organism or a potential human
50 pathogen causing gastrointestinal disease [5, 6]. Previous published papers have examined
51 blastocystiasis in association with acute or chronic diarrhea, particularly in
52 immunocompromised hosts, or even immunocompetent hosts such as children, pregnant
53 women, geriatric adults, or risky hosts, such as travelers, homosexuals, animal keepers, and
54 immune-suppressive patients [7-12]. Other symptoms probably included generalized
55 abdominal pain, nausea, vomiting, flatulence and anorexia [13] with non-specific and varied
56 symptoms of *Blastocystis* spp., resulting in misunderstanding of its potential pathogenicity
57 [14]. However, the presence of *Blastocystis* spp. in feces sometimes did not involve
58 gastrointestinal symptoms among infected cases [15, 16].

59 In Thailand, information about *Blastocystis* infection and distribution of the disease remains
60 scanty [17]. Some reports on the prevalence of blastocystiasis varied between 0.19-45.5%
61 [18-20]. The present study investigated the prevalence of *Blastocystis* infection among
62 asymptomatic, apparently healthy Thai people in rural, suburban, and urban areas of
63 Thailand. A large number joined the program, with differing genders, ages, occupations, and
64 residential locations, and included an evaluation of laboratory findings and parameters for
65 *Blastocystis*-infected participants.

66 2. MATERIALS AND METHODS

67 2.1 Study Population

68 A retrospective study was conducted among 5 groups of apparently healthy Thais (total of
69 6,537 participants) who joined the annual health check-up program serviced by Faculty of
70 Tropical Medicine, Mahidol University, during a three-year period (2010-2013) composed of
71 2,266 Border Patrol-policemen (Subdivision 12) from Sa Kaeo Province; 1,300 staff/lecturers
72 from Silpakorn University, Nakhon Pathom Province; 397 food handlers and cooks from
73 Silpakorn University, in Nakhon Pathom Province; 258 workers at the Home for Mentally
74 Handicapped Childcare, Ban Rajawadee, Nonthaburi Province; and 2,316 staff from the

Intercity Motorway Division, Department of Highways in Bangkok. Nonthaburi and Nakhon Pathom Provinces are both in the central part of the country, about 20 and 56 kilometers from Bangkok, respectively. Sa Kaeo Province is located in Eastern Thailand, about 245 kilometers from Bangkok. All participants were requested to collect fecal samples as one of the laboratory tests for routine health check-up.

2.2 Physical Examination

All participants were asked about abnormal diarrhea (at least 3 loose feces during a 24-hour period), abdominal pain, and any underlying disease. Routine physical examinations were conducted of all physical systems, particularly the gastrointestinal system. The height and weight of each participant were measured to determine nutritional status and obesity, compared with standard body-mass indices (BMI).

2.3 Fecal Examination

A total of 3,940 fecal samples from participants who enrolled in annual health check-up programs were examined to identify *Blastocystis* spp. by routinely direct simple smear in 0.85% normal saline solution and 1% Lugol's iodine solution for double examination to increase yields of detection.

2.4 Laboratory Investigations

Basic laboratory tests-complete blood count, urinalysis, chest X-ray, liver function test: aspartate transaminase (AST), alanine transaminase (ALT), alkaline phosphatase (ALP)-were performed for any remarkable changes involving blastocystiasis.

3. RESULTS

A total of 3,940 of 6,537 apparently healthy Thai people (60.3%) from various areas of Thailand were recruited into the study, people residing and working in urban Bangkok, suburban Nonthaburi and Nakhon Pathom Provinces, and rural Sa Kaeo Province (Fig. 1). The participants who were all adults (age range 26-59 years: (mean 42), joined the annual health check-up program serviced by Faculty of Tropical Medicine, Mahidol University during a 3-year period, and provided their feces for examination (Table 1). Only 40 of 3,940 (1.0%) were found to have the vacuolar form of *Blastocystis* spp. in their feces, by 0.85% normal saline solution and 1% Lugol's iodine solution (Fig. 2): no obvious clinical signs were noted in the physical examinations (male:female ratio, 2.3:1.0).. The enrolled participants were grouped into 5 groups by gender, age, occupations, and residential location, with significant difference in prevalence ($p < 0.001$; $\chi^2 = 66.3$). Most cases of blastocystiasis (47.5%: 19/40) were found among the age group 41-50 years. There were significant differences among the infected participants by occupations and residential location. Border Patrol-policemen (Subdivision 12) who had resided and worked in rural areas of Sa Kaeo Province, workers in the Home for Mentally Handicapped Childcare, Ban Rajawadee, in Nonthaburi Province, and staff/lecturers at Silpakorn University, Nakhon Pathom Province,

had significantly higher prevalent rates than those who had resided and worked in the Intercity Motorway Division Department of Highways in Bangkok ($P<0.001$: Table 1). However, food handlers and cooks at Silpakorn University showed no significant difference in *Blastocystis* infection rates from those who resided and worked in Bangkok ($P>0.001$).

Physical examinations found no obvious abnormalities. Initial laboratory tests-white blood cell count, hematocrit value, eosinophil count, SGOT, SGPT, ALP, and urinalysis-were analyzed (Table 2). Chest x-rays were unremarkable. The presence of *Blastocystis* infection was analyzed in relation to nutritional status according to body mass index (BMI) classified by WHO (2000) (Table 3): no nutritional involvement was found, since all participants were categorized as normal to overweight except only one female participant was underweight.

4. DISCUSSION

The prevalence of *Blastocystis* spp. varied among countries, and even within the same country, according to the population, targeted, area studied, and diagnostic tool used [21]. Among 6,537 apparently healthy Thai people, 2,266 (34.7%) came from rural areas in Sa Kaeo Province, while 4,271 (65.3%) resided in urban Bangkok and suburban Nonthaburi and Nakhon Pathom Provinces. Overall, *Blastocystis* spp. were detected in 40 fecal samples (male:female ratio, 2.3:1.0) from a total of 3,940 apparently healthy Thai people (1.0%). There was a significantly higher prevalence of *Blastocystis* infection among men who have sex with men (MSM) [22]. In this study, the vacuolar form of *Blastocystis* spp. was the only stage found in the fecal examinations, which was similar to the previous studies [12, 23]. The prevalence of *Blastocystis* infection was rather low because there were remaining 2,597 (39.7%) undiagnosed cases as fecal samples were not collected for routinely fecal examination. While, fecal examination using the direct simple smear technique was a simple and efficient tool for screening for intestinal parasites [24], it was often ignored because of the inconvenience of collecting feces, accompanied by unpleasant and embarrassed sensitivities. Hence, awareness should be raised about the importance of fecal examination to prevent and control parasitological diseases, and so that special attention is given to fecal analysis for pathogens that can easily spread disease to others by disease carriers. *Blastocystis* spp. is truly an enteropathogen as it sometimes presents in symptomatic patients. Although, there is sometimes no treatment, with recovery and improvement from self-limiting watery diarrhea, most infected patients are asymptomatic carriers [4, 25]. The different genotypes of *B. hominis* obtained from patients with IBS had insignificantly different pathogenic potential for degree of inflammation in sigmoidoscopic changes. However, severe inflammation mostly predominated among patients harboring genotype I isolates, while genotype II was not detected among symptomatic infected individuals. Those harboring genotypes III and IV exhibited both pathogenic and nonpathogenic strains [26].

There is general agreement among proponents from both sides that *Blastocystis* spp. is very frequently found in the feces of cases without GI symptoms [5]. However, this does not rule out the possibility that, under certain circumstances, infection with large numbers of *Blastocystis* spp. could also cause severe diarrhea. Hence, the role *Blastocystis* spp. as a

significant pathogen remains unelucidated, since no morphologic or physiologic evidence of pathology has been attributed to the organism, such as invaded tissues or toxins. It is somewhat of a challenge for clinicians to treat patients with diarrhea in who are *Blastocystis*-positive by fecal examinations. Metronidazole is the most effective drug for *Blastocystis* infection and is recommended in the same dose as used for treatment of amebiasis [27, 28]. However, another paper classified *Blastocystis* spp as the common nonpathogenic organism found in feces [29]. In this study, *Blastocystis*-positive individuals showed no obvious clinical signs or symptoms, and they did not correspond with the intensity of organisms found in feces: this was similar to a report by Horiki *et al.* (1997) [30]. In another study, no significant association was found between *Blastocystis* subtype and clinical features [31]. This findings supported the contention that *Blastocystis* spp. was not a serious pathogen and rarely caused clinical symptoms, since no invasion of host tissues was detected. Some reports showed that only a few specific *Blastocystis* subtypes of caused symptoms [32]. Some studies suggested that pathogenicity was related to a few specific subtypes as well as density of *Blastocystis* infection [7, 28, 32].

Each year, healthy Thai people are exposed to *Blastocystis* infection during their daily lives, by consuming contaminated food or water [18, 33, 34]. Not all Thais undergo systematic or routine health check-ups, which frequently depend on the health-welfare service provided by each institute/organization, or personal initiative and expense. The prevalence of *Blastocystis* infection among Border Patrol-policemen (Subdivision 12) in rural areas of Sa Kaeo Province, near border between Thailand and Cambodia (2.75%), was significantly higher than among staff of the Intercity Motorway Division, Department of Highways, in urban Bangkok (0.13%), $P<0.001$. The Border Petrolpolicemen were at higher risk of infection due to greater exposure to a contaminated environment, particularly in border areas with temporary migrant accommodation and inappropriate, unhygienic sanitation, or worse, none. Similarly, a high prevalence of *Blastocystis* infection was also reported in an army base in Chonburi Province and Myanmar immigrant workers in the Thai food industry [18, 35, 36]. The cystic form of *Blastocystis* spp. was viable for more than 1 or 2 month(s) in water at 25°C or 4°C, respectively. Contamination with only a small number of cysts could cause infection [37], and being resistant to standard water chlorination [38] can contaminate water or food; this was possibly the only infectious form involved in spreading *Blastocystis* infection. The transmission route was still unclear; however, the main transmission route was likely fecal-oral with the cyst form [39] from human to human, water-borne transmission and close contact with animals [40, 41]. In addition, those who resided and worked in the Home for Mentally Handicapped Childcare, Ban Rajawadee, Nonthaburi Province, and staff/lecturers at Silpakorn University, Nakhon Pathom Province, suburban areas near Bangkok, had significant higher prevalent rates of *Blastocystis* infection, at 4.2%, and 2.6%, than the staff of the Intercity Motorway Division, Department of Highways, in Bangkok, $P<0.001$. A previous study was conducted in Pathum Thani Province, a suburban area near Bangkok with a high prevalence of *Blastocystis* infection among orphans [19]. Hence, the residential locations of participants was probably a factor involved in disease transmission, if appropriate sanitation was lacking, and many reservoir hosts and insect vectors, such as

houseflies or cockroaches, were present to act as transporters of various parasitic diseases to humans [42].

In addition to residential location, occupation was another factor related to differences in the prevalence of *Blastocystis* infection. A comparison of *Blastocystis* infection among staff/lecturers and food handlers/cooks in the same work area of Silpakorn University, Nakhon Pathom Province, showed significantly greater prevalence among the staff/lecturers group than the food handlers/cooks group. There was no significant difference between food-handlers and cooks (0.3%) and workers in the Intercity Motorway Division, Department of Highways, in Bangkok, $P < 0.001$. Food handlers and cooks were more likely to contract and transmit *Blastocystis* spp. via fecal-oral route, with a high prevalent rate of 35.5%, including the most frequent symptoms of diarrhea, abdominal pain, and flatulence. Generally, those wishing to work as food handlers and cooks need to possess a verified health certificate showing freedom from enteric pathogens. The prevalence of *Blastocystis* infection among this group was rather low, because they were strictly monitored by disease control authorities, with annual health check-ups according to the university's food-safety policy. Although, this study did not focus on the prevalence of *Blastocystis* infection among mentally retarded children, the prevalence was high among workers who took close care of them. Blastocystiasis might spread between care-givers and vulnerable children, raising concern that mentally retarded children could easily become infected since they could not take adequate care of themselves and use appropriate hygienic behaviors independently. Previous studies found a high prevalence (45.2%) of *Blastocystis* infection among children in an orphanage in Pathum Thani Province, Thailand [19]. Zoonotic transmission was a possible source of *Blastocystis* infection among girls in a girls' home in Thailand who did not wash their hands after coming into contact with pets [43].

Initial laboratory results revealed no abnormal values correlating with *Blastocystis* infection, except for raised eosinophil count, which was also symptomatic of many other parasitic infections (Table 2). Abnormal values might be related to heavy infection and obvious clinical features. Hypoalbuminemia was reportedly associated with blastocystiasis-related diarrhea [44]. Moreover, an attempt was made to use 29-kDa *B. hominis* antigen as a marker for pathogenicity and diagnosis to differentiate between symptomatic and asymptomatic blastocystiasis [45].

5. CONCLUSION

Although *Blastocystis* spp. in otherwise healthy Thai people did not appear to be serious, since clinical manifestations were not evident: they presented mostly in overweight participants of both genders (males: 50.0%; females: 41.2%) (Table 3). However, the immunodeficient or immunocompetent, such as infants, children, geriatric men, pregnant women, and immunosuppressive patients, were at higher risk of diarrhea and clinical intestinal symptoms, since their immune systems were not normal or fully developed. Effective fecal screening health check-ups, and infection-eradication strategies, need to be rigorously enforced to control and prevent pathogens that may not be evident in

asymptomatic carriers, who can spread infections to others. The improvement of personal hygiene behaviors is important for reducing the risk of *Blastocystis* infection. Information derived from other studies would help us understand the transmission of this disease. Therefore, different occupations in different residential locations would yield more information on the risk of exposure to this disease.

CONCENT

Informed consent was obtained from all the study participants.

ETHICAL APPROVAL

This study was approved by the Ethics Committee on human rights related to Research Involving Human Subjects, Mahidol University (MURA2013/239).

REFERENCES

1. Tan KS, Singh M, Yap EH. Recent advances in *Blastocystis hominis* research: hotspots in terra incognita. *International Journal for Parasitology*. 2002;32:789-804.
2. Tan KS. New insights on classification, identification, and clinical relevance of *Blastocystis* spp. *Clinical Microbiology Reviews*. 2008;21:639-665.
3. Stensvold CR, Nielsen HV, Molbak K, Smith HV. Pursuing the clinical significance of *Blastocystis*-diagnostic limitations. *Trends in Parasitology*. 2009c;25:23-29.
4. Janarthanan S, Khoury N, Antaki F. An unusual case of invasive *Blastocystis hominis* infection. *Endoscopy Journal*. 2011;43:185-186.
5. Franklin AN and Harold WB. Basic clinical parasitology. Sixth edition. Appleton & Lange, Norwalk, Connecticut. 1994. p.55.
6. Leder K, Hellard ME, Sinclair MI, Fairley CK, Wolfe R. No correlation between clinical symptoms and *Blastocystis hominis* in immunocompetent individuals. *Journal of Gastroenterology and Hepatology*. 2005;20:1390-1394.
7. Zuel-Fakkar NM, Abdel-Hameed DM, Hassanin OM. Study of *Blastocystis hominis* isolates in urticaria: a case-control study. *Clinical and Experimental Dermatology*. 2011;36:908-910.
8. Chandramathi S, Suresh K, Anita ZB, Kuppusamy UR. Infections of *Blastocystis hominis* and microsporidia in cancer patients: are they opportunistic? *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 2012;106:267-269.
9. El Deeb HK, Salah-Eldin H, Khodeer S. *Blastocystis hominis* as a contributing risk factor for development of iron deficiency anemia in pregnant women. *Parasitology Research*. 2012;110:2167-274.

- 267 10. Tutanc M, Silfeler I, Ozgur T, Motor VK, Kurtoglu AI. The case of Henoch-
268 Schonlein Purpura associated with *Blastocystis hominis*. *Turkiye Parazitoloji Dergisi*.
269 2013;37:135-138.
- 270 11. Verma R and Delfanian K. *Blastocystis hominis* associated acute urticaria. *The*
271 *American Journal of the Medical Sciences*. 2013;346:80-81.
- 272 12. Basak S, Rajurkar MN, Mallick SK. Detection of *Blastocystis hominis*: a controversial
273 human pathogen. *Parasitology Research*. 2014;113:261-265.
- 274 13. Shah M, Tan CB, Rajan D, Ahmed S, Subramani K, Rizvon K, Mustacchia P.
275 *Blastocystis hominis* and *Endolimax nana* co-infection resulting in chronic diarrhea in
276 an immunocompetent male. *Case Reports in Gastroenterology*. 2012;6:358-364.
- 277 14. Jones MS, Ganac RD, Hiser G, Hudson NR, Le A, Whipps CM. Detection of
278 *Blastocystis* from stool samples using real-time PCR. *Parasitology Research*.
279 2008;103:551-557.
- 280 15. Rostami Nejad M, Nazemalhosseini Mojarad E, Dabiri H, Nochi Z, Pourhoseingholi
281 MA, Sahebekhtari N, Habibi M, Zali MR. A case-control study of *Blastocystis*
282 *hominis* among Iranian population. *East African Journal of Public Health*.
283 2010;7:101-104.
- 284 16. Surangsirat S, Thamrongwittawatpong L, Piyaniran W, Naaglor T, Khoprasert C,
285 Taamasri P, Mungthin M, Leelayoova S. Assessment of the association between
286 *Blastocystis* infection and irritable bowel syndrome. *Journal of the Medical*
287 *Association of Thailand*. 2010;93Suppl6:119-124.
- 288 17. Pipatsatitpong D, Rangsin R, Leelayoova S, Naaglor T, Mungthin M. Incidence and
289 risk factors of *Blastocystis* infection in an orphanage in Bangkok, Thailand. *Parasites*
290 *and Vectors*. 2012;5:37.
- 291 18. Taamasri P, Mungthin M, Rangsin R, Tongupprakarn B, Areekul W, Leelayoova S.
292 Transmission of intestinal blastocystosis related to the quality of drinking water. *The*
293 *Southeast Asian Journal of Tropical Medicine and Public Health*. 2000;31:112-117.
- 294 19. Saksirisampant W, Nuchprayoon S, Wiwanikit V, Yenthakam S, Ampavasiri A.
295 Intestinal parasitic infestations among children in an orphanage in Pathum Thani
296 province. *Journal of the Medical Association of Thailand*. 2003;86:263-270.
- 297 20. Kitvatanachai S and Rhongbutsri P. Intestinal parasitic infections in suburban
298 government schools, Lak Hok subdistrict, Muang Pathum Thani, Thailand. *Asian*
299 *Pacific Journal of Tropical Medicine*. 2013;6:699-702.
- 300 21. Yaicharoen R, Ngrenngarmert W, Wongjindanon N, Sripochang S, Kiatfuengfoo R.
301 Infection of *Blastocystis hominis* in primary schoolchildren from Nakhon Pathom
302 province, Thailand. *Tropical Biomedicine*. 2006;23:117-122.

- 303 22. Stark D, Fotedar R, van Hal S, Beebe N, Marriot D, Ellis JT, et al. Prevalence of
304 enteric protozoa in human immunodeficiency virus (HIV)-positive and (HIV)-
305 negative men who have sex with men from Sydney, Australia. The American Journal
306 of Tropical Medicine and Hygiene. 2007;76:549-552.
- 307 23. Yaicharoen R, Sripochang S, Sermsart B, Pidetcha P. Prevalence of *Blastocystis*
308 *hominis* infection in asymptomatic individuals from Bangkok, Thailand. The
309 Southeast Asian Journal of Tropical Medicine and Public Health. 2005;36 Suppl4:17-
310 20.
- 311 24. Babiker MA, Ali MS, Ahmed ES. Frequency of intestinal parasites among food
312 handlers in Khartoum, Sudan. Eastern Mediterranean Health Journal. 2009;15:1098-
313 10104.
- 314 25. Sekar U and Shanthi M. Blastocystis: Consensus of treatment and controversies.
315 Tropical Parasitology. 2013;3:35-39.
- 316 26. Fouad SA, Basyoni MM, Fahmy RA, Kobaisi MH. The pathogenic role of different
317 *Blastocystis hominis* genotypes isolated from patients with irritable bowel syndrome.
318 Arab Journal of Gastroenterology. 2011;12:194-200.
- 319 27. Editorial: *Blastocystis hominis*: commensal or pathogen? The Lancet. 1991;337:521-
320 522.
- 321 28. Fathy FM. A study on *Blastocystis hominis* in food handlers: diagnosis and potential
322 pathogenicity. Journal of the Egyptian Society of Parasitology. 2011;41:433-453.
- 323 29. Koshak EA and Zakai HA. A spectrum of pathogenic and non-pathogenic intestinal
324 parasites in pre-employment medical check-up for workers and their families. Journal
325 of Family and Community Medicine. 2009;10:47-53.
- 326 30. Horiki N, Maruyama M, Fujita Y, Yonekura T, Minato S, Kaneda Y. Epidemiologic
327 survey of *Blastocystis hominis* infection in Japan. The American Journal of Tropical
328 Medicine and Hygiene. 1997;56:370-374.
- 329 31. Jantermtor S, Pinlaor P, Sawadpanich K, Pinlaor S, Sangka A, Wilailuckana C,
330 Wongsena W, Yoshikawa H. Subtype identification of *Blastocystis* spp. isolated from
331 patients in a major hospital in northeastern Thailand. Parasitology Research.
332 2013;112:1781-1786.
- 333 32. Abdel-Hameed DM and Hassanin OM. Protease activity of *Blastocystis hominis*
334 subtype3 in symptomatic and asymptomatic patients. Parasitology Research.
335 2011;109:321-327.
- 336 33. Leelayoova S, Rangsin R, Taamasri P, Naaglor T, Thathaisong U, Mungthin M.
337 Evidence of waterborne transmission of *Blastocystis hominis*. The American Journal
338 of Tropical Medicine and Hygiene. 2004;70:658-662.

- 339 34. Leelayoova S, Siripattanapipong S, Thatthaisong U, Naaglor T, Taamasri P, Piyaraj P,
340 Mungthin M. Drinking water: a possible source of *Blastocystis* spp. subtype 1
341 infection in schoolchildren of a rural community in central Thailand. The American
342 Journal of Tropical Medicine and Hygiene. 2008;79:401-406.
- 343 35. Taamasri P, Leelayoova S, Rangsin R, Naaglor T, Ketupanya A, Mungthin M.
344 Prevalence of *Blastocystis hominis* carriage in Thai army personnel based in
345 Chonburi, Thailand. Military Medicine. 2002;167:643-646.
- 346 36. Nuchprayoon S, Samprasert V, Kaewzaaithim S, Saksirisampant W. Screening for
347 intestinal parasitic infections among Myanmar migrant workers in Thai food industry:
348 a high-risk transmission. Journal of Immigrant and Minority Health. 2009;11:115-
349 121.
- 350 37. Yoshikawa H, Yoshida K, Nakajima A, Yamanari K, Iwatani S, Kimata I. Fecal-oral
351 transmission of the cyst form of *Blastocystis hominis* in rats. Parasitology Research.
352 2004;94:391-396.
- 353 38. Zaki M, Zaman V, Sheikh NA. Resistance of *Blastocystis hominis* cyst to chlorine.
354 Journal of Pakistan Medical Association. 1996;46:178-179.
- 355 39. Yoshikawa H, Abe N, Iwasawa M, Kitano S, Nagano I, Wu Z, Takabashi Y. Genomic
356 analysis of *Blastocystis hominis* strains isolated from two long-term health care
357 facilities. Journal of Clinical Microbiology. 2000;38:1324-1330.
- 358 40. Hotez P. The other intestinal protozoa: enteric infections caused by *Blastocystis*
359 *hominis*, *Entamoeba coli*, and *Dientamoeba fragilis*. Seminars in Pediatric Infectious
360 Diseases. 2000;11:178-181.
- 361 41. Salim HR, Kumar GS, Vellayan S, Mak JW, Anuar AK, Init I, et al. *Blastocystis* in
362 animal handlers. Parasitology Research. 1999;85:1032-1033.
- 363 42. Maipanich W, Sa-nguankiat S, Pubampen S, Kusolsuk T, Lekka A. Intestinal
364 parasites isolated from house-flies in the tourist attraction areas in Thailand. The
365 Journal of Tropical Medicine and Parasitology. 2010;33:17-28.
- 366 43. Thathaisong U, Worapong J, Mungthin M, Tan-Ariya P, Viputtigul K, Sudatis A, et
367 al. *Blastocystis* isolates from a pig and a horse are closely related to *Blastocystis*
368 *hominis*. Journal of Clinical Microbiology. 2003;41:967-975.
- 369 44. Laodim P, Intapan PM, Sawanyawisuth K, Prasongdee TK, Laumaunwai P,
370 Maleewong W. Hypoalbuminemia as a predictor of diarrhea caused by *Blastocystis*
371 *hominis*. The Southeast Asian Journal of Tropical Medicine and Public Health.
372 2013;44:374-378.
- 373 45. Abou Gamra MM, Elwakil HS, El Deeb HK, Khalifa KE, Abd Elhafiz HE. The
374 potential use of 29 kDa protein as a marker of pathogenicity and diagnosis of

symptomatic infections with *Blastocystis hominis*. Parasitology Research. 2011;108:1139-1146.



Fig 1 Map of residing locations of participants: urban (Bangkok), suburban (Nonthaburi and Nakhon Pathom Provinces) and rural (Sa Kaeo Province) areas




Fig 2 Blastocystis vacuolar form with peripheral nuclei in the cytoplasm detected in 0.85% normal saline (A) and 0.1% Lugol's iodine solution (B) of fecal examination

Table 1

Basic participant data and distribution of *Blastocystis* infection (n = 40 cases) in rural, suburban and urban areas of Thailand

Age groups	Border Patrol-police Subdivision 12 (Sa Kaeo)		Home for Mentally Handicapped Childcare, Ban Rajawadee (Nonthaburi)		Silpakorn University (Nakhon Pathom)				Intercity Motorway Division, Department of Highways (Bangkok)	
					Staff and lecturers		Food handlers and cooks			
	M	F	M	F	M	F	M	F	M	F
21-30	3	0	0	1	0	1	0	0	1	0
31-40	1	0	0	1	1	2	0	0	1	1
41-50	16	0	0	1	0	1	0	1	0	0
51-60	4	0	1	2	0	1	0	0	0	0
No. of infected people	24	0	1	5	1	5	0	1	2	1
	2.8%		4.2%		2.6%		0.3%		0.1%	
No. of people undergoing fecal examination	872		143		235		374		2,316	
Z-test (p-values)	-7.71		-7.59		-5.81		-1.18			

M: Male; F: Female

Table 2

Initial laboratory parameters among *Blastocystis*-infected participants

Laboratory parameters	Normal range values	Number of patients (%)		Z-test (p-values)
		within normal limit	abnormal value	
WBC count (cells/mm ³)	4,800-10,800	94.9	5.1	5.62
Eosinophils (%)	2-3%	41.0	59.0	-1.12
Hct (%)	Male: 40-52 Female: 36-48	92.3	7.7	5.25
AST (unit/mL)	5-45	94.9	5.1	5.62
ALT (unit/mL)	5-45	84.6	15.4	4.38
ALP (IU/L)	40-150	97.4	2.6	5.88

WBC: white blood cell; Hct: hematocrit; AST: aspartate aminotransferase; ALT: alanine aminotransferase; ALP: alkaline phosphatase

Table 3

Number of *Blastocystis*-infected participants and nutritional status

Nutritional status (BMI)	No. of <i>Blastocystis</i> -infected participants (%)	
	M	F
<18.50 (underweight)	0.0 (0)	5.9 (1)
18.50-22.99 (normal weight)	33.3 (6)	35.3 (6)

23.00-24.99 (borderline overweight)	16.7 (3)	17.6 (3)
≥25.00 (overweight)	50.0 (9)	41.2 (7)
Total	18	17

395 BMI: body mass index, M: Male; F: Female

396