

Original Research Article

Separation of Palmitic and Palmitoleic Acid using Transition Metal Salts: Part 2

ABSTRACT

Fatty acids (FAs) play a major role in many physiological functions and offer various health benefits. Separation and isolation of fatty acids thus is of great interest. A set of experiments were conducted to evaluate the effectiveness of select transition metal salts in separating a saturated FA from mixture of C16:0 and C16:1 FAs. In an initial study, ZnSO₄, CuSO₄ and AgCl were able to cause significant precipitation of C16:0 FA from a mixture of C16:0 and C16:1 FAs in methanol. A follow-up study was conducted to ascertain the effect of ZnSO₄ concentration on the ability to separate individual FAs from a mixture of saturated and unsaturated FAs in methanol. Results indicated that a 30 mg/ml ZnSO₄ solution could remove 58% of C16:0 FA from a mixture containing both the saturated and the unsaturated FA forms. IR spectra suggests formation of a complex between zinc ion and the carboxylate group of the C16:0 FA, which in turn may cause precipitation. Computational chemistry calculations indicate that specific complex formation with the saturated FA might be due to more favorable thermodynamics between zinc ion - C16:0 FA complexation (with more negative value of Gibbs free energy of the reaction) than that between the metal ion and the unsaturated FA.

Keywords: *Palmitic acid, Palmitoleic acid, C16:0 FAME, C16:1 FAME, copper sulphate, silver nitrate, zinc sulphate, precipitation, complex formation*

1. INTRODUCTION

Fatty acids (FAs) play an important role in various physiological functions and contribute to many health effects. Palmitic acid has been proven to mediate hypothalamic insulin resistance [1]; Palmitoleic acid has been shown to boost insulin sensitivity thus improving hyperglycemia and hypertriglyceridemia [2]. Saturated FAs (SFA) are engaged in the synthesis of phospholipids and spingolipids, storage and production of energy, and lipid transport [3]. Poly-unsaturated fatty acids (PUFAs), especially the omega-3 fatty acids Eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) is reported to provide cardiovascular health benefits when integrated into the diet [4]. Many studies omega-3 fatty acids have shown that supplementation may render preventative and therapeutic effects on certain cancers including those of the prostate, colon, rectum, and breast, among others. [4] Therefore, isolation and separation of unsaturated fatty acids is of significant interest.

Various methods have been used for separation and isolation of saturated and unsaturated fatty acids: these include chromatography techniques such as gas chromatography, thin layer chromatography, and high pressure liquid chromatography; winterization; and urea complexation.¹

Silver ion chromatography is one of the most effective techniques used in fatty acid separation. This technique uses the property of olefinic compounds forming a complex with

transition metals (especially silver ions), resulting in separation of fatty acids based on the degree of unsaturation. An accepted model predicts formation of a σ -type bond between occupied 2p orbital of an olefinic double bond and the free 5s and 5p orbitals of the silver ion. Also there is a π – acceptor backbond between occupied 4d orbital of silver ion and the free antibonding 2p π^* orbital of the olefinic bond [5]. This property to form charge-transfer type complexes with transition metal ions has been widely adapted in high-performance liquid chromatography based olefin separation [6].

The property of silver ion complexation with olefinic double bonds has also been used to extract fatty acid ester derivatives from various organic solvents [7]. To give an example, separation of ethyl esters of eicosapentaenoic acid and docosahexaenoic acid had been attempted via circulating liquid membranes using AgCl as a carrier [8]. Another example involves the use of supercritical extraction (SC) to extract polyunsaturated fatty acid ethyl esters (PUFA-Et) by aqueous AgCl [9].

Another method to separate fatty acids uses the property of particular metal ions' ability of form precipitates with certain FAs. An example includes the use of alkali metal salts in combination with alkali metal hydroxides to precipitate out free fatty acids in the form of insoluble fatty acid salts [10].

Certain metallic ions demonstrate the ability to form precipitates with specific types of fatty acids and this property has been utilized to isolate one type of fatty acid from a mixture. For example, alkali metal salts along with alkali metal hydroxides have been used to precipitate out free fatty acids via the formation of insoluble fatty acids salts [10]. In another example, saturated straight chain aliphatic monocarboxylates have been separated using the property of Cu, Fe, Zn, Pb and Mg to form complexes with the monocarboxylates [11-13].

A method widely used for large scale separation of fatty acid is urea complexation. In this method, a homogenous solution of urea and free fatty acids/saturated fatty acids is formed in alcoholic solvent at 65°C. This is followed by cooling and separation of the resultant urea complex to room temperature [14]. Studies reveal that in a mixture containing both saturated and unsaturated FA forms, by using urea complexation, the SFA concentration was decreased to almost zero from an initial concentration of 4.5 % while the PUFA concentration was increased from an initial concentration of 33.5 % to 53.87 % [15]. The process is considered uncomplicated, quick, cheap, robust and environmental friendly [14, 16].

Winterization is another method adaptable for large scale separation of fatty acids. The method uses the difference of melting points of various fatty acids depending on the type and degree of unsaturation. Long-chain SFAs crystallize out at lower temperatures leaving PUFAs in liquid form [17]. For example, in biodiesel industry, winterization has been used to decrease saturated FAMES from 86.9% to 73.4% while increasing unsaturated FAMES from 12% to ~20% [18].

Currently used large-scale separation techniques such as urea complexation and winterization, besides being costly due to required temperature variations (E.G. in urea complexation, temperature of solution is increased to 65°C and then decreased to 4-24°C [19] whereas during winterization, temperature is decreased to -18°C[19]), are not quite selective. Chromatographic techniques are selective, but expensive (in terms of energy requirements) and difficult to scale-up. The essence of this study was to test methods that are scalable and energetically less costly. During preliminary studies, the ability of different metal salts to selectively precipitate saturated FAs from a mixture containing both saturated and unsaturated FAs dissolved in

methanol were evaluated [20]. Once a suitable salt was identified (in this case zinc sulphate), the effect of increasing ion concentration on selective precipitation of saturated (palmitic acid) from a mixture containing palmitic and palmitoleic acid in methanol was evaluated. Spectroscopic and computational studies were conducted to verify the composition and mechanistic aspects of formation of precipitates respectively.

2. EXPERIMENTAL DETAILS

2.1 Preliminary Studies

Preliminary studies were aimed at elucidating interaction of various transition metal salts with saturated and unsaturated fatty acids. For this purpose, 100 mg each of palmitic acid (saturated) and palmitoleic acid (unsaturated) were mixed in methanol. Approximately 0.17×10^{-3} moles of metal salts such as AgCl, CuSO₄, ZnSO₄, MnCl₂, FeSO₄, FeCl₃ and CoCl₂ (all salts were hydrated except for AgCl) were added individually to the solution of palmitic acid and palmitoleic acid in methanol. In the case of AgNO₃, CuSO₄ and ZnSO₄, immediate precipitate formation was observed as seen in photo 1, while in case of the other salts, no precipitation was observed (as seen in photo 2).




Photo 1: (Left to Right) a) AgCl b) CuSO₄ c) ZnSO₄ showing clear precipitation of saturated FA from a mixture containing both saturated and unsaturated FA




Photo 2: (Left to Right) a) Ferrous Sulphate b) Manganese Chloride c) Ferric Chloride d) Cobalt Chloride did not show any precipitation from a mixture containing both saturated and unsaturated FA

For metal salts showing immediate precipitation, studies were conducted to measure the decrease in the amount of fatty acids in the solution. Once AgCl , CuSO_4 and ZnSO_4 were added individually to the solution of FAs in methanol, the precipitate was allowed to settle for 24 hours, after which, the precipitate was filtered out and the supernatant was analyzed for the fatty acids concentration. FAs were first transesterified to FAMES using 4% solution of H_2SO_4 in methanol. The concentration of FAMES was estimated using a GC-FID. The procedure was repeated thrice from every metal salt. Figure 1 represents the results. The response of the study is measured as the difference is decrease of concentration of palmitoleic acid from the decrease in palmitic acid (decrease in concentration C16:0 – decrease in concentration of C16:1).

2.2 Optimization of ZnSO_4 Concentration

Preliminary studies indicated that ZnSO_4 preferentially precipitates palmitic acid from a mixture containing both palmitic and palmitoleic acids. Consequently, the present study was aimed at elucidating whether an increase in concentration of ZnSO_4 leads to increase in the formation of precipitates and if so, to what extent.

Hexadecanoic acid (C16:0) was obtained from Sigma Aldrich while 9-cis-hexadecanoic (C16:1) acid was purchased from Nu-Chek Prep. A 100 mg of each acid was added to 10 ml of methanol and the solution was mixed using a vortex mixer until the FAs were completely dissolved and a clear solution was obtained. To this solution, 50 mg of ZnSO_4 was added. Formation of an immediate precipitate was observed. The precipitate was allowed to form and settle for 24 hours. The precipitate was filtered and the supernatant was then transesterified to convert the remaining fatty acids in the solution to FAMES. For the transesterification process, the FA solution in methanol was mixed with 4% H_2SO_4 solution in methanol and heated at 120°C for 1 hour. The concentration of FAMES was estimated using

143 a GC-FID. The same procedure was repeated for different ZnSO₄ concentrations of 5, 10,
144 20, 30 and 40 mg/ml. Experiments at each concentration were run in triplicates.

145

146 **2.3 FTIR-ATR Analysis of Precipitate**

147

148 The precipitate obtained after adding ZnSO₄ to a mixture of C16:0 FA and C16:1 FA was
149 filtered and dried at 70°C to evaporate methanol present in the sample. Then the precipitate
150 was analyzed in a FTIR-ATR (ATR-Attenuated Total Reflectance mode) spectrometer to
151 obtain the spectra between 500-4000 cm⁻¹. Pure samples of palmitic acid and palmitoleic
152 acid were also analyzed using FTIR-ATR to compare their spectra to that of the precipitate.

153

154

155

156 **2.4 Computational Chemistry Calculations**

157

158 In order to explain the experimental results, theoretical calculations were performed.
159 Computational chemistry software, GuassView 5 and Gaussian 09 [21] were used to build,
160 optimize and find the energies of molecules and potential structures formed.

161 In the procedure it is mentioned that both fatty acids were completely mixed in methanol: this
162 means that the fatty acid molecules were ionized in solution, giving the carboxylate group of
163 the FA a net negative charge. The metal ions (in methanol) possess a net positive charge. It
164 was also observed from earlier studies that certain metal salts showed selective binding
165 preference with the saturated fatty acid forms. Accordingly, in order ascertain why certain
166 metals showed the tendency to form precipitates selectively with the saturated FA, following
167 computational chemistry calculations were undertaken:

168

- 169 a. Structures of palmitic acid, palmitoleic acid, a few of the metal ions used in the
170 experiment, and their corresponding "metal carboxylates" were optimized using the
171 method "Hartree-Fock" with basis set of "321-G" available in Gaussian 09;
- 172 b. Frequency calculations of the optimized structures were undertaken using the same
173 method and basis set, from which the sum of electronic and thermal free energies
174 (Standard Gibbs Free Energies) of ions and molecules were calculated; and
- 175 c. Assuming that positively charged metal ions formed a metal carboxylate by reacting
176 with the negatively charged fatty acid ion in methanol, the Gibbs free energies of
177 reactions of fatty acid ions with metals ions resulting in the formation of metal
178 carboxylate was calculated.

179

180 **3. RESULTS AND DISCUSSION**

181

182 **3.1 Preliminary Studies**

183 Figure 1 depicts the differences of the decrease of concentration of palmitoleic acid from the
184 decrease in palmitic Acid (decrease in concentration C16:0 – decrease in concentration of
185 C16:1).




Figure 1: Results of preliminary studies showing that metal salts selectively precipitate the saturated fatty acid (C16:0) more than the unsaturated fatty acid (C16:1)

From figure 1 it can be seen that copper, silver and zinc ions are effective in selectively precipitating out the saturated fatty acid from the solution. Considering cost and health benefits, it was decided to carry out further experiments with zinc [22] (sulphate).

3.2 Optimization of ZnSO_4 Concentration

This study was geared toward ascertaining whether the ZnSO_4 concentration has any impact on the amount of FAs that can be selectively separated. Figure 2 depicts how increased concentrations of ZnSO_4 lead to enhancement of the selective precipitation of the saturated FA (in comparison to unsaturated FA). The Y-axis in figure 2 is a measure of the difference in reduction of concentration of the unsaturated FA from that of the saturated FA. A positive difference indicates that the amount of C16:0 FA outweighs the amount of C16:1 in the solution.




Figure 2: Fatty acid precipitation with increasing concentration of ZnSO_4


An Analysis of variance (ANOVA) was conducted using Design-Expert (version 9.0.3.1) with concentration of ZnSO_4 as factors with 5, 10, 20, 30 and 40 mg/ml as levels. The ANOVA resulted in a p-value of 0.0002 - which is significant. This means that varying concentration of ZnSO_4 significantly affects the concentration difference between C16:1 FA and C16:0. From figure 2, it can be confirmed that increasing the ZnSO_4 concentration up to about 30 mg/ml leads to an increase in the relative amount of C16:0 FA precipitated. However, beyond 30 mg/ml ZnSO_4 concentration, the separatory effect seems to taper off. It can be clearly seen that the decrease in the concentration of C16:1 FA remains more or less constant at 10% whereas the reduction in concentration of C16:0 FA increases to a maximum of around 58%.

Another experimental model was analyzed where there were two factors : (a) Amount of ZnSO_4 added and (b) Type of fatty acid (which in turn has two levels : (i) C16:0 FA and (ii) C16:1). The response of this model was measured as the decrease in concentration of fatty acid after the addition of zinc sulphate. The ANOVA for this experimental model resulted in a p-value of < 0.0001. This means that the model is significant. The data for this experimental model is also represented in figure 2. Hence it can be surmised that (i) amount of ZnSO_4 added to the solution has a significant effect on the amount of precipitation and (ii) the saturated fatty acid selectively form a complex with zinc sulphate.

236
237
238

3.3 FTIR-ATR Analysis of Precipitate

Figure 4 depicts the spectra for C16:0 FA, C16:1 FA precipitates with zinc.



239
240
241

Figure 4: Spectra of C16:0 FA, C16:1 FA and Zinc Carboxylate precipitate

242
243
244
245
246
247
248
249
250
251

A stretch band above 3000 cm^{-1} indicates the presence of an alkene or “=C-H” bond [23], which can also be verified by observation of the differences in the spectra of C16:0 FA and C16:1 FA in figure 4 -- where there’s a distinct stretch band at 3003 cm^{-1} . The spectra for the zinc precipitate did not contain a stretch band above 3000 cm^{-1} -- indicating the absence of a carbon-carbon double bond in the precipitate. One distinct stretch band to be noted in the case of the precipitate is the one just above 1500 cm^{-1} -- which is indicative of the asymmetric carboxylate vibration in the IR spectra [$\nu_{\text{as}}(\text{COO}^-)$] [24]. It also confirms the presence of a zinc forming complex with the carboxylate group of the fatty acid [24] -- and in this case, a complex with the C16:0 FA.

252
253
254

3.4 Computational Chemistry Calculations

255
256
257
258
259
260
261
262
263
264

The Gibbs free energies of reactions (generated from computational chemistry calculations) are given in Table 1-4. Table 5 depicts information on the solubility of metal salts in methanol. The last column of each table gives the difference between the Gibbs Free energy of the reaction of the metal ion with the with palmitoleic acid and that with palmitic acid. If this value is negative, then it could be conjectured that the reaction of the metal ion with palmitic acid is more favorable than that with palmitoleic acid.

According to Table 1, negative Gibbs free energies for in column 6 indicate that Zn^{2+} has a tendency to complex with both C16:0 FA ion and C16:1 FA spontaneously. However, the

more negative value with palmitic acid indicates a preference for forming a complex with the metal ion when palmitic acid is mixed with palmitoleic acid. This result is consistent with the observations made experimentally.

Table 1: Computational calculations of Zn^{2+} – fatty acid chemistry

	Fatty acid ion (-1) (Ha)	Zn^{2+} (Ha)	Total Energy of Reactants (Ha)	Zinc Carboxylate (Ha)	Gibbs Free Energy (Ha)	Difference of Gibbs Free Energies (C16:0 - C16:1) (KJ/mol)
Palmitic acid	-807.778275	-1768.29	-3383.84747	-3384.88857	-1.041103	-3.257799999
Palmitoleic acid	-767.825167	-1768.29	-3303.94125	-3304.9811	-1.03985	

In the case for Cu^{2+} , as can be seen from table 2, Gibbs free energies for both reactions are negative - indicating that Cu^{2+} also has the potential to spontaneously form complexes with both forms of FAs. However, Cu^{2+} complexation with palmitic acid is favored in the presence of both forms of FAs since the Gibbs free energy with palmitic acid is more negative. This is again consistent with the results obtained experimentally.

Table 2: Computational calculations of Cu^{2+} – fatty acid chemistry

	Fatty acid ion (-1) (Ha)	Cu^{2+} (Ha)	Total Energy of Reactants (Ha)	Copper Carboxylate (Ha)	Gibbs Free Energy (Ha)	Difference of Gibbs Free Energies (C16:0 - C16:1) (KJ/mol)
Palmitic acid	-807.778275	-1630.13	-3245.68509	-3246.72107	-1.035988	-2.761200001
Palmitoleic acid	-767.825167	-1630.13	-3165.77887	-3166.8138	-1.034926	

Fe^{2+} also behaved consistent to the above cases: As can be seen from table 3, Gibbs free energies of Fe^{2+} complexing with both forms of FAs are negative indicating thermodynamic favorability. However, in this case, the reaction with palmitoleic acid is more favored. Nevertheless, experimentally, no precipitate was observed either with palmitic acid nor palmitoleic acid. This can be attributed to the fact that the solubility of ferrous sulphate is negligible in alcohols [25]. For both reactions to proceed, the availability of Fe^{2+} ions is essential.

Table 3: Computational calculations of Fe^{2+} – fatty acid chemistry

	Fatty acid ion (-1) (Ha)	Fe^{2+} (Ha)	Total Energy of Reactants (Ha)	Ferrous Carboxylate (Ha)	Gibbs Free Energy (Ha)	Difference of Gibbs Free Energies (C16:0 - C16:1) (KJ/mol)
Palmitic acid	-807.778275	-1255.39	-2870.94686	-2871.91519	-0.968329	67.8834

Palmitoleic acid	-767.825167	-1255.39	-2791.04064	-2792.03508	-0.994438	
------------------	-------------	----------	-------------	-------------	-----------	--

To validate the results, a calculation was run with select metal ions that did not form any precipitates experimentally. In the case of Co^{2+} , it can be seen from Table 4 that Gibbs free energies for formation of complexes with both forms of FAs are positive and thus non-spontaneous (even though cobalt chloride is soluble in methanol indicating readily available Co^{2+} ions [26]).


Table 4: Computational calculations of Co^{2+} – fatty acid chemistry

	Fatty acid ion (-1) (Ha)	Co^{2+} (Ha)	Total Energy of Reactants (Ha)	Cobalt Carboxylate (Ha)	Gibbs Free Energy (Ha)	Difference of Gibbs Free Energies (C16:0 - C16:1) (KJ/mol)
Palmitic acid	-807.778275	-1373.78	-2989.33515	-2871.76541	117.569746	78.1196
Palmitoleic acid	-767.825167	-1373.78	-2909.42894	-2791.88924	117.5397	

Table 5: Solubility of metal ions in methanol


ZnSO_4 (heptahydrate)	Soluble [27]
CuSO_4 (pentahydrate)	Soluble [28]
FeSO_4 (heptahydrate)	Insoluble / Sparingly Soluble [25]
CoCl_2 (hexahydrate)	Soluble [26]

Optimized structures of zinc pamitate and zinc palmitoleate are given in figure 5 and 6. The energy for an optimized molecule of zinc palmitate is -3385.811 Hatrees while that of zinc palmitoleate is -3305.797 Hatrees. Hence the complex between zinc and C16:0 is much more energetically stable that that with C16:1 FA. In figure 5 and 6, it is visible that the structure of zinc palmitate is linear when compared to that of zinc palmitoleate. Hence zinc palmitoleate molecules can stack themselves in a closely packed arrangement, promoting in precipitation and crystallization of the compound. These structural and energetic differences between the molecules of zinc pamitoleate and zinc palmitate could be the reason why zinc ions (and other metal ions) can easily form precipitates with the linear palmitic acid than with the skewed palmitoleic acid.



311

Figure 5: Optimized structure of zinc-palmitate



312

313

Figure 6: Optimized structure of zinc palmitoleate

314

315

316

317

4. CONCLUSION

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

REFERENCES

334

335

336

337

338

339

340

341

342

343

1. Benoit SC, et al. Palmitic acid mediates hypothalamic insulin resistance by altering PKC- θ subcellular localization in rodents. *The Journal of Clinical Investigation*, 2009. 119(9):2577-2589.
2. Yang ZH, Miyahara H and Hatanaka A. Chronic administration of palmitoleic acid reduces insulin resistance and hepatic lipid accumulation in KK-A(y) Mice with genetic type 2 diabetes. *Lipids in Health and Disease*. 2011. 10.
3. Spector A. Essentiality of fatty acids. *Lipids*. 1999. 34(1):S1-S3.
4. Shahidi F and Miraliakbari H. Omega-3 (n-3) fatty acids in health and disease: Part 1-cardiovascular disease and cancer. *Journal of medicinal food*. 2004. 7(4):387-401.

- 344 5. Morris LJ. Separations Of Lipids By Silver Ion Chromatography. *Journal of Lipid*
345 *Research*. 1966. 7(6):717.
- 346 6. Nikolova-Damyanova B. Retention of lipids in silver ion high-performance liquid
347 chromatography: Facts and assumptions. *Journal of Chromatography A*, 2009.
348 1216(10):1815-1824.
- 349 7. Teramoto M. et al. Extraction Of Ethyl And Methyl-Esters Of Polyunsaturated Fatty-
350 Acids With Aqueous Silver-Nitrate Solutions. *Industrial & Engineering Chemistry*
351 *Research*. 1994. 33(2): 341-345.
- 352 8. Teramoto M, Matsuyama H, and Nakai K. Separation of ethyl esters of
353 eicosapentaenoic acid and docosahexaenoic acid by circulating liquid membranes
354 using silver nitrate as a carrier. Uphill transport by use of temperature and solvent
355 dependencies of distribution ratio. *Separation Science and Technology*. 1996.
356 31(14):1953-1969.
- 357 9. Suzuki T et al. Extraction Of Polyunsaturated Fatty-Acid Ethyl-Esters From Aqueous
358 Silver-Nitrate Solution Using Supercritical Ethane And Ethylene And Modeling Of
359 The Extraction Process. *Kagaku Kogaku Ronbunshu*. 1994. 20(1): 97-104.
- 360 10. Awad A and Gray J. Methods to reduce free fatty acids and cholesterol in anhydrous
361 animal fat. Official gazette of the United States Patent and Trademark Office.
362 Patents. 2000. 1239(2).
- 363 11. Daloz D et al. Corrosion inhibition of rapidly solidified Mg-3% Zn-15% Al magnesium
364 alloy with sodium carboxylates. *Corrosion*. 1998. 54(6):444-450.
- 365 12. Hefter GT, North NA, Tan SH. Organic Corrosion Inhibitors in Neutral Solutions; Part
366 1 - Inhibition of Steel, Copper, and Aluminum by Straight Chain Carboxylates.
367 *Corrosion (Houston)*. 1997. 53(8):657-667.
- 368 13. Rocca E and Steinmetz J. Inhibition of lead corrosion with saturated linear aliphatic
369 chain monocarboxylates of sodium. *Corrosion Science*. 2001. 43(5): 891-902.
- 370 14. Hayes DG et al. Urea complexation for the rapid, ecologically responsible
371 fractionation of fatty acids from seed oil. *Journal of the American Oil Chemists*
372 *Society*. 1998. 75(10): 1403-1409.
- 373 15. Fei C, Salimon J, and Said . Optimisation of Urea Complexation by Box-Behnken
374 Design. *Sains Malaysiana*. 2010. 39(5): 795-803.
- 375 16. Wanasundara UN and Shahidi F. Concentration of omega 3-polyunsaturated fatty
376 acids of seal blubber oil by urea complexation: optimization of reaction conditions.
377 *Food Chemistry*. 1999. 65(1):41-49.
- 378 17. Shahidi F and Wanasundara UN. Omega-3 fatty acid concentrates: Nutritional
379 aspects and production technologies. *Trends in Food Science & Technology*. 1998.
380 9(6): 230-240.
- 381 18. Doğan TH. and Temur H. Effect of fractional winterization of beef tallow biodiesel on
382 the cold flow properties and viscosity. *Fuel*. 2013. 108(0): 793-796.
- 383 19. Mendes A., da Silva T, and Reis A. DHA concentration and purification from the
384 marine heterotrophic microalga *Cryptocodinium cohnii* CCMP 316 by winterization
385 and urea complexation. *Food technology and biotechnology*. 2007. 45(1):38-44.
- 386 20. Kulkarni S and Fernando S. Separation of Palmitic and Palmitoleic Acid and their
387 FAMES using Transition Metal Salts. *International Research Journal of Pure and*
388 *Applied Chemistry* 2014. 4(5): 551-561.
- 389 21. Frisch MJ et. al. Gaussian 09, Revision A.02. 2009. Gaussian, Inc. Wallingford CT.
- 390 22. Brown KH et al. Effect of supplemental zinc on the growth and serum zinc
391 concentrations of prepubertal children: a meta-analysis of randomized controlled
392 trials. *The American journal of clinical nutrition*. 2002. 75(6): 1062-1071.
- 393 23. Volland W. Organic Compound Identification Using Infrared Spectroscopy. 1999
394 [cited 2014 May 18].

- 395 24. Zeleňák, V, Vargová Z, Györyová K. Correlation of infrared spectra of zinc(II)
396 carboxylates with their structures. *Spectrochimica Acta Part A: Molecular and*
397 *Biomolecular Spectroscopy*, 2007. 66(2):262-272.
- 398 25. Oldberg O and Long H. *A Laboratory Manual of Chemistry: Medical and*
399 *Pharmaceutical*. 1891: WT Keener.
- 400 26. Cobaltous Chloride, Hexahydrate :Safety Data Sheet. LabChem.
- 401 27. DOSE/Dictionary of substances and their effects. Royal Society of Chemistry:
402 Cambridge, UK.
- 403 28. Haynes WM. *CRC Handbook of Chemistry and Physics* 92 ed. 2011, Boca Raton,
404 FL: CRC Press.
- 405