UNDER PEER REVIEW

Original Research Article

EFFECT OF ATRAZINE ON GERMINATION AND GROWTH PERFORMANCE OF WATER YAM (*Dioscorea alata*)

ABSTRACT

Effect of rates of application of atrazine on germination and growth performance of water yam minisetts was investigated at Ladoke Akintola University of Technology, Ogbomoso, Oyo State, Nigeria. The treatments in the field experiment were four (4) rates of atrazine i.e 100%, 50%, 25%, and 0% of the recommended dosage rate of 2.5kga.i/ha or 5litres/ha. A plot to which no herbicides was applied served as the control (0%). The experiment was arranged in a randomized complete block design with three replications. Result showed that though yam minisetts emergence was not hampered by the rates of application of atrazine, the growth parameters of the yam minisetts were significantly (p= 0.05) affected at early stage. Yam minisetts vine length and vine thickness were significantly (p=0.05) higher at 50% treatments compared to other treatments and control. Rates of application of atrazine however, did not affect the number of leaves of yam minisetts. The rates of application of atrazine did not significantly (p=0.05) affect the yield parameters of yam minisetts, though the heaviest weight was produced under 25% treatments. Similarly, the rates of application of atrazine were similar in their effect on weed control and were better than the control.

Key words: Yam minisetts, atrazine, dosage.

INTRODUCTION

Yams, *(Dioscorea spp.),* are important tuber crop and staple food for millions of people in many tropical and subtropical countries. It has become a cash crop in countries along the coast of West Africa (Egesi *et al,* 2005). They produce edible starchy storage tubers, which are of cultural, economic, and nutritional importance in the tropical and subtropical regions of the world (Coursey, 1967). Generally, yam tubers are boiled, roasted, baked, or fried. However, in some regions, the tubers are boiled and then pounded to glutinous dough called “fufu” Cooked yam tubers or their products are usually eaten in association with protein-rich sauces. It has also gained importance in pharmaceutical industries (Hahn, 1995). Nigeria accounts for about 70% of the world's production of yam, generating a global output of over 33 million metric tons (FAO,

UNDER PEER REVIEW

2000). *D, alata* is popular and prevalent in Abakaliki agro ecological zone of Ebonyi State, Nigeria where it is called "Mbala or Nvula" (Igbo names).

Traditionally, water yams are propagated vegetatively by means of tuber. Whole tuber called "seed" yam from l00g to 1500g are used as planting materials. Alternatively, larger tubers are often cut into approximately 200g pieces and used to establish the new crops (Onwueme, 1978; Okoli *et al*, 1982). A major constraint to yam production however, is the availability of the planting material. Since the tuber for which the crop is grown is equally needed as planting material, it becomes difficult to get enough material for planting since this constitute a competitive use for the tuber.

Furthermore, the conventional seed yam and or yam sett are usually very big thereby consuming a greater percentage of the total yam produced. The practical implication is that the seed represent the highest cost items, accounting for as much as 50 percent of operating expenses (Gray *et al*,1991) while planting material makes up 35 percent and 54 percent of the total cost of production (Osei, 1993).

The National Root Crop Research Institute in Nigeria developed a system termed the "minisett" technique which aims at overcoming the seed yam problem. A "minisett is defined as set less than one quarter of the minimum size (l00g) of yam sett often planted (Okoli *et al*, 1982). The technique involves cutting the selected "mother seed" yam into pieces 20g to 25g each, treating them in a fungicides or insecticides suspension and sprouting them in nursery beds or polyethylene bags. After three to four weeks, the sprouted minisetts are transplanted into the main fields. (Vernier *et al,* 1999)

However, a major constraint to minisetts production is weed control. This is because the tender yam seedling could not easily outcompete the accompany weeds. In addition, the tender nature of the seedlings makes the use of manual weeding using hoe weeding a delicate activity. Thus, a more effective and less cumbersome weeding using the herbicide becomes more appealing. It is in the light of this that this experiment was designed to evaluate the effect of atrazine, a popular pre-emergence herbicide, on the germination and growth performance of water yam minisett. The result is expected to enhance the success of rapid production of yam minisetts.

MATERIALS AND METHODS

UNDER PEER REVIEW

Field experiment was carried out at Ladoke Akintola University of Technology, Teaching and Research Farm, Ogbomoso, during the rainy seasons of 2011 and 2012.

The experiments were laid out in a randomized complete block design, with three replicates. The minisetts, 25g each, were treated with wood ash and cut the surfaces were allowed to dry for 24hrs before planting. The minisetts were planted in situ at 6 plants per bed at a spacing of 0.5m by 0.5m. The bed was dimensioned 2m x 2m. Grass mulch was applied on planting spots to conserve moisture and prevent excessive sunlight. Four rates of application were tested in the experiment namely; atrazine 100%, atrazine 50%, atrazine 25% and control treatment to which no herbicide was applied. Atrazine was applied to rain moistened soil using a knapsack sprayer model Cp3, which delivers at a rates of 2001/ha. To achieve the 100%, 50%, 25% of dosage rates of herbicides, 250ml, 125ml and 62.5ml of the herbicides were dissolved each in 10 litres of water. Stakes were provided for plant stands and the growing young vines were trained as soon they are long enough using threads. Data were collected on growth and yield parameters of water yam. Weed population density was estimated from 3 randomly placed $0.25m^2$ quadrat, while the wet weight of weeds harvested from the quadrats were measured on a sensitive scale. Mean data collected over the two years were subjected to analysis of variance (ANOVA) while the means were separated by DMRT at 5% probability level.

RESULT AND DISCUSSION

Effect of rates of application of atrazine on germination of yam minisetts is presented in the Table 1. There were no significant (p=0.05) differences among the treatments and the control at 3WAP. Atrazine did not have any effect on the germination of yam minisetts as both the atrazine treated and the untreated minisetts germinated at the same rate. At 6WAP, there were no significant differences on germination rate between atrazine 50%, Atrazine 25% and control. These were however significantly (p=0.05) higher than those of Atrazine 100%.

Effect of rates of application of atrazine on growth parameters of yam minisetts is presented in Table 2. There were no significant (p=0.05) effect of rates of application of atrazine on yam minisetts growth parameters across the period of data collection.

Similarly, application rates of atrazine did not have significant (p=0.05) effect on tuber yield of yam minisetts (Table 3). However, the heaviest tuber (1.3kg) was produced under 25%

UNDER PEER REVIEW

treatment (1.30kg) followed by the control (1.0kg). The smallest tuber (0.87kg) was produced by the 100% treatment.

The weed control effectiveness of the rates of application as measured by weed density and biomass is presented in Table 4. All the rates were superior to the control in effectiveness (p= 0.05). There were no significant difference (p= 0.05) between the rates in effectiveness.

The result from this experiment indicated that atrazine can safely be used for weed control in yam minisett production. The experimented rates did not have any significant negative effects on both the growth and tuber yield of water yam compared to the control. Thus, it could be safely recommended that the lowest rates (25%) be adopted for weed control in yam minisett production more so, when there were no significant differences between the weed control efficiencies of the rates. Similar finding had been reported by Olabode *et al* (2010). In addition, since atrazine had been listed as one of the most persistent herbicides (Akobundu 1987), an effective reduced dosage rate will help mitigate the problem of persistence of the herbicides in addition to added economic gains. Furthermore, since the use of herbicides does not involve using equipment like hoes and cutlasses that can easily injure or destroy the tender and fragile yam minisetts vines, the choice of this method of weed control aside reducing labour, drudgery and cost will guarantee safety of the yam minisett vines.

CONCLUSION

This experiment has shown that atrazine at the rates used in this experiment is adequate for use in the rapid multiplication of water yam tubers through minisetts. However, for economic and environmental concern, it is recommended that 25% of the recommended dosage rate of atrazine should be used for weed control in yam minisetts production.

UNDER PEER REVIEW

Table 1: Effect of rates of application of Atrazine on Germination of Yam Minisetts.

	%Germination	
Treatment	**3WAP**	**6WAP**
Atrazine l00%	16.7a	55.6b
Atrazine 50%	27.7a	77.8a
Atrazine 25%	33.3a	72.2a
Control	33.4a	77.8a

Mean followed by the same superscript along the column are not significantly different (p=0.05) by DMRT. WAP: Weeks After Planting

Table 2: Effect of rates of application of Atrazine on Vine length of yam Minisetts

	Vine length (cm)			
Treatment	**6**	**8**	**10**	**12**
	→	**WAP**	←	
Atrazine 100%	33. 1b	86.0a	122.6ab	147.0a
Atrazine 50%	58.1a.	98.0a	I41.4a	144.4a
Atrazine 25%	54.4a	71.4a	80.7b	135.9a
Control	46.8a	96.2a	124.6ab	126.4a

Mean followed by the same superscript along the column are not significantly different (p=0.05) by DMRT.

UNDER PEER REVIEW

Table 3: Effect of rates of application of Atrazine on Vine thickness of yam Minisetts

	Vine thickness (cm)			
Treatment	**6**	**8**	**10**	**12**
	→	**WAP**	←	
Atrazine 100%	0.5a	0.5a	0.5a	0.6ab
Atrazine 50%	0.6a	0.5a	0.6a	0.7a
Atrazine 25%	0.5a	0.5a	0.5a	0.6ab
Control	0.5a	0.5a	0.5a	0.5b

Mean followed by the same superscript along the column are not significantly different (p= 0.05) by DMRT.

Table 4: Effect of rates of application of Atrazine on number of leaves of yam Minisetts

	Number of leaves			
Treatment	**6**	**8**	**10**	**12**
	→	**WAP**	←	
Atrazine 100%	8.4a	24.0a	33.8a	49.0a
Atrazine 50%	11.8a	32.2a	37.5a	49,0a
Atrazine 25%	11. 8a	27.7a	32.7a	48.8a
Control	10.0a	24.7a	30.2a	52.2a

Mean followed by the same superscript along the column are not significantly different (p=0.05) by DMRT

UNDER PEER REVIEW

Table 5: Effect of rates of application of atrazine on weed density and weed biomass of yam minisetts

Treatment	Weed Density Unit/25m^2 8WAP	Weed Biomass g/25cm^2 8WAP
Atrazine 100%	10.3b	28.3b
Atrazine 50%	14.4b	33.3b
Atrazine 25%	15.0a	32.3b
Control	28.9a	79.7a

Mean followed by the same superscript along the column are not significantly different (p=0.05) by DMRT.

WAP: Weeks After Planting

Table 6: Effect of rates of application of Atrazine on the yield parameters of yam minisetts

Yield parameters			
Treatment	**Tuber Weight (kg)**	**Tuber length (cm)**	**Tuber Diameter (cm)**
Atrazine 100%	0.87a	21.20a	9.50a
Atrazine 50%	0.99a	24.90a	8.90a
Atrazine 25%	1.30a	24.90a	10.30a
Control	1.00a	25.30a	9.70a

Mean followed by the same superscript along the column are not significantly (p=0.05) different by DMRT

UNDER PEER REVIEW

REFERENCES

Akobundu, I.O. 1978. Weed control strategies for multiple cropping system of the humid and sub-humid tropics. Pp 80-95 IITA proceeding series 3.

Akobundu, I.O. 1987. Weed science in the tropics. John Wiley and sons, New York, 522pp.

Egesi, C.N., Ogunyemi, S. and Asiedu, R. 2005. Genotype x environment interaction in water yam *(Dioscorea alata)* yield trials in five yam-growing states in Nigeria. Journal of genetics and breeding 59: 81-88

Coursey, D. G. 1967. *Yams.* Longmans, London.230 pp.

Hahn, S. K. 1995. Yams: *Dioscorea* spp.(Dioscoreaceae). In: J. Smartt & N.W. Simmonds (Eds), Evolution of crop plants. Longman Scientific & Technical, UK. Pp. 112 - 120.

FAO 2000. FAOSTAT Agricultural data. Food and Agriculture Organisation of the United Nations. http//www.fao.org.

Onwueme I.C. 1978. The tropical tuber crops, Yams, Cassava, Sweet Potato and Cocoyam, xiv, 234p

Okoli O., LA Acha, H. Shiwachi, R. Asiedu, M.O Akoroda. 1982. Effect of Auxin on root Development in Yam *(Dioscorea rotundatd)* vine. Tropical Science vol 44: 80-84.

Gray, A., D. Crentsil, S. Gallat and S. Gogoe. 1997. Survey of yam trading practices and loss assessment in Techiman market. Project A0497; R6505. Chatham: Natural Resources Institute.

Osei, C. (1993). The yield of seed yams grown from mmisette as influenced by mulch in Rudat, H., *et al*., Improving farming systems in the interior savanna zone of Ghana: proceedings of the 2nd workshop on improving farming systems; Nyankpala April, 1990. Tamale: CRI/NAES.

UNDER PEER REVIEW

Olabode, O.S., Adesina G.O. and Babajide P.A. (2010). Weed control efficiency of reduced atrazine doses and its effect on soil organisms in maize *(zea mays* L) on field of South Western Nigeria. Journals of Tropical Agriculture 48 (1-2): 52-54

Vernier, P., N. Ericas, E. Aegbo, J. Hounhouigan, K. N'Kpenu and G. Orkwor. 1999. From biological control to a systems approach in post- harvest'. Proceedings of an IITA/GTZ meeting on integrated control of insect pests in rural maize stores, with particular reference to the Larger Grain Borer, and the future development of the post-harvest sector in sub-Saharan Africa. Eschborn: GTZ.