

Evaluations of Bentonite Plated Sulfur Application on Fe and Zn Bioavailability of Saline and Sodic Soil

A.R. Jafarnejadi^{1*}, A.Gholami¹

1- Soil science Department, Agricultural College, Ahvaz Branch, Islamiic Azad University

Corresponding author: arjafarnejady@gmail.com

Abstract

In saline and sodic soils, high pH strongly have affected on soil micronutrient availability. The study was conducted to evaluate using soil amendments (sulfur and manure) on some soil characteristics, iron and zinc concentrations of saline and sodic soil. A factorial randomized complete blocks design was considered with three replications. The main factor included organic matter (manure) as two levels (M_0 = without manure and M_1 = 40 tons manure/ha) and subplots were bentonite plated sulfur and Thiobacillus (five treatments) includes S_0 = control without S, S_1 = sulfur applying as soil gypsum requirement (1620kg/ha), S_2 = sulfur applying as double soil gypsum requirement (3240kg/ha), S_1T = S_1 +Thiobacillus treatment, S_2T = S_2 +Thiobacillus. The irrigation treatments were performed based on soil moisture monitoring plots. Also, the soil samples were taken in depth 0-30 cm and analyzed for determing soil pH, EC, SAR, soil Fe and Zn DTPA-extractable concentrations. The results showed that the manure and sulfur treatments were significant difference on soil EC, SAR and Zn DTPA-extractable. Also, the compare mean Duncan's treatments revealed that Zn DTPA-extractable concentrations was increased from 1.7 (M_0S_0) to 4.1 (M_1S_2) mg/kg ($p < 0.05$). Fe DTPA-extractable concentrations had increasing trend from 5.3 (M_0S_0) to 8.2 (M_0S_1T) mg/kg but the differences had not significant statistics ($p < 0.05$). The interactive effects of sulfur and organic matter application reduced soil SAR from 15.5 in the M_0S_0 treatment to 9.9 in the M_0S_2T treatment (in which sulfur was used together with Thiobacillus. Finally, organic manure and sulfur were recommended for increasing efficiency saline and sodic soils

Keywords: Soil amendments, Fe, Zn, Soil characteristics

Introduction

The soil salinity and drought are the most common environmental stress around the world including Iran (AKhani and Ghorbanli, 1993). The parts of natural and agricultural ecosystems are influenced under salt stress (Hafsi et al, 2007) and the soils were extended with difference's characteristics on the world (Rengasamy, 2005). Tóth et al (2008) reported that the area of salt affected soils is estimated at about one billion hectares. Sulfur application is amended for soil reclamation. The results explain the effect of the oxidizing bacteria, especially Thiobacillus (Mahjer milaniand Sistani, 1989).

Pot studies conducted in recent years at the Soil and Water Research Institute of Iran, suggest sulfur (if used together with Thiobacillus) has positive effects on soil physical, chemical, and biological properties (Besharati and Saleh rastin., 2006).

Sardinha et al., (2003) mentioned that organic and inorganic amendments consumption on saline soils can modify the effects of salinity on microbial activity and biochemical properties of soils. Nowadays, the large number of organic amendments such as mulch, manure and compost are used to improve saline and saline-sodic soils (Melero et al., 2007). Increased organic matter content can be a suitable method for solubilization of calcium carbonate and other minerals naturally occurring in soils (Minhas et al., 1995; Choudhary et al., 2002). In

saline or sodic soils, application of organic matter may increase leaching of sodium, reduce soil salinity, and increase soil water holding capacity and aggregate stability (Qadir et al., 2001). Moreover, Jalali and Ranjbar (2009) showed using organic amendments increased cation exchange capacity, saturated calcium (Ca), magnesium (Mg), and potassium (K) exchange sites, and prevented sodium from entering the exchange phase. When organic amendments are added to saline-sodic soils, more sodium is leached from soils, and soil properties such as infiltrability, water holding capacity, and aggregate stability are increased, while salinity and exchangeable sodium percentage declined (Qadir et al., 2001). Moreover, application of organic amendments reduces exchangeable sodium percentage (Qadir et al., 2003; Tejada et al., 2006). The researches have shown that the using of gypsum, compost and organic matter in saline soils increase leaching of sodium, soil permeability and are decreased soil exchangeable sodium and electrical conductivity (Tjada et al., 2006).

Considering sodic soils in Iran often contain calcium in the form of calcium carbonate, it is possible to use sulfur for amending these soils. The purpose of the study was to determine effective sulfur on soil micronutrients concentrations (Iron (Fe), Zinc (Zn)) and influenced soil pH and salinity in saline.

Materials and Methods

A factorial experiment using the randomized complete blocks design with three replications was conducted in saline-alkali soils 45 km on Ahvaz regions of South west Iran. There were 10 treatments. The each plots area was 4×2.5 (10 m^2). Manure levels (cow manure), 0 (M_0) and 40 t/ha (M_1), were applied to the main plots. Sulfur in the form of pelletized sulfur-bentonite mixture (made by Iran petrochemical national company) and *Thiobacillus thiooxidans* (5 treatments), including S_0 (the control with no sulfur), S_1 (sulfur as the gypsum requirement: 1620 kg/ha), S_2 (sulfur twice as much as the gypsum requirement: 3240 kg/ha), S_1T (S_1+T . *thiooxidans*), and S_2T (S_2+T . *thiooxidans*) were in the sub plots.

Gypsum requirement (GR) was calculated as fallowed equation1:

$$GR = \frac{(ESP_i - ESP_f) \times CEC \times BD \times D \times 8.61}{100f}$$

ESP_i : Initial ESP; ESP_f : Final ESP; CEC: Cation Exchange Capacity (me.100 /grsoil); BD: Bulk density (gr/cm^3); D: Soil depth (cm) and f: Gypsum convert coefficient to Sulfur (0.57).

The soil samples were taken in depths of 0-30 to determine initial soil properties, and water samples were taken to evaluate irrigation water quality in the region. After applying the treatments, the plots were irrigated. Irrigation interval during the experiment was determined by monitoring soil moisture. At the end of the experiment, soil samples were taken from depths of 0-30 cm in each treatment, and pH, soil salinity, sodium absorption ratio (SAR), and available iron (Fe) and Zn (Zinc) were measured. MSTAT-C software was applied for statistical analysis of the data and for the preparation of the variance analysis, and Duncan's multiple range test was employed for comparison of the means.

Results and discussion

Table 1, shows results obtained from analysis of soil samples taken in the region.

Table1: Soil chemical properties analysis

Depth (cm)	EC* (dS/m)	pH	OC (%)	SAR	P (mg/kg)	K (mg/kg)	TNV (%)	Sand (%)	Silt (%)	Clay (%)
0-30	33.6	8	0.23	27.7	2.2	253	50	44	36	20

*: EC: Electrical conductivity; pH: Soil acidity; OC: Organic carbone; SAR: Sodium absorption ratio; P: Phosphorous; K; Potassium; TNV: Total neutralization value;

Soil salinity (EC>4) and sodium absorption ratio (SAR>13) exceeded the standard limits (classification of salt-affected soils by US Salinity Laboratory) and, hence, the soil was saline-sodic. Moreover, the soil was classified as calcareous, and it had a medium texture.

Table 2 shows results observed in the analysis of water samples taken in the region.

Table2: Water chemical properties analysis

EC	pH	HCO ₃ ⁻	Cl ⁻	So ₄ ⁻²	T.A	Ca ⁺²	Mg ⁺²	Na ⁺	T.C
(μS/m)	-				me/l				
1990	7.9	4	11	5.4	20.4	6	4	10.4	20.4

T.A: Total Anions, T.C: Total Cations

Irrigation water salinity was in class C₃S₁ (salinity: 750-2250 μS/m and SAR<10) of Wilcox water quality table. Therefore, the water in the region was classified as very saline and of low sodium concentration.

Effects of treatments on soil properties

Results of ANOVA related to the applied treatments are presented in Table 3.

Table3: Analysis of Soil properties Variance

S.O.V	d.f	Salinity	pH	SAR	Fe	Zn
Replication	2	448.3	0.001	46.6	1.16	1.4
OM	1	2239.4*	0.3	77.1*	0.38	6.8*
Sulfur (S)	4	302.3	0.023	38.1	1.11	1.4
OM× S	4	285.3	0.034	8.5	3.8	0.73
Error	18	247.7	0.022	20.6	2.47	1.3

*- There is a significant difference between treatments at 0.05 levels

The individual effects of organic matter on salinity, SAR, and Zn showed statistically significant differences (p<0.05). However, the individual effects of sulfur on none of the studied properties were significantly different (p<0.05). Moreover, the interactive effects of organic matter and sulfur on the studied properties did not show statistically significant differences (p<0.05). Comparison of the means showed the individual effects of organic matter application on all of the studied properties (except Fe) were significant (p<0.05) (Table 4).

Table4: Comparison of the means soil properties of the study interactions between sulfur and organic matter (0-30 cm)

Treatment		EC (dS/m)	pH	SAR	Zn (mg/kg)	Fe (mg/kg)
M ₀	S ₀	21.6 b*	7.6 a	15.5 ab	1.7 a	5.3 a
	S ₁	24.5 b	7.53 abc	15 abc	1.93 ab	6.3 a
	S ₂	24.6 b	7.53 abc	11.6 ab	2.82 ab	7.1 a
	S ₁ T	16.7 b	7.57 ab	10.6 ab	2.00 ab	8.2 a
	S ₂ T	20.7 b	7.54 abc	9.9 b	2.7 ab	6.3 a
M ₁	S ₀	54.4 a	7.27 cd	19.4 a	3.7ab	6.1 a
	S ₁	46.3 ab	7.23 d	18.2 ab	2.87 ab	7.5 a
	S ₂	21.5 b	7.57 ab	11.9 ab	4.1 a	7.4 a
	S ₁ T	41.9 ab	7.3 bcd	13.1 ab	2.3 ab	6.4 a
	S ₂ T	39.4 ab	7.4 a-d	13 ab	3.1 ab	7.6 a

*: The similar letters has not significant differences at p<0.05, M₀: Without organic matter M₁: Using 40 (tons/ha); S₀: Without sulfur, S₁: Using sulfur as gypsum requirement, S₂: Using Sulfur as 2-fold gypsum requirement; T: Using Thiobacillus

When the rate of organic matter application was raised to 40 t/ha, salinity levels increased. This shows the organic matter used in the experiment was not of good quality. Application of various rates of sulfur together with Thiobacillus bacteria did not have significant effects on the studied properties ($p < 0.05$), but applying sulfur (individually or together with Thiobacillus) had positive effects on improving the studied properties (although these improvements did not cause statistically significant differences $p < 0.05$).


Interactive effects of treatments on soil properties

Table 4 revealed that the interactive effects of the applied treatments on the measured properties. The means of the interactive effects on salinity levels were different in the treatment, where organic matter was not applied as compared to treatments with various rates of sulfur application. All organic matter treatments had significantly increased on the mean salinity levels, this is due to organic matter used in the experiment was of high salt-containing and caused these changes in the soil. Furthermore, the trend of changes in salinity at both rates of organic matter application showed that the various sulfur application treatments caused reducing soil salinity. In treatments, where the organic matter (high-salt containing) was applied, these changes were more pronounced and salinity declined from 54.4 dS/m in the M_1S_0 to 21.5 dS/m in the M_1S_2 (Table 4).

Management of salinity and sodicity become more complicated when both occur in the same soil together. In research conducted in Iran, the positive effects of sulfur in reducing soil pH, salinity, and sodic, were greatest in Golestan province followed by Orumiyeh Province in the second place. However, sulfur application did not have these effects (or only had slight effects) in other provinces (Mohajer milani, 2000). A saline-sodic soil normally does not exhibit strong sodicity symptoms. The Ca can be replaced with sodium due to it has more valiant and ionic strength. Soil amendments are materials, such as gypsum, that directly supply soluble calcium for the replacement of exchangeable sodium, or other substances, such as or sulfur acid and or sulfur, that indirectly through chemical or biological action, make the relatively insoluble calcium carbonate commonly found in sodic soils, available for replacement of sodium.

The interactive effects of sulfur and organic matter application reduced soil SAR from 15.5 in the M_0S_0 treatment to 9.9 in the M_0S_2T treatment (in which sulfur was used together with Thiobacillus) (Table 4). Moreover, application of organic matter significantly increased SAR (as compared to treatments in which organic matter was not applied) because the organic matter was of the high salt-containing. Even under these conditions, the trend of changes in the studied properties was a descending, and sulfur application improved the sodic condition of the soil.

Application of sulfur and organic matter also caused changes in the chemical properties of the soil (such as its sodium and chlorine contents), and these changes were similar to those observed in salinity levels and in SAR. Applying organic matter and sulfur caused relative improvement in soil chemical properties and improved saline and sodic conditions of the soil (Table 4). The interactive effects of organic matter and sulfur on the levels of Fe and Zn availability are shown in Table 4, and Figure1 shows changes in the concentrations of Zn and Fe. Sulfur application significantly increased availability of nutrients so that Zn concentration increased from 1.7 in the control to 4.1 mg/kg in the M_1S_2 treatment.



Furthermore, changes in concentrations of nutrients were greater in treatments of organic matter application compared to treatments in which organic matter was not applied. Results shown in Table 4 indicate Fe concentrations followed a similar trend and increased from 5.3 in the M_0S_0 treatment to 8.2 mg/kg in the M_0S_1T treatment (but this increase was not statistically significant $p < 0.05$).

Figure 2 indicates changes in SAR and in concentrations of micronutrients (Fe and Zn).




Figure 2: Interactive effects organic matter and sulfur on SAR and Fe and Zn concentrations

SAR followed a descending trend when higher rates of sulfur were applied without Thiobacillus and declined from 15.5 in the M_0S_0 treatment to 9.9 in the M_0S_2T treatment, which was a significant reduction. With the decrease in SAR, concentrations of Zn and Fe followed an ascending trend. In other words, with soil sodic improved, soil fertility also followed an ascending trend of improvement. As shown in Table 4, SAR substantially increased with increased rates of organic matter application, and this increase in SAR can have adverse effects on concentrations of nutrients. However, with increased rates of sulfur application together with Thiobacillus, Zn and Fe became more available and salinity and sodic improved (in spite of the unfavorable saline and sodic conditions of the soil). This subject is due to Thiobacillus effects on sulfur (S). Thiobacillus converted S to So_4^{-2} and finally Acid sulfuric was created (Strong acid that could change soil pH and increased micronutrients such as Fe and Zn).

CONCLUSION

Saline and sodic soils are main challenge for agricultural plan on Khuzestan province, Iran. In saline and sodic soils, high pH strongly have affected on soil micronutrient availability. The results showed that the manure and sulfur with thiobacillus treatments were significant difference on soil EC, SAR and Zn DTPA-extractable. So, this is very effective to reduce micronutrient deficiencies (iron, zinc) in the production of healthy agricultural crop and food in saline and sodic soils.

Also, the compare mean Duncan's treatments revealed that Zn DTPA-extractable concentrations were increased. Finally, organic manure and sulfur were recommended for increasing efficiency saline and sodic soils. Also, other researches were recommended with organic matter different sources and sulfur with corps such as wheat and barley for studying micronutrients uptake by plants.

Acknowledgment

This article is extracted from research design on title” Evaluations pellets bentonite sulfur application on Fe and Zn bioavailability of saline and sodic soil” that supported as finance by Ahvaz Branch, Islamic Azad University, Iran. Hereby, authors have special thanks for this cooperation from research and technology vice president of Ahvaz Islamic Branch, Azad University, Iran.

References

1. Akhani, H., and Ghorbanli, M. 1993. A contribution to the halophytic vegetable and flora of Iran. P 35-44, In: H. leith and A.A. Al Massom (eds), Towards the rational use of high salinity tolerant plants, Kluwer Academic Publishers, Dordrecht.
2. Besharati, H. and Saleh rastin, N. 2006. Investigation effects Thiobacillus inoculation with Sulfur application on P availability increasing. Soil and water research Institute, Iran, Soil and water Journal. 12(1): 23-39.
3. Choudhary, O.P., Josan, A.S., Bajwa, M.S. 2002. Role of organic materials in mobilizing intrinsic calcium carbonate to ameliorate sodic irrigations. Proceedings of the 17th World Congress Soil Science, Symposia No. 34, Abstract Vol. III (Symposia 22–36), p. 1162.
4. Jalali M., F. Ranjbar. 2009. Effects of sodic water on soil sodicity and nutrient leaching in poultry and sheep manure amended soils. Geoderma 153. 194–204.
5. Hafsi, C., Lakhdar, A., Rabhi, M., Debez, A., Abdelly, C., and Ouerghi, Z. 2007. Interactive effects of salinity and potassium availability on growth, water status, and ionic composition of *Hordeum maritimum*. J. Plant Nutr. Soil Sci. 170: 469-473.
6. Melero, S., Madejon, E., and Herencia, J.C. 2007. Chemical and biochemical properties of a clay soil under dry land agriculture system as affected by organic fertilization. Eur J. Agron. 26: 327-334.
7. Minhas, P.S., Sharma, D.R., Singh, Y.P. 1995. Response to paddy and wheat to applied gypsum and FYM on an alkali water irrigated soil. J. Indian Soc. Soil Sci. 43: 452–455.
8. Mohajer milani, P and Sistani. H. 1989. Investigation effects sulfur different levels on P availability in Alfalfa cultivation. Soil and water research Institute, Iran, Soil and water Journal. 9(1): 42-63.
9. Qadir M, Schubert S, Ghafoor A, Murtaza G. 2001. Amelioration strategies for sodic soils: a review. Land Degrad Dev 12: 357–386.
10. Rengasamy, P. 2005. World salinisation with emphasis on Australia. Comp. Biochem. Phys. 141: 337-348.
11. Sardinha, M.T., Muller, H., Schmeisky, R., and Joergensen, G. 2003. Microbial performance in soils along a salinity gradient under acidic conditions. Appl. Soil Ecol. 23: 237-244.

12. Tejada, M., Garcia, C., Gonzalez, J.L., and Hernandez, M.T. 2006. Use of organic amendment as a strategy for saline soil remediation: influence on the physical, chemical and biological properties of soil. *Soil Bio. Biochem.* 38: 1413-1421.
13. Tóth, G., Montanarella, L., and Rusco, E. 2008. Updated map of salt affected soils in the European Union threats to soil quality in Europe, European Communities, Pp: 61-74.