

Original Research Article
**Anti-sickling and antibacterial activities of
Some Extracts from *Gardenia ternifolia* subsp.
jovis-tonantis (Welw.) Verdc. (Rubiaceae) and
Uapaca heudelotii Baill. (Phyllanthaceae).**

ABSTRACT

Aims: To evaluate the antisickling and antibacterial activities of *Gardenia ternifolia* and *Uapaca heudelotii*.

Study design: Evaluation of the antisickling and antibacterial activities of anthocyanins and organic acids extracted from *Gardenia ternifolia* and *Uapaca heudelotii*. *In vitro*

Place and Duration of Study: Faculty of Science, University of Kinshasa, between November 2012 and April 2013.

Methodology: The antisickling and antibacterial activities of anthocyanins and organic acids extracted from *Gardenia ternifolia* and *Uapaca heudelotii* were assessed using Emmel's and disc diffusion and micro-dilution methods respectively. The disc diffusion method was used to determine the antibacterial activity of extracts while the micro-dilution method was performed to determine their MIC and MBC.

Results: The present study revealed that anthocyanins and organic acids extracts from *G. ternifolia* and *U. heudelotii* possess antisickling and antibacterial activities. All tested extracts from *U. heudelotii* displayed interesting antisickling and antibacterial effects. At the extract dose of 6.25 µg/mL, the calculated normalization rates were 70% (for anthocyanins extract of *U. heudelotii*), 80% (for organic acids extract of *U. heudelotii*), 68% (for anthocyanins extract of *G. ternifolia*) and 72% (for organic acids extract of *G. ternifolia*). The reference bacterial strains *S. aureus* were more sensitive to anthocyanins extract: MIC = 31.25 (*U. heudelotii*) and 62.5 µg/mL (*G. ternifolia*) than the *E. coli* strains: MIC = 62.5 µg/mL (*U. heudelotii*) and 125 µg/mL (*G. ternifolia*).

Conclusion: This study provides a scientific basis for the antimicrobial and antisickling activities of anthocyanins and organic acids extracts from *Gardenia ternifolia* and *Uapaca heudelotii*. Isolation and purification of different phytochemicals may further yield significant antibacterial and antisickling new leads compounds.

Keywords: Sick cell disease, bacterial infections, Medicinal plants, drepanocyte normalization rate, MIC, Democratic Republic of the Congo.

Abbreviations: ACE: Anthocyanis extract, ATCC: American Type Culture Collection, DRC: Democratic Republic of the Congo, MBC: Minimum bactericidal concentration MHB: Mueller Hinton Broth, MIC: Minimum inhibitory concentration, OAE: organic acids extract, RBC: Red blood cell, SCD: Sick Cell Disease

1. INTRODUCTION

Microbial infections, especially those due to bacteria are recurrent pathologies of the Sick Cell Disease (SCD) and remain the leading cause of death in children patients (1, 2). Epidemiological data reveal that about 300,000 children are born each year in the world with pathological hemoglobin of which 70% are affected by SCD (3). Most of the patients die before the age of five years when they do not receive regular health care. This disease is endemic to tropical regions. In some African regions, carriers of S hemoglobin can reach 20% of the population with a prevalence of 25-30% in central Africa. In Democratic Republic of the Congo (DRC), more than one million of inhabitants or almost 2% of the population are sicklers.

Clinical management of SCD focuses mainly on prophylactic measures for alleviating the painful crises through administration of analgesics, antipyretics, oral antibiotics such as penicillin and the anticancer drug hydroxyurea as epigenetic modulator of fetal hemoglobin gene. Unfortunately, not only these treatments were not effective, they are also expensive for the poor African population and may present HIV / AIDS infection risk (4-6). Alternative strategy in the management of SCD is now focusing on the identification of the novel antisickling agents mainly that from medicinal plants. Indeed, traditional medicine continues to play a very significant role in the medical primary health care implementation in developing countries.

It is known that in SCD endemic regions, the use of herbs for the treatment of patients is a common practice. In previous studies of our research team, a number of medicinal plants traditionally used for the management of SCD in DRC were reported. These plant species have been scientifically validated for their beneficial effects in SCD condition including anti-sickling, anti-aggregating/anti-polymerization, radical scavenging (antioxidant) and anti-dehydrating effects. These activities are mainly due to anthocyanins and organic acids (7-10). In the present study, we hypothesized that the naturally occurring anthocyanins and organic acids of Congolese plants *G. tonatis* and *U. heudelotii* could also have antisickling as well as antibacterial properties.

Thus this research work was performed with the aim of evaluating the antisickling and antibacterial activities of anthocyanins and organic acids extracted from *Gardenia ternifolia* subsp. *jovis-tonantis* (Welw.) Verdc. (Rubiaceae) and *Uapaca heudelotii* Baill (Phyllanthaceae).

2. MATERIAL AND METHODS

2.1. Plant material collection and identification

The tested plant materials (leaves) used in this study were collected in Democratic Republic of the Congo during a field work in March 2011 and were authenticated by Mr B.L. Nlandu of the INERA (Institut National d'Etudes et Recherches Agronomiques). Vouchers specimens (*Gardenia ternifolia*: ref. N0 Ngb-BC002) and (*Uapaca heudelotii*: ref. N0 H. Breyné 2576) are on deposit at the INERA Herbarium of the Faculty of Science (University of Kinshasa).

2.2. Extraction and chemical screening

The dried and powdered plant material (10 g) was repeatedly extracted by cold percolation with 95% ethanol (EtOH) and water (100 mL, twice) for 48 hours. Chemical screening was performed in aqueous and organic extract according to as previously reported protocol (7, 12-14). Fractions were filtered and concentrated to dryness under reduced pressure using a rotary evaporator. Extraction of anthocyanins and organic acids was then done using 100 g of dried powdered plant material following an established protocol [4,15-18]. Anthocyanins extracts were then defatted by n-hexane and all extracts were stored at +4 °C.

2.3. Biological testing

2.3.1. In vitro antisickling bioassay

Blood samples used to assess the antisickling activity of the selected plant extracts were taken from known SCD patients attending the "Centre de Médecine Mixte et d'Anémie SS" located in Kinshasa, DRC. None of the patients had been transfused recently with Hb AA blood and all antisickling experiments were carried out with freshly collected blood. In order to confirm their SS nature, the above-mentioned blood samples were first characterized by Hemoglobin electrophoresis on cellulose acetate gel, as previously reported [1, 3-18]. They were found to be SS blood and were then stored at +4 °C in a refrigerator. An informed consent was obtained from all the patients participating in the study. All the research procedures have received the approval of Department of Biology Ethics Committee.

An aliquot of Hb S-blood was diluted with 150 mM phosphate buffered saline (NaH_2PO_4 30 mM, Na_2HPO_4 120 mM, NaCl 150 mM) and mixed with an equivalent volume of 2% sodium metabisulfite. A drop from the mixture was spotted on a microscope slide in the presence or absence of anthocyanins or organic acids extracts and covered with a cover slip. Paraffin was applied to seal the edges of the cover completely to exclude air (Hypoxia). Duplicate analyses were run for each extract. The red blood cells (RBCs) were analyzed by a computer assisted image analysis software (Motic Images 2000, version 1.3; Motic China Group Co LTD) and statistical data analysis were processed using Microcal Origin 7.1 package software as previously reported (1, 3-18).

2.3.2. Determination of antibacterial activity

- Microbial strains and media used

The activity of the plant samples was tested toward *Lactobacillus fermentum* (L. fermentum ATCC 9338), *Staphylococcus aureus* (S. aureus ATCC 25923), *Enterococcus faecalis* (E. faecalis ATCC 19433), *Salmonella typhimurium* (S. typhimurium ATCC 13311) and *Escherichia coli* (E. coli ATCC 25922). The tested strains were obtained from the

Comment [L1]: Select references used accurately.

84 American Type Culture Collection (ATCC, Rockville MD, USA). Mueller Hinton agar and Mueller Hinton Broth purchased
85 from Conda (Madrid, Spain) were used as media

86 - Disc diffusion method

87 The agar disc diffusion method was used to determine the antibacterial activity of plant extracts as follow: A 1 mL of
88 suspension of an 18 hours culture bacteria containing about 1×10^8 colony-forming units per milliliter (CFU/mL) were
89 spread on Mueller Hinton agar medium using sterile swabs. Filter paper discs (6 mm in diameter) were soaked in 10 μ L of
90 extracts and placed on the inoculated plates and allowed to dry for 30 min, then incubated at 37 °C for 24 hours. The
91 diameters of the inhibition zones were measured in millimeters (19). Two controls were included in the test: the first was a
92 control involving the presence of microorganisms but without the test extract sample and the last was two standard
93 antibiotics (gentamicin disc, 10 μ g/mL for gram-positive bacteria and ofloxacin disc, 20 μ g/mL for gram-negative
94 bacteria). Studies were performed in triplicate, and the developing inhibition zones were compared with those of reference
95 discs.

96 - Determination of Minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC)

97 The Minimum inhibitory concentration was determined by broth micro-dilution method as previously reported (1, 20). The
98 inocula of used microorganisms were prepared from 24 hours old broth cultures. The absorbance was read at 600 nm and
99 adjusted with sterile physiological solution to match that of a 0.5 McFarland standard solution. From the prepared
100 microbial solutions, other dilutions with sterile physiological solution were prepared to give a final concentration of 10^6
101 CFU/mL. Stock solutions of the extracts were prepared in 0.1% (v/v) aqueous tween 80 (Fisher chemicals) at
102 concentrations of 1 mg/mL. The two-fold serial dilutions in concentrations of the extracts were prepared in Mueller Hinton
103 Broth (MHB) (Conda, Madrid, Spain) to give final concentrations ranging from 250 to 1.95 μ g/mL.

104 An aliquot (10 μ L) of a 1×10^6 CFU/mL overnight culture was added to wells of a sterile 96-well micro-plate titer. The
105 positive control wells contained MHB+ bacteria suspension without plant extract while negative control wells contained
106 MHB only. The MIC was determined as the lowest plant extract concentration at which no growth were observed after 24
107 hours. MTT (30 μ L) in aqueous solution (0.01%) was used to evaluate the micro-organism viability. For MBC
108 determination, 10 μ L was taken from each well of complete inhibition of bacterial growth after incubation and spot
109 inoculated on freshly prepared MHB and incubated for 72 hours at 37 °C. The concentration at which no growth was
110 observed on subculture was determined as the MBC.

113 3. RESULTS AND DISCUSSION

116 3.1. Chemical screening

118 The results of chemical screening of *G. ternifolia* and *U. heudelotii* are presented in Table 1.

120 Table 1: Results of chemical screening of *G. ternifolia* and *U. heudelotii*


Chemical groups	Medicinal plant species (used parts)	
	<i>G. ternifolia</i> (leaves)	<i>U. heudelotii</i> (leaves)
Total polyphenols	+	+
Anthocyanins	+	+
Leuco-anthocyanins	+	+
Flavonoids	+	+
Gallic tannins	+	+
Catechic tannins	+	+
Coumarines	+	+
Quinones	+	+
Alkaloids	+	+
Saponins	+	+
Terpenes and steroids	+	+

122 It is deduced from the table 1 that the leaves of both *G. ternifolia* and *U. heudelotii* contain total polyphenols, catechic
123 tannins, alkaloids, saponins, terpenes, steroids, anthocyanins, leuco-anthocyanins, flavonoids, gallic tannins, coumarins
124 and quinines. Compounds which are present in these plant species, are well known for their broad spectrum of
125 pharmacological properties, including antimicrobial, antisickling and antioxidant activities (3-18, 21-24). Hence, the
126 presence of various secondary metabolites in these plants could justify their use as medicine in Africa.

Comment [L2]: Select the references used accurately

129 **3.2. Antisickling activity**


130
131 Figures 1a, b, c, and d show the optical micrograph phenotypes of SS blood alone (a) or treated with anthocyanins (b and
132 c) and organic acids crude extracts (d and e) extracted from the two selected plants.
133



134 1 (a)




136 1 (b)



138 1 (c)



1 (d)



1 (e)

Fig.1: Phenotype of SS red blood cells (SS RBCs) [NaCl 0.9% ; Na₂S₂O₅ 2%; X500]: (a): untreated SS RBCs (control); (b) SS RBCs treated with anthocyanins extract of *G. ternifolia* (20 µg/mL); (c): SS RBCs treated with anthocyanins extract of *U. heudelotii* (20 µg/mL); (d): SS RBCs treated with organic acids extract of *G. ternifolia* (20 µg/mL); (e): SS RBCs treated with organic acids extract of *U. heudelotii* (20 µg/mL).

Figure 1 shows that the control (a) contains in majority sickle-shaped RBCs, confirming the SS nature of the blood used. Mixed together with anthocyanins and organic acids extracts (b-d), the majority of SS RBCs are reversed normal-shape. At the extract dose of 6, 25 µg/mL, the calculated normalization rates were found to be 70% (for anthocyanins extract of *U. heudelotii*), 80% (for organic acids extract of *U. heudelotii*), 68% (for anthocyanins extract of *G. ternifolia*) and 72% (for organic acids extract of *G. ternifolia*) as it can be seen on the figure 2.




Fig.2: Evolution of the normalization rate of sickle erythrocytes with dose extracts from selected medicinal plant species (NaCl 0,9%; Na2S2O5 2%; X500). %Normalization rate = [number of drepanocytes of untreated SS blood (control) - number of drepanocytes of treated SS blood (sample)/ number of drepanocytes of untreated SS blood] ×100.

Our results indicated that anthocyanins and organic acids are the major antisickling agents of the tested plants. These results confirm those already reported by our research team on anthocyanins and organic acids from other plants used in Congolese folk medicine for the management of SCD (3-12). As previously reported, anthocyanins have the ability to interact with proteins (15). Interaction of these secondary metabolites with hemoglobin S could chemically compete with the polymerization of this abnormal hemoglobin thus preventing the sickling of sickle erythrocytes. In addition, anthocyanins (for which intestinal catabolism gives phenolic acids) are also known for their antioxidant properties and could affect the Fe³⁺/Fe²⁺ higher ratio in sickle cells and the stability of erythrocytes membrane by preventing the oxidation of membranes phospholipids (16).

3.3. Antibacterial activity

The antibacterial activity of extracts from selected plants against *L. fermentum*, *S. aureus*, *E. faecalis*, and *S. typhimurium* was determined. The results are shown in Table 2.

Table 2: Bacterial inhibitory growth effects of extracts from *G. ternifolia* and *U. heudelotii*

Reference bacterial strains	<i>Gardenia ternifolia</i>						<i>Uapaca heudelotii</i>					
	Anthocyanins extract			Organic acids extract			Anthocyanins extract			Organic acids extract		
	(µg/mL)			(µg/mL)			(µg/mL)			(µg/mL)		
	25	50	100	25	50	100	25	50	100	25	50	100
<i>L. fermentum</i>	-	+	+	+	+	+	-	-	+	-	+	+
<i>S. aureus</i>	-	+	+	-	-	+	+	+	+	+	+	+
<i>E. faecalis</i>	-	-	+	-	-	-	-	+	+	+	+	+
<i>S. typhimurium</i>	-	+	+	+	+	+	-	-	+	+	+	+
Activity (%)	58						75					

Legends - : Inactive extract (inhibition zone = 0 mm); +: Active extract (inhibition zone > 15 mm);
Positive control: gentamicin 10 µg/mL (inhibition zone = 19 mm), ofloxacin, 20 µg/mL (inhibition zone = 26 mm); %S: sensitivity percentage.

The antibacterial activity of extracts is closely dependent on the tested plants and the reference bacterial strains. Overall, *Uapaca heudelotii* was more active than *Gardenia ternifolia* (% average activity equal to 75 vs 58) (table 2). It should also be noted that *L. fermentum* and *S. typhimurium* showed inhibition zones to all *G. ternifolia* tested extracts while *S. aureus* and *E. faecalis* showed in contrary more less inhibition zones toward *U. heudelotii* extracts. Considering the strains globally, it should be noted that *E. faecalis* was the bacteria strain which has been showed less number of inhibition zones. *E. faecalis* is a Gram-positive bacterium belonging to the family of *Enterococcaceae* which is reported in the literature to be involved in drug resistance towards antibiotics such as aminoglycosides, cephalosporins, clindamycin, oxacillin and cotrimoxazole (25). However, it should be noted that at low dose (25 µg/mL), organic acids extracts of *U. heudelotii* inhibited the growth of this bacterium. This result showed that the secondary metabolites from the plants under study can serve as a source of new antibacterial agents. The results revealed also that *S. aureus* was the most sensitive strain.

3.4. Determination of Minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC)

The results of MIC and MBC are shown in Table 3.

Table 3: Minimum inhibitory concentration (MIC) and the minimum bactericidal concentration (MBC) of *Gardenia ternifolia* and *Uapaca heudelotii*.

Reference bacterial strains	MIC/MBC (µg/mL)			
	<i>Gardenia ternifolia</i>		<i>Uapaca heudelotii</i>	
	ACE	OAE	ACE	OAE
<i>E. coli</i> ATCC 25922	125/500	125/>500	62.5/250	125/>500
<i>S. aureus</i> ATCC 25923	62.5/250	125/>500	31.25/62.5	125/>500

Legends: ACE Anthocyanis extract; OAE Organic acids extract

The Gram positive bacterium *S. aureus* was more sensitive to Anthocyanis extract (ACE) than the Gram negative *E. coli*. (table 3). The higher sensitivity of Gram-positive bacteria could be attributed to their outer peptidoglycan layer which is not an effective permeability barrier. Gram-negative bacteria possessing an outer phospholipidic membrane carrying the structural lipopolysaccharide components make the cell wall impermeable to lipophilic solutes while porins constitute a selective barrier to hydrophilic solutes with an exclusion limit of 600 Da (20, 26). *S. aureus* is the bacteria most implicated in septicemia and osteomyelitis in SCD patients (2). Medicinal plant extracts displaying at the same time antibacterial (against *Staphylococcus aureus*) and antisickling effects could be useful in the management of Sickle cell disease. It has been reported that human gut microflora have the ability to metabolize anthocyanins into smaller phenolic acids which are more bioavailable and stable end-products (27). As anthocyanins metabolic stable end-products, they could have less antibacterial activity than their parents. The present study confirmed this hypothesis. As SCD is a chronic disease, using anthocyanins as medicinal foods or nutraceuticals would be a good approach instead of giving pharmaceutical products to sicklers during all their life. Such important nutraceuticals will contribute to solve health problem.

4. CONCLUSION

The present study provided evidence for the antibacterial and antisickling activities of anthocyanins and organic acids extracts from *Gardenia ternifolia* and *Uapaca heudelotii* and bring working data for future investigations that will lead to the characterization of chemical structure of anthocyanin and organic acid extracts from these plants.

ACKNOWLEDGMENTS

This research was founded by the International Foundation for Science (IFS, Stockholm, Sweden) and the Organization for the Prohibition of Chemical Weapons (OPCW) (IFS Research Grant N0 F/4921-2), research grant offered to Dr. Koto -te- Nyiwa Ngbolua.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

AUTHORS' CONTRIBUTIONS

Authors KN Ngbolua and DST Tshibangu wrote the first draft of the manuscript, PT Mpiana designed the study, SO Mihigo wrote the protocol, BK Mavakala and MC Ashande managed the analyses of the study and LC Muanyishay managed the literature searches. All authors read and approved the final manuscript

ETHICAL APPROVAL

All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable. All experiments have been examined and approved by the appropriate ethics committee

REFERENCES

1. Ngbolua KN, Mubindikila RN, Mpiana PT, Tshibangu DST, Ashande MC, Nzongola WXX, Baholy R, Fatiany PR. Phytochemical screening, Antibacterial and Antioxidant activities of *Anthocleista liebrechtsiana* Wild & T. Durand (Gentianaceae) originated from Democratic Republic of the Congo. Journal of Advancement in Medical and Life Sciences 2014; 1(3): DOI: 10.15297/JALS.V1I3.04.
2. Diagne I, Diagne-Gueye ND, Signate-Sy H, Camara B, Lopez-Sall P, Diack-Mbaye A, Sarr M, Ba M, Sow HD, Kuakivi N. Management of children with sickle cell disease in Africa: experience in a cohort of children at the Royal Albert Hospital in Dakar. Médecine Tropicale 2003; 63(4-5): 513-520.
3. Mpiana PT, Tshibangu DST, Shetonde OM, Ngbolua KN. *In vitro* antitrepanocytary activity (anti-sickle cell anaemia) of some Congolese plants. Phytomedicine 2007; 14, 192-195.
4. Mpiana PT, Mudogo V, Tshibangu DST, Kitwa EK, Kanangila AB, Lumbu JBS, Ngbolua KN, Atibu EK, Kakule MK. Antisickling Activity of Anthocyanins from *Bombaxpentadrum*, *Ficuscapensis*, *Zizyphusmucronata*: Photo-degradation effect. Journal of Ethnopharmacology 2008;120: 413-418.
5. Mpiana PT, Mudogo V, Tshibangu DST, Ngbolua KN, Shetonde OM, Mangwala PK, Mavakala BK. *In vitro* Antisickling Activity of Anthocyanins Extracts of a Congolese Plant: *Alchornea cordifolia* M. Arg. Journal of Medical Sciences 2007;7 (7): 1182-1186.
6. Mpiana PT, Mudogo V, Ngbolua KN, Tshibangu DST, Shetonde OM, Mbala BM. *In vitro* Antisickling Activity of Anthocyanins from *Ocimumbasilicum* L. (Lamiaceae). International Journal of Pharmacology 2007;3 (4): 371-374.
7. Ngbolua KN, Bishola TT, Mpiana PT, Mudogo V, Tshibangu DST, Ngombe KN, Tshilanda DD, Baholy R. *In vitro* antisickling and free radical scavenging activities of *Pentaclethra macrophylla* Benth. (Fabaceae). Journal of Advancement in Medical and Life Sciences 2014; 2(3): DOI: 10.15297/JALS.V1I2.03.
8. Ngbolua KN, Bishola TT, Mpiana PT, Mudogo V, Tshibangu DST, Ngombe KN, Ekutsu EG, Gbolo ZB, Kabena NO. Ethno-pharmacological survey, *in vitro* antisickling and free radical scavenging activities of *Carapa procera* DC. stem bark (Meliaceae). Nova Journal of Medical and Biological Sciences 2014; 2(2): 01-14.
9. Ngbolua KN, Bishola TT, Mpiana PT, Mudogo V, Tshibangu DST, Ngombe KN, Ekutsu EG, Tshilanda DD, Gbolo ZB, Mwanangombo DT, Fatiany PR, Baholy R. Ethno-botanical survey, *in vitro* antisickling and free radical scavenging activities of *Garcinia punctata* Oliv. (Clusiaceae). Journal of Advanced Botany & Zoology. 2014;

280 2(4): DOI: 10.15297/JABZ.V1I2.04.

281 10. Ngbolua KN, Mudogo V, Mpiana PT, Malekani MJ, Rafatro H, Urverg Ratsimamanga, Takoy L, Rakotoarimana
282 H, Tshibangu DST. Evaluation de l'activité anti-drépanocytaire et antipaludique de quelques taxons végétaux
283 de la République démocratique du Congo et de Madagascar. *Ethnopharmacologia* 2013; 50: 19-24.

284 11. Mpiana PT, Misakabu FM, Yuma PM, Tshibangu DST, Ngbolua KN, Muanyishay CL, Gbolo BZ, Misengabu NM,
285 Kayembe JS. Antisickling activity and physico-chemical stability of anthocyanins extracts from *Ipomoea*
286 *batatas* leaves. *Journal of Life Medicine* 2014; 2(1): 25-31.

287 12. Yuma PM, Mpiana PT, Bokota MT, Wakenge IB, Muanishay CL, Gbolo BZ, Mathina Di Mathina G, Tshibangu
288 DST, Ngbolua KN. Etude de l'activité anti-falcémiant et de la thermo et photo-dégradation des anthocyanes
289 de *Centella asiatica*, *Thomandersia hensii* et *Maesopsis eminii*. *International Journal of Biological & Chemical*
290 *Sciences* 2013;7(5): 1892-1901.

291 13. Mpiana PT, Kimbadi BL, Ombeni AM, Ngbolua KN, Tshibangu DST, Tshilanda DD, Misengabu NM, Muanyishay
292 CL, Muyisa SK. *In vitro* inhibitory effects and anti-sickle erythrocytes haemolysis of *Dicliptera colorata* C.B.
293 Clarke, *Euphorbia hirta* L. and *Sorghum bicolor* (L.) Moench. *Open Journal of Blood Diseases* 2013; 3: 43-48.

294 14. Mpiana PT, Dianzenza EN, Ngbolua KN, Tshibangu DST, Mbala BM, Shetonde OM, Atibu EK, Kakule MK,
295 Bokota MT. Antisickling properties, thermal and photochemical degradations of anthocyanins extract from
296 *Annonasenegalensis* (Annonaceae). *International Journal of Biological & Chemical Sciences* 2012; 6 (5):
297 2241-2251.

298 15. Mpiana PT, Ngbolua KN, Mudogo V, Tshibangu DST, Atibu EK, Tshilanda DD, Misengabu NM. Antisickle
299 erythrocytes haemolysis properties and inhibitory effect of anthocyanins extracts of *Trema orientalis*
300 (ULMACEAE) on the aggregation of human deoxyhemoglobin S *in vitro*. *Journal of Medical Sciences* 2011;11
301 (3): 129-137.

302 16. Mpiana PT, Ngbolua KN, Bokota MT, Kasonga TK, Atibu EK, Mudogo V. *In vitro* Effects of Anthocyanins Extracts
303 From *Justicia secunda* VAHL on the Solubility of Hemoglobin S and Membrane Stability of Sickle Erythrocytes,
304 *Blood Transfusion* 2010; 8: 248-254.

305 17. Mpiana PT, Bokota MT, Mbula JP, Ngbolua KN, Tshibangu DST, Atibu EK, Kwembe JTK, Kakule MK. Effet of
306 Anthocyanins Extracts from *Justicia matammensis* and *Justicia laxa* on Sickle cells. In: Anthocyanins,
307 Structure, Biosynthesis and Health benefits (Series: Biochemistry Research Trends, Nutritional and Diet
308 Research Progress); Noboru Motohashi (ed.), 2013; 111-124. NOVA PUBLISHERS, New York, USA.

309 18. Tshibangu DST, Shode FO, Koorbanally N, Mudogo V, Mpiana PT, Ngbolua KN. Antisickling triterpenoids from
310 *Callistemon viminalis*, *Meulaleuca bracteata* var. *Revolution Gold* *Syzygium guineense* and *Syzygium cordatum*.
311 The 14th NAPRECA Symposium and AAMPS Ethno-veterinary Medicine Symposium 8th-12th August 2011.
312 International Centre For Insect Physiology and Ecology (ICIPE): Kasarani, Nairobi, Kenya, pp.296-300 (YS
313 27).

314 19. Randrianirina AYO, Rajaonarivelo PJ, Rasondratovo B, Raharisololalao A, Moulis C, Mudogo V., Mpiana PT.,
315 Ethno-botanical survey, Chemical composition and *in vitro* Antimicrobial activity of essential oils from the root
316 bark of *Hazomalania voyroni* (Jum.) Capuron (Hernandiaceae). *Journal of Advancement in Medical and Life*
317 *Sciences* 2014 1(1): DOI: 10.15297/JALS.V1I1.11.

318 20. Ngbolua KN, Mubindukila RN, Mpiana PT, Masengo CA, Baholy R, Fatiany PR, Ekutsu GE, Gbolo ZB. *In vitro*
319 Assessment of Antibacterial and Antioxidant activities of a Congolese medicinal plant species *Anthocleista*
320 *schweinfurthii* Gilg (Gentianaceae). *J. of Modern Drug Discovery and Drug Delivery Research*. 2014;
321 3(3): DOI: 10.15297/JMDDR.V1I3.03.

322 21. Proestos C, Chorianopoulos N, Nychas GJ, Komaitis M. RP-HPLC analysis of the phenolic compounds of plant
323 extracts: Investigation of their antioxidant capacity and antimicrobial activity. *J Agr Food Chem*. 2005; 53:
324 1190-95.

325 22. Slobodnikov L, Kost'aloova D, Labudova D, Kotulova D, Kettmann V. Antimicrobial activity of *Mahonia aquifolium*
326 crude extract and its major isolated alkaloids. *Phytother Res*. 2004; 18: 674-76.

327 23. Chung KT, Wong TY, Wei CI, Hung YW, Lin Y. Tannins and human health: A review. *Crit Rev Food Sci Nutr*. ,
328 1998; 38: 421-64.

- 329 24. Mandal P, Babu SSP, Mandal NC. Antimicrobial activity of saponins from *Acacia auriculiformis*. *Fitoterapia* 2005;
330 76:462–65.
- 331 25. Martinko J, Madigan M. *Biologie des micro-organismes*, 11^e éd. Pearson Education, Paris, 2007
- 332 26. Kaur GJ, Arora DS. Antibacterial and phytochemical screening of *Anethum graveolens*, *Foeniculum vulgare* and
333 *Trachyspermum ammi*. *BMC Complementary and Alternative Medicine* 2009; 9: 30.
- 334 27. He J, Magnuson BA, Lala G, Tian Q, Schwartz SJ and Giusti MM. Intact Anthocyanins and Metabolites in Rat
335 Urine and Plasma after 3 Months of Anthocyanin Supplementation. *Nutrition & Cancer* 2004; 54(1): 3–12

286

337
338
339