

HISTOLOGICAL AND HISTOCHEMICAL EVALUATION OF ANTI-CADMIUM TOXICITY EFFECTS OF *MORINGA OLEIFERA* SEED OIL AND *ANACARDIUM OCCIDENTALE* NUT OIL IN THE HIPPOCAMPUS OF JUVENILE MALE WISTAR RATS

Abstract

The hippocampus is the part of the brain that is involved in memory formation, organization and storage, being a limbic system structure that is particularly important in forming new memories and connecting emotions and senses, such as smell and sound, to memories. Cadmium on the other hand is a metallic poison that affects the brain tissues as a whole producing seriously deleterious effects. It is believed that antioxidants could ameliorate the negative effects of toxic substances on body tissues. Since *Moringa oleifera* seed oil and *Anacardium occidentale* nut oil are natural edible oils which are greatly rich in antioxidants and medicinal phytochemicals, it is important to observe their potentials in helping the body to combat the effects of cadmium toxic effects particularly on the hippocampus. A total of 35 juvenile male Wistar rats were used for the study. They were the rats were randomly grouped into four A, B, C, D, E, F and G, each containing 5 animals. Group A, Group B and Group C animals were given 3ml per kg body weight of 0.9% NaCl solution intraperitoneally for 3 weeks; each rat in group B was induced intraperitoneally with a single dose of 2.5mg/kg body weight of 3CdSO₄.8H₂O, and in group C, each rat was given 5mg/kg body weight vitamin C dissolved in 0.9% NaCl solution and 6mg/kg body weight vitamin E dissolved in olive oil orally for 3 weeks. Group D animals were given 5mg/kg body weight vitamin C dissolved in 0.9% NaCl solution and 6mg/kg body weight vitamin E dissolved in olive oil orally for 3 weeks, after which each rat was induced intra-peritoneally with a single dose of 2.5mg/kg body weight of 3CdSO₄.8H₂O. Groups E, F and G were first given 2.5mg/kg body weight of 3CdSO₄.8H₂O, and after 48 hours, treatment began. Group E rats were treated with 4mg/kg body weight *Moringa Oleifera* seed oil each, group F rats were treated with 4mg/kg body weight *Anacardium occidentale* nut oil and group G rats were treated with 2mg/kg body weight *Moringa Oleifera* seed oil + 2mg/kg body weight *Anacardium Occidentale* nut oil. Cadmium intoxication has observable deleterious effects on brain tissues as a result of intra-peritoneal administration of cadmium sulphate. All agents used produced positive anti-cadmium effects, however to varying degrees. While all agents ameliorated the effects of cadmium on neuronal morphology and relative population cum distribution, moringa seed oil was the most potent in reducing the formation of senile plaques in the hippocampus. The potency combined effect of moringa oleifera seed oil and *Anacardium occidentale* nut oil in reducing or treating the oxidative damage caused by cadmium on the hippocampus is also observed. The natural oils were more potent than the selected regularly used antioxidants.

Key Words: Hippocampus, Cadmium Toxicity, Antioxidants, *Moringa oleifera*, *Anacardium occidentale*

Introduction

The hippocampus is believed to play important role in several functions of the body. The hippocampus is the part of the brain that is involved in memory formation, organization and storage (Squire, 1992; Eichenbaum and Cohen, 1993). Hippocampus is a limbic system structure that is particularly important in forming new memories and connecting emotions and senses, such as smell and sound, to memories. The hippocampus acts as a memory indexer by sending memories out to the appropriate part of the cerebral hemisphere for long-term storage and retrieving them when necessary (Squire, 1992; Eichenbaum and Cohen, 1993). Cadmium on the other hand is a metallic poison that affects the brain tissues as a whole producing seriously deleterious effects. It is believed that antioxidants could ameliorate the negative effects of toxic substances on body tissues. Since *Moringa oleifera* seed oil and *Anacardium occidentale* nut oil are natural edible oils which are greatly rich in antioxidants and medicinal phytochemicals, it is important to observe their potentials in helping the body to combat the effects of cadmium toxic effects particularly on the hippocampus.

The hippocampus is composed of multiple subfields. Though terminology varies among authors, the terms most frequently used are dentate gyrus and the cornu ammonis literally "Amun's horns", abbreviated CA. The dentate gyrus contains the fascia dentata and the hilus, while CA is differentiated into fields CA1, CA2, and CA3 (Chris, 2008). The horse shoe appearance of the hippocampus is caused by cell density differentials and the existence of varying degrees of neuronal fibers. The CA regions are also structured depth-wise in clearly defined strata (Philippe, 2007; Hiroaki, 2007; Annabelle, 2009). The alveus is the deepest layer and contains the axons from pyramidal neurons, passing on towards the fimbria/fornix, one of the major outputs of the hippocampus. Stratum oriens is the next layer superficial to the alveus (Annabelle, 2009). The cell bodies of inhibitory basket cells and horizontal trilaminar cells, named for their axons innervating the three layers-the oriens, pyramidal, and radiatum are located in this stratum (Annabelle, 2009). The basal dendrites of pyramidal neurons are also found here, where they receive input from other pyramidal cells, septal fibers and commissural fibers from the contra lateral hippocampus usually recurrent connections, especially in CA3 and CA2 regions (Annabelle, 2009).

Stratum pyramidale contains the cell bodies of the pyramidal neurons, which are the principal excitatory neurons of the hippocampus. This stratum tends to be one of the more visible strata to the naked eye (Annabelle, 2009). In region CA3, this stratum contains synapses from the mossy fibers that course through stratum lucidum. This stratum also contains the cell bodies of many interneurons, including axo-axonic cells, bistratified cells, and radial trilaminar cells (Chris, 2008). Stratum lucidum is one of the thinnest strata in the hippocampus and only found in the CA3 region. Mossy fibers from the dentate gyrus granule cells course through this stratum in CA3, though synapses from these fibers can be found in stratum pyramidale. Stratum radiatum like stratum oriens, contains septal and commissural fibers. It also contains Schaffer collateral fibers, which are the projection forward from CA3 to CA (Chris, 2008). Some interneurons that can be found in more superficial layers can also be found here, including basket cells, bistratified cells, and radial trilaminar cells (Chris, 2008). Stratum lacunosum is a thin stratum that too contains Schaffer collateral fibers, but it also contains perforant path fibers from the superficial layers of entorhinal cortex. Due to its small size, it is often grouped together with stratum moleculare into a single stratum called stratum lacunosum-moleculare (Chris, 2008). Stratum moleculare is the most superficial stratum in the hippocampus. Here the perforant path fibers form synapses onto the distal, apical dendrites of pyramidal cells (Chris, 2008). The hippocampal

sulcus or fissure is a cell-free region that separates the CA1 field from the dentate gyrus. Because the phase of recorded theta rhythm varies systematically through the strata, the fissure is often used as a fixed reference point for recording EEG as it is easily identifiable (Chris, 2008).

The dentate gyrus is composed of a similar series of strata namely; Polymorphic layer is the deepest layer of the dentate gyrus and is often considered a separate subfield (Annabelle, 2009). This layer contains many interneurons, and the axons of the dentate granule cells pass through this stratum on their way to CA3 (Annabelle, 2009). Stratum granulosum contains the cell bodies of the dentate granule cells. Stratum moleculare, outer third is where both commissural fibers from the contralateral dentate gyrus run and form synapses as well as where inputs from the medial septum terminate on the proximal dendrites of the granule cells. (Gross, 1993; Chris, 2008). Stratum moleculare, external two thirds is the deepest of the strata, sitting just superficial to the hippocampal fissure across from stratum moleculare in the CA fields. The perforant path fibers run through this stratum, making excitatory synapses with the distal apical dendrites of granule cells (Gross, 1993; Chris, 2008).

Cadmium is a soft, malleable, ductile, bluish-white divalent metal. It is similar in many respects to zinc but forms complex compounds (Holleman, 1985). Unlike other metals, cadmium is resistant to corrosion and as a result it is used as a protective layer when deposited on other metals. As a bulk metal, cadmium is insoluble in water and is not flammable; however, in its powdered form it may burn and release toxic fumes (CSEM, 2011). Cadmium cannot penetrate the adult blood brain barrier (BBB), although it might diffuse across the BBB with the help of a vehicle such as ethanol (Pal *et al.*, 1993). Cadmium can more effectively pass the BBB during the developmental stage in an organism and is more toxic in newborns (Sinha *et al.*, 2008; Wong and Klaassen, 1982). Cadmium has an LD50 of 5.7mg/Kg per body weight intraperitoneally on rats and once inside, it accumulates in different areas of the brain, induces lipid peroxidation and weakens the antioxidative defence (Sinha *et al.*, 2008; Gutierrez *et al.*, 1999). In battery workers Cd-induced oxidative stress was demonstrated to cause amyotrophic lateral sclerosis due to reduced brain SOD activity (Bar-sela *et al.*, 2001). Cadmium (0.4 mg CdAc₂/kg body weight) injected intraperitoneally to young albino rats for 30 days generated free radicals in the brain causing region-specific membrane changes, which in turn led to significant alterations in membrane fluidity, intracellular Calcium concentrations and phospholipid composition (Kumar *et al.*, 1996). It also resulted in a decreased GSH/GSSG ratio as well as activities of GR and glucose-6-phosphate-dehydrogenase (G6PDH) in various brain regions, although the decrease in GSH/GSSG was not seen in the hippocampus and midbrain (Shukla *et al.*, 1999). Cadmium-induced oxidative damage also induced enhanced lipid peroxidation and protein carbonylation in male Swiss albino mice that received 4 mg CdCl₂/kg body weight orally for three days (Sinha *et al.*, 2008). The oxidative impairment was characterised by increased ROS production, reduction of total thiols and the GSH pool together with an increase in GSSG level. In addition, also activities of antioxidant enzymes such as SOD, CAT, GST, GR, GPx and G6PDH were diminished. The authors also showed the protective abilities of taurine (single oral dose of 100 mg/kg body weight for five days before Cd treatment) and vitamin C (single oral dose of 100 mg/kg body weight for five days before Cd treatment) against oxidative impairment in brain tissue caused by Cd (Sinha *et al.*, 2008).

Moringa is a rich source of antioxidant (Chumark *et al.*, 2008) and has an LD50 of 340 – 400mg/Kg per body weight intraperitoneally on rats (Chumark *et al.*, 2008). It has been reported that aqueous extracts of leaf, fruit and seed of Moringa act as an antioxidant (Singh *et al.*, 2009). During a study reporting antioxidant property of freeze dried Moringa leaves from different

extraction procedures, it was found that methanol and ethanol extracts of Indian origin *Moringa* have the highest antioxidant activity with 65.1 and 66.8%, respectively (Lalas and Tsaknis, 2002; Siddhuraju and Becker, 2003). It was also reported that the major bioactive compounds of phenolics, such as quercetin and kaempferol are responsible for antioxidant activity (Bajpai *et al.*, 2005; Siddhuraju and Becker, 2003). During another study, quercetin and kaempferol have shown good antioxidant activity on hepatocyte growth factor (HGF) induced Met phosphorylation with IC50 value for 12 and ~6 $\mu\text{M/L}$, respectively (Labbe *et al.*, 2009). Another recent study comparing palm oil with *Moringa* seeds for their antioxidant potential found out that *Moringa* seed are superiors for radical scavenging (Ogbunugafor *et al.*, 2011).

The antioxidant properties and the effect on nitric oxide (NO) production of various extracts of leaves of *Anacardium occidentale*. The result suggested that the leaves of *A.occidentale* are a potent source of natural antioxidants (Jaiswal *et al* 2010). Cashew nut oil has an LD50 of 1230mg/kg body weight on rats intraperitoneally (Jaiswal *et al* 2010). Effects of immature cashew nut-shell liquid (*Anacardium occidentale*) against oxidative damage in *Saccharomyces cerevisiae* and inhibition of acetylcholinesterase activity. Study showed immature cashew nut shell liquid (iCNSL), a source of unsaturated long-chain phenols, and may have a potential role in protecting DNA against oxidative damage (Isao, 2005).

Materials and Methods

Moringa seed, 2000g , was produced from kuchikau town, Nassarawa state, Nigeria. 2000g of the powdered *Moringa oleifera* seed-oil was extracted by the following procedure. The seeds were heated inside an oven at temperature of 40⁰C and were later pounded with mortar and pestle to separate the chaff from the seeds. Finally the seed extract was grinded into powder with a grinding mill. The powdered extract of *Moringa Oleifera* was dissolved in 400ml at ambient temperature for two days and was later filtered. The aqueous extract was poured into molten mesh and was placed on an oil extractor machine, seed oil was removed at high temperature and pressure and oil extract was kept at -20⁰C before administration. 2000g of Cashew nut was produced from kuchikau town, Nassarawa state, Nigeria. 2000g of the powdered Cashew nut-oil was extracted by the following procedure. The seeds were heated inside an oven at temperature of 40⁰C and the seed extract was grinded into powder with a grinding mill. The extract was poured into molten mesh and was placed on oil extractor machine, seed oil was removed at high temperature and pressure and oil extract was kept at -20⁰C before administration.

The Cadmium Sulphate solution was prepared by dissolving the Cadmium Sulphate salt in normal saline solution. The Ascorbic acid (Vitamin C) was dissolved in normal saline solution and the Vitamin E (tocopherol) was dissolved in olive oil. All other chemicals were of the highest purity commercially available. The Cadmium Sulphate was administered at sub lethal dose according to the body weights of the rats intraperitoneally using insulin syringes. The Ascorbic acid and Alpha Tocopherol were administered orally using an oral cannula with a ball point at the tip.

A total of 35 juvenile male Wistar rats (100-200g) were used for the study. They were maintained under standard laboratory condition in the animal house (National Academy Press, 1996) the rats were randomly grouped into four A, B, C, D, E, F and G, each containing 5 animals. Group A, Group B and Group C animals were given 3ml per kg body weight of 0.9% NaCl solution intraperitoneally for 3weeks; each rat in group B was induced intraperitoneally with a single dose of 2.5mg/kg body weight of 3CdSO₄.8H₂O, and in group C, each rat was given 5mg/kg body weight vitamin C dissolved in 0.9% NaCl solution and 6mg/kg body weight

vitamin E dissolved in olive oil orally for 3 weeks. Group D animals were given 5mg/kg body weight vitamin C dissolved in 0.9% NaCl solution and 6mg/kg body weight vitamin E dissolved in olive oil orally for 3 weeks, after which each rat was induced intraperitoneally with a single dose of 2.5mg/kg body weight of $3\text{CdSO}_4 \cdot 8\text{H}_2\text{O}$, and after 48 hours, the rats were sacrificed. Groups E, F and G were first induced intraperitoneally with 2.5mg/kg body weight of $3\text{CdSO}_4 \cdot 8\text{H}_2\text{O}$, and after 48 hours, treatment began. Group E rats were treated with 4mg/kg body weight Moringa Oleifera seed oil each, group F rats were treated with 4mg/kg body weight Anacardium Occidentale nut oil and group G rats were treated with 2mg/kg body weight Moringa Oleifera seed oil + 2mg/kg body weight Anacardium Occidentale nut oil.

The animals were treated as for three (3) weeks and after every one week of treatment, the blood glucose levels of the rats were measured using a glucometer. Twenty four hours after the last administration, the rats were sacrificed by cervical dislocation. The skulls of the rats were then fractured open with forceps, the brains were removed into specimen bottles containing 40% formal calcium solution for histological and immunohistochemical analyses.

Data for oxidative enzyme tests and histochemical assays were statistically evaluated using SPSS 17.0 software (SPSS Incorporation, Chicago, USA) and were expressed in terms of Mean $\pm$ Standard Error of Mean. A value of $p < 0.05$ indicates a statistical significance between groups. All experimental recommendations provided in the guide for the care and use of laboratory animals was complied with (National Academy Press., 1996).

Table 1: Grouping of Animals

Group	Substance Administration
A	3ml of 0.9% NaCl solution per kg bw
B	Single dose of 2.5mg/kg bw $3\text{CdSO}_4 \cdot 8\text{H}_2\text{O}$
C	5mg/kg bw Vit. C + 6mg/kg bw Vit. E
D	5mg/kg bw Vit. C + 6mg/kg bw Vit. E + single dose 2.5mg/kg bw $3\text{CdSO}_4 \cdot 8\text{H}_2\text{O}$
E	4mg/kg bw of Moringa Oleifera seed oil
F	4mg/kgW of Anacardium Occidentale nut oil
G	2mg/kg bw of Moringa Oleifera seed oil + 2mg/kg bw of Anacardium Occidentale nut oil

Results




Figure A1: Histological demonstration of the frontal cortex tissue using the H&E staining technique [x400] showing normal neurons, glia and neuropil.




Figure A2: Histological demonstration using the Cresyl Fast Violet staining technique [x400]; showing numerous pyramidal cells and stellate cells as well as glia with normal morphology and normal pattern of distribution. Neuropil also appears normal



Figure A3: Histochemical demonstration of the frontal cortex tissue, using the Bielschowsky staining technique [x400] showing quite few senile plaques typical of a normal tissue.




Figure A4: Histochemical demonstration of Cathepsin D expression [x400]; showing normal expression of Cathepsin D in frontal cortex tissue




Figure A5: Histochemical demonstration of p53 expression [x400]; showing moderate expression of p53 in frontal cortex tissue.

GROUP A

Photomicrographs of the hippocampus of experimental animals administered 3ml of 0.9% NaCl solution per kg body weight as the control [N= Neurons; G = Glia; Senile plaque in being encircled].




Figure B1: Histological demonstration of the frontal cortex tissue of Group B animals using the H&E staining technique [x400] showing areas of clustered neurons with heterogeneous appearance.




Figure B2: Histological demonstration using the Cresyl Fast Violet staining technique [x400]; showing differential pattern of distribution of neurons and glia; area of densely packed predominantly pyramidal neurons and another area of prominent glia- predominantly astrocytes.



Figure B3: Histochemical demonstration of the frontal cortex tissue, using the Bielschowsky staining technique [x400] showing quite numerous senile plaques of various and randomly distributed.




Figure B4: Histochemical demonstration of Cathepsin D expression [x400]; showing quite expression of Cathepsin D in glia and relatively low expression in neurons.




Figure B5: Histochemical demonstration of p53 expression [x400]; neurons take on unusual morphology and pattern of p53 expression in the frontal cortex tissue.

GROUP B

Photomicrographs of experimental animals administered a single dose of 2.5mg/kg body weight of $3\text{CdSO}_4 \cdot 8\text{H}_2\text{O}$ at the onset of experiment [N= Neurons; G = Glia; Senile plaque in being encircled]




Figure C1: Histological demonstration of the frontal cortex tissue using the H&E staining technique [x400] showing normal neuron morphology and pattern of distribution; glia and neuropil.




Figure C4: Histochemical demonstration of Cathepsin D expression [x400]; showing relatively normal expression of Cathepsin D in frontal cortex tissue




Figure C2: Histological demonstration using the Cresyl Fast Violet staining technique [x400]; showing numerous neurons as well as glia with normal morphology and normal pattern of distribution. Neuropil also appears normal.




Figure C5: Histochemical demonstration of p53 expression [x400]; showing relatively moderate expression of p53 in the frontal cortex tissue.



Figure C3: Histochemical demonstration of the frontal cortex tissue, using the Bielschowsky staining technique [x400] showing relatively few senile plaques.

GROUP C

Photomicrographs of the hippocampus of experimental animals administered 5mg/kg body weight Vit. C + 6mg/kg body weight Vit. E throughout the duration of experiment [N= Neurons; G = Glia; Senile plaque in being encircled]




Figure D1: Histological demonstration of the frontal cortex tissue using the H&E staining technique [x400] showing heterogeneous neuron morphology




Figure D2: Histological demonstration using the Cresyl Fast Violet staining technique [x400]; showing heterogeneous neuron morphology and prominent astrocytes



Figure D3: Histochemical demonstration of the frontal cortex tissue, using the Bielschowsky staining technique [x400] showing quite large senile plaques.




Figure D4: Histochemical demonstration of Cathepsin D expression [x400]; showing heterogeneous neuron morphology and expression of Cathepsin D.




Figure D5: Histochemical demonstration of p53 expression [x400]; neurons take on heterogeneous morphology and relatively higher p53 expression in the neurons.

GROUP D

Photomicrographs of the hippocampus of experimental animals administered 5mg/kg body weight Vit. C + 6mg/kg bw Vit. E + single dose 2.5mg/kg body weight $3\text{CdSO}_4 \cdot 8\text{H}_2\text{O}$ [N= Neurons; G = Glia; Senile plaque in being encircled]




Figure E1: Histological demonstration of the hippocampus using the H&E staining technique [x400] showing neurons and glia




Figure E2: Histological demonstration using the Cresyl Fast Violet staining technique [x400]; showing numerous neurons as well as glia with normal morphology and normal pattern of distribution with normal neuropil



Figure E3: Histochemical demonstration of the hippocampus tissue, using the Bielschowsky staining technique [x400] showing quite few and small senile plaques




Figure E4: Histochemical demonstration of Cathepsin D expression [x400]; showing areas of enhanced expression of Cathepsin D in frontal cortex tissue



Figure E5: Histochemical demonstration of p53 expression [x400]; neurons express p53 relatively moderately

GROUP E

Photomicrographs of the hippocampus of experimental animals administered 4mg/kg body weight of *Moringa oleifera* seed oil throughout the duration of experiment



P 53




Figure F1: Histological demonstration of the frontal cortex tissue using the H&E staining technique [x400]; neurons and glia are observable




Figure F4: Histochemical demonstration of Cathepsin D expression [x400]; showing highly enhanced expression of Cathepsin D in the neurons




Figure F2: Histological demonstration using the Cresyl Fast Violet staining technique [x400]; neurons and glia are observable and prominent



Figure F5: Histochemical demonstration of p53 expression [x400]; neurons p53 expression in the cells is barely observable.



Figure F3: Histochemical demonstration of the frontal cortex tissue, using the Bielschowsky staining technique [x400] showing relatively abundant senile plaques.

GROUP F

Photomicrographs of the hippocampus of experimental animals administered 4mg/kgW of *Anacardium occidentale* nut oil throughout the duration of experiment




Figure G1: Histological demonstration of the hippocampus tissue using the H&E staining technique [x400]; neurons and glia are observable and prominent



Figure G2: Histological demonstration using the Cresyl Fast Violet staining technique [x400]; showing closely packed relatively large neurons



Figure G3: Histochemical demonstration of the frontal cortex tissue, using the Bielschowsky staining technique [x400] showing relatively few small-sized senile plaques.




Figure G4: Histochemical demonstration of Cathepsin D expression [x400]; showing relatively enhanced expression of Cathepsin D especially in the glia




Figure G5: Histochemical demonstration of p53 expression [x400]; p53 expression in neurons and glia is barely observable.

GROUP G

Photomicrographs of the hippocampus of experimental animals administered 2mg/kg body weight of *Moringa oleifera* seed oil + 2mg/kg body weight of *Anacardium occidentale* nut oil throughout the duration of experiment

Discussion

The present study evaluated the nature of a one-time single dose cadmium poisoning effects on the hippocampus of experimental animals; it also examined the individual effects of each of the oils- *Moringa oleifera* seed oil and *Anacardium occidentale* oil in ameliorating or neutralising such effects. The possible combined ameliorative effects of *Moringa oleifera* seed oil and *Anacardium occidentale* nut oil extracts on the destructive effects of cadmium on the hippocampus was also examined. The histological results showed that the control group A animals have normal features of a hippocampal tissue. Neurons and glia have normal morphology, characteristic of normal tissues. The expressions of Cathepsin D and p53 were also quite moderate and normal. These features could serve as adequate references with respect to other animal groups.

In group B which consists of animals induced with only cadmium showed features of deleterious effects of cadmium on the tissues. Heterogeneous morphology of neurons as well and unusually prominent glia- especially astrocytes are indications of toxic effects of cadmium on the tissue. More so, there could have been localized rather than randomly distributed damage of tissue damage as observed in Figure B2. Numerous senile plaque, quite more than usual for a normal tissue also support evidences of tissue damage. Glia expressed Cathepsin D quite prominently and at a quite high level; this also suggests the possibility of astrocytes reaction to the toxic effects of the substance on the tissue. The pattern of p53 expression was quite unusual and unconventional in Group B. all evidences- histological and histochemical, point to the toxic effects of the administered $3\text{CdSO}_4 \cdot 8\text{H}_2\text{O}$ on the hippocampus. This supports existing evidences of cellular destructive effects of cadmium (Yiin *et al.*, 1999). It has already been documented that cadmium causes cellular destruction, even cellular death at high concentration (Akerele, 1993).

Animals in group C which were treated with 5mg/kg body weight Vitamin C and 6mg/kg body weight Vitamin E showed normal features of a hippocampal tissue. The hippocampus is well populated by neurons and glia which are also normal in morphology. Relative to the control, the expressions of p53 and Cathepsin D are adequate and normal. Senile plaques are also few and do not show any characteristic of anomaly relative to the control. Vitamin E has potential to protect critical cellular structures against damage caused by oxygen free radicals and reactive products of lipid peroxidation; lipid peroxidation is prevented by vitamin E (Maydani, 1995).

Animals in group D which were treated with 5mg/kg body weight Vitamin C and 6mg/kg body weight Vitamin E and single dose 2.5mg/kg body weight $3\text{CdSO}_4 \cdot 8\text{H}_2\text{O}$. The hippocampus still show signs of altered hippocampal tissues due to toxic effects of cadmium. Neurons and glia are observable; neurons are however unusually heterogeneous and this suggests uncorrected effects of cadmium assault. The abnormally large senile plaques also point to tissue damage with serious molecular consequence. While the expression of Cathepsin D might be low, that of p53 is pronounced. Neuron morphology show little improvement over the untreated case in Group D; it is however obvious that it is not a case of complete correction or healing of the effects of cadmium intoxication.

Animals in group E which were induced with cadmium but treated with 2mg/kg body weight of *Moringa oleifera* seed oil showed improvement in cytoarchitecture of the hippocampus. Neurons

and glia as well as the neuropil are preserved and observable. Cathepsin D is however prominently expressed relative to the Group D especially that were treated with selected anti-oxidants, as well as the control Group. It is important to note that there are quite few and less prominent senile plaques. It is clear that there is an improvement in this tissue over the untreated Group B. It also appears that the extent of damages that lead to the formation of plaques is being greatly limited in this particular group. The heterogeneity of neuronal morphology is reduced [Figure E] related to Group B [Figure B]. There are evidences that *Moringa oleifera* seed oil effects against cadmium toxicity could have limited the extent of damage; also, the consequence excessive development of plaques as a result of such damage is prevented [especially when compared with the untreated Group B]. In terms of potency against the formation of plaques, Moringa seed oil appears better than the selected anti-oxidants in terms of prevention of excessive plaque formation. These features might be due to vitamin A, vitamin C, vitamin E and oxalic acid (Fuglie *et al.*, 1999). These constituents are said to have antioxidant effect, thus ability to remove superoxide and oxygen free radicals that are usually implicated in tissue destruction (Gustavo *et al.*, 2007). Ascorbic acid acts as a powerful hydrosoluble antioxidant in biological fluids and Moringa oleifera contains it (Fried *et al.*, 1990).

In group F animals which were induced with cadmium but treated with 2mg/kg body weight of *Anacardium occidentale* nut extract showed cytoarchitecture of the hippocampal tissue almost similar to those seen in group E animals which were induced with cadmium and treated with moringa oleifera. Neurons and glia are preserved. Cathepsin D is highly expressed; senile plaques are relatively smaller than in the untreated Group B, but more prominent than that of the Group E treated with Moringa oleifera seed oil. This is also expectedly due to its natural antioxidant property (Jaiswal *et al.* 2010). Study showed immature cashew nut-shell liquid (iCNSL), a source of unsaturated long-chain phenols, and may have a potential role in protecting against oxidative damage (Isao, 2005).

In group G which were induced with cadmium and treated with both 2mg/kg body weight of moringa oleifera seed oil and 2mg/kg body weight of *Anacardium occidentale* showed cytoarchitecture of the hippocampal tissue almost similar to those seen in group E and F animals due to their antioxidant properties but with more complete tissue repairs and regeneration. Neurons and glia are well preserved and they are prominent and well distributed. Senile plaques are very few and less prominent in a manner similar to the Group D treated with moringa seed oil only. This suggests that the moringa seed oil might be more potent in preventing the excessive formation of these plaques- more than other agents used. The combined effects of both oils in preserving neurons morphology however show possibility of greater potency.

Generally, cells in the hippocampus manifested some changes ranging from altered morphology and population densities as well as the cell's relative distribution. This implies that the activity of the hippocampus in memory formation and learning will be impaired and the role of the hippocampus that involved storage and retrieval of information will also be compromised. The findings in this study agree with the studies of Wolf *et al.*, (2009), who reported that rats exposed to high concentration of heavy metals, showed neurodegenerative changes in the hippocampus. Antioxidants exhibit positive effects that could ameliorate the damaging effects of cadmium on the hippocampus as seen in this investigation. The natural oils show improvement over the selected regularly used anti-oxidants, especially in limiting the extents of damage and the consequences in the hippocampus. Moringa seed oil produced better prevention of plaques especially, relative to *Anacardium occidentale*. It will be important to examine the roles of other

active phytochemicals in these oils in the differential potencies of the oils, as well as the mechanism of the effects.

Conclusion

It can be concluded from the present study that cadmium intoxication has observable deleterious effects on brain tissues as a result of intra-peritoneal administration of cadmium sulphate. All agents used produced positive anti-cadmium effects, however to varying degrees. While all agents ameliorated the effects of cadmium on neuronal morphology and relative population cum distribution, moringa seed oil was the most potent in reducing the formation of senile plaques in the hippocampus. The potency combined effect of moringa oleifera seed oil and *Anacardium occidentale* nut oil in reducing or treating the oxidative damage caused by cadmium on the hippocampus is also observed. The natural oils were more potent than the selected regularly used antioxidants.

References

- Agnello M, Filosto S, Scudiero R, Rinaldi AM and Roccheri MC.** (1993). *Cadmium induces an apoptotic response in sea urchin embryos*. Cell Stress Chaperones 12: 44-50.
- Akerele O.** (2007). *Nature Medical Bounty: don't throw it away*. World health forum 14, 390-395.
- Al, Krishnamurthy.** (2008). *Engineering Chemistry, 2/e*. pp. 82–83. ISBN 978-81-203-3666-7. Battery collection; recycling, nature protected". European Union. Retrieved November 4.
- Amaral, D; Lavenex P.** (2006). "Ch 3. *Hippocampal Neuroanatomy*". in Andersen P, Morris R, Amaral D, Bliss T, O'Keefe J. The Hippocampus Book. Oxford University Press. ISBN 9780195100273.
- Annabelle C.** (2009). *Rewarded Outcomes Enhance Reactivation of Experience in the Hippocampus Neuron*. 64:6, 910-921.
- Antonio MT, Benito MJ, Leret ML and Corpas I.** (1998). *Gestational administration of cadmium alters the neurotransmitter levels in newborn rat brains*. J Appl Toxicol 18: 83-88.
- Anwar F et al.** (2007). *Moringa oleifera: a food plant with multiple medicinal uses*. *Phytother Res.* (1):17-25. Source Department of Chemistry, University of Agriculture, Faisalabad-38040, Pakistan. anwar.gilani@aku.edu
- ATSDR** (2008). *Draft Toxicological Profile for Cadmium*. Atlanta, Georgia: US Department of Health and Human Services.
- ATSDR**, (2005). <http://www.atsdr.cdc.gov/clist.html>
- ATSDR.** (1999). *Toxicological Profile for Cadmium (Final Report)*. Atlanta, GA: Agency for Toxic Substances and Disease Registry. 434 pp.
- Au DW, Reunov AA and Wu RS,** (2001). *Reproductive impairment of sea urchin upon chronic exposure to cadmium. Part II: Effects on sperm development*. Environ Pollut 111: 11-20
- Bagchi, D., Vuchetich, P. J., Bagchi, M., Hassoun, E. A., Tran, M. X., Tang, L., and Stohs, S. J.** (1997). *Induction of oxidative stress by chronic administration of sodium dichromate and cadmium chloride to rats*. Free Radic. Biol. Med. 22, 471–478.
- Bailey et al.,** (1976), *Encyclopedia Britannica* 1993, Morton 1987, van Wyk 2005).

- Bertin G and Averbeck D.** (2006) *Cadmium: cellular effects, modifications of biomolecules, modulation of DNA repair and genotoxic consequences (a review)*. *Biochimie* 88: 1549-1559.
- Best PJ, White AM** (1999). *Placing hippocampal single-unit studies in a historical context*. (4):346–51. doi:10.1002/(SICI)1098-1063(1999)9:4<346::AID-HIPO2>3.0.CO;2-3. PMID 10495017.
- Bettinelli, M.; Baroni, U.; Pastorelli, N.** (1988). *Determination of arsenic, cadmium, lead, antimony, selenium and thallium in coal fly ash using the stabilised temperature platform furnace and Zeeman-effect background correction*. *Journal of Analytical Atomic Spectrometry* 3 (7): 1005–1011.
- Bliss T, Lomo T** (1973). *Long-lasting potentiation of synaptic transmission in the dentate area of the anaesthetized rabbit following stimulation of the perforant path*. *J Physiol* 232 (2): 331–56. PMID 4727084. PMC 1350458.
- Boyer P, Phillips JL, Rousseau FL, Ilivitsky S.** (2007). *Hippocampal abnormalities and memory deficits: new evidence of a strong pathophysiological link in schizophrenia*. *Brain Res Rev* 54 (1): 92–112. doi:10.1016/j.brainresrev.2006.12.008. PMID 17306884.
- Burke SN, Barnes CA** (2006). *Neural plasticity in the ageing brain*. *Nat Rev Neurosci* 7 (1): 30–40. doi:10.1038/nrn1809. PMID 16371948.
- Buxbaum, Gunter; Pfaff, Gerhard** (2005). *Cadmium Pigments*. Industrial inorganic pigments. Wiley-VCH. pp. 121–123. ISBN 978-3-527-30363-2.
- Buzsaki G** (1986). *Hippocampal sharp waves: their origin and significance*. *Brain Res.* 398 (2): 242–52. doi:10.1016/0006-8993(86)91483-6. PMID 3026567.
- Buzsaki G** (1989). *Two-stage model of memory trace formation: a role for noisy brain states*. *Neuroscience* 31 (3): 551–70. doi:10.1016/0306-4522(89)90423-5. PMID 2687720.
- Buzsaki G, Chen LS, Gage FH** (1990). *Spatial organization of physiological activity in the hippocampal region: relevance to memory formation*. *Prog Brain Res* 83: 257–68. doi:10.1016/S0079-6123(08)61255-8. PMID 2203100.
- Cantero, JL; Atienza M, Stickgold R, Kahana MJ, Madsen JR, Kocsis B** (November 26, 2003). *Sleep-dependent theta oscillations in the human hippocampus and neocortex*. *J Neurosci* 23 (34): 10897–903. PMID 14645485.
- Carballo, Rosa; Castiñeras, Alfonso; Domínguez-Martin, Alicia; García-Santos, Isabel; Niclós-Gutiérrez, Juan** (2013). "Chapter 7. *Solid state structures of cadmium complexes with relevance to biological systems*. In Astrid Sigel, Helmut Sigel and Roland K. O. Sigel. *Cadmium: From Toxicology to Essentiality*. Metal Ions in Life Sciences 11. Springer. pp. 145–189
- Case Studies in Environmental Medicine (CSEM)** (2011). *Cadmium*. Agency for Toxic Substances and Disease Registry. Archived from [the original](#) on 2011-06-06. Retrieved 2011-05-30.
- Chang, BS; Lowenstein DH** (2003). *Epilepsy* [92]. *N. Engl. J. Med.* 349 (13): 1257–66. doi:10.1056/NEJMr022308. PMID 14507951.
- Chiu YC, Algase D, Whall A, et al.** (2004). *Getting lost: directed attention and executive functions in early Alzheimer's disease patients*. *Dement Geriatr Cogn Disord* 17 (3): 174–80. doi:10.1159/000076353. PMID 14739541.
- Chumark P, Khunawat P, Sanvarinda Y, Phornchirasilp S, Morales NP, Phivthong-Ngam L, Ratanachamnong P, Srisawat P, Pongrapeeporn KU** (2008). *The in vitro and ex*

- vivo* antioxidant properties, hypolipidaemic and antiatherosclerotic activities of water extract of *Moringa oleifera* Lam. leaves. *J. Ethnopharmacol.* 116:439-446.
- Chung C.-P., Hsu HY, Chao AC, Chang FC, Sheng WY, Hu HH** (2006). *Detection of intracranial venous reflux in patients of transient global amnesia.* *Neurology* 66 (12): 1873–77
- Clayton L. Thomas.** (1989). *Taber's Cyclopedic Medical Dictionary.* F.A. Davis Company.
- Cooke SF, Bliss TV** (2006). *Plasticity in the human central nervous system.* *Brain* 129 (Pt 7): 1659–73. doi:10.1093/brain/awl082. PMID 16672292.
- Cotton, F. A.** (1999). *Survey of Transition-Metal Chemistry.* Advanced Inorganic Chemistry (6 ed.). John Wiley and Sons. p. 633. ISBN 0-471-19957-5.
- Diana RA, Yonelinas AP, Ranganath C** (2007). *Imaging recollection and familiarity in the medial temporal lobe: a three-component model.* *Trends Cogn Sci* 11 (9): 379–86. doi:10.1016/j.tics.2007.08.001. PMID 17707683.
- Duvernoy, HM** (2005). *Introduction [93]. The Human Hippocampus* (3rd ed.). Berlin: Springer-Verlag. p. 1. ISBN 3540231919.
- Eichenbaum H, Yonelinas AP, Ranganath C** (2007). *The medial temporal lobe and recognition memory [94].* 123–52. doi:10.1146/annurev.neuro.30.051606.094328. PMID 17417939. PMC 2064941.</ref>
- Eichenbaum, H; Cohen NJ** (1993). *Memory, Amnesia, and the Hippocampal System.* MIT Press.
- Eichenbaum, H; Otto TA, Wible CG, Piper JM** (1991). *Ch 7. Building a model of the hippocampus in olfaction and memory.* in Davis JL, Eichenbaum H,. *Olfaction.* MIT Press. ISBN 9780262041249.
- Ekstrom, AD; Kahana MJ, Caplan JB, Fields TA, Isham EA, Newman EL, Fried I** (2003). *Cellular networks underlying human spatial navigation [95]* (PDF). *Nature* 425 (6954): 184–88. doi:10.1038/nature01964. PMID 12968182.
- Eric R. Kandel, James H. Schwartz, and Thomas M. Jessel.** (1991). *Principles of Neural Science.* Elsevier, third edition,.
- EUR-Lex.** [Eur-lex.europa.eu](http://eur-lex.europa.eu) (April 18, 2011). Retrieved on June 5, 2011.
- Evans J and Hastings L.** (1992). *Accumulation of Cd(II) in the CNS depending on the route of administration: intraperitoneal, intratracheal, or intranasal.* *Fundam Appl Toxicol* 19: 275-278.
- FAO** (2008). FAOSTAT Data. *Food and Agriculture Organisation,* Rome.
- Fern, R., Black, J. A., Ransom, B. R., and Waxman, S. G.** (1996). *Cd(2p) induced injury in CNS white matter.* *J. Neurophysiol.* 76, 3264–3273.
- Fernandez N and Beiras R.** (2001). *Combined toxicity of dissolved mercury with copper, lead and cadmium on embryogenesis and early larval growth of the Paracentrotus lividus sea-urchin.* *Ecotoxicology* 10: 263-271.
- Finger, S** (2001). *Origins of Neuroscience: A History of Explorations Into Brain Function.* Oxford University Press US. ISBN 9780195146943.
- Friberg, L., Elinder, C. G., Kjellstrom, T., and Nordberg, G. F.** (1986). *Cadmium and Health: A Toxicological and Epidemiological Appraisal.* CRC Press, Boca Raton, FL.
- Fthenakis, V. M.** (2004). *Life cycle impact analysis of cadmium in CdTe PV production.* *Renewable and Sustainable Energy Reviews* 8 (4): 303. doi:10.1016/j.rser.2003.12.001.

- Fu, W; Sood S, Hedges DW** (2010). *Hippocampal volume deficits associated with exposure to psychological trauma and posttraumatic stress disorder in adults: A meta-analysis*. Progress in Neuro-Psychopharmacology and Biological Psychiatry **34** (7): 1181–1188.
- Fuglie, L.J.** (1999, 2000). *Moringa stem planting*.
- Garcia-Segura LM** (2009). *Hormones and Brain Plasticity*. Oxford University Press US. ISBN 9780195326611.
- Geraci F, Turturici G, Galli D, Cossu G, Giudice G and Sconzo G:** *Stress response in mesoangioblast stem cells*. Cell Death Diff 13: 1057-1063, 2006. Glucose 6 phosphate dehydrogenase manual of Radox laboratories, Ltd
- Goering, P. L., Waalkes, M. P., and Klaassen, C. D.** (1995). *Toxicology of cadmium*. In *Toxicology of Metals: Biological Aspects*. Handbook of Experimental Pharmacology, Vol. 115 (R. A. Goyer and M. G. Cherian, Eds.), pp. 189–213. Springer-Verlag, New York.
- Golberg, D. C.; et al.** (1969). *Trends in Usage of Cadmium: Report*. US NRC/NAS/NAE. pp. 1–3.
- Goto Y, Grace AA** (2008). *Limbic and cortical information processing in the nucleus accumbens* [96]. Trends Neurosci 31 (11): 552–8. doi:10.1016/j.tins.2008.08.002. PMID 18786735. PMC 2884964.
- Grant, C. A.; Sheppard, S. C.** (2008). *Fertilizer impacts on cadmium availability in agricultural soils and crops*. Human and Ecological Risk Assessment 14 (2): 210–228.
- Gray, JA.** (2000). *The Neuropsychology of Anxiety: An Enquiry into the Functions of the Septo-Hippocampal System*. Oxford University Press, P.44.
- Gray, JA; McNaughton N** (2000). *The Neuropsychology of Anxiety: An Enquiry into the Functions of the Septo-Hippocampal System*. Oxford University Press.
- Gross, Charles G.** (1993). *Hippocampus Minor and Man's Place in Nature: A Case Study in the Social Construction of Neuroanatomy*. Hippocampus 3 (4): 403–416
- Gutierrez-Reyes, E.Y.; Albores, A.; Rios, C.** *Increase of striatal dopamine release by cadmium in nursing rats and its prevention by dexamethasone-induced metallothionein*. Toxicology 1998, 131, 145–154.
- Habeebu SS, Liu J and Klaassen CD:** *Cadmium-induced apoptosis in mouse liver*. Toxicol Appl Pharmacol 149: 203-209, 1998.
- Hampel H, Bürger K, Teipel SJ, Bokde AL, Zetterberg H, Blennow K** (2008). *Core candidate neurochemical and imaging biomarkers of Alzheimer's disease*. Alzheimers Dement 4 (1): 38–48.
- Harrison PJ** (2004). *The hippocampus in schizophrenia: a review of the neuropathological evidence and its pathophysiological implications*. Psychopharmacology (Berl.) 174 (1): 151–62. doi:10.1007/s00213-003-1761-y. PMID 15205886.
- Hart, B. A., Lee, C. H., Shukla, G. S., Shukla, A., Osier, M., Eneman, J. D., and Chiu, J. F.** (1999). *Characterization of cadmium-induced apoptosis in rat lung epithelial cells: Evidence for the participation of oxidant stress*. Toxicology 133, 43–58
- Hetherington, L. E.; et al.** (2008). Production of Cadmium. World Mineral Production 2002–06. British Geological Survey. p. 15. Retrieved 2012-04-1
- Hiroaki. W** (2007). *Neural dynamics of the cognitive map in the hippocampus*. Cognitive Neurodynamics 1:2, 119-141
- Holleman, A. F.; Wiberg, E; Wiberg, Nils** (1985). *Cadmium*. Lehrbuch der Anorganischen Chemie, 91–100. Walter de Gruyter. pp. 1056–1057

- HSDB.** 2009. *Hazardous Substances Data Bank. National Library of Medicine.* <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB> and search on CAS number. Last accessed: 10/26/09.
- IARC.** (1993). *Cadmium and cadmium compounds. In Beryllium, Cadmium, Mercury and Exposures in the Glass Manufacturing Industry.* IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, vol. 58. Lyon, France: International Agency for Research on Cancer. pp. 119-239.
- IARC.** (1993). *Cadmium and cadmium compounds (Group1).* IARC monographs Vol. 58:1993. International Agency for Research on Cancer, Lyon. IPCS (1992). Cadmium. Geneva, World Health Organization, International Programme on Chemical Safety (Environmental Health Criteria 134; <http://www.inchem.org/documents/ehc/ehc/ehc134.htm>).
- IPCS** (1992). *Cadmium. Geneva, World Health Organization, International Programme on Chemical Safety (Environmental Health Criteria 134; <http://www.inchem.org/documents/ehc/ehc/ehc134.htm>).*
- IPCS** (2005–2007). *Cadmium, cadmium chloride, cadmium oxide, cadmium sulphide, cadmium acetate, cadmium sulphate.* Geneva, World Health Organization, International Programme on Chemical Safety (International Chemical Safety Cards 0020, 0116, 0117, 0404, 1075 and 1318; <http://www.who.int/ipcs/publications/icsc/en/index.html>).
- Isao Kubo *, Noriyoshi Masuoka, Tae Joung Ha, Kazuo Tsujimoto** (2005). *Antioxidant activity of anacardic acids. Department of Environmental Science, Policy and Management, University Of California, Berkeley, Ca 94720-3112, Usa.*
- Jaiswall Ys, Pa Tatke1*, Satish Y Gabhe, Ashok Vaidya** (2010). *Antioxidant Activity of Various Extracts of Leaves of Anacardium Occidentale (Cashew).* 1Department of Pharmaceutical Chemistry, C.U.Shah College of Pharmacy, S.N.D.T Women's University, Mumbai- 400049, India. 2Medical Research Centre, Kasturba Health Society, Vile Parle-(W), Mumbai- 400 056, India.
- James A. Duke** (1983). *Handbook of Energy Crops.* unpublished.
- Järup, L., Berglund, M., Elinder, C.G., Nordberg, G., Vahter, M.** (1998). *Health effects of cadmium exposure- a review of the literature and a risk estimate.* Scand J. Work Environ. Health 24 (Suppl 1), 1-5.
- Jed W. Fahey, Sc.D.** (2005). *Moringa oleifera: A Review of the Medical Evidence for Its Nutritional, Therapeutic, and Prophylactic Properties.* Part 1. December 1.
- Jeong EM, Moon CH, Kim CS, Lee SH, Baik EJ, Moon CK and Jung YS:** *Cadmium stimulates the expression of ICAM-1 via NF-kappaB activation in cerebrovascular endothelial cells.* Biochem Biophys Res Commun 320: 887-892, 2004.
- Jiao, Y.; Grant, C. A.; Bailey, L. D.** (2004). *Effects of phosphorus and zinc fertilizer on cadmium uptake and distribution in flax and durum wheat.* Journal of the Science of Food and Agriculture 84 (8): 777–785.
- Joels M** (2008). *Functional actions of corticosteroids in the hippocampus.* Eur J Pharmacol 583 (2-3): 312–321. doi:10.1016/j.ejphar.2007.11.064. PMID 18275953.
- Johnson, D.** (1973). *Cashew cultivation in Brazil. Agron. Mocamb.* Lourenco Marques 7(3):119–129.
- Klaassen CD and Liu J.** (1997) Role of metallothionein in cadmiuminduced hepatotoxicity and nephrotoxicity. Drug Metab Rev 29: 79-102.

- Klaassen, C. D., Liu, J., and Choudhuri S.** (1999). *Metallothionein: An intracellular protein to protect against cadmium toxicity*. Annu. Rev. Pharmacol. Toxicol. 39, 267–294.
- Kumar, R., Asic, K., Agarwal, K., and Seth, P. K.** (1996). *Oxidative stress mediated neurotoxicity of cadmium*. Toxicol. Lett. 89, 65–69.
- Kuruba R, Hattiangady B, Shetty AK** (2009). *Hippocampal neurogenesis and neural stem cells in temporal lobe epilepsy* [99]. Epilepsy Behav 14 Suppl 1: 65–73. doi:10.1016/j.yebeh.2008.08.020. PMID 18796338. PMC 2654382.
- Lalas S, Tsaknis J** (2002). *Extraction and identification of natural antioxidant from the seeds of the Moringa oleifera tree variety of Malawi*. JAOSC 79:677-683.
- Lee, Ching-Hwa; Hsi, CS** (2002). *Recycling of Scrap Cathode Ray Tubes*. Environmental Science and Technology 36 (1): 69 doi:10.1021/es010517q PMID 11811492.
- Lewis S** (1998). *Aetiology of transient global amnesia*. The Lancet 352 (9125): 397–9.
- Liu Y-P, Yang C-S and Tzeng S-F.** (2008). *Inhibitory regulation of glutamate aspartate transporter (GLAST) expression in astrocytes by cadmium-induced calcium influx*. J Neurochem 105: 137-150.
- Liu, J., Kadiiska, M. B., Corton, J. C., Qu, W., Waalkes, M. P., Mason, R. P., Liu, Y., and Klaassen, C. D.** (2002). *Acute cadmium exposure induces stress-related gene expression in wild-type and metallothionein-I/II null mice*. Free Radic. Biol. Med. 32, 525–535.
- Llewellyn TO.** (1994). *Cadmium. (Materials Flow)*. U.S. Department of Interior, Bureau of Mines, Information Circular 9380. <http://pubs.usgs.gov/usbmic/ic-9380/cadmium.pdf>.
- Lohr GW, Walker HD** (1974). *Glucose 6 phosphate dehydrogenase in method of enzymatic analysis* 3rd edition. Verlag Chemie- Wehnheim, pg636
- Lopez E, Arce C, Oset-Gasque MJ, Canadas S and Gonzales MP.** (2006). *Cadmium induces reactive oxygen species generation and lipid peroxidation in cortical neurons in culture*. Free Radic Biol Med 40: 940-951.
- Lopez, E., Figueroa, S., Oset-Gasque, M. J., and Gonzalez, M. P.** (2003). *Apoptosis and necrosis: Two distinct events induced by cadmium in cortical neurons in culture*. Br. J. Pharmacol. 138, 901–911.
- Lubenov EV, Siapas AG** (May 2009). *Hippocampal theta oscillations are travelling waves*. Nature 459: 534. doi:10.1038/nature08010. PMID 19448612.
- Maguire, EA; Burgess N, Donnett JG, Frackowiak RSJ, Firth CD, O'Keefe J** (1998). *Knowing Where and Getting There: A Human Navigation Network* [100]. Science 280 (5365): 921–24. doi:10.1126/science.280.5365.921. PMID 9572740.
- Maguire, EA; Gadian DG, Johnsrude IS, Good CD, Ashburner J, Frackowiak RS, Frith CD** (2000). *Navigation-related structural change in the hippocampi of taxi drivers* [101]. PNAS 97 (8): 4398–403. doi:10.1073/pnas.070039597. PMID 10716738. PMC 18253.
- Malenka RC, Bear MF** (2004). *LTP and LTD: an embarrassment of riches*. Neuron 44 (1): 5–21. doi:10.1016/j.neuron.2004.09.012. PMID 15450156.
- Meydani M.** (1995). *Vitamin E*. Lancet; 345: 170–5.
- Minami A, Takeda A, Nishibaba D, Takefuta S and Oka N.** (2001). *Cadmium toxicity in synaptic neurotransmission in the brain*. Brain Res 894: 336-339.
- Mistral HP, Fridovich J** (1972). *The role of superoxide anion in the antioxidation of epinephrine and a simple assay for SOD*. J. Biol. Chem. 247: 3170-5\
- Molnar G, Salanki J and Kiss T.** (2004). *Cadmium inhibits GABA-activated ion currents by increasing intracellular calcium level in snail neurons*. Brain Res 1008: 205-211.

- Morris, RGM; Garrud P, Rawlins JNP, O'Keefe J** (1982). *Place navigation impaired in rats with hippocampal lesions*. *Nature* 297 (5868): 681–83. doi:10.1038/297681a0. PMID 7088155.
- Morrow, H.** (2010). *Cadmium and Cadmium Alloys*. Kirk-Othmer Encyclopedia of Chemical Technology. John Wiley & Sons. pp. 1–36.
- Morton JF** (1991). *The horseradish tree, Moringa pterygosperma (Moringaceae)- A boon to arid lands?* *Econ. Bot.* 45:318-333.
- Moser, EI; Kropf E, Moser M-B** (2008). *Place Cells, Grid Cells, and the Brain's Spatial Representation System*. 69. doi:10.1146/annurev.neuro.31.061307.090723. PMID 18284371.
- Moser, EI; Moser M-B** (1998). *Functional differentiation in the hippocampus*. *Hippocampus* 8 (6): 608–19. doi:10.1002/(SICI)1098-1063(1998)8:6<608::AID-HIPO3>3.0.CO;2-7. PMID 9882018.
- Nadel L, O'Keefe J, Black A** (1975). *Slam on the brakes: a critique of Altman, Brunner, and Bayer's response-inhibition model of hippocampal function*. *Behav Biol* 14 (2): 151–62. doi:10.1016/S0091-6773(75)90148-0. PMID 1137539.
- Nakazawa K, McHugh TJ, Wilson MA, Tonegawa S** (2004). *NMDA receptors, place cells and hippocampal spatial memory*. *Nat Rev Neurosci* 5 (5): 361–72. doi:10.1038/nrn1385. PMID 15100719.
- Nishimura Y, Yamaguchi J-Y, Kanada A, Horimoto K, Kanemaru K, Satoh M and Oyama Y** , (2006). *Increase in intracellular Cd²⁺ concentration of rat cerebellar granule neurons incubated with cadmium chloride: cadmium cytotoxicity under external Ca²⁺-free condition*. *Toxicol In Vitro* 20: 211-216.
- Nollen EA and Morimoto RI.** (2002). *Chaperoning signalling pathways: molecular chaperones as stress-sensing 'heat shock' proteins*. *J Cell Sci* 115: 2809-2816.
- Nowinski, W.L.** (2001). *Computerized brain atlases for surgery of movement disorders*. *Semin. Neurosurg.* 12(2), 183–194
- OECD.** (1994). *Risk Reduction Monograph No. 5: Cadmium*. OECD Environment Monograph Series No. 104. OECD Environment Directorate, Paris.
- Ogbunugafor HA, Eneh FU, Ozumba AN, Igwo-Ezikpe MN, Okpuzor J, Igwilo IO, Adenekan SO, Onyekwelu OA** (2011). *Physico-chemical and antioxidant properties of Moringa oleifera seed oil*. *Pak. J. Nutr.* 10:409-414
- Okuda, B., Iwamoto, Y., Tachibana, H., and Sugita M.** (1997). *Parkinsonism after acute cadmium poisoning*. *Clin. Neurol. Neurosurg.* 99, 263–265.
- Paglia DE and Valentine WN** (1967). *Studies on the quantitative and qualitative characterisation of erythrocyte glutathione peroxidase*. *J. Lab. Clin. Med.* 70: 158-67
- Pal, R.; Nath, R., Gill, K.D.** (1993). *Influence of ethanol on cadmium accumulation and its impact on lipid peroxidation and membrane bound functional enzymes (Na⁺, K⁺)-ATPase and acetylcholinesterase) in various regions of adult rat brain*. *Neurochem. Int.* 1993, 23, 451–458.
- Pearce, J** (2001). The effects of telencephalic pallial lesions on spatial, temporal, and emotional learning in goldfish. *J Neurol Neurosurg Psychiatry* 71 (3): 351.
- Philippe Taupin** (2007). *The Hippocampus: Neurotransmission and Plasticity in the Nervous System*. Nova Pub
- Plachy J.** (2000). *Cadmium*. In *Minerals Yearbook, Vol. I, Metals and Minerals*. U.S. Geological Survey.

- Prull MW, Gabrieli JDE, Bunge SA** (2000). *Ch 2. Age-related changes in memory: A cognitive neuroscience perspective.* in Craik FIM, Salthouse TA. The handbook of aging and cognition. Erlbaum. ISBN 9780805829662.
- Scand. J. Work.** (1998) *Environ. Health* 24 suppl. 1:52p.
- Scoullou, Michael J.; Vonkeman, Gerrit H.; Thornton, Iain; Makuch, Zen** (2001). Mercury, Cadmium, Lead: Handbook for Sustainable Heavy Metals Policy and Regulation. Springer. ISBN 978-1-4020-0224-3.
- Seok SM, Park DH, Kim Y-C, Moon C-H, Jung Y-S, Baik EJ, Moon C-K and Lee SH,** (2006). *COX-2 is associated with cadmium-induced ICAM-1 expression in cerebrovascular endothelial cells.* Toxicol Lett 165: 212-220
- Shukla A, Shukla GS and Srimal RC.** (1996). *Cadmium-induced alterations in blood-brain barrier permeability and its possible correlation with decreased microvessel antioxidant potential in rat.* Hum Exp Toxicol 15: 400-405.
- Shukla, G.S.; Srivastava, R.S.; Chandra, S.V.** (1988). *Glutathione status and cadmium neurotoxicity— Studies in discrete brain-regions of growing-rats.* Fund Appl. Toxicol. 11, 229–235.
- Siddhuraju P, Becker K** (2003). *Antioxidant properties of various solvent extracts of total phenolic constituents from three different agroclimatic origins of drumstick tree (Moringa oleifera Lam.) leaves.* J. Agric. Food Chem. 51:2144-2155.
- Sinha, M.; Manna, P.; Sil, P.C.** (2008). *Cadmium-induced neurological disorders: Prophylactic role of taurine.* J. Appl. Toxicol. 2008, 28, 974–986.
- Skaggs WE, McNaughton BL, Permenter M, et al.** (2007). *EEG sharp waves and sparse ensemble unit activity in the macaque hippocampus* [108]. J Neurophysiol 98 (2): 898–910. doi:10.1152/jn.00401.2007. PMID 17522177.
- Smith C.J.E., Higgs M.S., Baldwin K.R.** (April 20, 1999). Advances to Protective Coatings and their Application to Ageing Aircraft. RTO MP-25. Retrieved May 29, 2011
- Smith DM, Mizumori SJ** (2006). *Hippocampal place cells, context, and episodic memory.* Hippocampus 16 (9): 716–29. doi:10.1002/hipo.20208. PMID 16897724.
- Squire, LR** (1992). *Memory and the hippocampus: a synthesis from findings with rats, monkeys, and humans.* 195–231. doi:10.1037/0033-295X.99.2.195.
- Squire, LR; Schacter DL** (2002). *The Neuropsychology of Memory.* Guilford Press.
- Stocks J, Domandy TL** (1971). *The autoxidation of human red cell lipid induced by hydrogen peroxide.* Br. J. Haematol. 20: 95 – 111
- Sutherland GR, McNaughton B** (2000). *Memory trace reactivation in hippocampal and neocortical neuronal ensembles.* Curr. Opin. Neurobiol. 10 (2): 180–86. doi:10.1016/S0959-4388(00)00079-9. PMID 10753801.
- The'venod F** (2009). *Cadmium and cellular signaling cascades: to be or not to be?* Toxicol Appl Pharm 238(3):221-239.
- Tolcin AC.** (2009a). *Cadmium.* In *Mineral Commodity Summaries.* U.S. Geological Survey. <http://minerals.usgs.gov/minerals/pubs/commodity/cadmium/mcs-2009-cadmi.pdf>.
- Tolcin AC.** (2009b). *Cadmium.* In *2008 Minerals Yearbook.* U.S. Geological Survey. <http://minerals.usgs.gov/minerals/pubs/commodity/cadmium/myb1-2008-cadmi.pdf>.
- Toxicol let.** (1990). Dec;54(92-3): 199-205
- Tsangaris GT and Tzortzatou-Stathopoulou F:** *Cadmium induces apoptosis differentially on immune system cell lines.* Toxicology 128: 143-150, 1998.

- UNEP (2008). *Draft final review of scientific information on cadmium. United Nations Environment Programme, Chemicals Branch* (http://www.chem.unep.ch/Pb_and_Cd/SR/Draft_final_reviews/Cd_Review/Final_UNEP_Cadmium_review_Nov_2008.pdf).
- Vanderwolf CH (2001). *The hippocampus as an olfacto-motor mechanism: were the classical anatomists right after all?*. Behav Brain Res 127 (1-2): 25–47. doi:10.1016/S0166-4328(01)00354-0. PMID 11718883.
- VanElzakker, MB; Fevurly RD, Breindel T, Spencer RL (2008). *Environmental novelty is associated with a selective increase in Fos expression in the output elements of the hippocampal formation and the perirhinal cortex*. [110].(12): 899–908. doi:10.1101/lm.1196508. PMID 19050162. PMC 2632843.
- W.H.O. (1999). *InorganicmercuryGeneva,Switzerland:WorldHealth Organization,International Programme on Chemical Safety*, Vol. 118. 168
- Chris M. (2008). *The hippocampus and memory: insights from spatial processing*. Nature Reviews Neuroscience 9:3, 182-194
- Waalkes, M. P. (2003). *Cadmium carcinogenesis*. Mutat. Res. 533, 107–120.
- Waalkes, M.P. (2000). *Cadmium carcinogenesis in review*. Jour. Inorganic Biochemistry 79: 240-244.
- Wang S, Hu P, Wang H-L, Wang M, Chen J-T, Tang J-L and Ruan D-Y, (2008). *Effects of Cd on AMPA receptor-mediated synaptic transmission in rat hippocampal CA1 area*. Toxicol Lett 176: 215-222.
- Wang Y, Fang J, Leonard SS, Krishna K and Rao M. (2004). *Cadmium inhibits the electron transfer chain and induces reactive oxygen species*. Free Radic Biol Med 36: 1434-1443.
- Waterston, W. and Burton, J. H. (1844). *Cyclopædia of commerce, mercantil law, finance, commercial geography and navigation*. p. 122.
- Webster, W. S., and Valois, A. A. (1981). *The toxic effects of cadmium on the neonatal mouse CNS*. J. Neuropathol. Exp. Neurol. 40, 247–257.
- WHO (2000). *Cadmium. In: Air quality guidelines for Europe*, 2nd ed. Copenhagen, World Health Organization Regional Office for Europe (http://www.euro.who.int/__data/assets/pdf_file/0005/74732/E71922.pdf).
- WHO (2007). *Health risks of heavy metals from long-range transboundary air pollution*. Copenhagen, World Health Organization Regional Office for Europe (<http://www.euro.who.int/document/E91044.pdf>).
- WHO (1992a). *Cadmium*. Environmental Health Criteria 134. World Health Organisation, International Programme on Chemical Safety (IPCS), Geneva, Switzerland.
- Willow J.H. Liu (2011). *Traditional Herbal Medicine Research Methods*. John Wiley & Sons, Inc. 101: 1140-1144.
- Wilson MA, McNaughton BL (1994). *Reactivation of hippocampal ensemble memories during sleep [111]*. Science 265 (5172): 676–79. doi:10.1126/science.8036517. PMID 8036517.
- Wong, K. L., and Klaassen, C. D. (1982). *Neurotoxic effects of cadmium in young rats*. Toxicol. Appl. Pharmacol. 63, 330–337.
- Yamagami K, Nishimura M and Sorimachi M, (1998). *Cd²⁺ and Co²⁺ at micromolar concentrations stimulate catecholamine secretion by increasing the cytosolic free Ca²⁺ concentration in cat adrenal chromaffin cells*. Brain Res 646: 295-298.