

Short Research Article

PAIN ASSESSMENT AND TREATMENT IN PEDIATRIC PATIENTS

ABSTRACT

Although there is a considerable number of pain assessment scales, only some of them could be used for practical intention. The Faces Pain Scale is used for children 4 to 6-7 years old, while the Visual Analog Scale (VAS) is used for children above this age.

A 7-months study from December 1, 2011 to May 31, 2012, was used to assess the acute pain among 177 children from birth to 13 years old in Obstetrics and Gynecology Hospital, University Hospital Centre, Pediatric Hospital and Outpatient Services in Tirana. For patients incapable to verbalize the pain intensity in a reliable way (neonates to < 4years old), it is used the hetero-evaluation. The hetero-evaluation is realized only with one assessment scale. While patients older than 4 years and < 6 years old were assessed using the hereto-evaluation and auto-evaluation scale according to the Faces Scale and VAS.

The auto-evaluation of the acute pain intensity is applied according to two different assessment scales, consisting in the same procedure, on the same patients. The assessment according to VAS is applied in two positions, horizontally and vertically. The results obtained from the different scales and the VAS positions are compared between one another. The NFCS and the Faces Scale are much more easily accessible by the nursing staff; while the multidimensional scales (OPS, CRIES, PIPP) were found more difficult to be used.

KEYWORDS: pain, pediatric patients, pain treatment, visual analogue scale.

INTRODUCTION

The medical science is becoming more and more sophisticated, but unfortunately there is more to do with the pain management. Adults are the first persons that benefited from the recent researches in this direction, mainly in the last two decades; meanwhile “uncommonly”, the pain is still untreated or insufficiently untreated among the children.^{[5][13][38]}

Physical pain is not just a transmission of electrical impulses in the corresponding sectors; it's considered as a “conflict” between the stimulant and the person that is suffering.^{[1][3][44]} Unrelieved pain may cause anxiety and stress, or in other cases even psychological and behavioral long term consequences.^[26]

Pain is an important part of the disease; it's the first evident symptom.^{[4][41]} Recently it is not considered anymore by the researchers and the doctors as a symptom, but as a specific disease, defined as “The Fifth Vital Sign”. Synchronal with this conception, there are other new notions like nervous system hypersensitivity caused by the acute pain and the rehabilitation, known for their important role due to major developments in the multimodal analgesia.

Last years the researches are mostly focused on pain assessments as the main element that precedes the treatment.^[48] There is used a specific number of assessments scales to measure the pain, mainly for the acute pain; but none of them fulfill at the same the criteria of validity,

reliability and applicability. Anyways, we can say that there are adopted some assessment scales according the researches or for daily use only.

Pain assessment is an essential element for the treatment. The success of a good analgesic management is based firstly in the valuation of the pain according to the scales.^{[27][48]}

Schematically, we can say that we use the behavioral scale like NFCS or CHEOPS or multidimensional scales (that combine the physiological markers like Pam or SpO₂ with the behavioral variations) like OPS, PIPP, CRIES, etc. to diagnose the children.^{[19][21]}

Pain assessment is based in two groups of measurement: the measurement of physiological parameters^[26] and the measurement of behavioral factors.^{[11] [18]} The physiological factors (like tachycardia, tachypnea, sweating, and the increase of arterial pressure) are always part of the pain, but they are not specific. Therefore, the behavioral scales are the main modalities used to assess the pain in neonates, infants and children under 4 years old that are not capable of verbal communication.^[12] These scales are based on facial expressions (crying or grimace), motor response, the vision and verbal response; for example the FLACC scale.

Although there is a considerable number of pain assessment scales, only some of them could be used for practical intention. We can use the FLACC scale for abovementioned age-group. The Faces Pain Scale is used for children 4 to 6-7 years old, while the Visual Analog Scale (VAS) is used for children above this age.^{[14][17][46]}

MATERIALS AND METHODS

A 7-months study from December 1, 2011 to May 31, 2012, was used to assess the acute pain among 177 children from birth to 13 years old in Obstetrics and Gynecology Hospital, University Hospital Centre, Pediatric Hospital and Outpatient Services in Tirana.

For patients, incapable to verbalize the pain intensity in a reliable way (neonates to <4years old), it is used the hetero-evaluation. The hetero-evaluation is realized only with one assessment scale.

While patients ≥ 4 years old and < 6 years old are assessed using the hereto-evaluation and auto-evaluation scale according to the Faces Scale and VAS.

The auto-evaluation of the acute pain intensity is applied according to two different assessment scales, consisting in the same procedure, on the same patients. The assessment according to VAS is applied in two positions, horizontally and vertically. The results obtained from the different scales and the VAS positions are compared between one another.

The auto-evaluation scales are combined as below:

- VAS/ NRS
- VAS/ Faces Scale

Pain is divided in 4 different levels, for all the scales:

LEVELS		NFCS	CRIES	OPS	NRS	VAS	Faces Scale	CHEOPS	PIPP
		0-10	0-10	0-10	0-10	0-10	0-10	4-13	0-21
1	Mild Pain	0 - < 3					2	4 - < 7	0 - < 7


2	Moderate Pain	$\geq 3 - < 5$	4	$\geq 7 - < 9$	$\geq 7 - < 11$
3	Intense Pain	$\geq 5 - < 7$	6	$\geq 9 - < 11$	$\geq 11 - < 17$
4	Maximum Pain	$\geq 7 - 10$	8 ose 10	$\geq 11 - 13$	$\geq 17 - 21$
Therapeutic Intervention Threshold		3/10	4/10	9/13	11/21

(According to Consensus Conference “*Évaluation et Stratégies de Prise en Charge de la Douleur Aiguë chez L’Enfant de 1 Mois à 15 Ans*”, ANAES, March 2000)


The population taken in consideration for the study is divided in 4 age-groups according to the suitable usage of the assessment scales, and it is represented in the table below.

AGE GROUP		HETERO-EVALUATION		AUTO-EVALUATION	
		Postoperative	Non postoperative	Postoperative	Non postoperative
I	0 to <1 month	CRIES	NFCS PIPP		
II	≥ 1 month to < 4 years old	OPS CRIES	CHEOPS		
III	≥ 4 years old to < 6 years old	OPS	CHEOPS	VAS Faces Scale	VAS Faces Scale
IV	≥ 6 years old			VAS NRS	VAS NRS

The assessment is applied among 177 patients, 40.1% of them females and 59.9% males.




The division according to the type of pain (postoperative/acute non postoperative) is represented below:



In the table below, it is represented the distribution of the patients according to age-group, associated with the graphical presentation:

GROUP I	GROUP II	GROUP III	GROUP IV
0 - < 1 month	≥ 1 month - < 4 years old	≥ 4 years old - < 6 years old	≥ 6 years old
22	114	17	24
12.43%	64.41%	9.60%	13.56%
177			

Scoring averages of different scales according to age-groups are represented in the table below. Since there are used two different positions of VAS (horizontal and vertical), all the information below is given separated according to the different variants (in which VASH is the horizontal variant and VASV is the vertical one).



The scoring system for the Face Scale, NFCS, VAS, OPS, CRIES and NRS is 0-10. While, for the CHEOPS (4-13) and PIPP (0-21) scales is applied the conversion of the average values to the 0-10 system.

DISCUSSION

According to the study's results, the percentage of postoperative patients with intense pain is 47.62% and with maximum pain is 52.38%. There are 5.77% non-postoperative patients with moderate pain, 55.77% with intense pain and 38.46% with maximum pain.^{[6][14]} In conclusion, all postoperative patients and 94.23% of non-postoperative children undergoing painful procedures experienced intense and maximum pain.^{[16][37][43]}

The average of the obtained results using all the scales varies in that way to show intense or maximum pain levels, as well.

Most of the researchers claim that if we compare the pain levels between males and females, the first ones are lower, but it is not considered as statistically significant ($p > 0.05$ for all the scales). However, we should take in consideration the fact that the auto-evaluation is applied among a small sample (24 patients-13.56%).^[15]

The previous studies indicate different levels of scoring, using the visual analog scale (VAS) in two different orientations-horizontal and vertical. Researchers argue that there are inequalities because of the difficulties among the children's spatial orientation, from 0-10 years old.^[20]

The NFCS and the Faces Scale are much more easily accessible by the nursing staff; while the multidimensional scales (OPS, CRIES, PIPP) were found more difficult to be used.^{[17][47]} This because of:

1. The considerable number of parameters to be assessed and saved in epidemiological files.

2. In many cases, there is not the required information about the levels of AP before and after the procedure.
3. Since there are required the reference values (the values of parameters before the surgical intervention or before the painful procedure), that could not be always documented.^[34]

The presence of parents during the auto-evaluation has played an important role in children's maximal collaboration, increasing in this way the obtained data validity.^{[9][33][45]}

TREATMENT

Pain treatment consists in providing a maximum pain relief and a minimum of side effects.^{[10][35][42]} The multimodal analgesia is considered to be as the most suitable method, based on the synergistic effects of different analgesics, decreasing the necessity of using high doses and the incidence of adverse effects of them.^{[2][7][21]}

Analgesics

Mild and moderate pain: could be used medicaments as paracetamol, NSAIDs (ibuprofen, naproxen, ketoprofen) and mild opioids (codeine, tramadol, dextropropoxyphene).

Average pain intensity: could be used opioids, accompanied with NSAIDs.

Intense pain: there are recommended major opioids (morphine, fentanyl, hydrocodone, oxycodone). NSAIDs could be used to increase the opioid's effects.^{[23][28][32]}

Local anesthetics could be used as peridural and intrathecal administration.

There are two types of analgesics: minor and major, or opioids and non opioids.^[30]

Acetaminophen (Paracetamol) is a minor analgesic that increases the effects of analgesia when it is used with another major analgesic. It is the most widely used analgesic among children for oral and rectal use. The optimal dose of acetaminophen it is not exactly defined, but usually it is used 10-15mg/kg, oral use. High doses of acetaminophen used in children with serious infections are considered as an indication for hepatic insufficiency. Therefore, it is suggested to be used for short period of times^[36].

Ibuprofen is a non steroid anti inflammatory drug used mainly to treat the pain and temperature among children.^[22] It is suggested a 15mg/kg dose for oral use to have an analgesic effect. However, for a repeated dose among 6 months to 12 years old children, it is suggested a dose of 10 mg/kg, every 6h oral use (maximum daily dose-40 mg/kg).^[36]

Naproxen has a longer half-life than ibuprofen, therefore it could be used every 8-12h. It is not known any information about the effect on infants. The usual or normal dose is considered 5-10 mg/kg oral use, every 8-12 h (maximum daily dose-20 mg/kg).^{[22][36]}

Opioid analgesics are used by the nursing staff under strict supervision of a doctor, because of the serious side effects, such as inhibition of respiration.^[29]

Opioids are used to treat pain among patients of different ages.^[22] If the dosage is correct, the analgesic effects among children could be easily distinguished.^{[7][8]} There are different routes of administration available for opioids, including oral, intravenous, rectal, transmucosal or transdermal administration.^[31] *It is important to avoid the intramuscular or subcutaneous route to pediatric patients.* Most of opioids are available in liquid forms and could be easily applied to children with difficulties in managing the opioid pills. Recently, to manage neoplastic pain treatment among children, it is used Fentanyl Transdermal Patch or other forms of Fentanyl.

Sedatives are mostly used for their calming or sleep-inducting effect to reduce pain.^[25]

The reasons why the pain could not be treated with opioids:

1. Side effects, such as respiratory depression.
2. Dependence.
3. Insufficient information possessed by the staff about these drugs.

Massage

Skin stimulation relieves pain, as well. There are used different techniques such as pressure, massage, vibration, heat and cold therapy, oily treatment which are considered to be very effective and safe. The activation of A β big fibers and the inhibition of A δ and C small fibers, which close the pain “gate”, can cause sedative effects.^{[7][8]}

Friction massage applied to the painful region could relax the muscles and decrease the tension. It is not recommended to the damaged skin since it could enlarge the damage. Skin irritation could be treated with menthol products, which contain methyl salicylate that could be absorbed, causing an analgesic effect in the affected area. Heat and cold therapy are use to decrease muscular pain.^{[37][39][42]} The immediate change of temperature, from heating to cold can cause analgesic effect that could last for hours. But, this is a method that can not be used within the first 24 hours to a traumatized area. Cold therapy reduces blood circulation causing positive effects, but preliminary we should take in consideration some advices about heat and cold therapy:

- To not use cold therapy to a hypo-vascular area
- To avoid extreme temperatures that can cause burn or freezing injuries
- To not apply heat therapy to a fresh wound because it might cause or increase bleeding
- To stop the application if the pain is increasing

Transcutaneous Electrical Nerve Stimulation (TENS)

TENS by definition covers the complete range of transcutaneously applied currents, with low voltage, transmitted to the affected area through some electrodes. It produces analgesics effects through A β fibers stimulation and A δ fibers inhibition.

Skin irritation could be a side effect caused by the electrode stickers. This could be avoided using hypoallergenic stickers, while redness could be avoided by cleaning the electrodes with soapy water. Contraindications of TENS include the application in eyes, forehead, mouth, neck. It is still not approved that TENS could perform safety during the very first months of pregnancy, but it is used to lower back part after it.^{[7][8]}

Biofeedback

It is a method that trains patients to control certain bodily processes that normally happen involuntarily, such as heart rate, respiratory rate and muscle tone. Relaxation decreases pain, decreasing the anxiety as well, and increasing pain control.

Fear and death could accelerate pain levels. Sharing emotionally patient’s experiences is as important as the treatment.^[24]

CONCLUSION

Assessment is the first step toward the appropriate postoperative and procedural acute pain management. The correct usage of assessment scales, according to a standard protocol ensures to truthfully measure pain intensity, being the “key to success” for the necessary information used to create the pain treatment modalities.

REFERENCES

1. Loeser JD, Treede RD. The Kyoto protocol of IASP Basic Pain Terminology. *Pain*. 2008;137:473–477.
2. Thienhaus O, Cole BE, Weiner RS. *Pain management: a practical guide for clinicians*. 6th ed. New York, NY: CRC Press; 2002. Classification of pain.
3. Ingelmo PM, Fumagalli R. Neuropathic pain in children. *Minerva Anestesiologica*. 2004;70:393–398.
4. Svendsen KB, et al. Breakthrough pain in malignant and non-malignant diseases: a review of prevalence, characteristics and mechanisms. *European Journal of Pain*. 2005;9:195–206.
5. Albertyn R, et al. Infant pain in developing countries; a South African perspective. In: Anand KJS, Stevens BJ, McGrath PJ, editors. *Pain in neonates and infants*. 3rd ed. Amsterdam: Elsevier; 2007. pp. 263–371.
6. Gaughan DM, et al. The prevalence of pain in pediatric human immunodeficiency virus/acquired immunodeficiency syndrome as reported by participants in the Pediatric Late Outcomes Study (PACTG 219) *Pediatrics*. 2002;109:1144–1152.
7. *Cancer pain relief and palliative care in children*. Geneva: World Health Organization; 1998.
8. Okpala I, Tawil A. Management of pain in sickle-cell disease. *Journal of Royal Society of Medicine*. 2002;95:456–458.
9. Hofmann M, et al. Posttraumatic stress disorder in children affected by sickle-cell disease and their parents. *American Journal of Hematology*. 2007;82:171–172.
10. Benjamin L. Pain management in sickle cell disease: palliative care begins at birth? *Hematology*. 2008;466–474.
11. von Baeyer CL, Spagrud LJ. Systematic review of observational (behavioral) measures of pain for children and adolescents aged 3 to 18 years. *Pain*. 2007;127:140–150.
12. Pillai Riddell R, Racine N. Assessing pain in infancy: the caregiver context. *Pain Research & Management*. 2009;14:27–32.
13. McGrath PJ, et al. Core outcome domains and measures for pediatric acute and chronic/recurrent pain clinical trials: PedIMMPACT recommendations. *The Journal of Pain*. 2008;9:771–783.

- 302 **14.** Cohen LL, et al. Evidence-based assessment of pediatric pain. *Journal of Pediatric*
303 *Psychology*. 2008;33:939–955.
- 304
- 305 **15.** Huguet A, Stinson JN, McGrath PJ. Measurement of self-reported pain intensity in children
306 and adolescents. *Journal of Psychosomatic Research*. 2010;68:329–336.
- 307
- 308 **16.** von Baeyer CL. Children's self-report of pain intensity: what we know, where we are headed.
309 *Pain Research & Management*. 2009;14:39–45.
- 310
- 311 **17.** Hicks CL, et al. The Faces Pain Scale – revised: toward a common metric in pediatric pain
312 measurement. *Pain*. 2001;93:173–183.
- 313
- 314 **18.** Crellin D, et al. Analysis of the validation of existing behavioral pain and distress scales for
315 use in the procedural setting. *Pediatric Anesthesia*. 2007;17:720–733.
- 316
- 317 **19.** Blount RL, Loiselle KA. Behavioural assessment of pediatric pain. *Pain Research &*
318 *Management*. 2009;14:47–52.
- 319
- 320 **20.** Stinson J, et al. Review of systematic reviews on acute procedural pain in children in the
321 hospital setting. *Acute Pain*. 2008;10:105–106.
- 322
- 323 **21.** Ramelet AS, et al. Clinical validation of the Multidimensional Assessment of Pain Scale.
324 *Pediatric Anesthesia*. 2007;17:1156–1165.
- 325
- 326 **22.** Franck LS, et al. Assessment of sickle cell pain in children and young adults using the
327 adolescent pediatric pain tool. *Journal of Pain and Symptom Management*. 2002;23:114–120.
- 328
- 329 **23.** Vetter TR. A primer on health-related quality of life in chronic pain medicine. *Anesthesia and*
330 *Analgesia*. 2007;104:703–718.
- 331
- 332 **24.** Palermo TM, et al. Evidence-based assessment of health-related quality of life and functional
333 impairment in pediatric psychology. *Journal of Pediatric Psychology*. 2008;33:983–996.
- 334
- 335
- 336 **25.** Long AC, Krishnamurthy V, Palermo TM. Sleep disturbances in school-age children with
337 chronic pain. *Journal of Pediatric Psychology*. 2008;33:258–268.
- 338
- 339 **26.** Blount RL, et al. Evidence-based assessment of coping and stress in pediatric psychology.
340 *Journal of Pediatric Psychology*. 2008;33:1021–1045.
- 341
- 342

27. Albertyn R, et al. Challenges associated with paediatric pain management in Sub Saharan Africa. *International Journal of Surgery*. 2009;7:91–93.
28. WHO Expert Committee on the Selection and Use of Essential Medicines. WHO model list of essential medicines for children: 2nd list (updated) March 2010. Geneva: World Health Organization; 2010.
29. Anand KJ, et al. Tolerance and withdrawal from prolonged opioid use in critically ill children. *Pediatrics*. 2010;125:1208–1225.
30. Berde CB, Sethna NF. Analgesics for the treatment of pain in children. *New England Journal of Medicine*. 2002;347:1542.
31. Challapalli V, et al. Systemic administration of local anesthetic agents to relieve neuropathic pain. *Cochrane Database of Systematic Reviews*. 2005;(4):CD003345.
32. De Lima L, et al. Potent analgesics are more expensive for patients in developing countries: a comparative study. *Journal of Pain & Palliative Care Pharmacotherapy*. 2004;18:59–70.
33. Ho IK, et al. Healthcare utilization and indirect burden among families of pediatric patients with chronic pain. *Journal of Musculoskeletal Pain*. 2008;16:155–164
34. Scoping document for the WHO treatment guidelines for chronic pain in children. Geneva: World Health Organization; 2008
35. International Narcotics Control Board. The report of the International Narcotics Control Board for 2008. New York, NY: United Nations; 2009
36. Merskey H, Albe-Fessard DG, Bonica JJ, et al. Pain terms: a list with definitions and notes on usage: recommended by the IASP Subcommittee on Taxonomy. *Pain* 1979;6:249-52
37. Annequin D., Canaoui P., Cloup C. Enquête multicentrique sur la douleur en réanimation pédiatrique In *Journées Parisiennes de Pédiatrie*, Paris, Flammarion Médecine Sciences, 1991:89-94
38. Mc Caffery M, Pasero C, eds. *Pain: Clinical Manual*, 2nd ed. St. Louis, MO: Mosby 1999.
39. Evaluation et stratégie de prise en charge de la douleur aiguë en ambulatoire chez l'enfant de 1 mois à 15 ans. www.anaes.fr
40. Stevens B, Taddio A, Ohlsson A, Einarson T. The efficacy of sucrose for relieving procedural pain in neonates: a systematic review and meta-analysis. *Acta Paediatr* 1997;86:837-42.

- 385
386
387
388
389
390
391

392
393
394
395
396
397

398
399
400

401
402
403

404
405
406

407
408
41. Stevens B, Johnston C, Petryshen P, Taddio A. Premature Infant Pain Profile: development and initial validation. Clin J Pain 1996;12:13-22
 42. Am. Acad. Pediatrics La prévention et la prise en charge de la douleur et du stress chez le nouveau-né Pediatrics 2000; 105: 454-461
 43. Goddard JM, Pickup SE. Postoperative pain in children. Anaesthesia 1996;51:588-90
 44. Fournier-Charrière E, Dommergues J. La douleur chez l'enfant : aspects spécifiques. La Revue du Praticien 1994 ; 44 :1925-1931.
 45. Gauvain-Piquard A. La décision de traiter un enfant pour sa douleur In : Cook J., Tursz A. in "L'enfant et la douleur, famille et soignants", Paris, Ed. Syros, 185, 1998:105-113
 46. Hicks CL, Von Baeyer C, Spafford PA, Van Korlaar I, Goodenough B. The Faces Pain Scale revised: toward a common metric in pediatric pain measurement. Pain 2001; 93:173-183.
 47. Ohnhaus EE & Adler R: Methodological problem in measurement of Pain: A Comparison Between VRS & VAS. Pain : 379,1975
 48. Pichard Leandri E. Evaluation de la douleur chez l'enfant. Rev Int Pediatr 1997 ; 28 : 20-25.