

Original Research Article

Pilot study of photoallergic reactions to UV filters among dental professionals

ABSTRACT (ARIAL, BOLD, 11 FONT, LEFT ALIGNED, CAPS)

Aims: To evaluate the incidence of photoallergic reactions to selected UV-filters among dental professionals (dentists, nurses and attendants) and dental patients.

Study design: In the study were included occupationally exposed to UV-filters containing dental materials dental professionals, and randomly chosen dental patients of different gender, age and occupations, the main inclusion criteria being the lack of occupational exposure to dental materials.

Place and Duration of Study: Department "Oral and Image Diagnostic", Faculty of Dental Medicine, Medical University of Sofia, between July 2014 and March 2015.

Methodology:

We included a total of 59 participants: 25 occupationally exposed dental professionals (15 women and 10 men; mean age 43.57 ± 8.76) and 34 dental patients (14 women and 20 men; mean age 40.41 ± 14.17). Skin photopatch testing with Benzophenone – 3, Benzophenone – 4, Para-aminobenzoic acid (PABA), BUTYL METHOXY-DIBENZOYL-METHANE, and 2,2-bis-[4-(2-hydroxy-3-methacryloxypropoxy)phenyl]-propane (BIS-GMA) (Chemotechnique Diagnostics) was performed, according to: Photopatch testing: a consensus methodology for Europe.

Results: Higher, but without statistical significance was the incidence of positive reactions to Benzophenone-3 and to Benzophenone-4 after UV-irradiation among women ($P=.23$; $P=.63$) and dental patients ($P=.75$; $P=.54$). A decrease of the incidence of positive skin patch test reactions to PABA was observed among the groups of men and dental patients, without statistical significance ($P=.63$, $P=.27$). Positive skin patch test reactions to BUTYL METHOXY-DIBENZOYL-METHANE were observed in the group of women and the one of dental professionals only after application of UV-irradiation; no statistical significance was revealed ($P=.08$). A slight increased incidence of positive skin patch test reactions after UV-irradiation was observed in all defined groups, but no statistical significance was established ($P=.40$; $P=.33$ - Table 3).

Conclusion: No statistically significant increased rate of sensitization to the studied photoallergens after UV-irradiation among dental professionals was observed. A slight photosensitizing action of Bis-GMA could be suggested, however, need further work to validate reliability.

Keywords: photosensitization, dental professionals, photopatch skin testing, Benzophenone – 3, Benzophenone – 4, Para-aminobenzoic acid, BUTYL METHOXY-DIBENZOYL-METHANE, BIS-GMA

1. INTRODUCTION

Photosensitisation is the term used for reactions dependent on the presence of certain photochemicals and the action of optical radiation in the 320–800 nm range. The adverse reaction is brought about by visible light or UV induced excitation involving free radicals or reactive oxygen species. The mechanism of reaction is similar to that of allergic contact dermatitis, but absorption of irradiation is necessary for the formation of antigens. Photoallergic reactions usually occur within 24–48 h [1].

Photosensitisers can be of endogenous origin such as porphyrins and flavins, or exogenous such as those derived from tar, vegetable and plant products, and fragrance materials. In addition, a number of commonly used drugs, e.g. antimicrobials, are potential photosensitisers

Modern techniques in dental cavity restorations, veneer- and orthodontic bonding and fissure sealing, depend on in situ polymerisation of monomers, most often brought about by light activation. UV absorbers in adhesives and other dental materials [2,3] absorb UV and visible light and can give rise to photosensitising reactions. In addition, leachables and degradation products from the dental materials as well as residuals from oral hygiene products and drugs may have radiation absorbing properties and, hence, may contribute to or be responsible for photoactivated reactions. It is of interest to discuss

potential adverse effects, such as photosensitivity reactions among dental professionals and dental patients.

The **purpose** of the present pilot study was to evaluate the incidence of photoallergic reactions to selected UV filters among dental professionals (dentists, nurses and attendants) and dental patients.

2. MATERIAL AND METHODS

2.1 STUDIED INDIVIDUALS

A total of 59 participants, divided into two groups, were included in the study: occupationally exposed dental professionals, and randomly chosen dental patients of different gender, age and occupations, the main inclusion criteria being the lack of occupational exposure to dental materials. Data regarding age and gender characteristics are presented in Table 1.

Table 1. General characteristics of the studied groups of individuals

Studied groups	Age (years) (M±SD)	Gender		Total
		Women n (%)	Man n (%)	
Dental patients	40.41 ± 14.17	14 (79,5)	20 (20.5)	34
Dental professionals	43.57 ± 8.76	15 (72.3)	10 (27.7)	25
Total	42.23 ± 16.3	29 (49.2)	30 (50.8)	59

The study was granted by the Medical University – Sofia, Grant № 48/2014, and was approved the Medical Ethics Board at the Medical University of Sofia. All participants were informed about the purpose of the study and gave their written informed consent.

2.2 SKIN PHOTOPATCH TESTING

Skin photopatch testing with the following photoallergens, included in the European Photopatch Baseline Series (EP-1000 - Chemotechnique Diagnostics) - Benzophenone – 3 (10.0% pet), Benzophenone – 4 (2.0% pet), PABA (10.0% pet), BUTYL METHOXY-DIBENZOYL-METHANE (10.0% pet), as well as with BIS-GMA was performed, according to Photopatch testing: a consensus methodology for Europe [4], by application of the allergens in IQ-Ultra hypoallergenic patches of Chemotechnique Diagnostics (IQ Chambers®, Vellinge, Sweden) on the mid upper back skin, avoiding 3-5 cm on either side of the vertebrae. Duplicate sets on left and right side of the back of the patient were applied for 48 h, after which both sets are removed. At this point, one set should be covered with an ultraviolet (UV) opaque material and the other irradiated with a calibrated metered broad-spectrum UVA source (UV-Therapy and Photodiagnosis system UV 236 PUVA) with a dose of 5 J/cm². Readings were recorded using the International Contact Dermatitis Research Group (ICDRG) scoring system with readings before irradiation, immediately after irradiation and also 48 and/or 72 h thereafter. Interpretation key based on recommendations by the ICDRG was applied (Table 2).

Table 2. Interpretation key of skin patch test results based on International Contact Dermatitis Research

Group

Symbol	Meaning
(-)	negative reaction
?	doubtful reaction
+	weak positive reaction (non-vesicular)
++	strong positive reaction (oedematous or vesicular)
+++	extreme positive reaction (ulcerative or bullous)
IR	irritant reaction

Lack of anti-allergic medication constituted a mandatory condition before placing the patches and during the testing.

2. 3 STATISTICAL METHODS

The statistics were calculated with SPSS 19.0. Available for cross-tabulation statistics were used - Fisher Exact Test for statistical significance, and two-sample t-test. Values of $P < 0.05$ were accepted as statistically significant.

3. RESULTS

Distribution by gender was uniform. The mean duration of occupational exposure of dental professionals to photosensitizers as ingredients of dental materials is 18 years, and 68 % of them were exposed up to 20 years.

Data concerning the incidence of positive skin patch test reactions to the tested substances, before and after UV-irradiation (distribution by gender and groups defined by occupational exposure in dentistry) are presented in (Table 3).

Table 3. Distribution of positive skin patch test reactions to tested substances, before and after UV-irradiation among the studied groups.

Tested substances		Positive reactions - distribution by gender		Positive reactions - distribution by studied groups	
		Men n (%)	Women n (%)	Dental professionals n (%)	Dental patients n (%)
Benzophenone – 3	before UV-irradiation	3 (10.0%)	4 (13.8%)	4 (16,0%)	3 (8,8%)

	after UV-irradiation	4 (13.3%)	7 (24.1%)	4 (16,0%)	7 (20,6%)
BIS-GMA	before UV-irradiation	6 (20.0%)	4 (13.8%)	6 (24,0%)	4 (11,8%)
	after UV-irradiation	7 (23.3%)	5 (17.2%)	7 (28,0%)	5 (14,7%)
PABA	before UV-irradiation	5 (16.7%)	4 (13.8%)	5 (20,0%)	4 (11,8%)
	after UV-irradiation	4 (13.3%)	4 (13.8%)	5 (20,0%)	3 (8,8%)
Benzophenone – 4	before UV-irradiation	5 (16.7%)	2 (6.9%)	3 (12,0%)	4 (11,8%)
	after UV-irradiation	4 (13.3%)	4 (13.8%)	3 (12,0%)	5 (14,7%)
BUTYL METHOXY-DIBENZOYL-METHANE	before UV-irradiation	3 (10.0%)	0 (0.0%)	0 (0.0%)	3 (8,8%)
	after UV-irradiation	2 (6.7%)	3 (10.3%)	1 (4,0%)	4 (11,8%)

Higher was the incidence of positive reactions to Benzophenone-3 after UV-irradiation among women (13.8% before vs 24.1% after UV-irradiation) and the group of dental patients (8.8% vs 20.6%), but no statistical significance was established ($P=.23$, $P=.75$).

Concerning the incidence of positive reactions to Benzophenone-4 after UV-irradiation, data were similar – it increased among women (6.9% vs 13.8%) and dental patients (11.8% vs 14.7%). However, no statistical significant differences were observed ($P=.63$, $P=.54$).

No reactions of photosensitization to PABA were observed in our study; furthermore, a decrease of the incidence of positive skin patch test reactions after UV-irradiation was observed among the groups of men and dental patients. No statistical significance was revealed ($P=.63$, $P=.27$).

Results from photopatch testing to BUTYL METHOXY-DIBENZOYL-METHANE are available in (Table 3). Interestingly, in the group of women and the one of dental professionals positive skin patch test reactions were observed only after application of UV-irradiation. Nevertheless, no statistical significance was revealed ($P=.08$).

Finally, due to the wide use of Bis-GMA as an ingredient of a number of dental materials, in the present pilot study we studied its possible photosensitizing action. A slight increased incidence of positive skin patch test reactions after UV-irradiation was observed in all defined groups, but no statistical significance was established ($P=.40$; $P=.33$ - Table 3).

4. DISCUSSION

Chemical ultraviolet (UV) filters have, over the last few decades, been increasingly used not only in conventional sunscreen products but also in many cosmetics and toiletries. Allergic contact dermatitis as well as photoallergic contact dermatitis reactions were well documented as a consequence of such use. We present the results from a pilot study on the incidence of photosensitivity reactions among dental professionals to the presented most common UV filters, applicable in dentistry as well.

Benzophenones are ultraviolet light filters that have been documented to cause adverse cutaneous reactions, including contact and photocontact dermatitis, contact and photocontact urticaria, and anaphylaxis. In recent years, they have become particularly well known for their ability to induce allergy and photoallergy. Topical sunscreens and other cosmetics are the sources of these allergens in most patients, but reports of reactions secondary to use of industrial products also exist. Benzophenones

as a group have been named the American Contact Dermatitis Society's Allergen of the Year for 2014 to raise awareness of both allergy and photoallergy to these ubiquitous agents [5].

While UV filters can cause both allergic and photoallergic contact dermatitis [6], the latter condition occurs more frequently. It should be emphasized that considering the widespread use of sunscreens, contact and photocontact dermatitis is uncommon. Photoallergic dermatitis from sunscreens is from organic UV filters, with benzophenone-3 (also known as oxybenzone) being the most common cause [7,8,9].

Benzophenone-3 (Fig 1) is used as UV-adsorber in topical sunscreens, moisturizers, shampoos, hair care products, lipsticks, lip balms, nail polish, etc. It is also common UV-adsorber and photostabilizer in dental composite materials, synthetic resins and other plastic materials [10,11].




Fig. 1. Chemical structure of Benzophenone-3

It is one of the most widely used organic UVA filters today, providing broad-spectrum ultraviolet coverage, including UVB and short-wave UVA rays. As a photoprotective agent, it has an absorption profile spanning from 270 to 350 nm with absorption peaks at 288 and 350 nm. Despite its photoprotective qualities, much controversy surrounds oxybenzone because of its possible hormonal and photoallergenic effects, leading many countries to regulate its use [12, 13].

The substance is currently regulated in the Cosmetics Directive in Annex VII, part 1 ("List of permitted UV filters which cosmetic products may contain") in a concentration up to maximum 10%. The regulation demands a warning on the label "contains oxybenzone" due to the photo-allergenic potential of the substance [14,15]. By the current dossier, submitted in December 2007, the applicants apply for a maximum allowed concentration up to 6%.

Benzophenone-4 (Sulisobenzene) is an ingredient in various sunscreen products, moisturizers, nail polish, lipsticks, lip balms, as well as in textiles, plastics, paints and cosmetics, protecting the skin from damage by UVB and short-wave UVA ultraviolet light (Fig 2) [16].




Fig. 2. Chemical structure of Benzophenone 4

Hughes and Stone (2007) investigated over a 3-year period whether any of selected chemical UV filters added to usual cosmetics/facial series produced positive patch test results in the absence of photostimulation. According to their results, benzophenone 4 produced significantly more positive patch test results than the other UV filters tested, and was the third most frequently positive result overall [17].

According to the results from the present study, the groups of women and of dental patients without occupational exposure in dentistry are with highest incidence of positive skin patch test reactions to Benzophenone-3 and Benzophenone-4 after UV-irradiation, without statistical significance. Generally, no risk of photosensitization to benzophenons in dentistry was established.

BUTYL METHOXY-DIBENZOYL-METHANE (Avobenzone) – (Fig. 3) is an oil soluble ingredient used in sunscreen cosmetics of the type creams, lotions, lipsticks, sun oils, etc., to absorb the full spectrum of UVA rays.




Fig. 3. Chemical structure of avobenzone

Its ability to absorb ultraviolet light over a wider range of wavelengths than many other sunscreen agents has led to its use in many commercial preparations marketed as "broad spectrum" sunscreens. Most sunscreens work by blocking UV rays and free radicals, but avobenzone works by absorbing the rays and converting them to energy that is less damaging to the skin. Avobenzone has an absorption maximum of 357 nm [18].

In our study, most intriguing were the results about skin patch test reactions to avobenzone after UV-irradiation. This substance seemed to play as a typical photosensitizer, since positive reactions were observed only after application of UV-irradiation the in the group of women and the one of dental professional. Since this is a pilot study, further investigations are needed before a more categorical statement about the role of exposure in dental practice for the incidence of sensitization to be given.

Para-aminobenzoic acid (PABA) – (Fig. 4), is an intermediate in the synthesis of folate by bacteria, plants, and fungi. Many bacteria, including those found in the human intestinal tract such as *E. coli*, generate PABA.




Fig. 4. Chemical structure of PABA

Patented in 1943, PABA was one of the first active ingredients to be used in sunscreen. It is a UVB absorber in wavelengths between 290 and 320 nm. In the past, PABA was widely used as a UV filter. Animal and in vitro studies in the early 1980s suggested PABA might increase the risk of cellular UV damage. [19]. On the basis of these studies, as well as problems with allergies and clothing discoloration, PABA fell out of favor as a sunscreen. However, its water-insoluble derivatives, such as padimate O are currently used in some products. Other uses include its conversion to specialty azo dyes and crosslinking agents.

Allergic reactions to PABA can occur. It is formed in the metabolism of certain ester local anesthetics, and many allergic reactions to local anesthetics are the result of reactions to PABA [20].

According to Gao et al (2014), who investigated the prevalence of photoallergic contact dermatitis reactions to different photoallergens among 4957 patients during a 7-year period, para-aminobenzoic acid was among the most predominant photoallergens [21]. Greenspoon et al (2013) completed a retrospective chart review of all patients who underwent photopatch testing between 2001 and 2010 in Canada. As most common relevant photoallergens they outlined benzophenone-3, PABA, and butylmethoxy-dibenzoylmethane [22].

Basing on the information above, in our study we studied the prevalence of photosensitization to PABA among the defined by us groups of individuals.

According to the results achieved in our pilot investigation PABA doesn't seem to be a significant photosensitizer among dental professionals, as well as among the control group. We didn't observe increase in positive reactions after UV-irradiation; furthermore, decrease of the incidence of positive skin patch test reactions among the groups of men and dental patients was found.

2,2-bis-[4-(2-hydroxy-3-methacryloxypropoxy)phenyl]-propane - Bis-GMA is a common monomer in composite fillings and fissure sealants, widely used in everyday dental practice. No data was found in the available literature concerning a possible action of this compound, so we decided to include Bis-GMA in our test series. The obtained results were intriguing. The sensitization rates to Bis-GMA were highest, if compared with the other tested allergens, in all defined by us groups. Nevertheless, as groups at risk of sensitization could be outlined those of dental professionals and men (sensitization rate 24% and 20% - Tabl. 3). Regarding the incidence of positive skin patch test reactions after UV-irradiation, it was slightly increased in all defined groups, without statistical significance. These findings are interesting and require further investigations, with more participants included.

5. CONCLUSION

A pilot study was performed to evaluate the incidence of photoallergic reactions to selected UV filters among dental professionals and dental patients. No statistically significant increased rate of sensitization to the studied photoallergens after UV-irradiation among dental professionals was observed. A slight photosensitizing action of Bis-GMA could be suggested, however, need further work to validate reliability.

CONSENT (WHERE EVER APPLICABLE)

All authors declare that written informed consent was obtained from the participants in the present study. A copy of the written consent is available for review by the Editorial office/Chief Editor/Editorial Board members of this journal.

ETHICAL APPROVAL (WHERE EVER APPLICABLE)

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

REFERENCES

- DeLeo V. Occupational phototoxicity and photoallergy. In: Kanerva L, Elsner P, Wahlberg JE, Maibach HI (eds) Handbook of occupational dermatology. Springer, Berlin, Heidelberg, New York, pp 314–324; 2000.
- Hickel RA, Folwaczny M Various forms of glass ionomers and compomers. In: Management alternatives for the carious lesion. Operative Dentistry Supplement. 2001;6:177–190.
- Van Meerbeek B, Vargas M, Inoue S, Yoshoda Y, Peumans M, Lambrechts P, Vanherle G. Adhesives and cements to promote preservation dentistry. In: Management alternatives for the carious lesion. Operative Dentistry Supplement. 2001;6:119–144.
- Bruynzeel DP, Ferguson J, Andersen K, Goncalo M, English J, Goossens A, et al. Photopatch testing: a consensus methodology for Europe. The European Taskforce for Photopatch Testing. JEADV. 2004;18, 679-682.
- Heurung AR, Raju SI, Warshaw EM. Benzophenones. Dermatitis. 2014;25(2):92-5.
- Fisher AA, Rietschel RL, Fowler JF, Fisher AA. Fisher's contact dermatitis. 6th ed. Hamilton (ON): BC Decker Inc.; 2008.
- European Multicentre Photopatch Test Study (EMCPPTS) Taskforce. A European multicentre photopatch test study. Br J Dermatol. 2012;166:1002-9.

8. Bryden AM, Moseley H, Ibbotson SH, Chowdhury MM, Beck MH, Bourke J, et al. Photopatch testing of 1155 patients: results of the U.K. Multicentre Photopatch Study Group. *Br J Dermatol*. 2006;155:737-47.
9. Zug KA, Warshaw EM, Fowler JF Jr, Maibach HI, Belsito DL, Pratt MD, et al. Patch-test results of the North American Contact Dermatitis Group 2005-2006. *Dermatitis*. 2009;20: 149-60
10. 2-HYDROXY-4-METHOXYBENZOPHENONE. National Library of Medicine (US), Division of Specialized Information Services. Accessed 11 May 2015. Available: <http://toxnet.nlm.nih.gov/cgi-bin/sis/search2/f?./temp/~2PfOus:1>.
11. SCCP/1201/08 Opinion on benzophenone-3. Accessed 11 May 2015. Available: <https://www.rinabook.com/new-read/781873/opinion-on-benzophenone-3.html>.
12. Burnett ME, Wang SQ. Current sunscreen controversies: A critical review. *Photodermatology, Photoimmunology & Photomedicine*. 2011;27(2):58–67. doi:10.1111/j.1600-0781.2011.00557.x
13. Calafat AM, Wong LY, Ye X, Reidy JA, Needham LL. Concentrations of the sunscreen agent Benzophenone-3 in residents of the United States: National Health and Nutrition Examination Survey 2003 – 2004. *Environ Health Perspectives*. 2008;116, 893-897.
14. Kiec-Swierczynska M, Krecisz B, Swierczynska-Machura D. Photoallergic and allergic reaction to 2-hydroxy-4-methoxybenzophenone (sunscreen) and allergy to cetyl alcohol in cosmetic cream. *Contact Dermatitis*. 2005;53:170-171. DOI: 10.1111/j.0105-1873.2005.0407b.x
15. Langan SM, Collins P. Photocontact allergy to oxybenzone and contact allergy to lignocaine and prilocaine. *Contact Dermatitis*. 2006;54:173-174. DOI: 10.1111/j.0105-1873.2005.0739f.x
16. Nohynek GJ, Schaefer H. Benefit and risk of organic ultraviolet filters. *Regul. Toxicol. Pharmacol*. 2001;33(3): 285–99. doi:10.1006/rtp.2001.1476.
17. Hughes TM, Stone NM. Benzophenone 4: an emerging allergen in cosmetics and toiletries? *Contact Dermatitis*. 2007;56(3):153-6.
18. Vielhaber G, Grether-Beck S, Koch O, Johncock W, Krutmann J. Sunscreens with an absorption maximum of > or =360 nm provide optimal protection against UVA1-induced expression of matrix metalloproteinase-1, interleukin-1, and interleukin-6 in human dermal fibroblasts. *Photochem Photobiol Sci*. 2006;5(3):275-82. doi:10.1039/b516702g
19. Osgood PJ, Moss SH, Davies DJG. The Sensitization of Near-Ultraviolet Radiation Killing of Mammalian Cells by the Sunscreen Agent Para-aminobenzoic Acid. *Journal of Investigative Dermatology*. 1982;79 (6):354–7. doi:10.1111/1523-1747.ep12529409.
20. Maki T, Takeda K. Benzoic Acid and Derivatives. *Ullmann's Encyclopedia of Industrial Chemistry*. Wiley-VCH; 2000. doi:10.1002/14356007.a03_555.
21. Gao L, Hu Y, Ni C, Xu Y, Ma L, Yan S, Dou X. Retrospective study of photopatch testing in a Chinese population during a 7-year period. *Dermatitis*. 2014;25(1):22-6. doi: 10.1097/DER.0000000000000008.
22. Greenspoon J, Ahluwalia R, Juma N, Rosen CF. Allergic and photoallergic contact dermatitis: a 10-year experience. *Dermatitis*. 2013;24(1):29-32. doi: 10.1097/DER.0b013e31827edc8b.