

Optimizing Diabetes by Herbal Drugs: A Review

ABSTRACT

Plant-based medicinal products have been acknowledged since ancient times and several medicinal plants and their products have been used to control diabetes in the traditional medicinal systems of many cultures worldwide. Several synthetic oral hypoglycemic agents are the primary forms of treatment for diabetes. However, prominent side-effects of such drugs are the main reason for an increasing number of people seeking alternative therapies that may have less severe or no side effects but little toxicological information exists concerning traditional antidiabetic plants.

Keywords: - Diabetes mellitus, Hypoglycemic, Plants

1. INTRODUCTION

Diabetes mellitus is a heterogeneous metabolic disorder characterized by altered carbohydrate, lipid and protein metabolism caused by insulin deficiency, often combined with insulin resistance [1]. It is considered as one of the five leading causes of death in the world [2]. About 150 million people are suffering from diabetes worldwide and it was almost five times more than the estimated ten years ago and this may be doubled by the year 2030 [3].

Hyperglycemia occurs because of uncontrolled hepatic glucose output and reduced uptake of glucose by skeletal muscle with reduced glycogen synthesis. When the renal threshold for glucose re absorption is exceeded, glucose spills over into the urine (glycosuria) and causes an osmotic diuresis (polyuria), which in turn, results in dehydration, thirst and increased drinking (polydipsia). Insulin deficiency causes wasting through increased breakdown and reduced synthesis of proteins.

2. Various forms of Diabetes mellitus [4]

I. General-genetic and other factors not precisely defined

a) Type 1 diabetes mellitus (T1DM OR IDDM).

- Type 1A – Auto-immune type 1 diabetes mellitus.
- Type 1B – Non-auto-immune or idiopathic type 1 diabetes mellitus.

b) Type 2 diabetes mellitus (T2DM OR NIDDM).

II. Specific defines gene mutations

a) Maturity onset diabetes of the young (MODY)

- MODY 1 (Hepatic nuclear fact 4- α gene mutations)
- MODY 2 (Glucokinase gene mutations)
- MODY 3 (Hepatic nuclear fact 1- α gene mutations)
- MODY 4 (Pancreatic determining fact x gene mutations)
- MODY X (Unidentified gene mutations(s))

b) Maternally inherited diabetes and deafness (MIDD)

c) Mitochondrial leucine t-RNA gene mutations

d) Insulin gene mutation

e) Insulin receptor gene mutations

III. Diabetes secondary to pancreatic disease

- Chronic pancreatitis
- Surgery

- Tropical diabetes (chronic pancreatitis associated with nutritional and or toxic factors)
- Gestational diabetes
- IV. Diabetes secondary to other Endocrinopathies
 - Cushing's disease
 - Glucocorticoid administration
 - Acromegaly
- V. Diabetes secondary to immune suppression
- VI. Diabetes associated with genetic syndromes eg: Prader–Willi syndrome
- VII. Associated with diabetes drug therapy
 - a) Drugs with Hypoglycemic effects
 - β-Adrenergic receptor antagonists,
 - Salicylates, Indomethacin, Clofibratemand ACE inhibitor.
 - b) Drugs with Hyperglycemic effects
 - Epinephrine, Glucocorticoid, β-adrenergic receptor agonist, Calcium channel blockers, Phenytoin and H₂ receptor blocker

Virtually all forms of diabetes mellitus are caused by a decrease in the circulating concentration of insulin (insulin deficiency) and a decrease in the response of peripheral tissues to insulin (insulin resistance). These abnormalities lead to alterations in the metabolism of carbohydrates, lipids, ketones and amino acids. The central feature of the syndrome is hyperglycemia characterized by a high blood glucose concentration.

Hyperglycemia may become toxic and fatal as a result of accumulation of non-enzymatically glycosylated products and osmotically active sugar alcohols such as sorbitol in tissues, the effects of glucose on cellular metabolism also may be responsible [5]. Diabetic ketoacidosis is an acute emergency which develops on prolonged hyperglycemia and develops in the absence of insulin because of an accelerated fat breakdown to acetyl CoA, which in the absence of aerobic carbohydrate metabolism is converted to acetoacetatae, β-hydroxybutyrate, that causes acidosis [6]. Hemoglobin undergoes glycosylation on its amino-terminal Valine residue to form the glucosyl Valine adduct of hemoglobin, termed hemoglobin A1c. The half-life on the modified hemoglobin is equal to that of the erythrocytes (about 120 days). Since the amount of glycosylated protein formed is proportional to the glucose concentration and the time of exposure of the protein to glucose, the concentration of hemoglobin, A1c in the circulation reflects the severity of the glycemic state [4].

2.1 Complications of the disorder [7]:

Most diabetes complications arise from prolonged exposure of tissues to elevated glucose levels. These complications develop as a consequence of the metabolic derangements in diabetes, often over many years. Many of these result in diseases of blood vessels, either large (macro vascular disease) or small (microangiopathy). Macro vascular disease consists of accelerated atheroma, which is much more common and severe in diabetic patients. Microangiopathy is a distinctive feature of diabetes mellitus and particularly affects the retina, kidney and peripheral nerves. Diabetes mellitus is the commonest cause of chronic renal failure, which itself represents a huge and rapidly increasing problem, the costs of which to society as well as to individual patients are staggering. Coexistent hypertension promotes progressive renal damage nephropathy and reduces the risk of myocardial infarction. Angiotensin-converting enzyme inhibitors or antagonists of angiotensin AT₁ receptors are more effective in preventing diabetic nephropathy than other antihypertensive drugs, perhaps because they prevent fibroproliferative actions of angiotensin- II and aldosterone. Although the underlying mechanism of diabetic complications is unclear much attention has been focused on the role of oxidative stress, which contributes to the pathogenesis of different diabetic complications.

Traditional treatments for diabetes may create a feeling of improved well-being without necessarily reducing hyperglycemia. There have been unsubstantiated claims that certain plants can ameliorate complications of diabetes (Table No:1), but the objective assessment of complications is difficult and the placebo effect of natural remedies cannot be discounted. No controlled studies of microvascular, macrovascular, or neuropathic complications have been undertaken with traditional therapies. A plant that is thirst quenching or increases sympathetic tone might gain a traditional reputation as efficacious without providing any long-term benefits to the underlying malady. Members of the *Allium* family, particularly garlic, are traditionally considered to give strength and combat arterial vascular

disease [8]. Garlic reduces polydipsia and weight loss in severely streptozocin-induced diabetic mice without improving glycemic control [9]. Onion has been reported to lower free fatty acid concentrations in healthy subjects, and seeds of *T. foenumgraecum* (fenugreek) reduced cholesterol levels in diabetic dogs [10, 11]. Other traditional antidiabetic plants have been anecdotally claimed to possess hypolipidemic properties.

Table No 1. Antidiabetic plants traditionally considered as efficacious in treatment of diabetes-associated complications

Complications	Plants
Polydipsia	<i>Panax ginseng</i> , <i>Polygonatum humile</i> , <i>Polygonatum macropodum</i> , <i>Polygonatum officinale</i>
Emaciation	<i>Allium sativum</i> , <i>Cichorium intybus</i>
Atherosclerosis	<i>Allium cepa</i> , <i>Allium sativum</i> , <i>Lycium chinensis</i> , <i>Panax ginseng</i> , <i>Taraxacum officinale</i> , <i>Trigonella foenumgraecum</i>
Retinopathy	<i>Daucus carota</i> , <i>Lycium chinensis</i> , <i>Taraxacum officinale</i> , <i>Vaccinium myrtillus</i>
Nephropathy	<i>Lycium chinensis</i>
Impotence	<i>Ceiba petandra</i> , <i>Coriandrum sativum</i> , <i>Crocus sativa</i> , <i>Panax ginseng</i> , <i>Papaver somniferum</i>

2.2 Insulin Therapy:

Insulin is the mainstay for treatment of virtually type-1 DM and many type-2 diabetes mellitus patients. When necessary, insulin may be administered intravenously or intramuscularly; however, long-term treatment relies predominantly on subcutaneous injection of the hormone. Subcutaneous administration of insulin differs from physiological secretion of insulin in at least two major ways: The kinetics do not produce the normal rapid rise and decline of insulin secretion in response to ingestion of nutrients, and the insulin diffuses into the portal circulation; the direct action of secreted insulin on hepatic metabolic processes is thus eliminated.

3. Screening methods of antidiabetic agents [12, 13]:

Before the advent of insulin and oral hypoglycemic drugs, the major form of treatment involved the use of plants. More than 400 plants have been recommended and recent investigations have affirmed the potential value of some of these treatments. The hypoglycemic and/or anti-hyperglycemic effect of several plants used as anti-diabetic remedies have been confirmed and the mechanisms of their actions are being studied. Chemical studies directed at the isolation, purification and identification of the substances responsible for the anti diabetic activity are also being conducted various diabetogens (Table No: 2) have been discovered by the researchers to assess the anti-diabetic potentials of the huge herbal resources.

Induction of experimental diabetes in the rat by particularly destroying the pancreatic beta cells is the most convenient method employed now a day and the most commonly used diabetogenic agents are alloxan and streptozotocin.

4. Herbal drugs used in the treatment of diabetes:

In the recent times many traditionally used medicinally important plants were tested for their anti-diabetic potential by various investigators in experimental animals. Several synthetic oral hypoglycemic agents that are used as conventional medicine today have a natural plant origin, eg: metformin was derived from the plant, *Galega officinalis* (Goat's Rue or French Lilae) which was a common remedy for diabetes [14, 15]. The synthetic hypoglycemic agents used in clinical practices have serious side effects like hematological effects, coma, disturbs the functions of liver and kidney. In addition, they are not suitable for use during pregnancy. Compared with synthetic drugs, drugs derived from plants are frequently considered to be less toxic with fewer side effects. Therefore, the search for more effective and

safer antidiabetic agent has become an area of active research. Most commonly used plants in herbal formulations in the treatment of diabetes are given in Table No: 3. The list of poly herbal formulation used is shown Table No: 4. The use of natural sweetener alternative to table sugar, artificial sweeteners like barley malt, maple syrup, rice syrup, fructose (fruit sugar) and the herb stevia are utilized more today. Stevia is an excellent natural alternative to artificial sweeteners. It is very concentrated, so only a small amount is used and it doesn't have the side effects that artificial sweeteners have. Aloe Vera, cinnamon, ginseng, onion, yam and ginseng powder are used as prediabetic remedy [71].

Table No 2. List of various diabetogens and doses used in diabetic study.

S.No.	Diabetogen	Dose	Animal
1	Alloxan	65-150 mg/kg	Rat
2	Streptozotocin	40-60 mg/kg	Rat
3	Cyclosporin A	40 mg/kg	Rat
5	Dehydro ascorbic acid	650 mg/kg	Rat
6	Methyl alloxan	53 mg/kg	Rat
7	Ethyl alloxan	50-130 mg/kg	Rat
8	Dexamethasone	2-5 mg/kg	Rabbit
9	Sodium diethyl Dithiocarbonate	0.5 –1 g/kg	Rabbit
10	Uric acid	1 g/kg	Rabbit

Table No 3. Plants reported having antidiabetic activity.

Plants	References
<i>Aegle Marmelos</i>	16
<i>Aerva lanata</i>	17
<i>Aegle marmelos</i>	18
<i>Allium cepa</i>	19
<i>Allium sativum</i>	20
<i>Alpinia galanga</i>	21
<i>Aporosa lindleyana</i>	22
<i>Averrhoa bilimbi</i>	23
<i>Azadirachta indica</i>	24
<i>Barleria lupulina</i>	25
<i>Bauhinia candicans</i>	26
<i>Berberis aristata</i>	27
<i>Boerhaavia diffusa</i>	28,29
<i>Boswellia glabra</i>	30
<i>Bougainvillea spectabilis</i>	31
<i>Caesalpinia bonducella</i>	32
<i>Curcuma longa</i>	33
<i>Eugenia jambolina</i>	34
<i>Fraxinus excelsior</i>	35
<i>Gymnema Sylvestre</i>	36
<i>Holostemma ada kodien</i>	72
<i>Momordica charantia</i>	37,38
<i>Ocimum gratissimum</i>	39
<i>Panax ginseng</i>	40
<i>Pterocarpus marsupium</i>	41
<i>Syzygium cumini</i>	42
<i>Talinum cuneifolium</i>	73
<i>Terminalia catappa</i>	43
<i>Terminalia chebula</i>	44
<i>Urtica dioica</i>	45
<i>Vinca rosea</i>	46
<i>Zizypus jujube</i>	47

Table No 4. Some of the polyherbal formulations used in treatment of Diabetes mellitus.

166

Formulation	Reference
Ayushan- 82	48
Diabecon (D- 400)	49
Dianex	50
Diarun plus65	51
MA- 471	52

167

168

169

5. The mechanism involved in the treatment of diabetes

170

The present treatment of diabetes is focused on controlling and lowering blood glucose to a normal level.

171

The mechanisms may be

172

- To stimulate cell of pancreatic islet to release insulin;

173

- To resist the hormones which rise blood glucose;

174

- To increase the number or rise the potency and sensitivity of insulin receptor sites to insulin;

175

- To decrease the leading-out of glycogen;

176

- To enhance the use of glucose in the tissue and organ;

177

- To clear away free radicals, resist lipid per oxidation and correct the metabolic disorder of lipid and protein;

178

- To improve microcirculation in the body.

179

180

Anti-diabetic medicinal plants undoubtedly have a significant effect on the lowering of blood sugar, but their mechanism of action is yet to be elucidated. The first evidence that the natural products have insulin potentiating activity was reported in 1929 by Glazer and Halpern. There are several mechanisms through which these herbs act to control the glucose level. They are more or less similar actions to the synthetic drugs. The mechanism of action of herbal anti-diabetics could be grouped as:

181

182

183

184

- Stimulation of insulin secretion (*Allium sativum* [20], *Allium cepa* [19],

185

- *Panax ginseng* [40])

186

- Inhibition in renal glucose reabsorption (*Fraxinus excelsior* [35])

187

- Stimulation of glycogenesis and hepatic glycolysis (*Momordica charantia* [38])

188

- Protective effect on the destruction of the beta-cells (*Thea sinensis* [53])

189

- Improvement of digestion and reduction of blood sugar and urea (*Aegle marmelos* [18])

190

- Prevents pathological conversion of starch to glucose (*Eugenia jambolina* [34], *Tinospora cordifolia* [34], *Pterocarpus marsupium* [41])

191

- Increasing the use of glucose by tissues and effect on adrenergic receptors (*Panax*

192

- Ginseng* [40], *Allium sativum* [20], *Allium cepa* [19])

193

- Potentiates the action of exogenously injected insulin (*Nelumbo nucifera* [54], *Rubia cordifolia* [55])

194

- Cortisol lowering activities (*Boerhaavia diffusa* [28,29], *Ocimum sanctum* [56])

195

196

197

Over 150 plant extracts some of the active principles, including flavonoids are known to be used for the treatment of diabetes [25, 57-59]. Huff and Howell reported that Quercetin stimulate insulin release and enhance calcium uptake from isolated cells, which suggest a place for flavonoids in NIDDM [60, 61]. The presence of 3' and 4' -OH in the B-ring and double bonds between C₂ and C₃ in flavones and flavonols are important factors for binding enzyme glycogen phosphorylase[62] which is a key enzyme in the regulation of glycogen metabolism and catalyzes a degradative phosphorylysis of glycogen to glucose.

198

199

200

201

202

203

204

205

6. Toxic Effects of Herbs:

206

207

Little toxicological information exists concerning traditional antidiabetic plants. Use of the plants over many centuries and sometimes as regular constituents of the diet might be expected to reveal any obvious detrimental side effects through the cumulative knowledge of personal experiences. Nevertheless, patients are prone to overindulge in natural treatments, believing their efficacious

208

209

210

reputation to imply safety, but chronic consumption of large amounts of traditional remedies must always be regarded with caution.

Catharanthus roseus (periwinkle) is widely used as a traditional treatment for NIDDM. Many major alkaloids isolated from this plant, including leurosine, vindoline, vindolinine, and catharanthine, recorded as mild hypoglycemic effect within 2-5 h in healthy rats, but none was sufficiently potent to encourage further investigation because of cytotoxic and neurological effects of *Catharanthus* alkaloids [63].

Ficus bengalensis (banyan tree), used traditionally in Asia for the treatment of DM, but studies on rats observed gross behavioral, neurologic, autonomic toxic effects [64]. A central nervous stimulation and hypertensive effect of ephedrine from *Ephedra* species and central nervous stimulation by *Panax* species has limited their usage in the treatment of diabetes [63, 40].

Unripe fruits of *Blighia sapida* (ackee fruit). A traditional treatment for diabetes in Central America and Africa. Hypoglycins (aminopropylpropionic acid derivatives) are effective in healthy and diabetic animals and humans, promoting glucose use and inhibiting gluconeogenesis secondary to the inhibition of long-chain fatty acid oxidation [65]. However, their toxic effects have precluded further development because it induces neuroglycopenia [65]. Other noncultivated traditional antidiabetic mushrooms such as *C. comatus* can accumulate heavy metals with toxic consequences if consumed in excess [63]. The poisonous effects of *A. phalloides* are probably due to neuroglycopenia after hepatic glycogen depletion and hepatic necrosis [66].

A study in South Africa recorded cases of spontaneous hypoglycemia, hepatic and renal necrosis after consumption of herbal medicines like *Adenia gummifera* (imfulwa), *Chenopodium Ambrosioides*, *Spilanthes Mauritania*, *Iboza riparia* [67]. A case-study report noted that the hypoglycemic effect of *M. charantia* was additive to that of chlorpropamide, and *V. myrtillus*, and synthalin apparently reduced insulin requirements [68, 69, 70].

7. Conclusion

This is an exciting time in medical and diabetic history. Diabetes is world's fast emergent metabolic disorder and a knowledge of this disorder increases, similar more appropriate therapies. Traditional plant medicines are used throughout the world for diabetes. So a study such medicine might offer a natural key to unlock a researcher in the future.

References

1. Rang and Dale. Pharmacology. International print-opera limited, Noida, UP, 2005:385.
2. Vats V, Yadav S P, Grover JK. Ethanolic extract of *Ocimum sanctum* leaves partially attenuates streptozocin induced alteration in glycogen content and carbohydrate metabolism in rats. J Ethnopharmacology. 2004; 90:155.
3. Gosh R, Sharachandra KH, Rita S, Thokchom IS. Hypoglycemic activity of *Ficus hispida* (bark) in normal and diabetic albino rats. Indian journal pharmacology 2004; 36:222.
4. Laurance LB, John SL, Keith LP. Goodman and Gilman's: The pharmacological basis of therapeutics. Mc Graw-Hill medical publishing division. New Delhi. 11th ed. 2006:1620.
5. Brownlees M. The pathological implications of protein glycation. Clin. Invest. Med. 1995; 18:275.
6. Satyanarayana U, Chakrapani U. Biochemistry. Book and allied (p) Ltd. Kolkata. 3rd ed. 2006:294.
7. Larmer J. Insulin and oral hypoglycemic drugs, glucogan .In: Gilman AG, Goodman LS, Rall TW, Murad F, Editors. The pharmacological basis of therapeutics. 7th Edition New York: Macmillan Publishing; 1985:1490.
8. Day C. The *Allium* Alliance. Nutr Food Sci 1984; 90:20.
9. Swanston-Flatt SK, Day C, Bailey CJ, Flatt PR. Traditional plant treatments for diabetes: studies in normal and streptozotocin diabetic mice. Diabetes Res Clin Pract. In press
10. Augusti KT, Benaim ME. Effect of essential oil of onion (allyl propyl disulphide) on blood glucose, free fatty acid and insulin levels of normal subjects. Clin Chim Acta 1975; 60:121.
11. Valette G, Sauvaire V, Baccou JC, Ribes G: Hypocholesterolemic effect of fenugreek seeds in dogs. Atherosclerosis 1984; 50:105.
12. Srinivasan K, Ramarao R. Animal models in type 2 diabetes research: An overview. Indian J. Med. Res. 2007; 125:451.

- 267 13. Bell HR, Hye RJ. Animal models for diabetes mellitus: physiology and pathology. J Surg Res. 1983;
268 35:433.
- 269 14. Oubre AY, Carlson TJ, King SR, Reaven GM. From plant, to patient: an ethnomedical approach to
270 the identification of new drugs for the treatment of NIDDM. Diabetologia 1997; 40:614.
- 271 15. Gloria YY, David ME, Kaptchuk TJ, Russell SP. Systematic Review of Herbs and Dietary
272 Supplements for Glycemic Control in Diabetes. Diabetes Care 2003; 26:1277.
- 273 16. Sharmila U, Kshama KS, et al. A study of hypoglycemic and antioxidant activity of *Aegle marmelos* in
274 alloxan induced diabetic rats. Indian J Physiol Pharmacol 2004; 48:476.
- 275 17. Vetrivelvan T, Jagadeeshan M. Antidiabetic activity of alcoholic extract of *Aerva lenata* (L) Juss. Ex
276 Schultes in rats. J Ethnopharmacol.2002; 80:103.
- 277 18. Das AV, Padayatti PS, Paulose CS. Effect of leaf extract of *Aegle marmelose* (L.)Correa ex Roxb. on
278 histological and ultrastructural changes in tissues of streptozotocin induced diabetic rats. Indian J Exp
279 Biol 1996; 34:341.
- 280 19. Kumari K, Mathew BC and Augusti KT. Antidiabetic and hypolipidemic effects of S-methyl cysteine
281 sulfoxide isolated from *Allium cepa* Linn. Indian J Biochem Biophys1995; 32: 49-54.
- 282 20. Augusti KT and Sheela CG. Antiperoxide effect of S-allyl cysteine sulfoxide, an insulin Secretagogue,
283 in diabetic rats. Experientia 1996; 52: 115.
- 284 21. Akthar MS, Khan MA, Malik MT. Hypoglycemic activity of *Alpinia galangal* rhizome and its extracts in
285 rabbits. Fitoterapia 2002; 73:623.
- 286 22. Jayakar B, Suresh B. Antihyperglycemic and hypoglycemic effect of *Aporosa lindleyana* in normal
287 and alloxan induced diabetic rats. Journal of Ethnopharmacology, 2003: 84:247.
- 288 23. Pusparaj P, Tan CH, Tan BHK. Effects of *Averrhoa bilimbi* leaf extract on blood glucose and lipids in
289 streptozotocin diabetic rats. J Ethnopharmacology.2000; 72:69.
- 290 24. Gupta S, Kataria M, et al. The protective role of extracts of neem seeds in diabetic caused by
291 streptozotocin in rats. J Ethnopharmacology.2004; 90:185.
- 292 25. Suba V, Murugasen R, et al. Antidiabetic potential of *Barleria lupulina* in rats. Fitoterapia 2004; 75: 1.
- 293 26. Fuentes O, Arancibia AP, Alarcon J. Hypoglycemic activity of *Bauhinia candicans* in diabetic induced
294 rabbits. Fitoterapia 2004; 75:527.
- 295 27. Bhupesh CS, Kamal S, et al. Antidiabetic activity of steam bark of *Berberies aristata* D.C. in alloxan
296 alloxan induced diabetic rats. The internet journal of pharmacology 2008; 6:1.
- 297 28. Pari L and Amamath Satheesh M: Hypoglycemic activity of *Boerhaavia diffuse* L. Effect on hepatic
298 key enzymes in experimental diabetes. J Ethnopharmacology 2004; 91: 109.
- 299 29. Satheesh MA and Pari L: Antioxidant effect of *Boerhavia diffusa* L. In tissues of alloxan induced
300 diabetic rats. Indian J Exp Biol 2004; 42: 989.
- 301 30. Kavitha JV, Rosario JF, et al. Hypoglycemic and other related effects of *Boswellia glabra* in alloxan
302 induced diabetic rats. Indian J Physiol Pharmacol 2007; 51:29.
- 303 31. Senapati AK, Dash GK, Ghosh T, Christina AJM. A study on antiinflammatory and hypoglycemic
304 activity of *Bougainvillea spectabilis*. Indian J. Nat. Prod. 2006; 22:3.
- 305 32. Kannur DM, Hukkeri VI, Akki KS. Antidiabetic activity of *Caesalpinia bonducella* seed extracts in rats.
306 Fitoterapia 2006; 77:546.
- 307 33. Ali Hussain HEM. Hypoglycemic, hypolipidemic and antioxidant properties of combination of
308 Curcumin from *Curcuma longa*, Linn. and partially purified product from *Abroma augusta* linn. In
309 streptozocin induced diabetes. Indian J of clinical biochemistry 2002; 17:33.
- 310 34. Grover JK, Vats V and Rath SS: Anti-hyperglycemic effect of *Eugenia jambolana* and *Tinospora*
311 *cordifolia* in experimental diabetes and their effects on key metabolic enzymes involved in
312 carbohydrate metabolism. J Ethnopharmacology 2000; 73:461.
- 313 35. Eddouks M, Maqhrani M. Phlorizin-like effect of *Fraxinus excelsior* in normal and diabetic rats. J
314 Ethnopharmacology 2004; 94:149.
- 315 36. Sugihara V, Nojima H, et al. Antihyperglycemic effects of *Gymema sylvestra* leaves in streptozocin
316 induced mice. J Asian Nat. Prod Res. 2000; 12:321.
- 317 37. Virdi J, Sivakami S, et al. Anti hypoglycemic effects of three extracts from *Momordi charantia*. J
318 Ethnopharmacology 2003; 88:107.
- 319 38. Sarkar S, Pranava M and Marita R. Demonstration of the hypoglycemic action of *Momordica*
320 *charantia* in a validated model of diabetes. Pharmacol Res 1996; 33:1.

39. Aguiyi JC, Obi CI, Gang SS, Igweh AC. Hypoglycemic activity of *Ocimum gratissimum* in rats. *Fitoterapia* 2000; 71:444.
40. Park EY, Kim HJ, Kim YK, Park SU et al. Increase in Insulin Secretion Induced by Panax ginseng Berry Extracts Contributes to the Amelioration of Hyperglycemia in Streptozotocin-induced Diabetic Mice. *J Ginseng Res*, Vol. 2012; 36:153.
41. Manickam M, Ramanathan M, Jahromi MA, et al. Antihyperglycaemic activity of phenolics from *Pterocarpus marsupium*. *J Nat Prod* 1997; 60: 609.
42. Kumar A, Iyavaranan R, et al. Antidiabetic activity of *Syzygium cumini* and its isolated compounds against streptozotocin induced diabetic rats. *Journal of medicinal plant research* 2008; 2:246.
43. Mansoor SA, Swamy VBM, Kumar P, Danapal R. Antidiabetic activity of *Terminalia catappa* Linn. Leaf extracts in alloxan induced diabetic rats. *Iranian J pharmacology and therapeutics* 2005; 4:36.
44. Sabu MC, Ramadasan K. Antidiabetic activity of medicinal plants and its relationship with their antioxidant property. *J Ethnopharmacology* 2002; 81:155.
45. Mohamed B, Fatima ZM, et al. Antihyperglycemic activity of the aqueous extract of *Urtica dioica*. *Fitoterapia* 2003; 74:677.
46. Sumana G, Suryawanshi SA. Effect of *Vinca rosea* extracts in treatment of alloxan induced diabetic male albino rats. *Indian j Exp. Biology* 2001; 39:748.
47. Aydin E, Fahrettin K, Hulusi A, Husseyin U, Yalcin T, Muzaffer U. Hypoglycemic effect of Zizyphus jujube Leaves. *J Pharm Pharmacol* 1995:4772.
48. Chowday DP, Dua M, Kishore B, Kishore P. Hypoglycemic effect of code formulation: Ayshu-82. *Journal of research in Ayurveda and Siddha*, 1998; 19:107.
49. Dhawan D, Bandhu HK, Singh B, Singh A, Nagpal JP. Effect of D-400 (A herbal formulation) on the regulation of glucose metabolism in diabetic rats. *Indian journal pharmacology* 1996; 28:224.
50. Mutalik S, Chetana M, Sulochana B, Devi PU. Effect of Dianex, a poly herbal formulation on experimental induced diabetic mellitus. *Phytother Res*. 2005; 19:409.
51. Senthilvel G, Jagadeesan M, et al. Effect of polyherbal formulation (Diarunal Plus) on streptozotocin induced experimental diabetes. *International journal of tropical of Tropical Medicine* 2006; 1:88.
52. Sircar AR, Ahuja RC, et al. Hypoglycemic, hypolipidemic and general beneficial effects of an herbal mixture MA-471. *Alternative Therapies in Clinical practice* 1996; 3:26.
53. Donga JJ, Surani VS, Sailor GU, Seth AK. A systematic review on natural medicine used for therapy of diabetes mellitus of some Indian medicinal plants. *An Int. J. Phar. Sci.* 2011; 2:36.
54. Pulkot KM, Kakali S, Pal M, Saha BP Effect of *Nelumbo nucifera* rhizome extract on blood sugar level in rats. *J Ethnopharmacology* 1997;58:207.
55. Rahul SS, Kishor SJ, Abhay KS. hypoglycaemic activity of roots of *Rubia cordifolia* in normal and diabetic rats. *Pharmacologyonline* 2007; 1:162.
56. Rai V, Iyer U and Mani UV: Effect of Tulasi (*Ocimum sanctum*) leaf powder supplementation on blood sugar levels, serum lipids and tissues lipids in diabetic rats. *Plant Foods Hum Nutr* 1997; 50: 9.
57. Meiselman HL, Halpern BP, Dateo GP. Reduction of sweetness judgement by extracts from the leaves of *Ziziphus jujube* *Physiology and Behavior* 1976; 17:313.
58. Choi JS, Yokozawa T, Oura H. Improvement of hyperglycemia and hyperlipidemia in streptozotocin – diabetic rats by methanolic extract of *Prunus davidiana* stems and its main component, prunigen. *Planta Med* 1991; 57; 208.
59. Ermenisoglu A, Kelestimur F, Koker AH, et al. Hypoglycemic effect of *Ziziphus Jujube* leaves. *J Pharm pharmacology* 1995; 47:72.
60. Hiff CS, Howell SL. Effect of epicatechin on rats isolated langerhans. *Diabetes* 1984; 33:291.
61. Tapas AR, Sarkar DM, Kakde RB. Flavonoids as nutraceuticals: A review. *Tropical journal of Pharmaceutical research*. 2008; 7:1089.
62. Atsushi K, Norio N, et al. Structural activity relationships of flavonoids as potential inhibitors of glycogen phosphorylase. *J. Agric. Food. Chem.* 2008; 56:4469.
63. Clifford JB, Caroline D. Traditional plant medicine as treatment for diabetes. *Diabetes care* 1989; 12:553.
64. Rakesh KS, Shikha M, Dolly J, Prashant KR, Geeta W. Antidiabetic effects of *Ficus bengalensis* aerial roots in experimental animals. *J Ethnopharmacology* 2009; 123:110.

- 374 65. Bressler R, Corriedor C, Brendel K. Hypoglycin and hypoglycin-like compounds. Pharmacol
375 Rev1969; 212:105.
- 376 66. Potron M. Champignons et diabete. Concours Med 1956; 36:3795.
- 377 67. Neame PB, Pillay VKG. Spontaneous hypoglycemia, hepatic and renal necrosis following the intake
378 of herbal medicines. S Afr Med J 1964; 38:729.
- 379 68. Allen FM. Blueberry leaf extract: physiologic and clinical properties in relation to carbohydrate
380 metabolism. JAMA 1927; 89:1577.
- 381 69. Wilder RM, Allan FN. Synthalin, blueberry leaf extract and glukohormet. JAMA 1928; 38:254.
- 382 70. Aslam M, Stockley IH. Interaction between curry ingredient (karela) and drug (chlorpropamide).
383 Lancet 1979; 1:607.
- 384 71. Jeffrey IM, Elisem B. Nutritional Strategies for the Diabetic/Prediabetic Patient. CRC Press 2006: 268.
- 385 72. Yasodha KJ, Raghu D, Jayaveera KN, Ravindra K. Anti-diabetic activity of Alcoholic extract of
386 *Holostemma ada kodien* in rats. The internet journal of Endocrinology. 2009; 5:2.
- 387 73. Yasodha KJ, Rasheed AS, Jayaveera KN, Ravindra K, Srikar A. Anti-diabetic activity of Alcoholic
388 extract of *Talinum cuneifolium* in rats. Pharmacologyonline 2008; 2:63-73