

1 Original Research Article

2 Self-Medication Practice among Secondary School Students in
3 Ekiti-State, Southwest Nigeria
4

5 Abstract

6 *Aims:* This study investigated the practice of self-medication and the types of drugs
7 students self-medicate as well as factors influencing practice of self-medication
8 among secondary school students in Ekiti State, Southwest Nigeria.
9 *Methods:* The study employed a descriptive survey design using a combination of
10 simple random and stratified sampling techniques to select 500 respondents to form
11 the study group.
12 *Instrumentation:* A pre-tested structured questionnaire was used to collect
13 information from respondents.
14 *Data Analysis:* SPSS 17.0 was used to analyse the data. The research questions were
15 answered using descriptive statistics of frequency and percentages. Inferential
16 statistics of t test and Analysis of Variance (ANOVA) were used to test the difference
17 in the practice of self-medication between age groups and sex at 95% confidence
18 level.
19 *Major Findings:* The study revealed that 80.4% of the respondents practised self-
20 medication. Also, analgesic 38.8% topped the type of drug that students self-
21 medicated. On the reasons adduced for self-medication, “I know what to do always”
22 has the highest response of 60.4%. The t cal 0.82<t tab 1.96 at P=0.05 denotes that
23 that there is no significant difference between male and female in the practice of self-
24 medication. Also, a df 2; 477, f cal 3.544>f tab 3.01 at P=0.05 signifies a significant
25 difference between age groups in the practice of self-medication.
26 *Conclusion:* The practice of self-medication was found to be prevalent among
27 secondary school students with. Analgesic was the drug most self-medicated and
28 significant difference exists between the age groups in the practice of self-medication.
29 It was recommended that, health educators should organize health campaigns and
30 sensitization programme in secondary schools to enlighten students on the health
31 implication of self-medication.
32 **Keywords:** self-medication, gender, age, practice, students
33

34 Introduction

35 Self-care behaviour in form of self-medication is a global phenomenon [1] and it has
36 become an issue under debate in health care [2]. Evidence has shown that self-
37 medication is not restricted to a region or race since both developing and developed
38 countries are experiencing the menace of inappropriate self-medication [3, 4]. The
39 variations in the level and patterns of self-medication among developing and
40 developed nations may be due to the disparities in the cultural, economic, health
41 care system and policies in each nation. Self –medication as a behaviour could be

42 classified as responsible or non-responsible and as a result, the behaviour cannot be
43 considered entirely harmful [5].

44 Responsible self-medication entails using approved and available medicine in a safe
45 and effective way as directed though, without prescription [6]. The types of drugs
46 used are indicated for a self-recognisable condition following initial medical
47 diagnosis, which means that users have previous knowledge of the dose, time and
48 side effect of the overdose of the drug. Responsible self-medication is possible in
49 the developed nations because of high quality of education, accessibility to health
50 information, safety and quality health care including government policies on health
51 coupled with the health seeking behaviour and sceptical expert knowledge [10].

52 Non-responsible self-medication is the use of drugs in the treatment of self-
53 diagnosed ailments or symptoms of diseases without supervision or prescription by
54 a physician [7]. It is characterised by indiscriminate use of medicine for the
55 management of ailment many of which have resulted into intoxication [8]. Major
56 problem of non-responsible self-medication is the lack of clinical evaluation by
57 medical professional which can result into missed diagnosis and delay in appropriate
58 treatment [9], drug resistance, use of expired drugs, wrong doses and prolonged
59 duration of usage [4]. The act of non-responsible self-medication seems to be
60 prevalent in the developing countries because of the poor level of literacy, scarcity
61 of health information and non-implementation of government policies on health
62 issues [11].

63 In Nigeria, high degree of irresponsible self-medication is practised [12]. The most
64 common drugs used without prescription include antimalarial, analgesics,
65 antipyretics [4], antibiotics and cough syrup [13]. The sources of the drugs people
66 use without prescription, sometimes are from pharmacy and left over drugs from
67 previous medication [14]. Students have reasonable information about drugs
68 through print media, pharmacy shops, medicine dealers, family and friends [15] or
69 patent medicine shops [13]

70 The factors influencing self-medication remains issues of intense debate in
71 academic discourse [16]. Un-controlled drugs in open market and poor drug
72 regulatory practices [17] contribute to self-medication. Many Nigerians, most times,
73 trivialised ailments such as headache, fever, cough, throat infection, common cold
74 and stomach ache [18] while, some do perceive some ailments to be too mild to
75 necessitate medical consultation [19] resulting into self-medication to manage such
76 ailments. Other factors like demographic factors have been found to influence self-
77 medication. For instance, gender, age, sex and social role were discovered to have
78 influence on self-medication [16]. Studies revealed higher prevalence of self-
79 medication among males than females while some found no strong association
80 between gender [18, 17].

81 Easy availability and accessibility to healthcare facilities and trained physicians
82 remain a cornerstone for reducing the practice of self-medication [20, 21]. The fact
83 remains, that many rural communities have little or no access to modern medical

84 care because of the uneven distribution or inadequate health facilities [22]. This
85 may encourage the practice of self- medication.
86 The practice of self-medication among secondary school students seems to be on
87 the increase. Many literatures exist on prevalence of self-medication among
88 university students [1] and among workers of tertiary institutions [16] but it was
89 observed that there is a relative dearth of literature on self-medication among
90 secondary school students. Therefore, this study investigated the practice of self-
91 medication among students of secondary schools, the types of drugs students self-
92 medicate and the factors influencing the practice. Lastly, the differences in the
93 practice of self-medication between gender as well as age group were also
94 determined.

96 **Methods**

98 **Study setting:** The study was conducted among secondary school students in Ekiti
99 State, Southwest Nigeria.

100 **Study Design:** Descriptive research design of the survey type was used to conduct
101 the study among students in rural and urban secondary schools in Ekiti State,
102 Southwest Nigeria.

103 **Sampling techniques:** A total of 500 students were selected using simple random
104 and stratified sampling techniques. Ekiti State was stratified into urban and rural
105 towns, after which simple random sampling was used to select two towns each from
106 each stratum and two secondary schools from each town. The schools were
107 stratified into junior secondary (JSS) and senior secondary schools (SSS). Due to the
108 difference in school population, proportionate sampling was used to select
109 respondents. Finally, 290 were selected from JSS and 210 from SSS making a total of
110 500 respondents.

111 **Instrumentation:** A pre tested, structured questionnaire was designed from related
112 studies. The questionnaire was in two parts. Part A elicited information on the
113 demographic variables of respondents and Part B was used to collect the data on
114 practice, reasons for practice of self-medication and types of drugs students self-
115 medicate. The first question was if students engage in self-medication followed
116 questions to determine the reasons for the practice of self-medication. The items
117 were measured with a two- point scale of “sometimes” and “Never”. The content
118 validity of the questionnaire was determined by experts in health education. Pilot
119 testing of the instrument on 20 respondents using test retest method of reliability
120 gave reliability co-efficient of 0.73. The researchers sought verbal consent from the
121 school principals before administering the questionnaire and selected respondents.
122 Copies of the questionnaire were administered to the respondents by the
123 researchers with the support of trained research assistants after explaining the
124 items in the questionnaire. The voluntary return of the completed questionnaire by
125 respondents signified consent and were included in the study

126 **Statistical Analysis:** Data were analysed using SPSS 17.0. Descriptive statistics of
127 frequency and percentage were used to answer the research questions. Inferential
128 statistics was used to find the differences in the practice of self-medication using t
129 test between male and female and Analysis of Variance (ANOVA) between the age
130 groups. The formulated hypotheses were tested at P= 0.05 level of significance.

131 **Results**

132 The response rate for the administered questionnaire was 96% because only 480
133 copies of questionnaire were fully completed and returned out of 500.

134
135 Table 1 shows that 58.3% respondents are 10-14 years, 15-19 years 39.6% while,
136 those in 20-24 years were 2.1%. Also, 50.4% were schooling in the urban areas and
137 49.6% in the rural areas. In addition, 45% were males and 55% were females.
138 Finally, 57.1% of the respondents were in junior secondary school (JSS) and 42.9%
139 were in senior secondary school (SSS).

Comment [U1]: Describe as demographic data then mention table

Variable		F	%
Age	10-14	280	58.3
	15-19	190	39.6
	20-24	10	2.1
	Total	480	100.0
Sex	Male	216	45.0
	Female	264	55.0
	Total	480	100.0
Location	Urban	242	50.4
	Rural	238	49.6
	Total	480	100.0
Level	JSS	274	57.1
	SSS	206	42.9
	Total	480	100.0

141 **Table 1: Distribution of the demographic variables of respondents**

Comment [U2]: It should be in top.

142
143 Table 2 reveals that 80.4% indulged in the act of self-medication while 19.6% of the
144 respondents did not practice self-medication. The breakdown of self-medication
145 practice among respondents were as follows; those within 10-14 years 44.6%, 15-19
146 years 34.2% and 20-24 years 8 1.6% including male 35.4% and female 45.5%

Comment [U3]: Same like before

Variable		Yes		No	
		F	%	F	%
Age	10-14	214	44.6	66	13.8
	15-19	164	34.2	26	5.4
	20-24	8	1.6	2	0.4
	Total	386	80.4	94	19.6
Sex	Male	170	35.4	46	9.6
	Female	216	45	48	10
	Total	386	80.4	94	19.6

Table 2: Frequency distribution of practice of self-medication

Comment [U4]: In top

Table 3 shows the analysis of the types of drugs the 386 respondents self-medicate. Analgesics 38.8% topped the list, followed by antibiotics 17.9% and antipyretics 14.2%. Others such as sedatives, blood supplement and appetite stimulant has 1.2% responses each.

Comment [U5]: Same As Before

Types of Drugs	F	%
Analgesic	186	38.8
Antibiotics	86	17.9
Antipyretics	68	14.2
Sedatives	6	1.2
Antimalarial	28	5.9
Blood Supplement	6	1.2
Appetite stimulant	6	1.2
Total	386	80.4

Table 3: Frequency distribution of the types of drugs students self-medicate

Comment [U6]: In top

Table 4 shows the reasons respondents engage in self-medication. I know what to do always, 60.4% of the respondents sometimes engaged in self-medication because they perceived that they know what to do. This was closely followed by “Ailment/disease is not serious to warrant going to the hospital” 58.3% and “using left over drugs drugs applied on previous health problem” 57.1%. The item with least response is lack of money to pay the doctor’s bill 41.7%.

Comment [U7]: Same As Before

	Sometimes		Never	
I engage in self-medication because:	F	%	F	%
Ailment/disease is not serious to warrant going to hospital	280	58.3	200	41.7
Using left over of drugs applied on previous health problem	274	57.1	206	42.9
Lack of money to pay for doctor's bill	176	36.7	304	63.3
Health facility too far away to my residence	222	46.3	258	53.7
Long duration of waiting to see doctor	200	41.7	280	58.3
No physician to prescribe drug for me	204	42.5	276	57.5
I know what to do always	290	60.4	190	39.6

Table 4: Frequency Distribution and Percentages responses on the reason for practicing self-medication

Table 5 shows that $t_{cal} 0.82 < t_{tab} 1.96$ $P=0.05$. The finding signifies that there is no significant difference between male and female in the practice of self-medication. The stated hypothesis that there is no significant difference between male and female in the practice of self-medication is thus not rejected.

	N	$\bar{X}$	SD	df	t cal	t tab	Remark
Practice Male	216	1.79	.410	478	0.82	1.96	NS
Female	264	1.82	.386				

$P=0.05$

Table 5: t test analysis of Gender and practice of Self-Medication

Table 6 shows the df 2;477, $F_{cal} 3.54 > F_{tab} 3.01$ at $P=0.05$. The fact that F_{cal} calculated is greater than the table value denotes that the stated hypothesis that there is no significant difference between age groups of respondents in the practice of self-medication is significant and thus not accepted.

Practice	SS	df	MSS	F cal	F tab	Sig.
Between Group	1.107	2	.553	3.544*	3.01	S
Within Group	74.485	477	.156			
Total	75.592					

$P=0.05$

Table 6: Analysis of Variance showing the difference between age and practice of Self-Medication

Discussion

This study discovered a high prevalence 80.4% of self-medication practice among the respondents. This finding is in line with previous similar study where a higher prevalence 97.8% was discovered among students of secondary school [23] as well as the submission that self-medication is prevalent among individuals in many developing countries of the world [11]. Nonetheless, a study on practice of self-medication revealed a lower prevalence of 11.9% [5]. The discrepancy in the prevalence could be because self-medication practice was limited to the preceding three months while this study did not limit self-medication to a particular period. Finding analgesics to be the drugs most used for self-medication agreed with a study conducted in Islamabad, Pakistan where analgesics were the most class of medicine used [24]. The similarities could be linked with the method of obtaining drugs in Nigeria which is similar to what happens in Pakistan where almost every pharmacy sells drug without demanding for prescription. On the contrary, a study carried out among adult population in Southwest Nigeria, discovered that antibiotics is the most common drugs used by students [13]. The difference could be attributed to the stage of development and exposure because the population in this study are younger generation with more female than male with most of them just attaining the stage of puberty. Findings in studies revealed that 84% of girls used analgesics without prescription to relieve themselves of menstrual pains [25] and over 67% girls take palliative drugs medicine for their menstrual pains without prescription [26]. The need to reduce dysmenorrhea among the girls may increase the use of analgesic. The higher number of female who are just entering into puberty in this study may be attributed to the increase in the prevalence of self-medication as well as using analgesics.

This survey showed that the most common reasons students self-medicated was “I know what to do always” this was closely followed by “ailment/disease is not serious to warrant going to hospital” and “using left over of drugs applied on previous health problem”. These findings are almost similar to the discoveries that students perceive some ailments to be too minor to necessitate medical consultation [19, 24] but disagreed with the finding that non-availability and accessibility to healthcare facilities increased the practice of self-medication [20]. Since, more than 50% of the respondents did not self-medicate because of the reasons relating to lack of money to pay doctors bill, health facility too far to residence, waiting time to see doctor or no physician to prescribe drugs, shows that the act of self-medication among the students was motivated by their attitude. The perception of knowing what to do could result into misplaced confidence which can cause inappropriate self-medication that would expose the respondents to risks associated with the practice of self-medication [27]

The present study discovered that there is no significant difference between male and female students in the practice of self-medication. This finding disagreed with the significant difference that was found between male and female in the practice of self-medication [23] and no strong association between gender and the practice of self-

239 medication [17] in similar studies. The slightly higher mean among female shows that
240 female practices self-medication than male negates the findings that showed a higher
241 prevalence among male than female [17, 18] but agreed with similar finding with higher
242 proportion of females 64.3% self-medicating than males 45.7% males [28]. The reason
243 for this finding could be linked to the fact that adolescent female may need to relieve
244 pains more than their male counterparts and male on the other hand, has the tendency
245 to trivialize illness or neglect mild illness.
246 The significant difference between age groups of respondents in the practice of self-
247 medication disagreed with the assertions in previous studies. For instance, no
248 significant difference was observed between different age groups and self-
249 medication [29] and significant relationship were found to exist between age and
250 the practice of self-medication. [16,18]. The variations in this study and others could
251 be as a result of the age of the respondents in this study because many of the
252 respondents at the time of this study were experiencing major physiological,
253 emotional and psychological changes.

254
255 **Conclusion**

256 The practice of self-medication was found to be prevalent among secondary school
257 students. Respondents self-medicated analgesic more than antibiotics. The major
258 reason identified for self-medicating was the perception that they know what to do
259 always followed by the feeling that the ailment/disease was not serious enough to
260 necessitate going to hospital. The danger in this perception is that serious ailment
261 could be trivialised and delay in the proper management of such ailment could be
262 fatal. There was no significant difference between males and females in the practice
263 of self-medication. Finally, a significant difference was found to exist between age in
264 the practice of self-medication. Based on the findings, health educators should
265 organize health campaign and sensitization programme in secondary schools to
266 enlighten students on the negative health implications of self-medication. Also,
267 secondary school students should be encouraged on the need to make use of health
268 care facilities

269
270 **Inclusion and Exclusion Criteria**

271 The final classes in Junior and Senior secondary schools were included because they
272 are regarded to be able to interpret the questionnaire than those in junior classes
273 All those who completed the questionnaire were also included.

274 **Limitations**

275 Students in private secondary schools were not included in the study hence
276 generalization of this study would not be related to them.

277 **Contribution of Authors**

278 Awosusi AO initiated the study, wrote the introduction and designed drafted the
279 questionnaire from related studies. Actively participated in the administration and
280 collation of the questionnaire and data analysis

Comment [U12]: In result section

Comment [U13]: In last part of Discussion

Konwea PE selected the respondents; pilot tested the instrument, participated in data administration and interpreted the data analysis for the study including the discussion of the study.

Acknowledgement

The researchers thank the research assistants who helped in the collection of data, the respondents for their co-operation and the school principals for the consent to conduct the study in their schools

Conflict of Interest: None declared

REFERENCES

1. Danshana B. Self-Medication: A Current Challenge. *Journal of Basic Clinical Pharmacy*. 2014; 5(1): 19-23
2. Almasdy DS. Self-Medication Practice with Non-Prescription Medication among University Students: a review of literature. *Archives of Pharmacy Practice*. 2011; 2(3): 95-100
3. Sarahrood S, Arzi A. Self-Medication with Antibiotics, Is it a Problem among Iranian College Students in Tehran? *J Biological Sciences*. 2009; 9: 829-32
4. Geissler PW, Nokes K, Prince RJ, Achieng OR, Aagaard-Hansen J, Ouma JH. Children and Medicine: Self Treatment of Common Illness among Luo School Children in Western Kenya. *Journal of Social Science and Medicine*. 2000; 50(112): 1771-1783
5. Kalaiselvi SS, Ganesh K, Archana R. Prevalence of self-medication practices and its associated factors in urban Puducherry, India. *Perspec Clin Res* 2014; 5(1): 32-36
6. WHO. Essential Medicines and Health Products . Information Portal. 2014; Apps.who.int/medicinedocs/en/cl/cl4.1/clmd
7. Ruiz ME. Risk of Self-Medication Practices. *Current Drug Safety*. 2010; 5(4): 315-23
8. Galato D, Galafassi L, Alano GM, Trauthman SC. Responsible Self- Medication: Review of the process of Pharmaceutical Attendance. *Brazilian Journal of Pharmaceutical Sciences* . 2009; 45(4): 625-633
9. Hamej MJ, Odhacha A, Roberts JM. Malaria Control in Buogoma District Kenya: A survey of Home Treatment of Children with Fever, Bed Net Use and Antenatal Clinics. *Bulletin of World Health Organization*. 2001; 79: 1014-1023
10. Talevi A. The New Patient and Responsible Self-Medication Practices: a critical review. *Current Drug Safety*. 2010; 5(4) 342-53.
11. Awad AI, Eltayeb IB. Self-medication Practices with Antibiotics and antimalarial among Sudanese Undergraduate University Students. *Annals of Pharmacotherapy*. 2007; 41:1249-1255

Comment [U14]: I actually do not know whether Journal Wants like this or not. But to my opinion this part should not appear in article

- 12.Olayemi OJ, Olayinka BO, Musa AJ. Evaluation of Antibiotic Self-Medication: Pattern amongst undergraduate students of ABU (Main Campus) Zaria. *J Appl Science Res.* 2010; 2(1): 35-38
13. Afolabi AO. Factors Influencing the Pattern of Self-Medication in an adult Nigerian Population. *Annals of African Medicine.* 2008; 7(3): 120-127
- 14.Vanden EJ, Marcus R, Hadler JL, Imhoff B, Vugia DJ, Cieslak PR, Zell E, Deneen V, McCombs KG, Zansky SM, Hawkins MA, Besser RE. Consumers Attitude and Use of Anti-Microbial Drugs . *Emerging Infectious Diseases.* 2003; 9: 1128-35
- 15.William IL. Self-Medication. *J National Medical Association.* 1978; 70(10): 731-732
- 16.Osemene KP, Lamikara A. A Study of the Prevalence of Self-Medication Practice among University Students in Southwestern Nigeria. *Trop J Pharmaceutical Res.* 2012; 11(4): 683-689
- 17.Fadare JO, Tamuno I. Antibiotic Self-Medication among University Undergraduate in Northern Nigeria. *Journal of Public Health Epidemiology.* 2011; 3(5): 217-220
- 18.Emmanuel A, Daniel G, Achema G, Afor B, Onyejekwe G, Gimba SM. Self-Medication Practice among Undergraduate Nursing Students of University of Jos, Nigeria. *Nigerian Journal of Pharmaceutical Sciences.* 2011; 10(2): 22-26.
- 19.Omolase CO, Adeleke EO, Afolabi AT. Self-Medication amongst General Out-Patients in a Nigerian Community Hospital. *Ann Ibadan Post Graduate Medicine.* 2007; 5(2): 64-67
- 20.Kumar N, Kanchan T, Unnikrishnan B, Rekha T, Mithia P. Perceptions and Practises of Self-Medication among Medical Students in Coastal South India. *PLoS ONE.* 2013; 8(8): e72247.doi:10.1371/journal.pone.0072247
- 21.Bamgboye EA, Amoran OE, Yusuf OB. Self-Medication Practices among Workers in a Tertiary Hospital in Nigeria. *African Journal of Medicine and Medical Sciences.* 2006; 35(4): 411-415
- 22.Arikpo G, Eja M, Enyi-Idoh K. Self-Medication in Rural Africa. *The Internet Journal.* 2009; 11(1):
- 23.Al-Hussaini M, Mustafa S, Ali S. Self-medication among Undergraduate Medical Students in Kuwait with reference to the Role of the Pharmacist. *Journal of Research Pharmacy Practice.* 2014; 3(1): 23-27
- 24.Aqeel T, Shabbir A, Basharat H, Bukhari M, Mobin S, Shahid H, Waqar SA. Prevalence of self-medication among urban and rural population of Islamabad, Pakistan. *Trop J Pharm Res.* 2014; 13(4): 627-633.
- 25.Sadiq MA, Salih AA. Knowledge and practice of adolescent females about menstruation in Baghdad. *Journal of General Practice.* 2013; 2(1):138-142
- 26.Poureslami M, Osati-Ashitani F. Assessing knowledge, attitudes and behaviour of adolescent girls in suburban districts of Tehran about dysmenorrhoea and menstrual hygiene. *Journal of International women's Studies.* 2002;3: 51-61

363 27. James H, Handu SS, Khalid AJ, Khaja A, Otoom S, Sequeira RP. Evaluation of
364 the knowledge, attitude and practice of self-medication among first year
365 medical students. Med Princ Pract. 2006; 15: 270-275
366 28. DeMoraes ACF, Delaporte TRM, Molena-Fernades CA, Falcao MC. Factors
367 associated with medicine use and selfmedication are different in adolescents.
368 Clinics. 2011; 66(7): 1149-55
369 29. Calvalho RS, Kara-Jose N, Temporini ER, Kara-Junior N, Noma-Campos R. Self-
370 medication: initial treatment used by patients seen in an ophthalmic
371 emergency rooms. Clinics (Sao Paulo, Brazil). 2009; 64(8):735-741