Original Research Article

Safety Evaluation of Giant African land Snails *(Archachatina* marginata) Haemolymph on HematologicalandBiochemical Parameters of Albino Rats.

Abstract

The safety of *Archachatina* maginata haemolymph on biochemical and hematological indices of albino rats was investigated. Twelve white albino rats were grouped into 3(A-C) of 4 animals each. Group A rats serve as control groups and received no treatment, while groups B and C received 2 and 4 ml/kg of the haemolymph, respectively, for ten days. Haemolymph administration resulted in significant increase (P<0.05) in the serum activities of aspartate aminotransferase (AST), alkaline phosphatase (ALP), alanine aminotransferase (ALT), total protein, albumin, total bilirubin and creatinine but significant decrease(P<0.05) the level of urea and chloride than the control rats. While serum bicarbonate, sodium, potassium, body weight gain, organ/body weight ratios were not significantly (P>0.05) altered. However, there was significant (P <0.05) decrease in the kidney/body weight ratio when compared with the control values. The haemolymph also produce a significant (p< 0.05) dose dependent changes in erythrocyte, leucocyte and platelete indices when compared with control value. The results indicated changes in the investigated haematological and some biochemical parameters showing a more pronounced effect on the liver function than the renal function tests and at a high dose than the low dose .Hence, caution should be taken when using the haemolymph of *Archachatina maginata* for therapeutic purpose because it may also have some liver or tissue membrane damaging effect.

Keywords:*Archachatina marginata*, Haemolymph, Biomarker enzyme, Liver, Kidney Haematological and Biochemical parameters

1. Introduction

Medicinal uses of natural product are gaining popularity in developing countries as has been estimated that 80%of people on this planet still rely on their traditional *materi amedica*(medicinal plants and other materials) for their primary health care needs. Emphasizes however has been laid that safety should be overriding criteria in the selection of herbal medicine for use in health care [1].

Scientific validations are being made globally to get evidences for traditionally used natural products. The active ingredients of natural products are chemicals that are similar to those in purified medications, but natural products lack defined dose and potency data and have the same potential to cause serious adverse effects [2].Whilst the literature documents severe toxicity resulting from the use of herbs, the potential toxicity of wild animal and their products has not been recognized. One of the wild animals claimed to be used in folklore medicine without recourse to its safety is *Archachatina* maginataand it haemolymph.

The Giant African land snails (*Archachatina marginata*) are soft bodied animals that belong to the phylum Mollusca, class Gastropoda, family Achatinidae and order Pulmonata [3].They are distributed through out the world in the tropical and sub-tropical regions where most of the genera of the family are confined to Africa [4]. They serve as wildlife dietary protein sources in Nigeria and some parts of Africa [5].

According to Abere and Lameed [6], snails are useful in the treatment of arteriosclerosis and other heart-related diseases due to their low cholesterol level. They also have pain reducing

effect and prevent loss of blood during labour as well as in the treatment of small pox [7].Previous biological studies reported that the isolated mucin motifs from *A. marginata* exhibited rectal absorption enhancing properties for the administration of insulin in rats [8], and produce marked and consistent blood glucose lowering effect [9]. There are also reports that the haemolymph of *A. marginata* exhibit pharmacological values for hypertensive patients, in thepreparations of vaccines, for treatment of kidney diseases, tuberculosis, diabetes, circulatory and stomach disorders [10]. Recently, we also reported that *A. marginata* haemolymph exhibited a significant protective effect against CCL4 induce liver damage [11].

Despite these pharmacological reports and wide spread use of the snail and its haemolymph, information on the safety or toxicity appears scanty. Therefore, this study was set out to assess the safety of the *Archachatina maginata* haemolymph on the biochemical parameters of organ damage. The organs (liver and kidney) have been carefully selected because of their role in the detoxification, excretion and absorption of the haemolymph. The haematological study was also carried out to determine it effect on blood component and corroborate it with alterations in the biochemical parameters

2. MATERIALS AND METHODS

2.1 Snail Collection

African Giant Snails (*Archachatina maginata)* within weight ranges of 150-210g were purchased from Kure market, Minna.Niger State. They were housed in a plastic vessel containing a layer of moist humus soil. The container was covered with a wire mesh for proper ventilation and also to prevent the snails from crawling out. The snails were fed *ad libitum* on fresh pawpaw leaves (*Carica papaya*).

2.2 Assay kit

The assay kits for AST, ALT, ALP total protein and albumin were products of Randox LaboratoriesLtd, United Kingdom. The urea and assay kits were obtained from QuinicaClinicaAplicada S.A., Spain. All other reagents used were of analytical grade and were prepared in distilled water.

2.3 Experimental Animals.

A total of twelve (12) white albino rats (*Rattus novergicus*) of both sex weighing between 120 and 200g were obtained from Small Animal Holding Unit of the Department of Biochemistry,Federal University of Technology Minna. The rats were kept in clean plastic cages and maintained under standard laboratory conditions (temperature: 22±3°C; photoperiod: 12 h natural light and 12 h dark; humidity: 40-45%).The animals were maintained on standard animal feeds (Bendel feeds and flour mills, Edo state, Nigeria) and tap water *ad libitum*. The principles governing the use of laboratory animals as laid out by the Federal university of Technology, Minna Committee on Ethics for Medical and Scientific Researchand also existing internationally accepted principles for laboratory animal use and care as contained in the Canadian Council on Animal Care Guidelines and Protocol Review [12] were duly observed.

2.4 Haemolymph Collection

The apex of the snails was opened by method adopted from Akinloye and Olorode[13].The haemolymph was drained into a clean conical flask and stored in refrigerator.

2.5 Experimental Design

Twelve (12) were randomly divided into three groups of 4 rats each as follows: Each group was separately housed.

(i) Group A— serve as control groups and received no treatment.
(ii) Group B— was intraperitoneally administered with 2ml/kg of African Giant Snails (*Archachatina maginata)* haemolymph for 10 days.
(iii) Group C— was intraperitoneallyadministered with 4ml/kg of African Giant Snails (*Archachatina maginata)*haemolymph for 10 days

2.6 Collection of Blood, serum and organs

Collection of sample for hematologicaland biochemical analyses was as described by Yakubu et al., [14]. At the end of the ten days treatment, the animals were denied their feeds but still had water *ad libitum* for 24h before they were sacrificed under ether anaesthesia. The blood was collected into sample bottles containing EDTA for haematological analyses. Another 4 mL of the blood was collected into a clean, dry centrifuge tubes. The blood sample was allowed to stand for 10 minutes at room temperature and then centrifuged at 1000rpm for 15minutes to get the serum. The animals were thereafter quickly dissected and the liver, kidneys spleen and heart were removed, cleaned and weighed.

2.7 Determination of Haematological Parameters

The haematological components including haemoglobin (Hb), haematocrite (HCT), red blood cells (RBC), mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH), mean corpuscular haemoglobin concentration (MCHC), white blood cells (WBC), Granulocyte count (GRA) lymphocytes (LY), platelate count (PLT) Mean platelate volume (PCT) Plateletecritandplatelete distribution weight were determined using the automated haematologic analyzer SYSMEX KX21, a product of SYSMEX Corporation, Japan employing the methods described by Dacie and Lewis [15].

2.8 Determination of Biochemical Parameters

The biochemical analyses were determined for alkaline phosphatase (ALP) based on methods of Tietz, [16], Aspartate transaminase (AST) and alanine transaminase (ALT) as described by Reitman and Frankel, [17]. The serum total protein concentration was estimated by biuret method as described by Gornall et al., [18], albumin as described by Doumas et al., [19], The procedure of Blass et al., 1974 [20], was used to determine serum creatinine concentration while the serum urea concentration was determined by the method of Veniamin and Vakirtzi [21]. The method of Evelyn and Malloy [22], was employed to determine the serum bilirubin concentration of the samples. While Na+, K+, bicarbonate and chloride ions concentrations were determined by flame photometry [16].

2.9 Determination of body weight and organ/body weight ratios.

The body weights of the rats were determined on day 1, and 10 of the experiment and the weight gains were computed. Organ-body weight ratios were computed by expressing the absolute weight of the organs to the body weight of the animals as described below.

Weight gain= Final weight of rat (g) - Initial weight of rat (g)

Organ-body weight ratios= organ weight (g)/body weight (g) x 100

2.10 Statistical Analysis

Data were analyzed using statistical package for social science (SPSS) version 16 and presented as means ± SEM. Comparisons between different groups was done using Analysis of Variance (ANOVA) and Duncan's Multiple Range Test (DMRT). Values of $P<0.05$ were considered as statistically significant as described by Mahajan, [23].

3. Results

3.1 Hematological parameters

Table 1 illustrates the changes in hematological indices in rats following the repeated administration of *Archachatina maginata* haemolymph. The RBC and HCT were significantly ($p < 0.05$) lowered in rats administered 4 ml/kg body weight haemolymph than the control rats, however, rats administered 2 ml/kg body weight showed no significant difference in RBC and HCT as compare with control. Furthermore all haemolymph treated rat showed a significant ($p < 0.05$) increase in HGB, MCV, MCH, RDH and decrease MCHC than the control rats. However, HGB, MCV and PDW count were higher in 2 ml/kg haemolymph administered rats than those administered 4 ml/kg.

The haemolymph also produce a dose dependent significant ($p < 0.05$) increase in WBC, LY and GRA as compared with the control values. While no significant ($p > 0.05$) difference was observed in PLT, MPV, PCT and PDW in rats administered 2 ml/kg haemolymph as compare with the control value, rats administered 4 ml/kg show a significant decrease in these parameters than the control rats.

Table 1: Haematological parameters of rats administered *Achachatina maginata* haemolymph.

Parameters	Control rats	Rat dose with 2 ml/kg Haemolymph	Rats dose with 4ml/kg Haemolymph
WBC (x 10^9 /L)	23.80±2.01 ^{a}	85.10±4.05^b	120.06±3.52^c
GRA (%)	29.30±3.15^a	69.00±3.15^b	73.30±4.06^b
LY(x 10^9 /L)	14.30±1.14^a	25.40±1.22^b	30.90±1.10^c
RBC (x 10^{12} /L)	2.54±0.21 ^{b}	2.44±0.23^b	1.97±0.02^a
HCT (L/L)	0.20±0.11^b	0.26±0.05^b	0.16±0.05^a
HGB(g/l)	126.00±2.07^a	370.06±7.02^c	296.03±5.10^b
MCH(pg)	49.80±1.11^a	151.70±2.59^b	150.40±6.61^b
MCHC(g/L)	646.86±23.01^b	41.70±0.00^a	33.01±0.00^a
MCV(FL)	77.05±4.31^a	105.96±2.17^c	84.00±2.17^b
RDW(FL)	42.60±1.46^a	74.00±2.17^c	47.30±5.19^a
PLT(x 10^9 /L)	295.45±3.12^b	288.98±2.11^b	173.04±4.22^a
MPV(FL)	23.76±2.11 ^{b}	24.06±1.10^b	12.90±0.42^a
PCT(L/L)	0.41±0.05 ^{b}	0.39±0.005^b	0.21 005^a
PDW(%)	2.54±0.21 ^{b}	2.44±0.23^b	1.97±0.02^a

Values are mean ± SEM of 4 determinations. Values along the same row with different superscripts are significantly different ($p < 0.05$).WBC=White Blood Cells; GRA= granulocyte count: LY= lymphocytes; RBC=Red Blood Cells; HCT= haematocrite; HGB=Haemoglobin; MCH= mean corpuscular haemoglobin, MCHC=Mean Corpuscular Haemoglobin Concentration MCV=Mean Corpuscular Volume; PLT= platelate count; MPV= Mean platelate volume; PCT=Plateletecrit; PDW=platelete distribution weight.

3.2 Biochemical parameters

The effects of administration of 2 and 4 ml/kg of haemolymph on the activities of some biomarker enzymes, level of total protein, albumin and total bilirubin of rats are shown in Table 2. There were significant ($P < 0.05$) increases in serum activities of biomarker enzymes of rats administered with the haemolymph at dose of 2 and 4 ml/kg except for ALP whose activity was significantly lower in 2 ml/kg haemolymph administered rat as compared with control rats. The serum activities of ALP and ALT were significantly ($P < 0.05$) higher in rats receiving 4 ml/kg of haemolymph than those receiving 2 ml/kg. However, no significant difference ($P > 0.05$) was observed in AST activities between the two treatment doses.

The *Archachatina maginata* haemolymph also significantly increased (P<0.05) the level of total protein, albumin and total bilirubin as compared with the control rats. There were no significant (P>0.05) difference in the level of albumin and total bilirubin between the 2 doses of the haemolymph administered, however the level of total protein was significant (P< 0.05) higher at dose of 4 ml/kg than 2 ml/kg.

Table 2: Liver function indices of rats administered *Archachatina maginata* haemolymph.

Parameters	Control rats	Rat dose with 2 ml/kg Haemolymph	Rats dose with 4ml/kg Haemolymph
Total bilirubin(umol/l)	8.80±0.37[a]	11.80±0.21[b]	10.43±0.20[b]
Albumin (g/dl)	2.90±0.05[a]	3.30±0.11[b]	3.80±0.01 [b]
Total protein (g/L)	60.90±0.15[a]	68.80±1.23[b]	71.90±2.05[c]
AST(U/L)	36.0±0.27[a]	67.00±2.23 [b]	67.00±0.34 [b]
ALT(U/L)	12.0±0.15[a]	25.20±0.05 [b]	67.00±1.09[c]
ALP(IU/L)	75.30±1.08[b]	56.40±0.98[a]	90.30±1.08[c]

Values are mean ± SEM of 4 determinations. Values along the same row with different superscripts are significantly different (p < 0.05).

3.3Serum Electrolyte, Urea andCreatinine

Table 3 illustrates the changes in the serum electrolyte, urea and creatinine in rats following the administration of *Archachatina maginata* haemolymph. The level of urea and chloride were significantly (P<0.05) lower while the level of creatinine increased in animals treated with 2 and 4 ml/kg of haemolymph than the control rats. However, serum bicarbonate sodium and Potassium level was not significantly (P>0.05) altered.

Table 3: Kidney function indices of rats administered *Archachatina maginata* haemolymph.

Parameters	Control rats	Rat dose with 2 ml/kg Haemolymph	Rat dose with 4 ml/kg Haemolymph
Urea(mmol/L)	4.36±0.20[c]	2.30±0.44[a]	3.90±0.01 [b]
Creatinine(µmol/L)	104.90±0.11[a]	118.97±2.34[b]	125.98±2.05[c]
Sodium (MEq/L)	162.76±3.23[a]	168.97±3.98[a]	163.09±3.64[a]
Potassium(mmol/L)	8.60±0.41[a]	8.20±0.19[a]	7.90±0.97[a]
Chloride(mmol/L)	101.97±4.59[c]	65.45±2.35[a]	95.05±4.56[b]
Bicarbonate(mmol/L)	22.03±0.31[a]	21.60±0.09[a]	23.80±1.05[a]

Values are mean ± SEM of 4 determinations. Values along the same row with different superscripts are significantly different (p < 0.05).

3.4 Body Weight and Organs/Body Ratio

The effects of administration of 2 and 4 ml/kg of *Archachatina maginata* haemolymphon body weight and organs/body ratio of albino rat are shown in Table 4 and 5 respectively. Body weight gain of the rats administered *Archachatina maginata* haemolymph for 10days were not significantly (P>0.05) different from those of the control rats (Table 4). Similarly the computed liver, spleen and heart body weight ratios of the rats were not significantly (P>0.05) different from those of the control rats (Table 5). However, there was significant (P < 0.05) decrease in the kidney/body weight ratio as compared with the control values.

Table 4: weight changes of rats administered *Archachatina maginata* haemolymph.

Parameters	Initial weight (G)	final weight (G)	Weight gain(G)
Control rats	112.70±3.47	126.10±3.59	13.45±2.12[a]
Rat dose 2ml//kg haemolymph	172.40±3.03	186.14±4.54	13.74±1.03[a]
Rats dose 4ml//kg haemolymph	200.40±4.21	214.50±4.40	14.10±2.10[a]

Values are mean ± SEM of 4 determinations. Values along the same row with different superscripts are significantly different ($p < 0.05$).

Table 5: Organ/body weight ratios of rats administered *Archachatina maginata* haemolymph.

Parameters	Control rats	Rats dose with 2ml//kg Haemolymph	Rat dose with 4ml//kg Haemolymph
Liver	3.330±0.110[a]	3.688±0.001[a]	3.418±0.201[a]
Kidney	0.703±0.001[b]	0.5931±0.031[a]	0.607±0.015[a]
Heart	0.321±0.003[a]	0.376±0.004[a]	0.359±0.00[a]
Spleen	0.309±0.015[a]	0.399±0.012[a]	0.338±0.010[a]

Values are mean ± SEM of 4 determinations. Values along the same row with different superscripts are significantly different ($p < 0.05$).

4. Discussion

Hematological parameters generally provide information on deleterious effects of foreign components on the blood and also explain blood-related functions of chemical compounds [24]. White blood cells defend the body against infections or any foreign body. The significant increase on WBC and factors relating to it (WBC, LY and GRA) recorded in this research may indicate immunological response to the constituents of the haemolymph and this augmented the production of more WBC, LY and GRA to improve the health status of the animals. The decreases in RBC and HCT observed in haemolymph treated rats indicate anemic condition which could be attributed to the destruction of the erythroblast, thereby, limiting their synthesis. The decrease in the MCHC in this study was an indication of erythrocytes swelling, however, the increase RDW and MCV reflect hemolytic anemia and cytotoxic effect the haemolymph [25].

The biochemical indices monitored in the serum of rats are useful 'markers' for assessment of tissue damage. The measurement of activities of various enzymes in the body fluids plays a significant role in disease investigation and diagnosis [26], assault on the organs/tissues and to a reasonable extent the toxicity of the drug [27]. Biomarker enzymes can also indicate tissue cellular damage caused by chemical compounds long before structural damage that can be picked by conventional histological techniques [28]. The transaminase (ALT and AST) are 'markers' of liver damage and can thus be used to assess liver cytolysis with ALT being a more sensitive biomarker of hepatotoxicity than AST [29]. Consequently, in the present work marked increase in ALT and AST observed after treatment with haemolymph may have occurred as a result of metabolism of the haemolymph by the liver and hepatic tissue turn-over, as response by the body system towards overcoming stress induced by the test substance.

Alkaline phosphatase is a "*marker*" enzyme for the plasma membrane and endoplasmic reticulum [30]. It is often employed to assess the integrity of plasma membrane [28] and

endoplasmic reticulum. The reduction in the enzyme activities after 2 ml/kg administration of the haemolymph may be attributed to either loss of membrane component (including alkaline phosphatase) into the extracellular fluids [31] or inactivation of the enzyme molecule in situ [32] or depletion of important molecules required by the enzyme for maximum activity [27].This reduction in ALP activities can constitute threat to the life of cells that are dependent on a variety of phosphate group for their vital process e.g. synthesis of major membrane phospholipids, (phosphatidylethanolamine and phosphatidylcholine) since there may be less availability of the phosphate group which consequently affect membrane fluidity and decrease the permeability of the epithelial cells [33]. The increase in the activity of ALP after 4 ml/kg administration of the haemolymph could be a consequence of activation of the enzyme or an increase in the rate of the synthesis of the enzyme induced by the higher dose of components of the haemolymph. This increase could also indicate an enhanced functional activity of the organs (kidney and small intestine) [34], since there will be an increase in the rate of ion transport across the cell membranes in these tissues.

The concentrations of total protein, albumin, total bilirubin, urea, creatinine and electrolytes are useful 'markers' of secretory, synthetic and excretory functioning of the liver and kidney [35] The observed increase in the total protein, albumin and total bilirubin content suggests a compromise of the synthetic ability of the liver arising from the administration of the haemolymph. The haemolymph might have increased the functional activity of the liver by interfering with the equilibrium in the rate of synthesis and destruction, removal or clearance of total protein, albumin and total bilirubin from the system of the animals. A similar finding has been previously reported in rats administered aqueous extract of *Crateva adansonii* [36]. Such increase in total protein could, however, lead to dehydration which is detrimental to cellular homeostasis. This will negatively affect the metabolic activities of the liver and consequently the health of the animals.
The kidneys regulate the excretion of urea and reabsorption of electrolytes into the blood. Filtration occurs at the glomeruli while reabsorption takes place in the renal tubules [37]. When there is compromise of normal glomerular function, substances normally cleared by the kidneys such as urea and creatinine accumulate in the biological fluid. The significant decrease in serum urea following the administration of the haemolymph for 10 days may be due to decreased protein catabolism (as observed in table 2) or renal dysfunction [38], while the significant increase in creatinine content of the serum may be attributed to compromise of the renal functional capacity. The haemolymph might have either interfered with creatinine metabolism leading to increased synthesis or the tissue might have compromised all or part of its functional capacity of tubular excretion [39].

The absence of an effect on the levels of serum bicarbonate sodium and Potassium suggests that the normal excretion of this electrolytes by the kidney was not been adversely affected by the haemolymph. Furthermore, the significant decreases in the levels of chloride ions concentration following the administration of haemolymph suggest that some aspects of tubular functioning as it relates to chloride have been compromised.
The treatment of the rats with 2 and 4 ml/kg of the haemolymph for 10 days in this study does not affect the body weight gain of the animals (Table 4). Organ body weight ratios are normally investigated to determine whether the size of the organ has changed in relation to the weight of the whole animal. The absence of an effect on the computed liver/body, spleen/body and

heart/body weight ratios suggest that the haemolymph did not cause any form of swelling, atrophy and hypertrophy on the organs [40]. However, the significant reduction in kidney/body weight ratio following the 10 days administration of the haemolymph may be attributed to tissue necrosis since the organ is concerned with the excretion of foreign substances [27].

5. CONCLUSION

In conclusion, the results of this study indicate that administration of *Archachatina maginata* haemolymph at the doses of 2 and 4 ml/kg caused changes in the hematological profile and some biochemical parameters of liver and kidney dysfunction with a more pronounced effect on the liver function than the renal function tests and at a high dose than the low dose. Hence, caution should be taken when using the haemolymph of *Achachatina maginata* for therapeutic purpose because it may also have some liver or tissue membrane damaging effect.

Ethical Clearance

Ethical Clearance was given by Federal University of Technology, Minna/Nigerian Ethical Review Board (CUERB) in accordance with internationally accepted principles for laboratory animal use and care as contained in the Canadian Council on Animal Care Guidelines and Protocol Review**.**

REFERENCES

1. WHO. Legal Status of Traditional Medicine and Complementary/ Alternative medicine: A world wide review. WHO Publishing 1, 2001.
2. Jowell T. Herbal Medicines. House of Commons official report (Hansard). 1999; 26: 426
3. Uwalaka RE, Ahaot EO.Poverty Reduction through Snail Farming in Nigeria Inter. J. Vet. Sci., 2013;2(3): 81-84.
4. Pilsbry HA. Family Archatinidae. Manual Onchol 1904; 16: 205- 325.
5. Ohimain EI, Philip CAN, Japhet Y, Akinnusi B, Oyenike FA. Performance and Carcass Characteristics of Giant African Land Snails Fed *Alchornea cordifolia* Leaf Meal In Replacement For Soybean MEAL, Inter, J. Res. In Agric and Food Sci, 2014;1(6):1-4
6. AbereSA, and LameedGA. The medicinal utilization of snails in some selected states in Nigeria. *In*: JC Onyekwelu, VAJ Adekunle and DO Oke (eds.) Proceeding of the first National conference of the Forests and Forest Products Society (FFPs) held in Akure, Ondo State between 11th and 18th of April, 2008. pp: 233-237
7. Akinnusi O. Introduction to snail and snail farming. Triola Publishing Company, Abeokuta. 2002; P70.
8. Fred-Jaiyesimi AA, Awobajo O.Biochemical and Histopathological Investigation Of TheEffect of *Archachatinamarginata*Haemolymph In Normal Rats.Inter. J. of Appl. Bio and Pharm. Tech., 2011;2(4):30-33.
9. Adikwu MU. Evaluation of snail mucinmitifs as rectal absorption enhancer for insulin in non-diabetic rat models. Bio Pharm Bull. 2005;28(9): 1801 -4
10. Ademolu RO, Idowu AB, Agbelusi OM. Effects of stocking density on growth and haemolymph biochemical value of *Archachatinamarginata*. Afr. J Biotech. 2006; 8(12): 2908 – 10
11. Bashir L, ShittuOK, Prince CO, Asmau AN. and Aisha MI. Evaluation of Antioxidant Activity of Giant African Snail (*Archachatinamaginata)*Haemolymph in CCl_4- Induced Hepatotoxixity in albino Rats. Brit J Pharm Res.2015 (in press)

12. CCAC, Canadian Council on Animal Care Guidelines and Protocol Review., 1997.
13. Akinloye OA,Olorode BR. Effect of different feeding conditions on the performance, Haemolymph Biochemical and mineral value of African Giant land Snail. *Archachatinamarginata*. J. of Agric and Environ. 2000; 1: 143 – 147
14. Yakubu MT, Bilbis LS, Lawal M, Akanji MA. Effect of repeated administration of sildenafil citrate on selected enzyme activities of liver and kidney of male albino rats. Nig J. of Pure and Appl. Sci., 2003; 18: 1395- 1400.
15. Dacie, JV, Lewis SM, Practical Haematology, 7th edn. Churchill Livingston. Edingburgh, 1991.
16. Tietz NW. Clinical Guide to Laboratory Tests. 3rd Edition, 1995, Pp. 286 – 288
17. Reitman S, Frankel SA. colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases.American Journal of Clinical Pathology. 1957; 28:56-63
18. Gornall AC, Bardawill CJ, David MM. Determination of serum protein by means of biuret reaction. Journal of Biol.Chem, 1949; 177:751-756
19. Doumas BT, Watson WA, Biggs HG. Albumin standards and measurement of serum album with bromocresol green. Clinical Chemistry Acta, 1971; 31: 87.
20. Blass KG, Thierbert RJ, Lam LK. A study of the mechanism of the Jaffe reaction. J.CliniChem and ClinBiochy, 1974; 12:336-343.
21. Veniamin MP Vakirtzi C. Chemical basis of the carbamidodiacetylmicromethod for estimation of urea, citrulline and carbamyl derivatives. Clin. Chem, 1970; 16:3 – 6.
22. Evelyn KA,Malloy HT. Micro determination of oxyhaemoblobin, methaemoglobin and sulphaemoglobin in a single sample of blood. J. Biol. Chem., 1938; 126, 655-661.
23. Mahajan BK. Significance of differences in means. In: Methods in Biostatistics for Medical and Research Workers, 6th edition. New Delhi: JAYPEE Brothers Medical Publishers. 1997; Pp. 130-155
24. Betancourt-Alonso MA, Orihuela A, Aguirre V, Vázquez R, Flores-Pérez I. Changes in behavioural and physiological parameters associated with *Taeniapisiformis* infection in rabbits (*Oryctolaguscuniculus*) that may improve early detection of sick rabbits. World Rabbit Sci., 2011; 19: 21-30.
25. Tekie H, kin TB, Aklas G, Tan SM, Slit M. the evaluation of red cell distribution width in chronic hemodialysis patient. Int j nephrol, 2014;754370
26. MalomoSO.Toxicological implication of ceftriaxone administration in rats. Nig. J. Biochem.Mol. Biol., 200015(1): 33-38.
27. Yakubu MT, Bilbis LS, Lawal M, Akanji MA. Effect of repeated administration of sildenafil citrate on selected enzyme activities of liver and kidney of male albino rats. NigJ.Pure Appl. Scie., 2003; 18:1395- 1400.
28. Akanji MA, Olagoke OA, Oloyede OB. Effect of chronic consumption of metabisulphite on the integrity of the rat kidney cellular system. Toxicol., 1993; 81: 173-179.
29. Pramyothin P, Samosorn P, Poungshompoo S, Chaichantipyuth C. The protective effects of Phyllanthusemblica Linn. extract on ethanol induced rat hepatic injury. J. Ethnophar.,2006; 107:3, 361–364.
30. Wright PJ, Plummer DT. The use of urinary enzyme measurement of direct renal damage caused by nephrotoxic compound. Bioch. Pharma., 1974; 23, 65–73.

31. Malbica JO, Hart LG.. Effect of adenosine triphosphate (ATP) and some anti-inflammatory agents on purified lysosomal fraction having high acid phosphatase and labile glucouronidase activity. Bioch.Pharma., 1971; 20:2017–2022
32. Umezawa H. Hooper IR. Aminoglycoside Antibiotic. Springer Verk Berlin; Hadelberg, New York, 1982.
33. Yakubu MT, Akanji MA, Oladiji AT. Aphrodisiac potentials of the aqueous extract of Fadogiaagrestis (Schweinf. Ex Hiern) stem in male albino rats. Asian Journal of Androl., 2005; 7: 399-404
34. Brain RT, Kay HD. Kidney phosphatase II: The enzyme in disease.Bioch. Jour., 1927; 21: 1101 – 1108.
35. Yakubu MT, Musa IF.Liver and Kidney Functional Indices of Pregnant Rats Following the Administration of the Crude Alkaloids from Sennaalata (Linn. Roxb) Leaves. Irani. Jour.Toxicol., 2012;6:16, 615-625
36. Salau AK, AkanjiMA, Yakubu, MT.Safety Evaluation of Aqueous Extract of Cratevaadansonii Leaves on SelectedTissues of Rats. Fountain Journal of Natural and Applied Sciences: 2013; 2(1): 17 – 28
37. Mayne PD. Clinical Chemistry in Diagnosis and Treatment. 6th Ed. Lloyd-Luke (Medical Books) Ltd 2005.p 185
38. Whelton A, Watson AY, Rock RC. In: Burtis, C.A., Ashwood, E.R. (Eds.), Tietz Textbook of Clinical Chemistry. W.B. Saunders Company, London, 1994; 1528–1531.
39. Zilva JF, PanmallPR, Mayne PD. Clinical Chemistry in Diagnosis and Treatment, 5th edition, England Clays Ltd., St. Ives Plc., England, 1991.
40. Amresh GR, Singh PN, Rao CV. Toxicological screening of traditional medicine Laghupatha (Cissampelospareira) in experimental animals. J.Ethnopharm., 2008; 116: 454-460.