

Original Research Article

Experimental evaluation of analgesic and antioxidant effects of hydromethanolic extract of *Dioscorea dumetorum* tuber.

Author details: *Celestine O. Ukwueze¹, Samuel O. Onoja², Maxwell I. Ezeja².

¹Department of Veterinary Surgery and Theriogenology, College of Veterinary Medicine, Micheal Okpara University of Agriculture, Umudike, Abia, Nigeria

²Department of Veterinary Physiology, Pharmacology, Biochemistry and Animal Health and Production, College of Veterinary Medicine, Micheal Okpara University of Agriculture, Umudike, Abia, Nigeria
ukwezecele@yahoo.com

ABSTRACT

Aims: The hydromethanolic extract of *D. dumetorum* was evaluated for antinociceptive and antioxidant activities to validate its uses in traditional medicine for these purposes.

Methods: The antioxidant activity of *D. dumetorum* was done in vitro using 2, 2-diphenyl-1-picrylhydrazyl (DPPH) photometric assay. The antinociceptive effect of the extract were investigated using acetic acid-induced writhing reflex and tail flick models with aspirin and pentazocine as reference drugs respectively.

Result: The extract (200 mg/kg) produced a comparable effect to aspirin (100 mg/kg) in the acetic acid induced abdominal writhing reflex. In tail flick respond model, the extract produced significant ($P < 0.05$) dose-dependent increase in pain reaction time (PRT) in the treated rats when compared to the distilled water treated rats. The extract (200 mg/kg) and pentazocine (3 mg/kg) produced 62.86 and 62.52% increase in PRT, respectively when compared to the negative controls. Similarly, DPPH radical scavenging activity of *D. dumetorum* extract produced concentration-dependent increase in the percentage antioxidant activity in the DPPH photometric assay. The extract at 400µg/ml concentration produced antioxidant activity comparable to the standard.

Conclusion: The results of our study confirmed the uses of *D. dumetorum* in traditional medicine and provided evidence that tuber extract of *D. dumetorum* might indeed be potential sources of natural antinociceptive and antioxidant agents.

Keywords: *Dioscorea dumetorum*, antinociceptive, antioxidant, pentazocine, aspirin

INTRODUCTION:

Food is very essential to man and animal as it prevents and reduces several risk factors of diseases and enhances mental and physical well-being [1]. One of such food is *Dioscorea dumetorum* (Bitter yam or trifoliate yam). *Dioscorea dumetorum* occurs in both wild and cultivated forms. It is mainly cultivated in Tropical Africa with Nigeria and Cameroon as the major producers [2]. It is a tuber crop with fleshy edible parts which can be yellow or white and occurs naturally in clusters hence its name cluster yam. It has different names among different ethnic groups in Nigeria. It is called “una” by the Igbos of Southeast Nigeria, *Kosanrogo* in

Hausa, and *Esuru* in Yoruba [1, 2, 3]. The *D. dumetorum* has high nutritive value, containing high crude protein (11.4%), crude fibre (2.03%), carbohydrate (77.55%) and energy (361.90kcal/100g), vitamins, minerals and amino acid [4, 5]. The phytochemical profile has been reported containing, fatty acids, phenols, sterols, aldehydes, ketones, alcohols, hydrocarbons, esters, amines and alkaloid [3, 6].

There are edible and non edible *D. dumetorum*. The edible varieties are cultivated while non edible varieties grow in wild and are not normally eaten except at times of food scarcity. However, the non edible *D. dumetorum* are highly utilized in traditional medicine treatment of some illness such diabetes mellitus, schistosomiasis, jaundice, malaria, as a topical anesthetic and also applied to suppurating abscesses [1, 7, 8]. The antidiabetic, antioxidant and antimicrobial effects of *D. dumetorum* have been reported [2, 8, 9, 10, 11]. These effects have been attributed to some compounds found in the tuber such as 9, 12, Octadecadienoic acid (Linoleic acid) and, 3, 5-Di-t-butyl phenol [6].

However, the analgesic property of the *D. dumetorum* has not been extensively studied owing to the little scientific information on the antinociceptive effects of the plant. This necessitated the design of this study to evaluate the antinociceptive effects of the hydromethanolic extract of *D. dumetorum* in order to validate its uses in traditional medicine for this purpose.

MATERIALS AND METHODS

Fresh tubers of non edible strain of *D. dumetorum* were harvested in the wild from Umuida in Enugu-Ezike, Enugu State, Nigeria.

Preparation of the Plant Extract

The tuber was sliced into smaller pieces, dried and reduced into a coarse powder using the manual grinder (Corona China). The cold maceration method was used for the extraction of coarse powder of *D. dumetorum* (86 g) using 80% methanol in a Winchester bottle for period of 48 h at room temperature. Filtration was done using Whatman No. 1 filter papers, funnel and conical flask. The filtrate was kept in a hot oven at 40°C to enhance the evaporation of the methanol and concentrate the filtrate. The *D. dumetorum* extract (DDE) was stored in a refrigerator at 4°C throughout the duration of this study [12]. The weight of *D. dumetorum* recovered was 13.3 g, and the percentage yield was 15.5% w/w dry matter. The extract was dark brown in color and solid in consistency.

Experimental Animals

Thirty five male Wistar Albino rats (120-140 g) and thirty male mice (28-35 g) were used in the study. The animals were obtained from the laboratory animal unit of the Department of Veterinary Physiology, Pharmacology, Biochemistry, Animal Health and Production; Michael Okpara University of Agriculture, Umudike. They were maintained in accordance with the recommendations of the “guide for the care and use of laboratory animals” [13]. They were kept in clean dusted aluminum cage at 6 animals per cage and were fed *ad libitum* with standard commercial pelleted feed (Vital Feed®) with free access to clean drinking water, kept at normal environmental temperature and natural light/darkness daily cycle. They were allowed 2 weeks to acclimatize before the commencement of the experiment. The animal experiment protocol was approved by the Michael Okpara University of Agriculture Animal Ethics Committee.

Oral Acute Toxicity Study

The oral acute toxicity test of *D. dumetorum* extract (DDE) was determined according to the OECD guideline No. 425 (acute oral toxicity - up-and-down-procedure). Briefly, a group of five rats were dosed orally with 2000 mg/kg of *D. dumetorum* and were observed for 48 h for signs of toxicity and death [14].

In vitro Antioxidant Activities of the *D. dumetorum* Extract

The free radical scavenging activity of DDE was analyzed by the 2,2-diphenyl-1-picrylhydrazyl (DPPH) photometric assay [12]. This was done using 2 ml of the extracts at different concentrations (25, 50, 100, 200 and 400 µg/ml) each mixed with 0.5 mM DPPH (in 1 ml of methanol) in a cuvette and incubated for 30 min in the dark at room temperature. The absorbance at 517 nm was taken after the incubation. The concentrations were prepared in triplicates, and the percentage antioxidant activity (AA) calculated as follows:

$$\% \text{ antioxidant activity (AA)} = 100 - \left(\frac{[\text{absorbance of sample} - \text{absorbance of blank}] \times 100}{\text{absorbance of control}} \right)$$

A volume of 1 ml of methanol plus 2 ml of the extract was used as the blank while 1 ml of the 0.5 mM DPPH solution plus 2 ml of methanol was used as a negative control. Ascorbic acid (vitamin C) was used as a reference standard [15].

Effect of *D. dumetorum* on Acetic Acid-induced Abdominal Writhing in Mice

The acetic acid-induced writhing test measures abdominal constrictions together with stretching of the hind limbs resulting from intraperitoneal (i.p.) injection of 0.7% acetic acid (10mL/kg). This was carried out according to the procedures described by Vale *et al.* [16]. Five Groups (A-E) of mice consisting of 6 mice each were fasted for 12 h but given adequate amount of water. Group A mice received distilled water (10 ml/kg) and served as negative control. Group B mice received aspirin (100 mg/kg) orally (positive control), while Groups C-E mice received 50, 100 and 200 mg/kg of DDE by oral administration, respectively. After 45 min, the mice received 10 ml/kg of 0.7% acetic acid intraperitoneally. Then, the number of writhing or abdominal stretches produced in each mouse was counted for 30 min.

Effects of *D. dumetorum* on Tail Flick Response in Rats

The experiment was carried out by measuring tail withdrawal time from hot water [17]. Thirty rats were randomly divided into 5 Groups (A-E) of 6 rats each and fasted for 12 h. The rats were treated as follows; Group A served as negative control and received 10 ml/kg distilled water (10mg kg) orally, Group B served as positive control and received pentazocine (3mg/kg) intraperitoneally while Group C-E received DDE (50, 100 and 200 mg/kg, respectively) orally. One hour post-drug treatment about 3 cm of the tail of each rat was dipped into a water bath containing hot water maintained at temperature of $50 \pm 1^\circ\text{C}$. The time taken for the rat to flick the tail, known as pain reaction time (PRT) was recorded for all the rats.

Statistical Analysis

Statistical analysis was done using SPSS, version 17.0. Data obtained were presented as mean $\pm$ standard error of the mean and analyzed using one-way Analysis of Variance. The variant mean was separated by least significant difference of the different groups. Significance was accepted at the level of $P < 0.05$.

RESULTS

Oral Acute Toxicity Study

Neither clinical signs of toxicity nor death was recorded following oral administration of the *D. dumetorum* extract at higher dose of 2000 mg/kg at the end of 48 hours period.

Effects of *D. dumetorum* on Acetic Acid-induced Writhing reflex in Mice. The *D. dumetorum* methanolic extract and aspirin produced high inhibition to acetic acid –induced writhing reflex in mice. The DDE (50 mg/kg, 100 mg/kg and 200 mg/kg) produced significant ($P < 0.05$) dose-dependent decrease in the mean number of abdominal constriction in the treated group of mice when compared to the negative control. However, The DDE (200 mg/kg) produced a comparable effect ($P < 0.05$) to aspirin (100 mg/kg) in the acetic acid induced abdominal writhing reflex (Table 1).

Effects of *D. dumetorum* on Tail Flick Response in Rats

In tail flick respond model, the extract produced significant ($P < 0.05$) dose-dependent increase in pain reaction time (PRT) in the treated rats when compared to the distilled water treated rats. The extract (200 mg/kg) and pentazocine (3 mg/kg) produced 62.86 and 62.52% increase in PRT, respectively when compared to the negative controls (Table 2).

Effect of *D. dumetorum* on DPPH Radical Scavenging

The result of DPPH radical scavenging activity of DDE is showed in figure 1. The DDE produced concentration-dependent increase in the percentage antioxidant activity (AA) in the DPPH photometric assay. The increase in concentration from 25µg/ml to 400µg/ml produced an appreciable increase in the percentage AA.

Table 1: Effects of *Dioscorea dumetorum* extract on acetic acid induced writhing reflex in mice

Treatment	Mean number of writhing $\pm$ SEM	% inhibition
Distilled water 10 ml/kg	125.5 $\pm$ 0.78	-
aspirin 100 mg/kg	30.75 $\pm$ 0.62*	75.5
DDE 50 mg/kg	44.44 $\pm$ 0.94*	64.59
DDE 100 mg/kg	39.14 $\pm$ 1.40*	68.81
DDE 200 mg/kg	18.75 $\pm$ 0.81*	85.06

* $P < 0.05$ compared with the negative control (distilled water), SEM: Standard error of mean
DDE: *Dioscorea dumetorum* extract,

Table 2: Effects of *Dioscorea dumetorum* extract on tail flick response in rats

Treatment	Mean PRT $\pm$ SEM (Sec)	% inhibition of pain
Distilled water 10 ml/kg	5.87 $\pm$ 0.20	-
pentazocine 3 mg/kg	9.54 $\pm$ 2.36*	62.52

DDE 50 mg/kg	5.80 ± 0.56	-1.19
DDE 100 mg/kg	8.22 ± 0.35	40.03
DDE 200 mg/kg	9.56 ± 1.07*	62.86

*P<0.05 compared with the negative control (distilled water), SEM: Standard error of mean

DDE: *Dioscorea dumetorum* extract, PRT: Pain reaction time

175




Figure 1: Effect of DDE on DPPH radical scavenging

DISCUSSION AND CONCLUSION

The DDE was well tolerated by the rats, as no death and clinical signs of toxicity were recorded at the high dose of 2000 mg/kg. This indicates that the LD50 is greater than 2000mg/kg dose.

The acetic acid-induced writhing reflex is a sensitive procedure in screening peripherally acting analgesics (18). Intra-peritoneal injection of acetic acid induces an abdominal constriction response which is thought to involve local peritoneal receptors [18, 19]. Acetic acid induces writhing by direct stimulation of acid sensitive receptors or through the release of pain mediators such as histamines, serotonin, and prostaglandins [20]. Prostaglandins sensitize the pain receptor [12]. In this study, the number of writhing movements during a 30min observation in the control group was 125.5 ± 0.78. The aspirin (reference drug) gave a 75.5% inhibition of writhing in the animals while the 200mg/kg dose gave 85.06% inhibition showing that its effect is higher and comparable to that of the reference drug. Active substances against pain may either interfere with pain mediator systems or act on the central nervous system (CNS) by blocking the pain influx

transmission [20]. Aspirin (acetylsalicylic acid) inhibit the production of prostaglandin in CNS and body by inhibiting the cyclooxygenase (COX) enzymes [21]. The observed effects in the present study suggest that the extract may have good peripheral pain relieving effect through inhibitory effect on pain mediator systems.

In addition, the effect of the extract on deep pain was tested using tail flick method, a nociception model for the evaluation of the deep analgesic effects of drugs [12]. The extract caused a significant dose-dependent increase in the PRT which is highly comparable to that of reference drug. The reference drug, Pentazocine is a synthetic opioid agonist-antagonist that produces narcotic analgesia and sedation by an interaction with κ -receptors [12, 22]. The DDE may have acted on the central nervous system (CNS) by blocking the pain influx transmission and provide relief to deep pain sensation which may have been mediated through an increase in the pain threshold in the central nervous system [12, 20].

The efficacies of antioxidants are often associated with their ability to scavenge free radicals [23]. A dose- dependent relationship was found in DPPH radical scavenging effect of DDE. The effect increased as the concentration increased (fig. 1). It suggested that further increase in the concentration of the extract may compare well with the standard drug. Sakthidevi and Mohan [23] reported high DPPH radical scavenging effect of *Dioscorea alata* at concentration of 1000g/kg. The positive DPPH test suggested that DDE was scavenger of free radicals and potential antioxidant. This can be attributed to some of *D. dumetorum* phytoconstituents, such as phenols, flavonoids and tannin. Phenolic compounds are well known as antioxidant and scavenging agents for free radicals associated with oxidative damage [7, 23].

In conclusion, this study has shown that, the methanol extract of *D. dumetorum* showed significant antinociceptive effect and moderate antioxidant activity. This justified its use in traditional medicine as a topical anaesthetic. However, more work is required for the isolation and characterization of the active principles responsible for these activities.

REFERENCES

1. Ogbunugafor HA, Ilodigwe EE, Ajaghaku DL, Ezekwesili CN, Okafor CS, Ajuzieogu CF, et al. *Dioscorea dumetorum*-Fed Rats Exhibited Decreased Body Weight, Blood Glucose, and Insulin in STZ-induced Diabetes. *Functional Foods in Health and Disease*. 2014; 4(2):87- 97
2. Bhogireddy N, Naga AVK, Ramesh B, Pradeep KM, Reddy OVS, Gaddaguti V, et al. Anti-inflammatory and anti-diabetic activities with their other ethnomedicinal properties of the plants. *J. Med. Plants Studies*. 2013;1(5): 587- 96
3. Alozie YE, Lawal OO, Umoh IB, Akpanabiatu, MI. Fatty acid composition of *Dioscorea dumetorum* (Pax) varieties. *Afri. J. Food Agri. Nutri. Dev*. 2010; 10: 2956-66.
4. Polycarp D, Afoakwa EO, Budu AS, Otoo E. Characterization of chemical composition and anti-nutritional factors in seven species within the Ghanaian yam (*Dioscorea*) germplasm . *Intern. Food Res. J*. 2012;19 (3): 985-992

- 246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
5. Alozie Y, Akpanabiatu MI, Eyong EU, Umoh IB, Alozie G. Amino Acid Composition of *Dioscorea dumetorum* Varieties. Pakistan J.Nutri. 2009; 8 (2): 103-105.
 6. Ezeocha VC, Ojmelukwe PC, Onwuka GI. Effect of cooking on the nutritional and phytochemical components of trifoliate yam (*Dioscorea dumetorum*). Glo. Adv. Res. J. Biochem. Bioinform. 2012; 1(2): 26-30.
 7. Sonibare MA, Abegunde RB. In vitro antimicrobial and antioxidant analysis of *Dioscorea dumetorum* (Kunth) Pax and *Dioscorea hirtiflora* (Linn.) and their bioactive metabolites from Nigeria. J. Appl. Biosci. 2012. 51: 3583– 90
 8. Fasaanu OP, Oziegbe M, Oyedapo OO. Investigations of activities of alkaloid of trifoliate yam (*dioscorea dumetorum*, (kunth) pax. Ife J. Sci. 2013; 15(2): 251-62.
 9. Iwu MM, Okunji CO, Ohiaeri GO, Akah P, Corley D. Hypoglycaemic activity of dioscoretine from tubers of *Dioscorea dumetorum*. Planta Medica. 1990; 56: 264-267.
 10. Kelmanson JE, Jager AK, Van Staden J. Zulu medicinal plants with antibacterial activity. Journal of Ethnopharmacology. , 2000; 69: 241-246.
 11. Chan YC, Hsu CK, Wang MF, Su TY. A diet containing yam reduces the cognitive deterioration and brain lipid peroxidation in mice with accelerated senescence. Int. J. Food Sci Tech. 2004; 39: 99-107.
 12. Onoja SO, Ukwueze CO, Ezeja MI, Udeh NE. Antinociceptive and antioxidant effects of hydromethanolic extract of *Bridelia micrantha* stem bark. J. Exp Integr Med. 2014 ; 14 (4) 273-277
 13. DHHS. Guide for the Care and Use of Laboratory Animals. Washington, DC: Institute of Laboratory Animal Resources Commission on Life Sciences, National Research Council, National Academy Press; 1985
 14. OECD. Guidelines for the Testing of Chemicals, Acute Oral Toxicity - Up-and-Down-Procedure 425. [Adopted Mar 2008].
 15. Iwalewa EO, Adewale IO, Aiwo BJ, Arogundabe T, Osinowo A, Daniyan OM, *et al.* Effects of *Harungana madagascariensis* stem bark extract on the antioxidant markers in alloxan induced diabetic and carrageenan induced inflammatory disorders in rats. J Complement Integr Med 2008; 5(1):1-18.
 16. Vale ML, Marques JB, Moreira CA, Rocha FA, Ferreira SH, Poole S, *et al.* Antinociceptive effects of interleukin-4, -10, and -13 on the writhing response in mice and zymosan-induced knee joint incapacitation in rats. J Pharmacol Exp Ther 2003;304: 102-8.

- 292
293 17. Adzu B, Amos S, Wambebe C, Gamaniel K. Antinociceptive activity of *Zizyphus spina-*
294 christi root bark extract. *Fitoterapia* 2001;72:344-50
295
296 18. Adedapo AA and Orherhe VE. Antinociceptive and Anti-Inflammatory Studies of the
297 Aqueous Leaf Extract of *Carica Papaya* in Laboratory Animals. *Asian j. Exp. Biol. Sci.*
298 2013; 4 (1):89-96
299
300 19. Chakraborty A, Devi RKB, Rita S, Sharatchandra K, Singh TI. Preliminary studies on
301 anti-inflammatory and analgesic activities of *Spilanthes acmella* in experimental animal
302 models. *Indian J Pharmacol.* 2004; 36 (3): 148-50.
303 20. Anaga AO, Onehi EV. Antinociceptive and anti-inflammatory effects of the methanol
304 seed extract of *Carica papaya* in mice and rats. *Afr J Pharm Pharmacol* 2010; 4:140-4.
305
306 21. Vane JR, Botting RM. The mechanism of action of aspirin. *Thrombo Res.* 2003; 110:
307 255–8.
308
309 22. Martin B. Opioid and non-opioid analgesic. In: Charles CR, Robert SE, editors. *Modern*
310 *Pharmacology*. 4th ed. Boston: Little Brown and Company; 1994; 431-50.
311
312 23. Sakthidevi G, Mohan VR. Total Phenolic, Flavonoid Contents and In vitro Antioxidant
313 Activity of *Dioscorea alata* l. Tuber. *J. Pharm. Sci. & Res.* 2013; 5(5)115 - 9
314
315
316