

Optimize Diabetes by Herbal Medicine: A Review

Yasodha Krishna Janapti

Department of Pharmaceutical Chemistry, Jawaharlal Nehru Technological University Hyderabad, Sri
Sarada College of Pharmacy, Ananthram (viL), Bhongiri (M), Nalgonda (Dt), Andhra Pradesh, India -
508116

E- Mail: krishna.yasodha@gmail.com,

Yasodha krishna

suniljunapudi@gmail.com

ABSTRACT

Plant-based medicinal products have been acknowledged since ancient times and several medicinal plants and their products have been used to control diabetes in the traditional medicinal systems of many cultures worldwide. Several synthetic oral hypoglycemic agents are the primary forms of treatment for diabetes. However, prominent side-effects of such drugs are the main reason for an increasing number of people seeking alternative therapies that may have less severe or no side effects but little toxicological information exists concerning traditional antidiabetic plants. The present paper is an attempt to list the plants with anti-diabetic and related beneficial effects originating from different parts of the world and polyherbal formulations. History has shown that medicinal plants have been used in traditional healing around the world for a long time to treat diabetes; this is because such herbal plants have hypoglycemic properties and other beneficial properties, as reported in scientific literature. There are some such plants described in this review, which clearly shows the importance of herbal plants and polyherbal formulations in the treatment of diabetes mellitus. The effects of these plants may delay the development of diabetic complications and provide a rich source of antioxidants that are known to prevent/delay different diseased states.

Keywords: - Diabetes mellitus, Hypoglycemic, Plants

1. INTRODUCTION

Diabetes mellitus is a heterogeneous metabolic disorder characterized by altering carbohydrate, lipid and protein metabolism caused by insulin deficiency, often combined with insulin resistance [1]. It is considered as one of the five leading causes of death in the world [2]. About 150 million people are suffering from diabetes worldwide and it was almost five times more than the estimated ten years ago and this may be doubled by the year 2030 [3]. Despite considerable progress in the treatment of diabetes by oral hypoglycemic agents, search for newer drugs continues because the existing synthetic drugs have several limitations. The herbal drugs with antidiabetic activity are yet to be commercially formulated as modern medicines, even though they have been acclaimed for their therapeutic properties in the traditional systems of medicine. The plants provide a potential source of hypoglycemic drugs because many plants and plant derived compounds have been used in the treatment of diabetes. Many Indian plants have been investigated for their beneficial use in different types of diabetes and reports occur in numerous scientific journals. Ayurveda and other traditional medicinal system for the treatment of diabetes, describe a number of plants used as herbal drugs. Hence, they play an important role as an alternative medicine due to less side effects and low cost. The active principles present in medicinal plants have been reported to possess the pancreatic beta cells re-generating, insulin releasing and

fighting the problem of insulin resistance. Hyperglycemia is involved in the etiology of development of diabetic complications. Hypoglycemic herbs increase insulin secretion, enhance glucose uptake by adipose or muscle tissues and inhibit glucose absorption from the intestine and glucose production from liver [4]. Insulin and oral hypoglycemic agents like sulphonylureas and biguanides are still the major players in the management, but there is a quest for the development of more effective anti-diabetic agents [74].

Hyperglycemia occurs because of uncontrolled hepatic glucose output and reduced uptake of glucose by skeletal muscle with reduced glycogen synthesis. When the renal threshold for glucose reabsorption is exceeded, glucose spills over into the urine (glycosuria) and causes an osmotic diuresis (polyuria), which in turn, results in dehydration, thirst and increased drinking (polydipsia). Insulin deficiency causes wasting through increased breakdown and reduced synthesis of proteins.

2. VARIOUS FORMS OF DIABETES MELLITUS [4]

I. General-genetic and other factors not precisely defined

a) Type 1 diabetes mellitus (T1DM OR IDDM).

- Type 1A – Auto-immune type 1 diabetes mellitus.
- Type 1B – Non-auto-immune or idiopathic type 1 diabetes mellitus.

b) Type 2 diabetes mellitus (T2DM OR NIDDM).

II. Specific defines gene mutations

a) Maturity onset diabetes of the young (MODY)

- MODY 1 (Hepatic nuclear factor 4- α gene mutations)
- MODY 2 (Glucokinase gene mutations)
- MODY 3 (Hepatic nuclear factor 1- α gene mutations)
- MODY 4 (Pancreatic determining factor x gene mutations)
- MODY X (Unidentified gene mutations(s))

b) Maternally inherited diabetes and deafness (MIDD)

c) Mitochondrial leucine t-RNA gene mutations

d) Insulin gene mutation

e) Insulin receptor gene mutations

III. Diabetes secondary to pancreatic disease

- Chronic pancreatitis
- Surgery
- Tropical diabetes (chronic pancreatitis associated with nutritional and or toxic factors)
- Gestational diabetes

IV. Diabetes secondary to other Endocrinopathies

- Cushing's disease
- Glucocorticoid administration
- Acromegaly

V. Diabetes secondary to immune suppression

VI. Diabetes associated with genetic syndromes eg: Prader-Willi syndrome

VII. Associated with diabetes drug therapy

a) Drugs with Hypoglycemic effects

β -Adrenergic receptor antagonists,
Salicylates, Indomethacin, Clofibrate and ACE inhibitor.

b) Drugs with Hyperglycemic effects

Epinephrine, Glucocorticoid, β -adrenergic receptor agonist, Calcium channel blockers, Phenytoin and H_2 receptor blocker

Virtually all forms of diabetes mellitus are caused by a decrease in the circulating concentration of insulin (insulin deficiency) and a decrease in the response of peripheral tissues to insulin (insulin resistance). These abnormalities lead to alterations in the metabolism of carbohydrates, lipids, ketones and amino acids. The central feature of the syndrome is hyperglycemia characterized by a high blood glucose concentration.

Hyperglycemia may become toxic and fatal as a result of accumulation of non-enzymatically glycosylated products and osmotically active sugar alcohols such as sorbitol in tissues, the effects of glucose on cellular metabolism also may be responsible [5]. Diabetic ketoacidosis is an acute emergency which develops on prolonged hyperglycemia and develops in the absence of insulin because of an accelerated fat breakdown to acetyl CoA, which in the absence of aerobic carbohydrate metabolism is converted to acetoacetatae, β -hydroxybutyrate, that causes acidosis [6]. Hemoglobin undergoes glycosylation on its amino-terminal Valine residue to form the glucosyl Valine adducts of hemoglobin, termed hemoglobin A1c. The half-life on the modified hemoglobin is equal to that of the erythrocytes (about 120 days). Since the amount of glycosylated protein formed is proportional to the glucose concentration and the time of exposure of the protein to glucose, the concentration of hemoglobin, A1c in the circulation reflects the severity of the glycemic state [4].

2.1 COMPLICATIONS OF THE DISORDER [7]:

Most diabetes complications arise from prolonged exposure of tissues to elevated glucose levels. These complications develop as a consequence of the metabolic derangements in diabetes, often over many years. Many of these result in diseases of blood vessels, either large (macro vascular disease) or small (microangiopathy). Macro vascular disease consists of accelerated atheroma, which is much more common and severe in diabetic patients. Microangiopathy is a distinctive feature of diabetes mellitus and particularly affects the retina, kidney and peripheral nerves. Diabetes mellitus is the commonest cause of chronic renal failure, which itself represents a huge and rapidly increasing problem, the costs of which to society as well as to individual patients are staggering. Coexistent hypertension promotes progressive renal damage nephropathy and reduces the risk of myocardial infarction. Angiotensin-converting enzyme inhibitors or antagonists of angiotensin AT₁ receptors are more effective in preventing diabetic nephropathy than other antihypertensive drugs, perhaps because they prevent fibroproliferative actions of angiotensin- II and aldosterone. Although the underlying mechanism of diabetic complications is unclear much attention has been focused on the role of oxidative stress, which contributes to the pathogenesis of different diabetic complications.

Traditional treatments for diabetes may create a feeling of improved well-being without necessarily reducing hyperglycemia. There have been unsubstantiated claims that certain plants can ameliorate complications of diabetes (Table No:1), but the objective assessment of complications is difficult and the placebo effect of natural remedies cannot be discounted. No controlled studies of microvascular, macrovascular, or neuropathic complications have been undertaken with traditional therapies. A plant that is thirst quenching or increases sympathetic tone might gain a traditional reputation as efficacious without providing any long-term benefits to the underlying malady. Members of the *Allium* family, particularly garlic, are traditionally considered to give strength and combat arterial vascular disease [8]. Garlic reduces polydipsia and weight loss in severely streptozocin-induced diabetic mice without improving glycemic control [9]. Onion has been reported to lower free fatty acid concentrations in healthy subjects, and seeds of *T. foenumgraecum* (fenugreek) reduced cholesterol levels in diabetic dogs [10, 11]. Other traditional antidiabetic plants have been anecdotally claimed to possess hypolipidemic properties.

Table No 1. Antidiabetic plants traditionally considered as efficacious in treatment of diabetes-associated complications

Complications	Plants
Polydipsia	<i>Panax ginseng</i> , <i>Polygonatum humile</i> , <i>Polygonatum macropodum</i> , <i>Polygonatum officinale</i>
Emaciation	<i>Allium sativum</i> , <i>Cichorium intybus</i>
Atherosclerosis	<i>Allium cepa</i> , <i>Allium sativum</i> , <i>Lycium chinensis</i> , <i>Panax ginseng</i> , <i>Taraxacum officinale</i> , <i>Trigonella foenumgraecum</i>
Retinopathy	<i>Daucus carota</i> , <i>Lycium chinensis</i> , <i>Taraxacum officinale</i> , <i>Vaccinium myrtillus</i>
Nephropathy	<i>Lycium chinensis</i>
Impotence	<i>Ceiba petandra</i> , <i>Coriandrum sativum</i> , <i>Crocus sativa</i> , <i>Panax ginseng</i> , <i>Papaver somniferum</i>

2.2 INSULIN THERAPY:

Insulin is the mainstay for treatment of virtually type-1 DM and many type-2 diabetes mellitus patients. When necessary, insulin may be administered intravenously or intramuscularly; however, long-term treatment relies predominantly on subcutaneous injection of the hormone. Subcutaneous administration of insulin differs from physiological secretion of insulin in at least two major ways: The kinetics do not produce the normal rapid rise and decline of insulin secretion in response to ingestion of nutrients, and the insulin diffuses into the portal circulation; the direct action of secreted insulin on hepatic metabolic processes is thus eliminated.

3. SCREENING METHODS OF ANTIDIABETIC AGENTS [12, 13]:

Before the advent of insulin and oral hypoglycemic drugs, the major form of treatment involved the use of plants. More than 400 plants have been recommended and recent investigations have affirmed the potential value of some of these treatments. The hypoglycemic and/or anti-hyperglycemic effect of several plants used as anti-diabetic remedies have been confirmed and the mechanisms of their actions are being studied. Chemical studies directed at the isolation, purification and identification of the substances responsible for the anti diabetic activities are also being conducted various diabetogens.

Induction of experimental diabetes in the rat by particularly destroying the pancreatic beta cells is the most convenient method employed now a day and the most commonly used diabetogenic agents are alloxan and streptozotocin.

4. HERBAL DRUGS USED IN THE TREATMENT OF DIABETES:

In the recent times many traditionally used medicinally important plants were tested for their anti-diabetic potential by various investigators in experimental animals. Several synthetic oral hypoglycemic agents that are used as conventional medicine today have a natural plant origin, eg: metformin was derived from the plant, *Galega officinalis* (Goat's Rue or French Lilae) which was a common remedy for diabetes [14, 15]. The synthetic hypoglycemic agents used in clinical practices have serious side effects like hematological effects, coma, disturbs the functions of liver and kidney. In addition, they are not suitable for use during pregnancy. Compared with synthetic drugs, drugs derived from plants are frequently considered to be less toxic with fewer side effects. Therefore, the search for more effective and safer antidiabetic agent has become an area of active research. Most commonly used plants in herbal formulations in the treatment of diabetes are given in Table No: 2. The list of poly herbal formulation used is shown Table No: 3. The use of natural sweetener alternative to table sugar, artificial sweeteners like barley malt, maple syrup, rice syrup, fructose (fruit sugar) and the herb stevia are utilized more today. Stevia is an excellent natural alternative to artificial sweeteners. It is very concentrated, so only a small amount is used and it doesn't have the side effects that artificial sweeteners have. Aloe Vera, cinnamon, ginseng, onion, yam and ginseng powder are used as prediabetic remedy [71].

Table No 2. Plants reported having antidiabetic activity.

Plants	References
<i>Abelmoschus moschatus</i> Medik	85
<i>Abrus precatorius</i> L.	82
<i>Acacia arabica</i>	82
<i>Achyranthes aspera</i> L.	85
<i>Achyrocline satureioides</i>	86
<i>Acosmium panamense</i> Schott.	85
<i>Aegle Marmelos</i>	16
<i>Aerva lanata</i>	17
<i>Aegle marmelos</i>	18
<i>Agrimony eupatoria</i> L.	87
<i>Ajuga iva</i> L. Schreber	85

<i>Allium cepa</i>	19,82
<i>Allium sativum</i>	20,82
<i>Alpinia galangal</i>	21
<i>Aloe vera and Aloe barbadensis</i>	82
<i>Andrographis paniculata</i> Burm	88
<i>Annona squamosa</i> L	89
<i>Aporosa lindleyana</i>	22
<i>Artemisia herba-alba</i> Asso	90
<i>Artemisia dracunculus</i> L	91
<i>Astragalus membranaceus</i> Bunge	85
<i>Averrhoa bilimbi</i>	23
<i>Azadirachta indica</i>	24
<i>Barleria lupulina</i>	25
<i>Bauhinia candicans</i>	26,92
<i>Bauhinia forficata</i> Link	85
<i>Berberis aristata</i>	27
<i>Bidens pilosa</i> L	93
<i>Biophytum sensitivum</i>	94
<i>Bixa orellana</i> L	95
<i>Boerhaavia diffusa</i>	28,29
<i>Boswellia glabra</i>	30
<i>Bougainvillea spectabilis</i>	31
<i>Brassica nigra</i> (L) Koch	96
<i>Bryonia alba</i> L	97
<i>Bumelia sartorum</i> Mart	98
<i>Caesalpinia bonducella</i>	32
<i>Cajanus cajan</i> (L) Millsp	99
<i>Carum carvi</i>	85
<i>Casearia esculenta</i> Roxb	100
<i>Cassia auriculata</i> L	101
<i>Catharanthus roseus</i>	102
<i>Chamaemelum nobile</i>	103
<i>Cichorium intybus</i> L	104
<i>Clausena anisata</i> (Willd) Benth	105
<i>Coriandrum sativum</i> L	106
<i>Combretum lanceolatum</i>	81
<i>Cuminum cyminum</i> L	107
<i>Curcuma longa</i>	33
<i>Cuminum nigrum</i> L	108
<i>Cyamopsis tetragonoloba</i> (L) Taubert	109
<i>Dioscorea dumetorum</i>	110
<i>Eclipta alba</i> (L) Hassk	111
<i>Embllica officinalis</i> Gaertn	112
<i>Enicostema littorale</i> blume	113
<i>Eugenia jambolina</i>	34
<i>Ficus bengalensis</i> L	114
<i>Fraxinus excelsior</i>	35
<i>Garcinia kola</i> Heckel	115
<i>Gongronema latifolium</i> Endl	116
<i>Gymnema Sylvestre</i>	36
<i>Helicteres isora</i> L	117,118
<i>Holostemma ada kodien</i>	72
<i>Hypoxis hemerocallidea</i> conn Corm	119
<i>Inula racemosa</i> Hook	85
<i>Lagerstroemia speciosa</i> (L) Pers	120
<i>Lepidium sativum</i> L	121

<i>Mangifera indica</i>	82
<i>Momordica charantia</i>	37,38,82
<i>Morinda lucida</i> Benth	85
<i>Myrcia uniflora</i> Barb	122
<i>Nigella sativa</i> L	123
<i>Ocimum gratissimum</i>	39
<i>Ocimum sanctum</i> L	124,125
<i>Origanum vulgare</i> L	126
<i>Otholobium pubescens</i> L	127
<i>Paeonia lactiflora</i> Pall	128
<i>Panax ginseng</i>	40
<i>Pterocarpus marsupium</i>	41
<i>Phyllanthus amarus</i> Schum & Thonn	129
<i>Psidium guajava</i> L.	130,131
<i>Pterocarpus marsupium</i> Roxb	132,133
<i>Retama raetam</i>	134,135
<i>Salacia reticulate</i> W	136
<i>Spergularia purpurea</i>	137
<i>Suaeda fruticosa</i> (SF) Euras	138
<i>Syzygium cumini</i>	42
<i>Talinum cuneifolium</i>	73
<i>Tamarindus indica</i> L	139
<i>Telfaria occidentalis</i> Hook	85
<i>Terminalia catappa</i>	43
<i>Terminalia chebula</i>	44
<i>Trigonella foenum graecum</i>	82
<i>Tinospora cordifolia</i>	82
<i>Urtica dioica</i>	45
<i>Vinca rosea</i>	46
<i>Zizypus jujube</i>	47
<i>Sphenocentrum jollyanum</i>	75
<i>Ocimum canum</i>	76
<i>Jatropha curcas</i>	77
<i>Lawsonia inermis</i>	78
<i>Ficus hispida</i>	79
<i>Pergularia daemia</i>	80
<i>Wattakaka volubilis</i>	82

Table No 3. Some of the Polyherbal formulations used in treatment of Diabetes mellitus.

Formulation	Reference
<i>Acanthopanax senticosus</i>	83
<i>Aavaraiyathi churnam</i>	84
<i>Alnus hirsuta</i>	83
Ayushan- 82	48
Diabecon (D- 400)	49
Dianex	50
Diarun plus65	51
<i>Rosa davurica</i>	83
<i>Panax schinseng</i>	83
MA- 471	52

5. THE MECHANISM INVOLVED IN THE TREATMENT OF DIABETES

The present treatment of diabetes is focused on controlling and lowering blood glucose to a normal level. The mechanisms may be

- To stimulate cell of pancreatic islet to release insulin;
- To resist the hormones which rise blood glucose;
- To increase the number or rise the potency and sensitivity of insulin receptor sites to insulin;
- To decrease the leading-out of glycogen;
- To enhance the use of glucose in the tissue and organ;
- To clear away free radicals, resist lipid per oxidation and correct the metabolic disorder of lipid and protein;
- To improve microcirculation in the body.

Anti-diabetic medicinal plants undoubtedly have a significant effect on the lowering of blood sugar, but their mechanism of action is yet to be elucidated. The first evidence that the natural products have insulin potentiating activity was reported in 1929 by Glazer and Halpern. There are several mechanisms through which these herbs act to control the glucose level. They are more or less similar actions to the synthetic drugs. The mechanism of action of herbal anti-diabetics could be grouped as:

- Stimulation of insulin secretion (*Allium sativum* [20], *Allium cepa* [19], *Panax ginseng* [40])
- Inhibition in renal glucose reabsorption (*Fraxinus excelsior* [35])
- Stimulation of glycogenesis and hepatic glycolysis (*Momordica charantia* [38])
- Protective effect on the destruction of the beta-cells (*Thea sinensis* [53])
- Improvement of digestion and reduction of blood sugar and urea (*Aegle marmelos* [18])
- Prevents pathological conversion of starch to glucose (*Eugenia jambolina* [34], *Tinospora cordifolia* [34], *Pterocarpus marsupium* [41])
- Increasing the use of glucose by tissues and effect on adrenergic receptors (*Panax Ginseng* [40], *Allium sativum* [20], *Allium cepa* [19])
- Potentiates the action of exogenously injected insulin (*Nelumbo nucifera* [54], *Rubia cordifolia* [55])
- Cortisol lowering activities (*Boerhaavia diffusa* [28,29], *Ocimum sanctum* [56])

Over 150 plant extracts some of the active principles, including flavonoids are known to be used for the treatment of diabetes [25, 57-59]. Huff and Howell reported that Quercetin stimulate insulin release and enhance calcium uptake from isolated cells, which suggest a place for flavonoids in NIDDM [60, 61]. The presence of 3' and 4' -OH in the B-ring and double bonds between C₂ and C₃ in flavones and flavonols are important factors for binding enzyme glycogen phosphorylase[62] which is a key enzyme in the regulation of glycogen metabolism and catalyzes a degradative phosphorylysis of glycogen to glucose.

6. TOXIC EFFECTS OF HERBS:

Little toxicological information exists concerning traditional antidiabetic plants. Use of the plants over many centuries and sometimes as regular constituents of the diet might be expected to reveal any obvious detrimental side effects through the cumulative knowledge of personal experiences. Nevertheless, patients are prone to overindulge in natural treatments, believing their efficacious reputation to imply safety, but chronic consumption of large amounts of traditional remedies must always be regarded with caution.

Catharanthus roseus (periwinkle) is widely used as a traditional treatment for NIDDM. Many major alkaloids isolated from this plant, including leurosine, vindoline, vindolinine, and catharanthine, recorded as mild hypoglycemic effect within 2-5 h in healthy rats, but none was sufficiently potent to encourage further investigation because of cytotoxic and neurological effects of *Catharanthus* alkaloids [63].

Ficus bengalensis (banyan tree), used traditionally in Asia for the treatment of DM, but studies on rats observed gross behavioral, neurologic, autonomic toxic effects [64]. A central nervous stimulation and hypertensive effect of ephedrine from *Ephedra* species and central nervous stimulation by *Panax* species has limited their usage in the treatment of diabetes [63, 40].

Unripe fruits of *Blighia sapida* (ackee fruit). A traditional treatment for diabetes in Central America and Africa. Hypoglycins (aminopropylpropionic acid derivatives) are effective in healthy and diabetic animals and humans, promoting glucose use and inhibiting gluconeogenesis secondary to the inhibition of long-chain fatty acid oxidation [65]. However, their toxic effects have precluded further development because it induces neuroglycopenia [65]. Other noncultivated traditional antidiabetic mushrooms such as *C. comatus* can accumulate heavy metals with toxic consequences if consumed in excess [63]. The poisonous effects of *A. phalloides* are probably due to neuroglycopenia after hepatic glycogen depletion and hepatic necrosis [66].

A study in South Africa recorded cases of spontaneous hypoglycemia, hepatic and renal necrosis after consumption of herbal medicines like *Adenia gummifera* (imfulwa), *Chenopodium Ambrosioides*, *Spilanthes Mauritania*, *Iboza riparia* [67]. A case-study report noted that the hypoglycemic effect of *M. charantia* was additive to that of chlorpropamide, and *V. myrtillus*, and synthalin apparently reduced insulin requirements [68, 69, 70].

7. CONCLUSION

This is an exciting time in medical and diabetic history. Diabetes is world's fast emergent metabolic disorder and a knowledge of this disorder increases, similar more appropriate therapies. Traditional plant medicines are used throughout the world for diabetes. So a study such medicine might offer a natural key to unlock a researcher in the future.

ACKNOWLEDGEMENTS

I am very much thankful to the Sri Sarada College of Pharmacy, for his guidance, kind help and constant encouragement at every step during the progress of my work.

CONSENT

Not applicable.

ETHICAL APPROVAL

Not applicable

COMPETING INTERESTS

Author was declared that no competing interests exist.

ABBREVIATIONS

Not applicable

References

1. Rang and Dale. Pharmacology. International print-opera limited, Noida, UP, 2005:385.
2. Vats V, Yadav S P, Grover JK. Ethanolic extract of *Ocimum sanctum* leaves partially attenuates streptozocin induced alteration in glycogen content and carbohydrate metabolism in rats. *J Ethnopharmacology*. 2004; 90:155.
3. Gosh R, Sharachandra KH, Rita S, Thokchom IS. Hypoglycemic activity of *Ficus hispida* (bark) in normal and diabetic albino rats. *Indian journal pharmacology* 2004; 36:222.
4. Laurance LB, John SL, Keith LP. Goodman and Gilman's: The pharmacological basis of therapeutics. Mc Graw-Hill medical publishing division. New Delhi. 11th ed. 2006:1620.
5. Brownlees M. The pathological implications of protein glycation. *Clin. Invest. Med.* 1995; 18:275.
6. Satyanarayana U, Chakrapani U. Biochemistry. Book and allied (p) Ltd. Kolkata. 3rd ed. 2006:294.

7. Larmer J. Insulin and oral hypoglycemic drugs, glucogan .In: Gilman AG, Goodman LS, Rall TW, Murad F, Editors. The pharmacological basis of therapeutics. 7th Edition New York: Macmillan Publishing; 1985:1490.
8. Day C. The Allium Alliance. Nutr Food Sci 1984; 90:20.
9. Swanston-Flatt SK, Day C, Bailey CJ, Flatt PR. Traditional plant treatments for diabetes: studies in normal and streptozotocin diabetic mice. Diabetes Res Clin Pract. In press
10. Augusti KT, Benaim ME. Effect of essential oil of onion (allyl propyl disulphide) on blood glucose, free fatty acid and insulin levels of normal subjects. Clin Chim Acta 1975; 60:121.
11. Valette G, Sauvaire V, Baccou JC, Ribes G: Hypocholesterolemic effect of fenugreek seeds in dogs. Atherosclerosis 1984; 50:105.
12. Srinivasan K, Ramarao R. Animal models in type 2 diabetes research: An overview. Indian J. Med. Res. 2007; 125:451.
13. Bell HR, Hye RJ. Animal models for diabetes mellitus: physiology and pathology. J Surg Res. 1983; 35:433.
14. Oubre AY, Carlson TJ, King SR, Reaven GM. From plant, to patient: an ethnomedical approach to the identification of new drugs for the treatment of NIDDM. Diabetologia 1997; 40:614.
15. Gloria YY, David ME, Kaptchuk TJ, Russell SP. Systematic Review of Herbs and Dietary Supplements for Glycemic Control in Diabetes. Diabetes Care 2003; 26:1277.
16. Sharmila U, Kshama KS, et al. A study of hypoglycemic and antioxidant activity of Aegle marmelos in alloxan induced diabetic rats. Indian J Physiol Pharmacol 2004; 48:476.
17. Vetrichelvan T, Jagadeeshan M. Antidiabetic activity of alcoholic extract of Aerva lenata (L) Juss. Ex Schultes in rats. J Ethnopharmacol. 2002; 80:103.
18. Das AV, Padayatti PS, Paulose CS. Effect of leaf extract of Aegle marmelose (L.) Correa ex Roxb. on histological and ultrastructural changes in tissues of streptozotocin induced diabetic rats. Indian J Exp Biol 1996; 34:341.
19. Kumari K, Mathew BC and Augusti KT. Antidiabetic and hypolipidemic effects of S-methyl cysteine sulfoxide isolated from Allium cepa Linn. Indian J Biochem Biophys 1995; 32: 49-54.
20. Augusti KT and Sheela CG. Antiperoxide effect of S-allyl cysteine sulfoxide, an insulin Secretagogue, in diabetic rats. Experientia 1996; 52: 115.
21. Akthar MS, Khan MA, Malik MT. Hypoglycemic activity of Alpinia galangal rhizome and its extracts in rabbits. Fitoterapia 2002; 73:623.
22. Jayakar B, Suresh B. Antihyperglycemic and hypoglycemic effect of Aporosa lindleyana in normal and alloxan induced diabetic rats. Journal of Ethnopharmacology, 2003: 84:247.
23. Pusparaj P, Tan CH, Tan BHK. Effects of Averrhoa bilimbi leaf extract on blood glucose and lipids in streptozotocin diabetic rats. J Ethnopharmacology. 2000; 72:69.
24. Gupta S, Kataria M, et al. The protective role of extracts of neem seeds in diabetic caused by streptozotocin in rats. J Ethnopharmacology. 2004; 90:185.
25. Suba V, Murugasen R, et al. Antidiabetic potential of Barleria lupulina in rats. Fitoterapia 2004; 75: 1.
26. Fuentes O, Arancibia AP, Alarcon J. Hypoglycemic activity of Bauhinia candicans in diabetic induced rabbits. Fitoterapia 2004; 75:527.
27. Bhupesh CS, Kamal S, et al. Antidiabetic activity of steam bark of Berberies aristata D.C. in alloxan alloxan induced diabetic rats. The internet journal of pharmacology 2008; 6:1.
28. Pari L and Amamath Satheesh M: Hypoglycemic activity of Boerhaavia diffusa L. Effect on hepatic key enzymes in experimental diabetes. J Ethnopharmacology 2004; 91: 109.
29. Satheesh MA and Pari L: Antioxidant effect of Boerhavia diffusa L. In tissues of alloxan induced diabetic rats. Indian J Exp Biol 2004; 42: 989.
30. Kavitha JV, Rosario JF, et al. Hypoglycemic and other related effects of Boswellia glabra in alloxan induced diabetic rats. Indian J Physiol Pharmacol 2007; 51:29.
31. Senapati AK, Dash GK, Ghosh T, Christina AJM. A study on antiinflammatory and hypoglycemic activity of Bougainvillea spectabilis. Indian J. Nat. Prod. 2006; 22:3.
32. Kannur DM, Hukkeri VI, Akki KS. Antidiabetic activity of Caesalpinia bonducella seed extracts in rats. Fitoterapia 2006; 77:546.
33. Ali Hussain HEM. Hypoglycemic, hypolipidemic and antioxidant properties of combination of Curcumin from Curcuma longa, Linn. and partially purified product from Abroma augusta linn. In streptozotocin induced diabetes. Indian J of clinical biochemistry 2002; 17:33.

34. Grover JK, Vats V and Rathi SS: Anti-hyperglycemic effect of *Eugenia jambolana* and *Tinospora cordifolia* in experimental diabetes and their effects on key metabolic enzymes involved in carbohydrate metabolism. *J Ethnopharmacology* 2000; 73:461.
35. Eddouks M, Maqhrani M. Phlorizin-like effect of *Fraxinus excelsior* in normal and diabetic rats. *J Ethnopharmacology* 2004; 94:149.
36. Sugihara V, Nojima H, et al. Antihyperglycemic effects of *Gymema sylvestra* leaves in streptozocin induced mice. *J Asian Nat. Prod Res.* 2000; 12:321.
37. Virdi J, Sivakami S, et al. Anti hypoglycemic effects of three extracts from *Momordi charantia*. *J Ethnopharmacology* 2003; 88:107.
38. Sarkar S, Pranava M and Marita R. Demonstration of the hypoglycemic action of *Momordica charantia* in a validated model of diabetes. *Pharmacol Res* 1996; 33:1.
39. Aguiyi JC, Obi CI, Gang SS, Igweh AC. Hypoglycemic activity of *Ocimum gratissimum* in rats. *Fitoterapia* 2000; 71:444.
40. Park EY, Kim HJ, Kim YK, Park SU et al. Increase in Insulin Secretion Induced by *Panax ginseng* Berry Extracts Contributes to the Amelioration of Hyperglycemia in Streptozotocin-induced Diabetic Mice. *J Ginseng Res*, Vol. 2012; 36:153.
41. Manickam M, Ramanathan M, Jahromi MA, et al. Antihyperglycaemic activity of phenolics from *Pterocarpus marsupium*. *J Nat Prod* 1997; 60: 609.
42. Kumar A, Iivarasan R, et al. Antidiabetic activity of *Syzygium cumini* and its isolated compounds against streptozocin induced diabetic rats. *Journal of medicinal plant research* 2008; 2:246.
43. Mansoor SA, Swamy VBM, Kumar p, Danapal R. Antidiabetic activity of *Terminalia catappa* Linn. Leaf extracts in alloxan induced diabetic rats. *Iranian J pharmacology and therapeutics* 2005; 4:36.
44. Sabu MC, Ramadasan K. Antidiabetic activity of medicinal plants and its relationship with their antioxidant property. *J Ethnopharmacology* 2002; 81:155.
45. Mohamed B, Fatima ZM, et al. Antihyperglycemic activity of the aqueous extract of *Urtica dioica*. *Fitoterapia* 2003; 74:677.
46. Sumana G, Suryawanshi SA. Effect of *Vinca rosea* extracts in treatment of alloxan induced diabetic male albino rats. *Indian j Exp. Biology* 2001; 39:748.
47. Aydin E, Fahrettin K, Hulusi A, Husseyin U, Yalcin T, Muzaffer U. Hypoglycemic effect of *Zizyphus jujube* Leaves. *J Pharm Pharmacol* 1995:4772.
48. Chowday DP, Dua M, Kishore B, Kishore P. Hypoglycemic effect of code formulation: Ayshu-82. *Journal of research in Ayurveda and Sidda*, 1998; 19:107.
49. Dhawan D, Bandhu HK, Singh B, Singh A. Nagapal JP. Effect of D-400 (A herbal formulation) on the regulation of glucose metabolism in diabetic rats. *Indian journal pharmacology* 1996; 28:224.
50. Mutalik S, Chetana M, Sulochana B, Devi PU. Effect of Dianex, a poly herbal formulation on experimental induced diabetic mellitus. *Phytother Res.* 2005; 19:409.
51. Senthilvel G, Jagadeesan M, et al. Effect of polyherbal formulation (Diarunal Plus) on streptozocin induced experimental diabetes. *International journal of tropical of Tropical Medicine* 2006; 1:88.
52. Sircar AR, Ahuja RC, et al. Hypoglycemic, hypolipidemic and general beneficial effects of an herbal mixture MA-471. *Alternative Therapies in Clinical practice* 1996; 3:26.
53. Donga JJ, Surani VS, Sailor GU, Seth AK. A systematic review on natural medicine used for therapy of diabetes mellitus of some Indian medicinal plants. *An Int. J. Phar. Sci.* 2011; 2:36.
54. Pulok KM, Kakali S, Pal M, Saha BP Effect of *Nelumbo nucifera* rhizome extract on blood sugar level in rats. *J Ethnopharmacology* 1997;58:207.
55. Rahul SS, Kishor SJ, Abhay KS. hypoglycaemic activity of roots of *Rubia cordifolia* in normal and diabetic rats. *Pharmacologyonline* 2007; 1:162.
56. Rai V, Iyer U and Mani UV: Effect of *Tulasi (Ocimum sanctum)* leaf powder supplementation on blood sugar levels, serum lipids and tissues lipids in diabetic rats. *Plant Foods Hum Nutr* 1997; 50: 9.
57. Meiselman HL, Halpern BP, Dateo GP. Reduction of sweetness judgement by extracts from the leaves of *Zizyphus jujube* *Physiology and Behavior* 1976; 17:313.
58. Choi JS, Yokozawa T, Oura H. Improvement of hyperglycemia and hyperlipidemia in streptozocin – diabetic rats by methanolic extract of *Prunus davidiana* stems and its main component, prunigen. *Planta Med* 1991; 57; 208.
59. Ermenisoglu A, Kelestimur F, Koker AH, et al. Hypoglycemic effect of *Zizyphus Jujube* leaves. *J Pharm pharmacology* 1995; 47:72.
60. Hiff CS, Howell SL. Effect of epicatechin on rats isolated langerhans. *Diabetes* 1984; 33:291.

61. Tapas AR, Sarkarkar DM, Kakde RB. Flavonoids as nutraceuticals: A review. Tropical journal of Pharmaceutical research. 2008; 7:1089.
62. Atsushi K, Norio N, et al. Structural activity relationships of flavonoids as potential inhibitors of glycogen phosphorylase. J. Agric. Food. Chem. 2008; 56:4469.
63. Clifford JB, Caroline D. Traditional plant medicine as treatment for diabetes. Diabetes care 1989; 12:553.
64. Rakesh KS, Shikha M, Dolly J, Prashant KR, Geeta W. Antidiabetic effects of Ficus bengalensis aerial roots in experimental animals. J Ethnopharmacology 2009; 123:110.
65. Bressler R, Corridedor C, Brendel K. Hypoglycin and hypoglycin-like compounds. Pharmacol Rev1969; 21:105.
66. Potron M. Champignons et diabete. Concours Med 1956; 36:3795.
67. Neame PB, Pillay VKG. Spontaneous hypoglycemia, hepatic and renal necrosis following the intake of herbal medicines. S Afr Med J 1964; 38:729.
68. Allen FM. Blueberry leaf extract: physiologic and clinical properties in relation to carbohydrate metabolism. JAMA 1927; 89:1577.
69. Wilder RM, Allan FN. Synthalin, blueberry leaf extract and glukohormet. JAMA 1928; 38:254.
70. Aslam M, Stockley IH. Interaction between curry ingredient (karela) and drug (chlorpropamide). Lancet 1979; 1:607.
71. Jeffrey IM, Elisem B. Nutritional Strategies for the Diabetic/Prediabetic Patient. CRC Press 2006: 268.
72. Yasodha KJ, Raghu D, Jayaveera KN, Ravindra K. Anti-diabetic activity of Alcoholic extract of Holostemma ada kodien in rats. The internet journal of Endocrinology. 2009; 5:2.
73. Yasodha KJ, Rasheed AS, Jayaveera KN, Ravindra K, Srikar A. Anti-diabetic activity of Alcoholic extract of Talinum cuneifolium in rats. Pharmacologyonline 2008; 2:63-73
74. Kavishankar GB, Lakshmidevi N, Mahadeva Murthy S, Prakash HS, Niranjana SR. Diabetes and medicinal plants-A review. Int J Pharm Biomed Sci 2011, 2(3), 65-80
75. Alese MO, Adewole OS, Ijomone OM, Ajayi SA, Alese OO. Hypoglycemic and Hypolipidemic Activities of Methanolic Extract of Sphenocentrum jollyanum on Streptozotocin-induced Diabetic Wistar Rats. European Journal of Medicinal Plants 2014, 4(3): 353-364.
76. Alok Kumar Dash, Jhansee Mishra, Deepak Kumar Dash. In vivo Antioxidant Activity and Anti Hyperglycemic Relevant Enzyme Inhibition Properties of Petroleum Ether Extract of Traditionally Processed Ocimum canum Leaves. British Journal of Pharmaceutical Research 2014, 4(9): 1031-1040.
77. Momoh Johnson, Longe Adeteju Olufunmilayo, Campbell Charles Adegboyega, Omotayo Mutiat Adetayo. Evaluation of Antidiabetic and the Effect of Methanolic Leaf Extract of Jatropha curcas on Some Biochemical Parameters in Alloxan-induced Diabetic Male Albino Rats. European Journal of Medicinal Plants 2014, 4(12): 1501-1512.
78. Oyesola O. Ojewunmi, Tope Oshodi, Omobolanle I Ogundele, Chijioke Micah, Sunday Adenekan. In vitro Antioxidant, Antihyperglycaemic and Antihyperlipidaemic Activities of Ethanol Extract of Lawsonia inermis Leaves. British Journal of Pharmaceutical Research 2014, 4(3): 301-314.
79. Ravichandra VD, Padmaa M. Paarakh. Evaluation of Anti-diabetic Potentials of Methanol Extract of Ficus hispida Linn Leaves Against Alloxan Induced Diabetic Rats. British Journal of Pharmaceutical Research 2014, 4(3): 315-324, 2014
80. Sijuade AO, Omotayo OO, Oseni OA. Hypoglycemic Effect of Methanolic Extract of Pergularia daemia in Alloxan-Induced Diabetic Mice. British Journal of Pharmaceutical Research 2014, 4(22): 2614-2621.
81. Carlos Roberto Porto Dechandt, Damiana Luiza Pereira De Souza, et al. Changes in the Oxidative Stress Biomarkers in Liver of Streptozotocin-diabetic Rats Treated with Combretum lanceolatum Flowers Extract. British Journal of Pharmaceutical Research 2014, 4(20): 2340-2356.
82. Chandraprakash Dwivedi, Swarnali Dasgupta. Antidiabetic Herbal Drugs and Polyherbal Formulation Used For Diabetes A Review. The Journal of Phytopharmacology 2013; 2 (3): 44-51
83. Weicheng Hu, Jin-Hee Yeo, Yunyao Jiang, Seong-Il Heo, Myeong Hyeon Wang. The antidiabetic effects of an herbal formula composed of Alnus hirsuta, Rosa davurica, Acanthopanax senticosus and Panax schinseng in the streptozotocin-induced diabetic rats. Nutrition Research and Practice 2013;7(2):103-108.

84. 1Anbu N, Musthafa M D,Velpandian V. Anti-Diabetic Activity of Polyherbal Formulation Aavaraiyathi churnam in Alloxan Induced Diabetic Rats. International Journal of Toxicological and Pharmacological Research 2012-13; 4(4): 77-80
85. Kavishankar GB, Lakshmidhevi N, Mahadeva Murthy S, Prakash HS, Niranjana SR. Diabetes and medicinal plants-A review. Int J Pharm Biomed Sci 2011, 2(3), 65-80
86. Kadarian C, Broussalis AM, Miño J, Lopez P, Gorzalczany S, Ferraro G, Acevedo C..Hepatoprotective activity of *Achyrocline satureioides* (Lam) D. C. Pharmacol Res 2002, 45, 57-61.
87. Gray AM, Flatt PR.Actions of the traditional anti-diabetic plant, *Agrimony eupatoria* (agrimony): effects on hyperglycaemia, cellular glucose metabolism and insulin secretion. Br J Nutr 1998, 80, 109-114.
88. Dandu AM, Inamdar NM. Evaluation of beneficial effects of antioxidant properties of aqueous leaf extract of *Andrographis paniculata* in STZinduced diabetes. Pak J Pharm Sci 2009, 22, 49-52.
89. Panda S, Kar A. Antidiabetic and antioxidative effects of *Annona squamosa* leaves are possibly mediated throughquercetin-3-O-glucoside. Biofactors 2007, 31, 201-210.
90. Khazraji SM, Shamaony LA, Twaij HA.Hypoglycaemic effect of *Artemisia herba alba*. Effect of different parts andinfluence of the solvent on hypoglycemic activity. J Ethnopharmacol 1993, 40, 163-166.
91. Ribnický DM, Poulev A, Watford M, Cefalu WT, Raskin I. Poulev A, Watford M, Cefalu WT, Raskin I. Antihyperglycemic activity of Tarralin,an ethanolic extract of *Artemisia dracuncululus* L. Phytomedicine 2006, 13, 550-557.
92. Fuentes O, Arancibia-Avila P, Alarcón J.Hypoglycemic activity of *Bauhinia candicans* in diabetic induced rabbits. Fitoterapia 2004, 75, 527-532.
93. Chang SL, Chang CL, Chiang YM, Hsieh RH, Tzeng CR, Wu TK, Sytwu HK, Shyur LF, Yang WC. Polyacetylenic compounds and butanol fraction from *Bidens pilosa* can modulate thedifferentiation of helper T cells and prevent autoimmune diabetes in non-obese diabetic mice. Planta Med 2004, 70, 1045-1051.
94. Puri D. The insulinotropic activity of a Nepaiese medicinal plant *Biophytum sensitivum*: preliminary experimentalstudy. J Ethnopharmacol 2001, 78, 89-93.
95. Russell KR, Omoruyi FO, Pascoe KO, Morrison EY. Hypoglycaemic activity of *Bixa orellana* extract in the dog. Methods Find Exp Clin Pharmacol 2008, 30, 301-305.
96. Anand P, Murali KY, Tandon V, Chandra R, Murthy PS.Preliminary studies on antihyperglycemic effect of aqueous extract of *Brassica nigra* (L.) Koch in streptozotocin induced diabetic rats. Indian J Exp Bioi 2007, 45, 696-701.
97. Karageuzyan KG, Vartanyan GS, Agadjanov MI, Panossian AG, Hoult JR.Restoration of the disordered glucose-fatty acid cycle in alloxandiabetic rats bytrihydroxyoctadecadienoic acids from *Bryonia alba*,a native Armenian medicinal plant. Planta Med 1998, 64, 417-422.
98. Naik SR, Barbosa Filho JM, Dhuley JN, Deshmukh V. Probable mechanism of hypoglycemic activity of bassic acid, a natural product isolated from *Bumelia sartorum*. J Ethnopharmacol 1991, 33, 37-44.
99. Amalraj T, Ignacimuthu S. Hypoglycemic activity of *Cajanus cajan* (seeds) in mice. Indian J Exp Biol 1998, 36, 1032-1033.
100. Prakasam A, Sethupathy S, Pugalendi KV. Antihyperglycaemic effect of *Casearia esculenta* root extracts in streptozotocin-induceddiabetic rats. Pharmazie 2002, 57, 758-760.
101. Gupta S, Sharma SB, Singh UR, Bansal SK, Prabhu KM. Elucidation of mechanism of action of *Cassia auriculata* leaf extract for its antidiabeticactivity in strepto zotocin-induced diabetic rats. J Med Food 2010, 13, 528-534
102. Nammi S, Boini MK, Lodagala SD, Behara RB. The juice of fresh leaves of *Catharanthus roseus* Linn, reduces blood glucose in normal and alloxan diabetic rabbits. BMC Complement Altern Med 2003, 2, 3-4.
103. Eddouks M, Lemhadri A, Zeggwagh NA, Michel JB. Potent hypoglycaemic activity of the aqueous extract of *Chamaemelum nobite* in normaland streptozotocin-induced diabetic rats. Diabetes Res Clin Pract 2005, 67, 189-195.
104. Pushparaj PN, Low HK, Manikandan J, Tan BK, Tan CH. Anti-diabetic effects of *Cichorium intybus* in streptozotocin-induced diabetic rats. J Elhnopharmacol 2007,111, 430-434.
105. Ojewole JA. Hypoglycaemic effect of *Clausena anisata* (Willd) Hook methanolic root extract in rats. J Ethnopharmacol 2002, 81, 231-237.

106. Eidi M, Eidi A, Saeidi A, Molanaei S, Sadeghipour A, Bahar M, Bahar K. Effect of coriander seed (*Coriandrum sativum* L.) ethanol extract on insulin release from pancreatic beta cells in streptozotocin-induced diabetic rats. *Phytother Res* 2009, 23, 404-406.
107. Jagtap AG, Patil PB. Antihyperglycemic activity and inhibition of advanced glycation end product formation by *Cuminum cyminum* in streptozotocin induced diabetic rats. *Food Chem Toxicol* 2010, 48, 2030-2036.
108. Ahmad M, Akhtar MS, Malik T, Gilani AH. Hypoglycaemic action of the flavonoid fraction of *Cuminum nigrum* seeds. *Phytother Res* 2000, 14, 103-106.
109. Mukhtar HM, Ansari SH, Ali M, Bhat ZA, Naved T. Effect of aqueous extract of *Cyamopsis tetragonoloba* Linn, beans on blood glucose level in normal and alloxan-induced diabetic rats. *Indian J Exp Biol* 2004, 42, 1212-1215.
110. Iwu MM, Okunji CO, Ohiaeri GO, Akah P, Corley D, Tempesta MS. Hypoglycaemic activity of dioscoretine from tubers of *Dioscorea dumetorum* in normal and alloxan diabetic rabbits. *Planta Med* 1990, 56, 264-267.
111. Ananthi J, Prakasam A, Pugalendi KV. Antihyperglycemic activity of *Eclipta alba* leaf on alloxan-induced diabetic rats. *Yale J Biol Med* 2003, 76, 97-102.
112. Nampoothiri SV, Prathapan A, Cherian OL, Raghu KG, Venugopalan VV, Sundaresan A. In vitro antioxidant and inhibitory potential of *Tetralinia bellerica* and *Emblica officinalis* fruits against LDL oxidation and key enzymes linked to type 2 diabetes. *Food Chem Toxicol* 2010.
113. Maroo J, Vasu VT, Gupta S. Dose dependent hypoglycemic effect of aqueous extract of *Encostemma littorale* blume in alloxan induced diabetic rats. *Phytomedicine* 2003, 10, 196-199.
114. Kumar RV, Augusti KT. Antidiabetic effect of a leucocyanidin derivative isolated from the bark of *Ficus bengalensis* Linn. *Indian J Biochem Biophys* 1989, 26, 400-404.
115. Adaramoye OA, Adeyemi EO. Hypoglycaemic and hypolipidaemic effects of fractions from kolaviron, abiflavonoid complex from *Garcinia Kola* in streptozotocin-induced diabetes mellitus rats. *J Pharm Pharmacol* 2006, 58, 121-128.
116. Ugochukwu NH, Babady NE. Antioxidant effects of *Gongronema latifolium* in hepatocytes of rat models of non-insulin dependent diabetes mellitus. *Fitoterapia* 2002, 73, 612-618.
117. Suthar M, Rathore GS, Pareek A. Antioxidant and Antidiabetic Activity of *Helicteres isora* (L.) Fruits. *Indian J Pharm Sci* 2009, 71, 695-699.
118. Gupta RN, Pareek A, Suthar M, Rathore GS, Basniwal PK, Jain D. Study of glucose uptake activity of *Helicteres isora* Linn, fruits in L-6 cell lines. *Int J Diabetes Dev Ctries* 2009, 29, 170-173.
119. Mahomed IM, Ojewole JA. Hypoglycemic effect of *Hypoxis hemerocallidea* conn (African potato) aqueous extract in rats. *Methods Find Exp Clin Pharmacol* 2003, 25, 617-623.
120. Judy WV, Hari SP, Stogsdill WW, Judy JS, Naguib YM, Passwater R. Antidiabetic activity of a standardized extract (Glucosol) from *Lagerstemia speciosa* leaves in Type II diabetics. A dose-dependence study. *J Ethnopharmacol* 2003, 87, 115-117.
121. Eddouks M, Maghrani M. Effect of *Lepidium sativum* L. on renal glucose reabsorption and urinary TGF-beta 1 levels in diabetic rats. *Phytother Res* 2008, 22, 1-5.
122. Olajide OA, Awe SO, Makinde JM, Morebise O. Evaluation of the antidiabetic property of *Morinda lucida* leaves in streptozotocin-diabetic rats. *J Pharm Pharmacol* 1999, 51, 1321-1324.
123. Pepato MT, Oliveira JR, Kettelhut IC, Migliorini RH. Assessment of the antidiabetic activity of *Myrcia uniflora* extracts in streptozotocin diabetic rats. *Diabetes Res* 1993, 22, 49-57.
124. Kaleem M, Kirmani D, Asif M, Ahmed Q, Bano B. Biochemical effects of *Nigella saliva* L seeds in diabetic rats. *Indian J Exp Biol* 2006, 44, 745-748.
125. Rai V, Iyer U, Mani UV. Effect of Tulasi (*Ocimum sanctum*) leaf powder supplementation on blood sugar levels, serum lipids and tissue lipid in diabetic rats. *Plant Food for Human Nutrition* 1997, 50, 9-16.
126. Vats V, Yadav SP, Grover JK. Ethanolic extract of *Ocimum sanctum* leaves partially attenuates streptozotocin induced alteration in glycogen content and carbohydrate metabolism in rats. *J Ethnopharmacol* 2004, 90, 155-160.
127. Lemhadri A, Zeggwagh NA, Maghrani M, Jouad H, Eddouks M. Antihyperglycemic activity of the aqueous extract of *Origanum vulgare* growing wild in Tafilalet region. *J Ethnopharmacol* 2004, 92, 251-256.

128. Hsu FL, Lai CW, Cheng JT. Antihyperglycemic effects of paeoniflorin and 8-debenzoylpaeoniflorin, glucosides from the root of *Paeonia lactiflora*. *Planta Med* 1997, 63, 323-325
129. Raphael KR, Sabu MC, Kuttan R. Hypoglycemic effect of methanol extract of *Phyllanthus amarus* Schum & Thonn on alloxan induced diabetes mellitus in rats and its relation with antioxidant potential. *Indian J Exp Biol* 2002, 40, 905-909.
130. Mukhtar HM, Ansari SH, Bhat ZA, Naved T, Singh P. Antidiabetic activity of an ethanol extract obtained from the stem bark of *Psidium guajava* (Myrtaceae). *Pharmazie* 2006, 61, 725-727.
131. Rai PK, Mehta S, Watal G. Hypolipidaemic & hepatoprotective effects of *Psidium guajava* raw fruit peel in experimental diabetes. *Indian J Med Res* 2010, 131, 820-824.
132. Mukhtar HM, Ansari SH, Ali M, Bhat ZA, Naved T. Effect of aqueous extract of *Pterocarpus marsupium* wood on alloxan-induced diabetic rats. *Pharmazie* 2005, 60, 478-479.
133. Jahromi MA, Ray AB. Antihyperlipidemic effect of flavonoids from *Pterocarpus marsupium*. *J Nat Prod* 1993, 56, 989-994
134. Maghrani M, Lemhadri A, Jouad H, Michel JB, Eddouks M. Effect of the desert plant *Retama raetam* on glycaemia in normal and streptozotocin-induced diabetic rats. *J Ethnopharmacol* 2003, 87, 21-25.
135. Maghrani M, Michel JB, Eddouks M. Hypoglycaemic activity of *Retama raetam* in rats. *Phytother Res* 2005, 19, 125-128.
136. Yoshino K, Miyauchi Y, Kanetaka T, Takagi Y, Koga K. Anti-diabetic activity of a leaf extract prepared from *Salacia reticulata* in mice. *Biosci Biotechnol Biochem* 2009, 73, 1096-1104.
137. Eddouks M, Jouad H, Maghrani M, Lemhadri A, Burcelin RL. Inhibition of endogenous glucose production accounts for hypoglycemic effect of *Spergularia purpurea* in streptozotocin mice. *Phytomedicine* 2003, 10, 594-599.
138. Benwahhoud M, Jouad H, Eddouks M, Lyoussi B. Hypoglycemic effect of *Suaeda fruticosa* in streptozotocin-induced diabetic rats. *J Ethnopharmacol* 2001, 76, 35-38.
139. Maiti R, Das UK, Ghosh D. Attenuation of hyperglycemia and hyperlipidemia in streptozotocin-induced diabetic rats by aqueous extract of seed of *Tamarindus indica*. *Biol Pharm Bull* 2005, 28, 1172-1176.