

Case Study**Hepatic Metastasis in a Nigerian with Differentiated****Thyroid Cancer- an Uncommon Presentation: Case Report.****ABSTRACT**

The papillary and follicular histological sub-types of thyroid cancer are referred to as differentiated thyroid cancers. When these tumors metastasize, they rarely do so to the liver. Hepatic metastasis is very uncommon in this group of cancers and has only been reported in a handful of cases. We present one such case in a Nigerian patient and note its unique presentation and the need for careful attention to arrival at definitive diagnosis in cases of suspected primary liver cell cancers which are common in this part of the world.

Keywords: Follicular Thyroid Cancer; Differentiated Thyroid cancer; Hepatic Metastasis.

1. Introduction

Furthermore, thyroid carcinomas constitute less than 1% of all human cancers. The annual incidence world-wide ranges from 0.5 to 10 cases per 100,000 populations [1]. The median age at diagnosis is 45–50 years with two to four times more frequent in women than men [1]. Papillary and follicular thyroid cancers, together, are referred to as differentiated thyroid cancer (DTC). Differentiated thyroid carcinomas are relatively rare despite common incidence of thyroid nodules [2]. Liver metastasis from DTC is quite unusual, with a reported frequency of 0.5% [3]. Metastatic liver involvement from DTC is nearly always multiple or diffuse and usually found along with other distant metastases including the lungs, bones and the brain [4]. What makes the case presented here an interesting one is the fact that thyroid cancer metastases to liver are rare, and this represents a novel documentation of same from our region of Africa.

2. Case Report.

A 56-year old Nigerian lady presented with complaints of a 6 month history of abdominal swelling and weight loss. She had felt malaise with a significant drop in appetite over the same period. Her past history was significant for a thyroidectomy which was carried out 7 years earlier on account of what was then thought to be a non-malignant thyroid mass.

Examination findings included non-tender hepatomegaly, were no obvious peripheral stigmata of chronic liver disease. She had a well healed scar of a previous thyroidectomy and there were palpable lymph nodes in the deep cervical lymph node group. The nodes were firm, immobile and none-tender. Investigations carried out revealed deranged thyroid function tests (elevated T_3 and Thyroid Stimulating Hormone), markedly elevated alkaline phosphatase and γ glutamyltransferase (γ GT), with other liver function tests being normal. Screens for hepatitis B and C were negative.

A chest X-ray results showed bilateral pulmonary metastatic lesions. An abdominal ultrasound (USS) and computed tomography (CT) scan revealed a huge intrahepatic mass in segments 4a, 4b and 5 with lobulated margins and heterogeneous enhancement post contrast(Figure 1).




Figure 1: Computed tomography image of the abdomen showing a huge intra-hepatic mass in segments 4a, 4b and 5 with lobulated margins and heterogeneous enhancement post contrast.

Normal α fetoprotein level and colonoscopy results were followed by a USS-guided biopsy of the liver mass which histology were reported as sections containing thyroid and NOT hepatic tissue. It further revealed variously sized thyroid follicles lined by cuboidal cells (Figure 2). A whole body scintigraphy revealed normal tracer bio-distribution in the thyroid gland with abnormal uptake in the lungs and liver indicative of multiple functioning metastases in these latter organs.




Figure 2: Photomicrographs of needle biopsy from liver mass showing tumor fragments consisting exclusively of thyroid follicles lined by cuboidal epithelial cells with hyperchromatic nuclei (Left-H&E $\times$ 100, Right $\times$ 400).

A diagnosis of distant (lungs and hepatic) metastasis from a thyroid cancer was made and the patient has since been referred for expert radio-oncologic assessment and management.

3. Discussion

Liver metastasis from DTC is quite rare, with a reported frequency of 0.5% [3]. As of 2012, a review of literature by Hong-Jun Song et al, reported that only ten cases have been documented in the literature [4]. Three were males and seven females with an average age of about 63 years

(range from 32 to 85 years). Histologically, the primary tumor was identified as papillary in four patients, follicular in five, and Hürthle cell thyroid cancer in one patient. Our patient presented herein, fits well the typical epidemiological descriptors as she is 56 years old.

Unfortunately, the histology of the primary tumor of this patient is unknown as pathology was not carried out on the thyroid gland when it was removed 7 years ago. Hence, we cannot comment about the primary histological variant. This is because transformation of histopathologic types may occur in liver metastases from DTC [4]. This phenomenon has been documented in two of the patients in the 2012 review where the primary tumors were follicular and papillary but their liver metastases were a unique entity described as follicular variant of papillary thyroid carcinoma.

Metastatic liver involvement from DTC is nearly always multiple or diffuse and is usually found along with other distant metastases [4] as was in this index patient who had accompanying lung metastasis at presentation.

The patients seldom present with symptoms referable to the *de facto* liver involvement except when the masses were huge [4]. This was also mirrored in our report. The survival rate after liver metastases is poor; however, Song et al opines that this may not be attributable to the liver metastases alone as multiple other vital organs are concurrently involved by this time [4].

4. Conclusion

The case report highlights the possibility of such rare occurrence of hepatic metastasis from DTC. To the best of our knowledge, this represents a novel documentation of this association from our environment. Because of the possible confusion with a primary liver cell carcinoma

which is relatively common in our region of the world, the usefulness of histological diagnosis of doubtful intra-hepatic lesions is also highlighted.

CONSENT

All authors declare that written informed consent was obtained from the patient (or other approved parties) for publication of this case report and accompanying images.

References

1. Schlumberger MJ. Papillary and follicular thyroid carcinoma. *N Engl J Med* 1998;338: 297–306.
2. Keeston Jones M. Management of papillary and follicular thyroid cancer. *J RSoc Med* 2002;95(7):325–326.
3. Salvatori M, Perotti G, Rufini V, et al. Solitary liver metastasis from Hürthle cell thyroid cancer: a case report and review of the literature. *J Endocrinol Invest* 2004;27(1):52–56.
4. Song H-J, Xue YL, Qiu ZL, et al. Uncommon metastases from differentiated thyroid carcinoma. *Hell J Nucl Med* 2012;15(3):233–240.