

Formulation and physical characterization of a novel sustained-release ophthalmic delivery system for sparfloxacin: the effect of the biological environment

Abd EL – Gawad H. Abd EL – Gawad¹, Osama A. Soliman¹, Susan A. Barker², and Germeen N.S. Girgis^{1*}

Department of Pharmaceutics, Faculty of Pharmacy, Mansoura University, 35516, Mansoura, Egypt.

²School of Pharmacy, University of East Anglia, Norwich Research Park, Norwich, Norfolk, NR4 7TJ, UK

ABSTRACT

Aims: The purpose of this study was to prepare and evaluate ocular thermosetting gels containing sparfloxacin-HP- β -CyD. The formulations based mainly on pluronic F127 alone or combined with other polymers.

Study design: Experimental study.

Place and Duration of Study: School of Pharmacy, University of East Anglia, Norwich, Norfolk, UK, between July 2010 , January 2011.

Methodology: Sparfloxacin, the more effective fluoroquinolone antibiotic used in ocular infection was prepared in form of 1:2 HP- β -CyD spray dried complex. The complex was characterized by several techniques like DSC, FTIR and SEM. In this study, the possibility of incorporation of this complex into thermosetting gels was investigated using pluronic F127 alone or combined with other polymers. All placebo and medicated formulations were characterized physically (colour, clarity, pH, and drug contents). Rheological tests, texture analysis were performed in both non physiological conditions, and after dilution with the simulated tear fluid, to estimate biological environment effect on thermosetting gels properties.

Results: Sparfloxacin-HP- β -CyD complex was successfully prepared and characterized by several techniques, all confirmed complete inclusion between sparfloxacin and HP- β -CyD. The prepared complex showed a significant improvement in the aqueous solubility of sparfloxacin compared to untreated drug or the physical mixture. Ocular thermosetting gels containing sparfloxacin-HP- β -CyD complex were prepared simply by cold method using 20% pluronic F127 alone or 15% pluronic F127 combined with other polymers. All preparations were found colourless or yellow, clear, acceptable pH, and drug content. The rheology and texture analysis studies revealed that, formulations exhibited suitable gelling temperatures, gelling times, and appropriate strength to be used in ocular dosage forms. The biological environment highly affected gel properties according to the polymer type.

Conclusions: ophthalmic *in-situ* gelling formulations containing sparfloxacin-HP- β -CyD complex is considered promising delivery system. It showed good rheological and texture properties. The results confirmed the possibility of lowering pluronic F127 concentration by using other polymers in small concentrations.

Keywords:

Ocular, thermosetting gels, sparfloxacin, HP- β -CyD, pluronic F127, sodium alginate, chitosan, carbopol 980.

1. INTRODUCTION

Topical administration of eye drops is the ideal treatment for ocular diseases, especially when the drug must display a localized action (e.g., the cornea and/or anterior chamber). Unfortunately, in several cases, this treatment is not effective due to protective mechanisms of the human eye, short pre-corneal contact time, and corneal impermeability. And as a result, frequent dosing is usually needed leading to many side effects. Various ocular formulations have been developed in an effort to overcome the issues with conventional eye drop formulations.

Among them, the *in-situ* gel-forming formulations, which undergo phase transition from liquid to semisolid gel upon exposure to physiological environments, seem to be a promising tool. These formulations should be a free-flowing liquid at room temperature to allow easily reproducible administration into the eye as a drop, and then on exposure to physiological conditions; it is converted to the gel phase, and thus increasing the precorneal residence time of the delivery system and enhancing ocular bioavailability. According to the mechanism of gelation, *in-situ* forming gels are classified into three different types: the first type is temperature induced *in-situ* gel system (thermosetting gels), in which the gel is formed by increasing the temperature from room 25 °C to eye temp 34 °C polymers such as pluronic acid is working by this mechanism. The second is the pH-induced *in-situ* gel systems, in which the gels are formed increasing pH to 7.4, such as: carbopol 980, and chitosan. And finally, the third type is osmotically induced *in-situ* gel system in which the gels are formed by the presence of mono or divalent cations like sodium and calcium that found in tear fluids polymers working by this mechanism includes sodium alginate and gelrite (Basavaraj *et al.*, 2007).

Nowadays, *in-situ* gel forming systems are of great importance, having the combined advantage being patient convenient with favorable residence time for enhancing ocular bioavailability and for reducing systemic side effects. In addition it could be easily subconjunctivally injected as a method for periocular delivery system to prolong the drug release to the posterior eye segment tissues such as retina, and vitreous in order to treat the ocular diseases of these tissues.

Poloxamer 407 gives a colourless and transparent gel and is widely employed, but they suffer from a major drawback of having weak mechanical strength, which leads to rapid erosion (El Kamel, 2002). As a result, it requires higher concentration of about 25 to 30 % (W/V) to exhibit sol-gel phase transition at 37 °C when used alone. Gelation temperature can be adjusted within the required temperature range of 33-36 °C by modifying cross-linking agents, by mixing the different series of poloxamer, and by mixing with other polymers. An interesting approach focuses on using mixture of poloxamers with other polymers like carbopol, alginate, and chitosan to prepare *in-situ* gels of suitable characters. Carbopol, a mucoadhesive polymer, increases the formulation's mechanical strength, and also increases surface interaction with the ocular tissue and, consequently, contact time. Chitosan is a biodegradable polymer that has showed excellent ocular compatibility. It presents positively charged amine groups in its chemical structure that could interact with the negatively charged mucous layer, conferring a mucoadhesive characteristic.

Therefore, a combination of these polymers with poloxamer would be very promising for ocular administration, as the *in-situ* mechanical strength of the formulation would be higher than that of both polymers alone.

Sparfloxacin (Figure 1) is a new generation fluoroquinolone antibiotic, which effectively has been used to treat eye infections, it provides improved efficacy against important ocular pathogens like *Staphylococcus aureus*, *Streptococcus pneumoniae*, and *Staphylococcus epidermidis*. It is reported to be more active *in vitro* than ciprofloxacin against mycobacteria and gram-positive bacteria (Sweetman, 2009). However, its main drawbacks as a therapeutic molecule are its very low water solubility and its photosensitivity. In this study, we have investigated the potential of a sparfloxacin-HP- β -CyD inclusion complex to overcome these issues.

73




Figure (1): Chemical structure of sparfloxacin

Cyclodextrins are a family of cyclic oligosaccharides with a hydrophilic outer surface and a lipophilic central cavity, as indicated in Figure 2. They have potential as carriers in drug delivery systems, for example improving the apparent solubility of poorly water-soluble drugs and reducing stability issues associated with photo-sensitive compounds (Chao *et al.*, 2003).




Figure (2): The structure of HP-β-CyD

Hydrophilic cyclodextrins, especially HP-β-CyD, have been shown to be non-toxic and well tolerated in the eye (Loftsson, and Jarvis, 1999).

In this study, we have incorporated the prepared sparfloxacin-HP-β-CD complex into a range of thermo-setting gels and assessed their physical performance, both in the "as prepared" state and in a simulated ocular bio-environment.

The aims of this work are to prepare and characterize an inclusion complex of sparfloxacin and hydroxypropyl-β-cyclodextrin (HP-β-CyD), to develop ophthalmic *in-situ* gelling formulations containing the sparfloxacin-(HP-β-CyD) complex by using poloxamer either alone or combined with other polymers. And to study the effect of the biological environment on the physical properties of the different prepared gel formulations.

2. METHODOLOGY

2.1 Materials

Sparfloxacin of purity 98 % for HPLC, 2-Hydroxypropyl-β-cyclodextrin, poloxamer (pluronic F127), chitosan medium molecular weight, and alginate sodium salt from brown algae (All

for analytical grades reagents) were purchased from Sigma Aldrich, UK. Sodium chloride, calcium chloride dihydrates and sodium bicarbonate were obtained from Fisher Scientific Ltd., UK. Carbomer 980 (Noveon) was obtained from Surfachem, UK. Deionised water was produced from an Elga still, Elga, UK. Syringe filters of 0.45 µm pore size were obtained from Sartorius stedim biotech GmbH, Germany.

2.2 Experimental

2.2.1 Sparfloxacin and HP-β-CyD inclusion complex

2.2.1.1 Preparation sparfloxacin-HP-β-CyD inclusion complex

The sparfloxacin-HP-β-CyD inclusion complex was prepared by spray-drying from a hydro-alcoholic solution containing both sparfloxacin and HP-β-CyD (1:2 molar ratio). Spray-drying was performed using a Mini Spray Dryer Büchi 290 (Büchi, Switzerland) with flow rate 5 ml/minute, inlet temperature 85°C, atomized air pressure 3 kg/cm²; process yield being circa 50 %.

2.2.1.2 Preparation of sparfloxacin-HP-β-CyD physical mixture (PM)

The physical mixture was prepared by the same method described by Williams *et al.*, 1998. This is simply by mixing both the drug and HP-β-CyD, using mortar and pestle in the same molar ratio (1:2) that has been used to prepare the spray dried complex.

2.2.1.3 Characterization of sparfloxacin, HP-β-CyD, the physical mixture and the inclusion complex

The drug alone, HP-β-CyD, drug-HP-β-CyD complex and physical mixture were characterized and evaluated by using the following methods:

Differential scanning calorimetry (DSC)

For the thermal analysis of sparfloxacin, HP-β-CyD, physical mixture and the prepared inclusion complex, a Q1000 MTDSC system (TA Instruments, Newcastle, DE) was used. Samples of circa 5 mg in aluminium pans were heated from 25°C to 300°C at 10°C/minute (Chao *et al.*, 2003).

Fourier transform infrared spectroscopy (FT-IR)

FT-IR studies were carried on a Bruker Optics IFS66/S spectrometer (Coventry, UK) that is equipped with a heated Golden Gate MkII ATR accessory (Specac Limited, Orpington, UK). Spectra were acquired over the range 4000 to 500 cm⁻¹.

Scanning electron microscopy (SEM)

Scanning electron microscopy was performed using a JEOL JSM 5900 LV (JEOL, Japan), mounted with a tungsten filament with an acceleration voltage of 5–20kV. The samples were mounted onto stubs using double-sided tape and were gold coated to a thickness of

147 circa 15 nm by a Polaron SC7640 sputter gold coater (Quorum Technologies). The imaging
148 process was performed in a high vacuum environment.

149 **2.2.1.4 Dissolution rate study of inclusion complex compared to untreated drug and** 150 **physical mixture**

151 Dissolution studies were carried out for sparfloxacin, physical mixture, and inclusion
152 complexes by the same method described by **Aleem et al., (2008)** with some modifications.
153 The experiment has been performed by using USP paddle type dissolution apparatus
154 adjusted at $37\text{ }^{\circ}\text{C} \pm 1\text{ }^{\circ}\text{C}$ and 100 rpm. The dissolution medium used was 500 ml of freshly
155 prepared simulated tear fluid which was maintained at pH 7.4. Simulated tear fluid (STF)
156 was prepared by NaCl (0.67 g), NaHCO_3 (0.2 g), $\text{Ca Cl}_2 \cdot 2\text{H}_2\text{O}$ (0.008 g), and purified water
157 to 100 g, pH 7.4 was adjusted. The large volume of dissolution medium was used in order to
158 maintain the sink conditions as the methods described by **Shikhar et al., 2011**.
159 10 mg of the drug or its equivalent weights of the physical mixture, or the inclusion complex
160 was added to the dissolution medium, samples of 5 ml were withdrawn at time intervals of 5,
161 10, 20, 30, 40, 50, 60, 90, and 120 min. The volume of dissolution medium was adjusted to
162 500 ml by replacing the sample withdrawn with 5 ml of fresh simulated tear fluid (STF) pH
163 7.4. The solutions were immediately filtered through $0.45\text{ }\mu\text{m}$ membrane filter, suitably
164 diluted and the concentrations of sparfloxacin in samples were determined
165 spectrophotometrically at 290 nm.

166 The results of dissolution were analyzed statistically using one way analysis of variance
167 (ANOVA) followed by Tukey-Kramer multiple comparisons test (**Petrie, 1987; Gad and Weil,**
168 **1989**) at the time required for the release of 50% of drug ($T_{50\%}$). Statistical calculations were
169 carried out using Instate - 2 computer programs (Graphpad software Inc., V2. 04 San Diego,
170 CA, USA).

171 **2.2.2 Preparation of ophthalmic thermosetting gels containing sparfloxacin-HP- β -CyD** 172 **inclusion complex**

173
174 Different concentrations of pluronic F127 with or without other polymers were used for the
175 preparation of four plain gels giving names FP1, FP2, FP3, and FP4. For the medicated
176 gels, the sparfloxacin-HP- β -CyD inclusion complex was added to give a final drug loading
177 equivalent to 0.3 %w/w sparfloxacin. The four medicated gels were coded by FM1, FM2,
178 FM3, and FM4, the composition of all plain and medicated formulations were represented in
179 Table 1

180 181 **2.2.2.1 Preparation of plain in-situ forming gels**

182
183 All the plain in-situ gels were prepared by the cold method with some modification in each
184 formula according to the different polymer used.

185 For FP1, a certain volume of double distilled water was cooled to 4°C using an ice bath, over
186 a magnetic stirrer. Then pluronic F127 was slowly added portion wise with continuous
187 stirring for one hour keeping the temperature at $4\text{ }^{\circ}\text{C}$. Then the volume was completed by
188 water, the preparation was kept in refrigerator for at least 24 hours to ensure the complete
189 dissolution of the polymer and get a clear transparent solution (**Gratieri et al., 2010**).

190
191 For FP2, the carbopol 980 solution was prepared by dispersing the calculated weight in
192 double distilled water with continuous stirring until completely dissolved. Then carbopol
193 pluronic solution was prepared by dispersing the weighed amount of pluronic F127 in the
194 previously prepared carbopol solution after cooling to $4\text{ }^{\circ}\text{C}$, then stirring for 1 hour in an ice

bath. The final preparation was kept in refrigerator for at least 24 hours to ensure the complete dissolution (Lin and Sung, 2000).

For FP3, the chitosan solution was initially prepared by dissolving the required weight in a solution of acetic acid 0.5% v/w with continuous stirring until completely dissolved. Then the chitosan solution was refrigerated and used as a solvent for the poloxamer dispersion for 1 hour at 4 °C. The solution was kept in refrigerator for at least 24 hours to ensure the complete dissolution (Varshosaz et al., 2008).

For FP4, The sodium alginate solution was prepared by dispersing the required weight in double distilled water, with continuous stirring till dissolved, and then was cooled to 4 °C; then pluronic was added with stirring for 1 hour using ice bath, kept the partially dissolved solution in refrigerator for at least 24 h till complete formation of *in-situ* gel (Liu et al., 2006).

2.2.2.2 Preparation of medicated *in-situ* gels

Medicated *in-situ* gels were prepared with the same plain *in-situ* gels compositions in addition to the incorporation of sparfloxacin-HP-β-CyD complex which was previously prepared by the spray drying technique. The four medicated gels are coded by FM1, FM2, FM3, and FM4.

In order to prepare the medicated formulations FM1, FM2, FM3, and FM4, the accurately weighted amount of the sparfloxacin-HP-β-CyD complex which equivalent to 0.3% concentration of sparfloxacin in all formulations, was slowly added to the plain solution with continuous agitation in ice bath at 4 °C till homogenous distribution then stored in screw capped amber glass vials for further studies at 4 °C in the refrigerator. In addition, for comparison, a control solution was prepared by the dissolution of the drug-HP-β-CD spray dried complex in double distilled water in the same concentration that equivalent to 0.3% sparfloxacin which has been used to prepare all other formulations (Varshosaz et al., 2008).

2.2.3 Physical characterization of thermosetting gels

The formulations were evaluated physically for colour, clarity, pH and drug content. The drug content was measured by taking 1 ml of each formulation diluting to 100 ml with distilled water in measuring flask. Then 5 ml was diluted to 25 ml with distilled water. The drug concentration was measured by the spectrophotometer at $\lambda = 293 \text{ nm}$ using standard calibration curve within the range of (1 - 30 µg/ml).

In addition both the rheological property and the texture profile of all formulations were estimated in details as in the following techniques.

2.2.4 Rheological studies

Rheological properties were measured using oscillatory rheometer (TA Instruments, UK). Different four rheological tests were performed in order to evaluate all the prepared *in-situ* gel properties and each test was repeated at least three times.

Data were presented as G' (the solid component) and G'' (the liquid component), with the transition (or gelling) point being the point at which the two components become equal in magnitude. Both the temperature at which the systems gelled ($T_{\text{sol/gel}}$) and the time the systems took to gel at 37°C were measured.

Table (1): Composition of thermosetting gels

FORMULATIONS	INGREDIENTS				
	PLURONIC F127	CARBOPOL 980	CHITOSAN	SODIUM ALGINATE	SPARFLOXACIN-HP- β -CYD COMPLEX EQUIVALENT TO SPARFLOXACIN
FP1	20 %	---	---	---	---
FP2	15 %	0.2 %	---	---	---
FP3	15 %	---	0.3 %	---	---
FP4	15 %	---	---	0.1 %	---
FM1	20 %	---	---	---	0.3 %
FM2	15 %	0.2 %	---	---	0.3 %
FM3	15 %	---	0.3 %	---	0.3 %
FM4	15 %	---	---	0.1%	0.3 %
CONTROL	----	---	---	---	0.3 %

247

248

249 **2.2.4.1 Flow test**

250 It was conducted to determine the type of flow of all samples to know either it is Newtonian
 251 or Non-Newtonian. The sample was equilibrated at 35 °C (temperature of eye surface) for at
 252 least 5 minutes before the continuous flow test was started, the flow test is destructive one,
 253 and so the sample must be discarded after each time.

254

255 **2.2.4.2 Strain sweep (LVR determination test)**

256 It is very important test used to determine LVR (linear viscoelastic region) which is defined
 257 as the region where the stress was directly proportional to strain while the storage modulus
 258 G' remained constant. It is used to know the maximum strain for the gel and to indicate that
 259 any strain above this LVR will cause destruction of three dimensional network of the gel .So
 260 any further tests applied for measurement of elastic properties should be made in this LVR.
 261 This test was made at 35 °C, sample was equilibrated at this temperature for at least 5

minutes before the test began, and a torque sweep over the range of 0.1 and 1000 $\mu\text{N.M}$. at frequency of 1 HZ.

2.2.4.3 Temperature sweep

It is used to determine the temperature of the phase transition for the gel formulations. The phase transition temperature is defined as the temperature at which the gel is converted from liquid state to solid state. The test was performed by first setting the initial temp at 10 $^{\circ}\text{C}$ for 5 min before the test was began, then the temp ramp was performed to be increased from 10 to 45 $^{\circ}\text{C}$ at a rate 1 $^{\circ}\text{C}$ per min with applied frequency of 1 HZ and torque of 10 $\mu\text{N.M}$ for FP1 or 1 $\mu\text{N.M}$ for FP2, FP3, and FP4. Each sample was performed at least three times, the phase transition temp is called $T_{\text{sol/gel}}$ and is determined from the obtained curves by the temperature at which the two moduli G' , G'' (Elastic, Viscous modulus) crossover or intersected each other.

2.2.4.4 Time sweep

It is used to determine the gelation time which is defined as the time after which the gel was formed. The initial temperature was set at 10 $^{\circ}\text{C}$ for 5 minutes before the test began, then the temp was raised to 35 $^{\circ}\text{C}$ and the test was made for 10 min at frequency 1HZ and torque of 10 $\mu\text{N.M}$ for FP1 or 1 $\mu\text{N.M}$ for FP2,FP3, and FP4.

2.2.5 Texture profile

Texture profile analysis of the pre-gelled formulations was conducted using TA.XT2 Texture Analyser (Stable Microsystems, UK) and the strength of the gel was taken to be the hardness in compression. The test was performed by placing 5 gm of each sample in plastic sample tube and holding at 35 $^{\circ}\text{C}$ overnight, then put under probe in a manner that the probe penetrate the gel into its centre, 11.5 cm diameter Perspex cylinder probe was used for the test with pre-test speed 0.5mm/s, test speed 1 mm/s, post test speed 5mm/s, force 0.1g, penetration distance 3mm, trigger force 0.5g, and the data acquisition rate 400 pps.

2.2.6 Effect of simulated *in-vivo* conditions

In order to evaluate the formulation performance after ocular instillation, the gels were mixed with simulated tear fluid at a ratio of 40:7 (Wei *et al.*, 2002). These mixtures were subjected to the rheological studies and the texture analysis. For comparison, samples diluted with purified water in the same ratio were also studied.

2.2.7 *In-vitro* release study using membranless method

For the determination of the release profile of sparfloxacin from the inclusion complex incorporated in different thermosetting gel formulations, a membrane-free experimental set-up was used to study the drug release pattern of gels in the eyes. This method was similar to that described by Ur-Rehman *et al.*, (2010) and Gupta *et al.*, (2010).

The method was carried out with some modification in order to use a dialyzer diffusion cell chamber of volume 1500 μl (Harvard apparatus, USA) fitted within the vessels of a paddle USP II dissolution apparatus (Copley apparatus, UK).

The method was performed by filling 1 ml of the each *in-situ* gel formulation in the dialyzer body with care to avoid air bubbles inside the preparation. Then, it was placed at 35 $^{\circ}\text{C}$ inside an oven for 30 mins to ensure the complete gelation. Then, it was immediately transferred with care to the vessels of the dissolution bath containing 300 ml of simulated tear fluid (STF) as dissolution medium of pH 7.4. The temperature and stirring rate were maintained at 35 $^{\circ}\text{C}$ and 25 rpm, respectively. Aliquots of 5 ml were withdrawn from the release medium at predetermined time intervals and then replaced with fresh STF solution heated to 35 $^{\circ}\text{C}$. The samples were filtered through 0.45 μm syringe filters and subjected to spectrophotometric

analysis to determine the sparfloxacin concentrations at λ_{max} 290 nm. Each experiment was repeated three times and the average was calculated.

3. RESULTS AND DISCUSSION

3.1 Preparation and characterization of sparfloxacin-HP- β -CyD complex

Spray drying method is presently one of the most exciting technologies for the pharmaceutical industry. It is an ideal process for complex formation where the end-product must comply with precise quality standards regarding to the particle size distribution, residual moisture content, bulk density and morphology.

Previously, it was found that with spray drying technique one can co-precipitate the drug with a polymer like cyclodextrins in a stable amorphous solid dispersion.

The prepared sparfloxacin-HP- β -CyD complex was characterized by several techniques as DSC, FTIR, and SEM.

3.1.1 Differential scanning calorimetry (DSC)

The thermal analysis has been reported as an important method to recognize and characterize CyDs complexes. When guest molecules were included inside CyDs cavities or in the crystal lattice, the peak corresponding to their melting point is generally shifted to another temperature. Also the peak intensity may be decreased or disappeared (Liu and Zhu 2006).

The DSC curves of pure components and the prepared complex are illustrated in Figure (3). It was observed that, the DSC curve of HP- β -CyD alone showed a very broad endothermic peak, between 38 °C and 118 °C, which attained a maximum at 86 °C which might be corresponding to dehydration process of cyclodextrin. This finding was in agreement with the results obtained by Wang *et al.*, (2011). The authors concluded that in case of HP- β -CyD owing to its amorphous nature, it showed a broad endothermic peak.

For the sparfloxacin alone, it displayed one sharp endothermic peak at about 262 °C corresponding to the melting point of the crystalline form of the drug followed by decomposition phenomena at slight higher temperature. This is similar to the results obtained by Gupta *et al.*, (2010).

In thermogram of the physical mixture of drug with HP- β -CyD, there is an unchanged broad band due to the HP- β -CyD dehydration, while the sharp endothermic drug peak was decreased in its intensity but did not completely disappeared, this may be due to partial inclusion of drug with cyclodextrin by the physical mixing process. On the other hand, there was no peak corresponding to the sparfloxacin observed in case of the sparfloxacin- HP- β -CyD spray dried complex. This may be explained by the complete inclusion of the drug inside the cyclodextrin cavity by this method.

Thus, these thermal behaviour changes showed an interaction between the drug and HP- β -CyD and indicated the possibility of inclusion complex formation by these techniques. The thermal analysis of spray dried system revealed the disappearance of the endothermic peak of sparfloxacin at 262 °C. While, there was no change in the the endothermic peak at 86 °C, this may be attributed to HP- β -CyD dehydration. This phenomenon may be indicative of complete inclusion complex formed (Jun *et al.*, 2007).

3.1.2 Fourier transform infrared spectroscopy (FT-IR)


The FT-IR spectra of sparfloxacin, HP- β -CyD, sparfloxacin and HP- β -CyD physical mixture, sparfloxacin- HP- β -CyD inclusion complex are illustrated in Figure (4). The characteristic peaks of sparfloxacin in FT-IR spectra showed, that is, C = O stretching around 1715 cm^{-1} , C = C at 1620 cm^{-1} , and -CH (1440 to 1500 cm^{-1}), Figure (4 a), while that of HP- β -CyD Figure (4 b) showed prominent absorption bands at 3418 cm^{-1} for O-H stretching vibrations, 2930 cm^{-1} for C-H stretching vibrations and 1154, 1085 and 1036 cm^{-1} for C-H, C-O stretching vibration (Al Omari *et al.*, 2009).

The FT-IR spectrum of the physical mixture Figure (4 c) did not differ significantly from those of the individual components. However, the FT-IR spectrum of the sparfloxacin- HP- β -CyD inclusion complex shows no peaks similar to the drug alone. This can be probably due to the inclusion complexation of sparfloxacin into the HP- β -CyD cavity.


3.2.3 Scanning electron microscopy (SEM)

The surface morphology of the powders for sparfloxacin, HP- β -CyD, sparfloxacin and HP- β -CyD physical mixture, and sparfloxacin - HP- β -CyD complex was assessed by SEM. As illustrated in Figure (5), sparfloxacin existed in plate-like crystal, whereas HP- β -CyD was observed as amorphous, cylindrical spheres. Regarding sparfloxacin and HP- β -CyD physical mixture, the characteristic sparfloxacin crystals were clearly observed, which were mixed with HP- β -CyD particles or adhered to their surface. In contrast, the sparfloxacin-HP- β -CyD spray dried complex appeared in the form of irregular particles in which the original morphology of both components disappeared and tiny aggregates of amorphous particles of irregular size were present. The comparison of these images revealed that, the inclusion complex was structurally distinct from the isolated pure components and the physical mixture of sparfloxacin and HP- β -CyD. The sizes and shapes of particles of spray dried complex were different from those of the sparfloxacin or HP- β -CyD alone; this confirmed the formation of the inclusion complex of sparfloxacin and HP- β -CD by using the spray dried technique. This was similar to the finding of Wang *et al.*, (2011); they proved that, the inclusion complex of trans-ferulic acid with HP- β -CyD was confirmed by SEM method.


A)



B)



C)



D)




Figure (3): DSC thermograms of A) HP-β-CyD, B) sparfloxacin, C) physical mixture and D) Spray dried complex




Figure (4): FT-IR spectra of a) sparfloracin, b) HP- β -CyD, c) physical mixture and d) spray dried complex




Figure (5): SEM micrographs of a) HP- β -CyD, b) sparfloracin, c) physical mixture and d) spray dried complex

3.3 Dissolution rate study

In-vitro dissolution studies for the untreated sparfloxacin, sparfloxacin - HP- β -CyD physical mixture, and sparfloxacin-HP- β -CyD spray dried complex were illustrated in Figure (6). As a general trend, there was an increase in the sparfloxacin dissolution rate from the HP- β -CyD physical mixture and spray dried complex compared to the untreated drug. This may be due to the increase of the drug solubility by using HP- β -CyD.

The untreated sparfloxacin showed a low water dissolution profile due to its low water solubility. From the curve, it was found that, the percent released of sparfloxacin alone after 5 minutes was 34.69%.

Sparfloxacin dissolution rate was improved when it was mixed physically with HP- β -CyD. It showed an increase in the percent released after 5 minutes to be 73%. So, there was more than double increase in the percent by mixing the drug with HP- β -CyD physically. This may be explained on the bases of the partial inclusion of sparfloxacin in the cavity of HP- β -CyD by physical mixing. This was confirmed by DSC results. These results were in agreement with **Badr-Eldin et al, (2008)**. They proved that, the dissolution rate of tadalafil (which is a poor water soluble and effective phosphodiesterase-5 inhibitor) was enhanced when mixed with CyDs due to the *in-situ* inclusion process between the drug and CyD.

In case of sparfloxacin-HP- β -CyD spray dried complex, it was observed that, there was a highly significant increase in the drug dissolution rate compared to the drug alone. It showed a marked increase of the dissolved percentages during the initial intervals of the dissolution process, to be 93 % after the first 5 min., 98% after 10 min. Finally it was reached 99.8 % after 60 min. This may be attributed mainly to complete inclusion of the sparfloxacin in HP- β -CyD cavity by the spray drying technique. The complete inclusion was confirmed before by DSC, FTIR results. This was in agreement with **Betageri and Makarla, (1995)**, who found that there was a marked increase in the dissolution rate of glyburide (a poorly water soluble oral hypoglycemic agent) which might be due to the formation of solid complex as a result of the complete inclusion. Also, they mentioned that, particle size was reduced to the molecular size when the carrier brought the drug into the dissolution medium, leading to fast dissolution.

Riekes et al., 2010 proved that, there was an improvement of amiodarone dissolution rate (water insoluble anti arrhythmic agent) by complexation with β -CyD using freeze drying and spray drying techniques.

The high dissolution rate of the formed drug-complex was explained by changing the crystalline form of the drug into the amorphous form as indicated by SEM results. This finding was also proved by **Dollo et al., (1999)**, and **Aleem et al., (2008)**.

Also, **Badr-Eldin et al, (2008)**, concluded that, there was a marked enhancement of tadalafil dissolution after complexation with CyDs, which was more than 75 % after the first 5 min.

The results of two-way ANOVA statistical analysis revealed that, there was extremely significant difference between the tested prepared physical mixture and HP- β -CyD complex compared to the untreated sparfloxacin at $P < 0.001$.

Therefore, sparfloxacin-HP- β -CyD spray dried complex prepared in a molar ratio 1: 2 could be chosen for further ocular formulations of the sparfloxacin. This is owing to an improved water solubility of the formed complex in comparison to the untreated drug.

3.4 Physical evaluation of the *in-situ* gels

The results are shown in Table (2). All indicated that, the formulations were colourless or yellow in colour and the clarity was satisfactory. In addition, the pH values of all formulations were within the acceptable range (4-9) which the eye can tolerate, without any irritation (**Gupta et al., 1986**). The ophthalmic products outside this range may cause irritation to the

eye. And finally the drug contents of all formulations were found to be in the acceptable range.




Figure (6): *In-vitro* dissolution study of sparflaxacin in simulated tear fluid $\pm$ S.D.

Table (2): Physical characters of in-situ gels

Formulation	Colour	Clarity	pH	Drug content
FP1	colourless	clear	7.06	--
FP2	colourless	translucent	4.2	--
FP3	colourless	clear	4.05	--
FP4	colourless	clear	6.85	--
FM1	pale yellow	translucent	7.01	98 %
FM2	yellow	translucent	5.6	97.23 %
FM3	yellow	clear	4.38	103.8 %
FM4	pale yellow	translucent	6.85	95.09 %

3.5 Rheological study in non-physiological conditions

All the results of four rheological tests for plain and medicated formulations are represented in Table (3).

3.5.1 Flow test

All the four placebo formulations exhibited non Newtonian behaviour with plastic flow for FP1, and pseudo plastic flow (shear thinning) for the other three *in-situ* gels, FP2, FP3, and FP4. According to **Gratieri et al., (2010)**, the administration of the ophthalmic preparation should have as little as possible of the pseudo plastic character of the pre-corneal formulae. The incorporation of the drug complex has limited effect on the type of the flow, so FM1, FM2, FM3, and FM4 have the same flow types as plain ones.

3.5.2 Linear viscoelastic region (LVR).

It was found that FP1 has LVR of 1 to 100 μ .N.m and all other plain preparations have LVR equal 1-10 μ .N.m so, FP1 is the most strong and stable formula this is due to the higher concentration of pluronic F127 (20 %) compared to others which contains only (15%). For FM1, FM3, FM4 the LVR was the same as FP1, FP3, FP4 but FM2, it was found to be 1 to 100 μ .N.m.

This may be explained by the presence of interaction between carbopol 980 and cyclodextrins group especially HP- β CyD leading to decomplexation of the free sparfloxacin from the cyclodextrin cavity. This was previously proved by Rodríguez-Tenreiro *et al.* (2004) who studied the interactions between carbopol and β -cyclodextrin or hydroxypropyl- β -cyclodextrin by differential scanning calorimetry (DSC) and FT-IR spectroscopy.

As a result of the separation of sparfloxacin from the complex which is less water soluble than the complex, leaving free H-bonding which make crosslinking between two polymers, three dimensional network was formed giving more stronger gel with higher LVR.

3.5.3 Temperature test

It is important to know that the ideal phase transition temperature ($T_{sol/gel}$) for the in-situ gel should be located between the average room temperature and 35 $^{\circ}$ C which is the eye temperature (**Gratieri et al., 2010**). If it is higher than this temperature the gel will not be formed in the eye and it will be drained like a liquid.

It was found that the $T_{sol/gel}$ for FP1 was 21.3 $^{\circ}$ C, and for the other three plain gels (FP2, FP3, and FP4) was around 26 $^{\circ}$ C. This indicates that in non physiological conditions $T_{sol/gel}$ is dependent mainly on the pluronic F127 concentration. The presence of other polymers like carbopol 980, chitosan or sodium alginate has no effect on $T_{sol/gel}$. This is in agreement with **Gratieri et al., (2010)**, they proved that, the presence of chitosan in different concentrations did not significantly interfere with the formulations $T_{sol/gel}$. Also this was confirmed by many authors who reported that, $T_{sol/gel}$ is dependent on the poloxamer concentrations; there is an inverse relationship between the $T_{sol/gel}$ and the poloxamer concentration (**Talasaz et al., 2008**).

For the medicated preparations, it was found that, the incorporation of sparfloxacin-HP- β -CyD complex in all medicated formulations causes an increase in $T_{sol/gel}$ either a slight increase for FM1 and FM2 or a large one in case of FM3 and FM4. This is in accordance with the results obtained by Kim *et al.*, (2002), they found that the addition of human epithelial growth factor-CyD complexes (rhEGF-HP- β -CyD) increased the gelation temperature of poloxamer gel used for ophthalmic delivery system. The author explained this result by a possible mechanism, that the binding force (H-bonding) of cross-linked reticular poloxamer

555 gel became weaker by replacing rhEGF/HP- β -CyD complexes in the gel matrix. In addition,
556 it was observed that, the effect of incorporation of sparfloxacin-HP- β -CyD complex into the
557 gel on the phase transition temperature was found to be greater in case of gels containing
558 low concentration of pluronic F127 (FM3, FM4) than that containing higher pluronic F127
559 concentration (FM1, FM2).

560

561

3.5.4 Time test

562 It is important to know the gelling time of all the prepared formulations, because if the gel
563 takes a long time to form in the eye, the drug may be drained out of the eye before gel
564 formation occurs, leading to decreased ocular bioavailability.

565 From the obtained rheograms, the gelling time was found to be in the order of $FP2 < FP1 <$
566 $FP4 < FP3$.

567

568 In addition, the presence of carbopol 980 lowers the gelling time to great extent more than
569 sodium alginate, this result was in accordance with that obtained by Yvonne, (2008).

570 On the other hand, the presence of chitosan increases the gelling time of the preparation.

571

572 In general, it was found that, all the prepared formulations either plain or medicated will form
573 in-situ gels within the first three minutes of their instillation.

574

575

3.6 Rheological studies in physiological condition.

576

577

578

3.6.1 Linear viscoelastic region (LVR)

579 From the obtained rheograms after the dilution, it was found that FP1 and FM1 showed the
580 same strength and the same LVR range after dilution either by water or STF, indicating their
581 stability. This may be due to higher concentration of pluronic F127 compared to all other
582 preparations.

583

584

3.6.2 Temperature test

585 From the results shown in Table (3), it was observed that, after dilution with the distilled
586 water all the eight formulations showed an increase in $T_{sol/gel}$. This may be explained by
587 dilution effect which resulted in reduction of polymer concentration in the formulae leading to
588 an increase in $T_{sol/gel}$.

589

590 Regarding dilution with STF, FM1, FP1, showed the same $T_{sol/gel}$ after dilution either by STF
591 or distilled water (27.5, 24.4). This indicating that, effect is only due to decrease in pluronic
592 F127 concentration by dilution and there was no effect of STF composition on $T_{sol/gel}$.

593

594 On the other hand, all other 6 formulations containing other polymer in combination with
595 pluronic F127, showed decrease in $T_{sol/gel}$ after dilution with STF compared to the dilution
596 with water, in the same used ratio of dilution and the same method. This is an indication of
597 the effective composition of STF in case of these preparations. This may be due to the effect
598 of presence of other polymers presence in combination with pluronic F127 in these
599 preparations. Sodium alginate forms a stable hydrogel in presence of divalent cations of STF
600 (Ca^{+2}) by the interaction between carboxylic functional group located on polymer chain and
601 cation (Cohen *et al.*, 1997).

602

603

604

605

606

607

608
609
610
611
612

Table (3): Results of rheology tests in both physiological and non-physiological conditions

Formulation	LVR (μ .N.m)	Breakdown stress (μ .N.m)	Phase Transition Temp. (°C)	Phase Transition Time (seconds)	Phase Transition Temp. (° C) after dilution with STF	Phase Transition Temp. (° C) after dilution with water
FP1	1 to 100	630.96	21.3	39	24.4	24.4
FP2	1 to 10	158.49	26.5	26.5	28.6	32.7
FP3	1 to 10	31.62	26.3	177	27.5	29.6
FP4	1 to 10	25.15	25.8	79	29.6	34.7
FM1	1 to 100	630.96	22.25	44	27.5	27.5
FM2	1 to 100	158.49	26.9	56	24.4	30.6
FM3	1 to 10	31.62	29.8	180	24.4	31.6
FM4	1 to 10	25.15	28.3	44	30.7	36.8

613
614
615

3.7 Texture analysis results

617

From the results represented in Table (4), it was found that among the prepared gels FP1, FM1 have the highest texture or hardness compared with the others, this finding was in agreement with the obtained of rheology study as they contain higher pluronic F127 concentration. So the order of hardness for plain gels was FP1 > FP2 > FP4 > FP3. Incorporation of the sparfloxacin-complex in all formulations causes an increase in their hardness except FM3 which became weaker after the incorporation of drug complex within the clear gel matrix (decrease from 2.9 to 1.2). This may be explained by that, FM3 is a transparent preparation due to the complete dissolution of sparfloxacin in acetic acid. Also, this could be explained by the drug trapping between the gel network structure leading to loss of the texture of the medicated formulation in case of chitosan-pluronic F127 in-situ forming gel (Varshosaz *et al.*, 2008).

629

The gel textures were decreased after dilution either by STF or water, while FP1 and FM1 were observed to be the highest in resistance to dilution effect compared with others, for example FP1 decreased in texture from 33.9 g to 20.5 g when mixed with STF and to 21.7 g when mixed with water. Also, FM1 decreased from 36.2 g to 16.5 g when diluted with STF and to 17.2 g when diluted with water. On the other hand FP2 texture was extensively decreased from 32.4 g to 1.1 g (STF) and to 1.2 g (water). This may be due to the higher concentration of the pluronic acid (20%) in both FP1, FM1 preparations.

637

This may be attributed to the fact that poloxamer is non-ionic polyoxyethylene–polyoxypropylene–polyoxyethylene triblock copolymer molecules that aggregate into micelles at 34 °C due to the dehydration of the polymer blocks with temperature. The gel formation is a result of micellar enlargements, and the gel is more rigid at higher poloxamer concentrations (Mansour *et al.*, 2008).

643

644

645

646
647
648

Table (4): Results of texture analysis test represented in g.

Formulation	Hardness	Hardness	Hardness
	(Without dilution)	(Diluted by STF)	(Diluted by water)
FP1	33.9±1.1	20.58±1.2	21.7±1.1
FP2	32.4±2.06	1.1±0.1	1.2±0.1
FP3	2.9±0.1	0.3±0.05	0.4±0.05
FP4	26.36±1.06	0.9±0.06	0.9±0.09
FM1	36.2±1.8	16.58±1.4	17.2±1.5
FM2	35.03±0.5	0.4±0.09	0.3±0.05
FM3	1.2±0.1	1.1±0.1	0.9±0.08
FM4	29.5±2.5	1.1±0.09	1±0.12

649
650

3.8 *In- vitro* drug release study with membranless method

651

The cumulative percentage of sparfloxacin released from all medicated *in-situ* gels and the control were illustrated in Figure. (7). All preparations were containing sparfloxacin-HP-β-CyD complex equivalent to 0.3 % sparfloxacin. In case of the control, it was found that the whole sparfloxacin was released in the dissolution medium almost instantaneously after the start of release experiment (it released over 95 % after only 15 minutes). However, the incorporation of the sparfloxacin-HP-β-CyD complex in the pluronic F 127 based thermosetting gels caused slowing of the drug release compared to the control at each time interval. This finding was in agreement with that obtained by **Nie *et al.*, (2011)**.

660

The results showed that, there were significant differences in release rate between the four *in-situ* gels containing sparfloxacin-HP-β-CyD complex. It indicated that, FM2 which contains carbopol 980- pluronic F127 *in-situ* gel showed the fastest drug release profile. The sparfloxacin released was about 80 % after only 1 hr, and reached 96 % after 6 hrs. While FM1 which containing pluronic F127 alone released sparfloxacin in slower rate with 48 % after 1 hr, and 84 % after 6 hrs.

667

The obtained results may be explained by the presence of HP-β CyD in the formulation. There may be an interaction between carbopol 980 and HP-β-CyD leading to depletion of carbopol 980 from the preparation, and replacement of the sparfloxacin from the cyclodextrin cavity. These results were in agreement with **Rodríguez-Tenreiro *et al.*, (2004)**, who confirmed the interactions between carbopol and β-CyD or HP-β-CyD by differential scanning calorimetry (DSC) and FTIR spectroscopy.

674


In case of FM3, which contain chitosan-pluronic F127 and sparfloxacin-HP-β-CyD complex, it showed a different profile, drug release curve follow biphasic pattern compared to FM1. With a higher rate during first hour, then nearly the same as FM1 which was about 48 % after 1 hr., and then followed by slower rate. This may be related to the drug trapped between the tortuous ways of the gels which takes longer time with a slower slop to be released (**Varshosaz *et al.*, 2008**).

681

For FM4 (sodium alginate-pluronic F127 *in-situ* gel containing sparfloxacin-HP-β-CyD complex), it showed a decrease in the drug release rate. There was only 30 % drug released after 1 hr. and approximately 85 % after 6 hrs. This may be due to presence of sodium

685

686 alginate polymer along with pluronic F127, it instantaneously formed gels upon its addition to
 687 simulated lachrymal fluid, leading to slow drug release.
 688 This was in agreement with Cohen *et al.*, 1997, who concluded that *in-situ* gelling alginate
 689 system is a promising drug carrier for prolonged ophthalmic delivery of pilocarpine.
 690
 691



692
 693
 694 **Figure (7): *In-vitro* release of sparfloxacin from in-situ gels containing sparfloxacin-**
 695 **HP-β-CyD complex by membranless method.**
 696
 697

700 4. CONCLUSION

702 The sparfloxacin-HP-β-CyD inclusion complex was successively prepared by spray-drying
 703 method and its characterization was investigated by different analytical techniques. The
 704 utilized techniques confirmed the complete formation of the complex in case of spray drying
 705 technique, while the partial or incomplete inclusion with the physical mixing method.
 706

707 It was proved from the *in-vitro* dissolution study of that there was a marked enhancement in
 708 the dissolution rate of the sparfloxacin in case of the spray dried complex. Also, there was an
 709 improvement in drug released from the physical mixture compared to the untreated

sparfloxacin. As a result, the more water soluble sparfloxacin-HP- β -CyD complex can be incorporated into ophthalmic thermosetting gels based mainly on pluronic F127 alone or combined with other polymers.

The rheological and textural studies showed that, for placebo preparations, a solution to gel phase transition temperature ($T_{sol/gel}$) is $26\text{ }^{\circ}\text{C} \pm 0.5\text{ }^{\circ}\text{C}$, but varying gelling times (from 27 to 177 sec.) and gel strengths (2.9 to 33.9 gm).

Incorporation of the drug complex increased $T_{sol/gel}$, but had variable effects on the gelling temperature and gel strength.

Dilution of placebo with water resulted in a significant increase in $T_{sol/gel}$, but dilution with STF showed only an intermediate effect. Gel strengths decreased dramatically when diluted, irrespective to the dilution medium. While for medicated gels, after dilution with water, showed the same results as for placebo gels, but dilution with STF showed striking differences.

It could be concluded that the physical evaluation of the *in-situ* gels indicates that, the biological environment has a great influence on the performance of the formulation. This particular interaction cannot be explained simply by dilution, but shows a complex relationship between the gel composition, the sparfloxacin-HP- β -CyD complex, and the biological environment.

In-vitro release results showed that, the four medicated formulations succeeded to prolong the drug release from the pluronic F127 based thermosetting gels compared to control. It was found that, FM2 has the fastest release and FM4 the slowest release profile .

The results revealed the possibility to get sustained release profile of sparfloxacin from a novel formulation approach using thermosetting gels containing sparfloxacin-HP- β -CyD inclusion complex. These results indicate that, this formulation approach could be developed and trialed in the treatment of intra-ocular infections with a reduced dosing frequency.

AKNOWLEDGEMENTS

The authors wish to thank the Egyptian Education Mission Sector for funding.

I acknowledge some important points made by reviewer related to prove complex formation by preparing the physical mixture after treating each component (Drug, CyD) separately with the same spray-dried procedure, also to run the dissolution study using this physical mixture, and finally the gel formulations in the present work can be prepared by using non-treated components (Drug and CyD) and compared by the obtained results of gel containing the spray dried complex.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Al Omari MM., Daraghmeih NH, El-Barghouthi MI, Zughul MB, Chowdhry BZ, Leharne SA, Badwana AA. Novel inclusion complex of ibuprofen tromethamine with cyclodextrins: Physico-chemical characterization. *J. Pharm. Biomed. Anal.* 2009;50:449–58.
- Aleem O, Kuchekar B, Pore Y, Late S. Effect of β -cyclodextrin and hydroxypropyl β -cyclodextrin complexation on physicochemical properties and antimicrobial activity of cefdinir. *J. Pharm. Biomed. Anal.* 2008;47(3):535-40.
- Badr-Eldin SM, Elkheshen SA, Ghorab MM. Inclusion complexes of tadalafil with natural and chemically modified β -cyclodextrins. I: Preparation and *in-vitro* evaluation. *Eur. J. Pharm. Biopharm.* 2008;70:819–27.
- Basavaraj K, Nanjawade FV, Manvi AS, Manjappa. *In situ*-forming hydrogels for sustained ophthalmic drug delivery. *J. Controlled Release.* 2007;122:119–34.
- Betageri GV, Makarla KR. Enhancement of dissolution of glyburide by solid dispersion and lyophilization techniques. *Int. J. Pharm.* 1995;126:155–60.
- Chao J, Li J, Meng D, Huang S. Preparation and study on the solid inclusion complex of sparfloxacin with HP-[β]-cyclodextrin. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy.* 2003;59(4):705-11.
- Cohen S, Lobe E, Trevigoda A, Peled Y. A novel in-situ forming ophthalmic drug delivery system from alginates undergoing gelation in the eye. *J. Controlled Release.* 1997; 44(2-3):201-8.
- Dollo G, Corre P, Chollet M, Chevanne F, Bertault M, Burgot J, Verge R. Improvement in solubility and dissolution rate of 1,2-dithiole-3-thiones upon complexation with β -cyclodextrin and its hydroxypropyl and sulfobutyl ether- 7 derivatives. *J. Pharm. Sci.* 1999; 88: 889–95.
- El-Kamel AH. In vitro and *in vivo* evaluation of Pluronic F127-based ocular delivery system for timolol maleate. *Int. J. Pharm.* 2002;241(1):47-55.
- Esclusa-Diaz M T, Guimaraens-Méndez M, Pérez-Marcos MB, Vila-Jato JL, Torres-Labandeira JJ. Characterization and *in-vitro* dissolution behaviour of ketoconazole/ β - and 2-hydroxypropyl- β -cyclodextrin inclusion compounds. *Int. J. Pharm.* 1996; 143(2):203-10.
- Gad SC, Weil CS. "Statistics for Toxicologists", in: "Principles and Methods of Toxicology", 2nd Ed., Hayes, A. W., ed., Raven press Ltd., New York; 1998.
- Gonnissen Y, Verhoeven E, Peeters E, Remon JP, Vervaet C. Coprocessing via spray drying as a formulation platform to improve the compactability of various drugs. *Eur. J. Pharm. Biopharm.* 2008;69(1):320-34.

804 Gratieri T, Gelfuso GM, Rocha EM, Sarmento VH, de Freitas O, Lopez RFV. A
805 poloxamer/chitosan in-situ forming gel with prolonged retention time for ocular delivery.
806 *Eur. J. Pharm. Biopharm.* 2010;75(2):186-93
807
808
809 Gupta SK, Uppal RK, Agnihotri S, Jain N. Need for monitoring strict quality control
810 measures in ophthalmic preparations. *East Pharmacist.* 1986;43–8.

811

812 Gupta H, Aqil M, Khar RK, Ali A, Bhatnagar A, Mittal G. Sparfloxacin-loaded PLGA
813 nanoparticles for sustained ocular drug delivery. *Nanomedicine: Nanotechnology,*
814 *Biology, and Medicine.* 2010;6(2):324-33.
815
816 Jun SW, Kim MS, Kim JS, Park HJ, Lee S, Woo JS., Wang H. Preparation and
817 characterization of simvastatin/hydroxypropyl- β -cyclodextrin inclusion complex using
818 supercritical antisolvent (SAS) process. *Eur. J. Pharm. Biopharm.* 2007;66(3):413-21.
819
820 Lin H, Sung KC. Carbopol/pluronic phase change solutions for ophthalmic drug delivery.
821 *J. Control. Release.* 2000;69:379–88.
822
823 Liu L, Zhu S. Preparation and characterization of inclusion complexes of prazosin
824 hydrochloride with β -cyclodextrin and hydroxypropyl- β -cyclodextrin. *J. Pharm. Biomed.*
825 *Anal.* 2006;40:122–27.
826
827 Loftssona T, Jārvinen T. *Adv. Drug. Deliv. Rev.* 1999;36:59–79.

828 Mansour M, Mansour S, Mortada ND, ElHady SSA. Ocular poloxamer-based
829 ciprofloxacin hydrochloride in-situ forming gels. *Drug. Dev. Ind. Pharm.* 2008;34(7):744-
830 52.
831
832 Nie S, Hsiao WW, Pan W, Yang Z. Thermoreversible pluronic® F127-based hydrogel
833 containing liposomes for the controlled delivery of paclitaxel: *In-vitro* drug release, cell
834 cytotoxicity, and uptake studies. *Int. J. Nanomedicine.* 2011;6(1):151-66.

835 Petrie A. Lecture notes on Medical statistics, 2nd Ed., Blackwell scientific publications,
836 Oxford, U.K.1987.

837

838 Riekes MK, Tagliari MP, Granada A, Kuminek G, Silva MAS, Stulzer HK. Enhanced
839 solubility and dissolution rate of amiodarone by complexation with β -cyclodextrin through
840 different methods. *Materials Science and Engineering: C.* 2010;30(7):1008-13.
841
842 Rodríguez-Tenreiro C, Alvarez-Lorenzo C, Concheiro Á, Torres-Labandeira J.
843 Characterization of cyclodextrincarbopol interactions by DSC and FTIR. *Journal of*
844 *Thermal Analysis and Calorimetry.* 2004;77(2):403-11.
845
846 Shikhar A, Bommana MM, Gupta SS, Squillante E. Formulation development of
847 Carbamazepine–Nicotinamide co-crystals complexed with β -cyclodextrin using
848 supercritical fluid process, *J. of Supercritical Fluids.* 2011;55:1070–78.
849
850 Sweetman S. Editor. Martindale: the complete drug reference. London: Pharmaceutical
851 Press.2009.

852 Talasaz AHH, Ghahremankhani AA, Moghadam SH, Malekshahi MR, Atyabi F,
853 Dinarvand R. In situ gel forming systems of poloxamer 407 and hydroxypropyl cellulose
854 or hydroxypropyl methyl cellulose mixtures for controlled delivery of vancomycin, J. Appl.
855 Polym. Sci. 2008;109:2369–74.
856
857 Ur-Rehman T, Tavelin S, Gröbner G. Effect of DMSO on micellization, gelation and drug
858 release profile of Poloxamer 407. Int. J. Pharma. 2010;394(1–2):92-8.

859 Van Ooteghem MM. In: Edman, P. (Ed), Biopharmaceutics of Ocular Drug Delivery,
860 CRC Press, Boca Raton pp. 27–41.1993.

861 Varshosaz J, Tabbakhian M, Salmani Z. Designing of a Thermosensitive
862 Chitosan/Poloxamer in-situ gel for ocular delivery of ciprofloxacin, The Open Drug
863 Delivery Journal. 2008;2:61-70.
864
865 Wang J, Cao Y, Sun B, Wang C. Characterisation of inclusion complex of trans-ferulic
866 acid and hydroxypropyl- β -cyclodextrin. Food Chemistry. 2011;124(3):1069-75.

867 Wei G, Xu H, Ding PT, Li SM, Zheng JM. Thermosetting gels with modulated gelation
868 temperature for ophthalmic use: The rheological and gamma scintigraphic studies. J.
869 Controlled Release. 2002;83(1):65-74.
870
871 Williams RO, Mahaguna V, Sriwongjanya M. characterization of an inclusion complex of
872 cholesterol and hydroxypropyl- β -cyclodextrin. Eur. J. Pharm. Biopharm;1998;46(3):355-
873 60.
874
875
876
877
878
879
880
881
882

883
884
885
886