



SDI FINAL EVALUATION FORM 1.1

PART 1:

Journal Name:	Ophthalmology Research: An International Journal
Manuscript Number:	2015_OR_18295
Title of the Manuscript:	Sudden loss of vision associated with use of systemic non-steroidal anti inflammatory drugs.
Type of Article:	Case Report

PART 2:

FINAL EVALUATOR'S comments on revised paper (if any)	Authors' response to final evaluator's comments
Valsalva retinopathy has also been described following weight lifting, coughing, vomiting, sexual activity, end stage labor, blowing musical instruments, and compressive injurie, etc. Lab studies? Did you do it? ...can be used to rule out predisposing risk factors, including sickle cell disease, anemia, idiopathic thrombocytopenic purpura, and other blood dyscrasias. The dose of NSAID is low and for few days (3), I don't agree with you with this association. You must do an exhaustive workup in this case before confirm this hypothesis.	I am aware that this could have been valsava retinopathy which is a differential diagnosis but this patient is a man who had no history of coughing, sneezing, weight lifting, vomiting and was not involved in any musical activity or injury. Sexual activity however cannot be ruled out. His laboratory tests are as stated in the article. Bleeding peptic ulcers have long been reported after the use of NSAIDS. It may be necessary to watch out for retinal bleeding following use of these drugs in ophthalmology