

Case Study**Conjunctival Histology in Long-Standing Esotropia****Abstract**

Background: Conjunctivae can be restrictive in long-standing strabismus. To date, there are no reports in the literature describing the histology of the conjunctiva in these patients.

Methods: Conjunctivae biopsies over the medial and lateral rectus were taken at the time of strabismus surgery in 3 patients with restrictive large angle long-standing esotropia.

Results: The conjunctiva overlying the medial rectus has a much more condensed, organized lamina propria compared to the conjunctiva of the contralateral lateral rectus. The medial conjunctiva has scattered small-diameter vessels and numerous clusters of plasmatic cells. The lateral conjunctiva has large blood vessels with occasional scattered plasmatic cells and the lamina propria presents a lax structure. Conclusions: These new findings reinforce the clinical understanding that conjunctiva can become more restrictive over the medial rectus in long-standing esotropia.

Keywords

Strabismus; esotropia; conjunctiva; histology

Introduction

It has long been recognized that the conjunctiva can be restrictive in long-standing strabismus [1,2]. Recessing the conjunctiva in such situations increases the amount of prismatic diopters corrected postoperatively[3,4].

The microscopic structure of the restrictive conjunctiva has not been described.

We present the conjunctival histology findings on 3 patients with large angle long-standing esotropia.

Material and Method

In this case series, we look at the histologic aspect of the conjunctiva of 3 patients that have had long-standing horizontal strabismus, specifically, esotropia. For this study, we define large angle long-standing esotropia as esotropia of 50 prismatic diopters (PD) or more and esotropia that has been persistent for more than 20 years. We excluded patients with previous strabismus surgery or conjunctiva pathology. Every patient included in the case series tested positive on forced duction test on attempted abduction. The surgical technique utilized was a conjunctiva limbal approach. Recession of the medial rectus with recession of the conjunctiva allowing a forced duction test on abduction to be negative and resection of the opposite lateral rectus. Biopsies of conjunctiva overlying the medial and lateral rectus muscles from the esotropic eye were taken at the time of surgery.

Results

The changes found on histology were consistent among all patients. On H&E staining, we found that the conjunctiva overlying the medial rectus have a much more condensed organized lamina propria compared to the conjunctiva of the contralateral lateral rectus muscle. The medial conjunctiva has scattered small-diameter vessels and numerous clusters of plasmatic cells (figure1). In contrast, the lateral conjunctiva has large, abundant blood vessels with occasional scattered plasmatic cells. The lamina propria itself is poorly organized in a lax structure (figure 2). The conjunctiva epithelium overlying the lateral rectus muscle has fewer cells in comparison to the normal looking epithelium of the medial conjunctiva.

Discussion

Several studies have addressed the role of recessing the conjunctiva to increase success rate in strabismus surgery[2,3,4]. Although fornix conjunctiva incision is favored by most surgeons for primary horizontal rectus muscle surgery [5], a limbal

approach has been shown to allow recession of restrictive conjunctiva along with medial rectus muscle recession. Studies have shown that recessing conjunctiva along with the medial rectus results in consistently increased correction of diopters postoperatively compared to recession of rectus alone [4]. The conjunctiva over the opposite resected rectus muscle is often loose and sometimes needs to be trimmed in order to have a good cosmetic appearance. The main histologic finding is on the lamina propria. We think that the “increased density” of the lamina propria over the medial rectus could explain the small-diameter vessels as well as the increased number of plasmatic cells. The “tight” lamina propria could make the plasmatic cells to be close to one another, not necessarily increasing the absolute number of cells. We also think that the dense lamina propria could “compress” the vessels making them look smaller in diameter. There have been no studies to date that look at the histologic changes on the overlying conjunctiva in strabismus. Our study is limited by our small cohort, but the histologic findings were consistent among all our patients.

Conclusion

This is the first study to look at the histologic changes in the conjunctiva overlying the rectus muscles of patients with long-standing large angle esotropia. Conjunctiva overlying the medial and lateral rectus muscles in long-standing esotropia undergo histologic changes that may affect the structure of the conjunctiva. These new findings reinforce the clinical understanding that conjunctiva can become more restrictive over the medial rectus in long-standing esotropia. This description warrants further studies regarding the role of the conjunctiva in long-standing strabismus.

All patients have consented to the submission of the case report for submission to the journal.

Informed consent was obtained from all individual participants included in the study.”

“All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.”

References

1. Cole JG, Cole HG. (1962) Recession of the conjunctiva in complicated eye muscle operations
Am J Ophthalmol 52:618-22
2. Mazow M. (1976) Recession of the conjunctiva in esotropia surgery
Int Ophthalmol Clin 16 (3): 91-5
3. Helveston E.M, Ellis F.D, Patterson J.H, Weber J (1978)
Augmented recession of the medial recti
Ophthalmology 85(5):507-11
4. Willshaw H.E, Mashhoudi N, Powell (1986)
Augmented medial rectus recession in the management of esotropia
Br J Ophthalmol 70(11): 840-843
5. Mikhail M, Verran R, Farrokhyar F, Sabri K. (2013)
Choice of conjunctival incisions for horizontal rectus muscle surgery-a survey of American Association for Pediatric Ophthalmology and Strabismus members.
J AAPOS 17:184-187

Fig 1

H&E 40X the conjunctiva over medial rectus in large angle long-standing Esotropia. Condensed organized lamina propria. scattered small-diameter vessels and numerous clusters of plasmatic cells

Fig 2

H&E 40X large blood vessels with occasional scattered plasmatic cells. The lamina propria itself is poorly organized in a lax structure. The conjunctiva epithelium overlying the lateral rectus muscle has fewer cells in comparison to the normal looking epithelium of the medial conjunctiva

Figure 1.





Figure 2



138