

## **Case Study**

### **Is The Diagnosis of Acute Posterior Multifocal Placoid Pigment Epitheliopathy with Fatal Cerebral Stroke Correct? A case report**

**Author details: Omer Takes<sup>1</sup>, Aylin Yaman<sup>2</sup>, F.Tolga Koroğlu<sup>3</sup>, Handan Guleryuz<sup>4</sup>, A.Osman Saatci<sup>2</sup>**

**1.Resident, Department of Ophthalmology,Dokuz Eylul University ,Izmir,Turkey**

**2.Professor, Department of Ophthalmology, Dokuz Eylul University ,Izmir,Turkey**

**3.Professor, Department of Pediatrics, Dokuz Eylul University ,Izmir,Turkey**

**4. Professor, Department of Radiology, Dokuz Eylul University ,Izmir,Turkey**

**[omer\\_takes@hotmail.com](mailto:omer_takes@hotmail.com) , [aylinyaman@gmail.com](mailto:aylinyaman@gmail.com) , [tolga.koroglu@deu.edu.tr](mailto:tolga.koroglu@deu.edu.tr) ,  
[handanguleryuz68@gmail.com](mailto:handanguleryuz68@gmail.com) ,  
[osman.saatci@deu.edu.tr](mailto:osman.saatci@deu.edu.tr)**

### **ABSTRACT**

**Aim:** To reiterate the fatal cerebral stroke in association with acute posterior multifocal placoid pigment epitheliopathy (AMPEE).

**Report of a Case:** A previously healthy 9-year-old girl developed acute visual loss in her left eye and was diagnosed to have unilateral acute posterior multifocal placoid pigment epitheliopathy (AMPEE) elsewhere and received systemic steroid treatment with topical steroid drops. She was seen a month later by us as no visual improvement was noted by the family. The patient was hospitalized for a routine systemic work-up as the diagnosis seemed unconvincing .Three days after the hospitalization, she suffered a cerebral stroke and died in the hospital despite extensive treatment in the intensive care unit.

**Conclusion:** The neurologic changes including cerebral stroke can be seen in cases with AMPEE and high suspicion should be present in clinicians whenever a case with AMPEE diagnosed.


Keywords: Acute multifocal placoid pigment epitheliopathy, choroiditis, neurologic complications, stroke

## 1. Introduction

Acute posterior multifocal placoid pigment epitheliopathy (AMPEE) is in the spectrum of white-dot syndrome complex characterized by multiple circumscribed, flat gray-white lesions at the level of the retinal pigment epithelium with an acute onset of visual deterioration, metamorphopsia, and scotomas [1-3]. During the course, initially observed multiple lesions may end up with various sized chorioretinal scars. Neurologic involvement associated with AMPEE is a rare entity and a variety of clinical presentations such as headaches, strokes, cerebral venous sinus thrombosis, seizures and central nervous system vasculitis can be encountered [4-13]. We hereby report a case with possible unilateral AMPEE having a fatal stroke during the course.

## 2. Case Report

A previously healthy 9-year-old girl experienced sudden visual loss in her left eye and was diagnosed to have unilateral AMPEE at another eye clinic out of town. Oral and topical steroids were commenced without any visual benefit. As no visual improvement was experienced the family demanded a second opinion and she was seen by us a month after the episode. On our examination, visual acuity was 20/20 in OD and light perception in OS. Left afferent pupillary defect was present. There were no cells in the anterior chamber and vitreous in OS while the right anterior segment was normal. The right fundus was normal, and the left optic disc looked pale and posterior pole looked pale with slightly constricted vessels. Her initial fundus pictures were compared to the newly acquired pictures (Figure 1a and 1b)



66

67

68

69 Figure 1a and b : a) Initial color picture showing the pale and slightly swollen optic disc  
 70 with palish looking posterior pole b) color picture taken a month after the episode  
 71 showing whitish optic nerve with slight chorioretinal atrophy at the posterior pole.


72

73

74 Fluorescein angiogram (FA) and optical coherence tomography (OCT) images were also  
 75 taken ( Figure 2a, 2b and 2c) and fluorescein angiographic pictures were also compared.

76

77



78

79

80 Figure 2a, b and c : a) Initial midvenous phase angiogram showing the widespread  
 81 hyperfluorescence at the deeper layers with disc leaking b) Composite midphase

angiogram taken a month after the episode showing the widespread window defects at the posterior pole c) Thinned retina

We felt that both the initial and recent fundus pictures of the left fundus were not compatible with the initial diagnosis of unilateral AMPEE and left ophtalmic artery occlusion might be in the differential diagnosis. Thereby, she was hospitalized for a meticulous systemic work up. During the systemic evaluation the patient suffered a syncope episode and developed status epilepticus. She was intubated and admitted to the intensive care unit. A magnetic resonance imaging study demonstrated frontal and left temporo-parieto-occipital cytotoxic edema consistent with ischemic infarcts (Figure 3a and b).

we deleted the question marks.




Figure 3a and 3 b : Cranial MRI, FLAIR axial image(a) and diffusion weighted axial(b) images showing bilateral cerebral acute infarction.

infarction

She was aggressively treated for brain edema and refractory status epilepticus in the pediatric intensive care unit. Nevertheless, the brain edema progressed to herniation subsequently and the patient died. The family refused an autopsy.

### 3. Discussion

Stroke related to AMPEE is most likely to occur simultaneously or within a few weeks of the ophthalmic disease presentation. Thus far five cases with AMPEE and cerebral vasculitis with eventual stroke related death were reported [4,5,8,10,13]. Wilson et al. [4] reported a 24-year old man with bilateral AMPEE and he was put on 40 mg/day oral prednisone. Almost a month later, he suddenly collapsed and hospitalized while he was on steroid taper. Radionuclide anterior cerebral perfusion study and EEG demonstrated brain death and the patient died. An autopsy was performed. While there was no evidence of widespread vasculitis, vasculitis changes were demonstrated in medium sized arterioles within the leptomeninges and a few smaller intracerebral vessels. Hammer et al. [5] described a 25-year-old woman who was six months postpartum. She received a diagnosis of bilateral AMPEE and was put on 40 mg/day oral prednisone. Three weeks later she developed hemiplegia. A computerized tomographic investigation showed diffuse cerebral infarction and the patient passed away due to cerebral herniation. De Vries et al. [8] reported a 23-year-old white man with bilateral ocular involvement who developed hemiparesis and hyperesthesia three days after the initial eye examination and receiving the diagnosis of AMPEE. MRI study of the brain showed left medial cerebral artery occlusion and narrowing of the right posterior cerebral artery with infarctions in these vascular territories and the patient died due to herniation. After reviewing the findings in the autopsy, the authors concluded that the cause was focal granulomatous vasculitis affecting the large cerebral arteries. El Sanhoury et al. [10] described a 53-year-old woman with an already established Crohn disease who received a diagnosis of bilateral AMPEE and put the patient on systemic steroid. Prednisone was tapered from 80 mg/day to 20 mg/day over five days. On the sixth day, she developed severe headache and unresponsiveness. She was intubated but three days later she died despite the administration of pulse steroid therapy. Neither MRI nor autopsy could be performed. Very recently, Tsang et al. [13] reported a previously healthy 29-year-old man who had received a diagnosis of bilateral AMPEE following a two month long flu-like symptoms and severe headaches. Approximately, three weeks later, he developed bilateral cerebral infarction and died 14 days after despite intravenous methylprednisolone and craniotomy for the management of raised intracranial pressure. At autopsy, multiple infarctions of varying ages within a 10 day period were detected.

So far

cerebral

All above mentioned five patients with AMPEE having a fatal course due to neurological problems had bilateral ocular involvement and their ages varied between 23 and 53 years. Present case was nine years old and had unilateral ocular involvement. Besides, the initial fundus appearance was not compatible with AMPEE according to our clinical judgement. However, as the family refused the autopsy we could not further comment on the diagnosis.

AMPEE should be considered among the causes of stroke and aseptic meningitis in adults. Severe neurological complications are difficult to predict at the time of ocular disease. Neuroimaging or other diagnostic procedures such as lumbar puncture and even cerebral angiography are not necessary for each and every patient with AMPEE unless there is a clue for neurological involvement. It is difficult to predict the potential neurologic complications when the initial diagnosis is established.

#### **4. Conclusion**

The neurologic changes including cerebral stroke can be seen in cases with AMPEE and high suspicion should be present in clinicians whenever a case with AMPEE diagnosed.

#### **References**

1. Quillen DA, Davis JB, Gottlieb JL, Blodi BA, Callanan DG, Chang TS, Equi RA, The white-dot syndromes. Am J Ophthalmol 2004;137:538-550.
2. Nussenblatt RB. White-dot syndromes. In, Uveitis, Fundamentals and clinical practice, Fourth ed. Eds. Nussenblatt RB and Whitcup SM, Mosby Elsevier USA 2010 pp 389-392.
3. Agarwal A. Acute posterior multifocal placoid pigment epitheliopathy. In, Gass Atlas of macular diseases, Fifth edition. ed. Agarwall A. Elsevier Saunders 2012 USA pp 954-959.

4. Wilson CA, Choromokos EA, Sheppard R. Acute posterior multifocal placoid pigment epitheliopathy and cerebral vasculitis. *Arch Ophthalmol* 1988;106:796–800.
5. Hammer ME, Grizzard WS, Travies D. Death associated with acute, multifocal, placoid pigment epitheliopathy. Case report, *Arch Ophthalmol* 1989;107:170–171.
6. Comu S, Verstraeten T, Rinkoff JS, Busis NA. Neurological manifestations of acute posterior multifocal placoid pigment epitheliopathy. *Stroke* 1996;27:996–1001.
7. O'Halloran HS, Berger JR, Lee WB, Robertson DM, Giovannini JA, Krohel GB, Meckler RJ, Selhorst JB, Lee AG, Nicolle DA, O'Day J. Acute multifocal placoid pigment epitheliopathy and central nervous system involvement: nine new cases and a review of the literature. *Ophthalmology* 2001; 108:861–868.
8. de Vries JJ, den Dunnen WF, Timmerman EA, Kruithof IG, De Keyser J. Acute posterior multifocal placoid pigment epitheliopathy with cerebral vasculitis: a multisystem granulomatous disease. *Arch Ophthalmol* 2006; 124:910-913.
9. Luneau K, Newman NJ, Srivastava S, Biousse V. A case of acute posterior multifocal placoid pigment epitheliopathy with recurrent stroke. *J Neuroophthalmol* 2009 ;29:111-118.
10. El Sanhoury A, Sisk RA, Petersen MR. Mortality from cerebral vasculitis associated with rapid steroid taper during treatment of acute posterior multifocal placoid pigment epitheliopathy. *Arch Ophthalmol* 2012 ;130 :935-937.
11. Thomas BC, Jacobi C, Korporal M, Becker MD, Wildemann B, Mackensen F. Ocular outcome and frequency of neurological manifestations in patients with acute posterior multifocal placoid pigment epitheliopathy (APMPPE). *J Ophthalmic Inflamm Infect* 2012;2:125–138.
12. Matamala JM, Fruerhake W, Verdugo R. Delayed recurrent stroke in a young patient with acute posterior multifocal placoid pigment epitheliopathy. *J Stroke Cerebrovasc Dis* 2013 ;22:e630-634.
13. Tsang BK1, Chauhan DS, Haward R, Whiteman I, Frayne J, McLean C. Fatal ischemic stroke complicating acute multifocal placoid pigment epitheliopathy: histopathological findings *J Neuroophthalmol* 2014 ;34(1):10-15.

205

206

207

208

209