

Direct Correlation Function of Hard Spherocylinder Fluids

Tooraj Ghaffary^{1*}, Mohammad Mehdi Gharahbeigi², Fatemeh Farmanesh³, Leila Shahmandi²

¹Department of Physics, Shiraz Branch, Islamic Azad University, Shiraz, Iran.

²Young Researchers club, Shiraz Branch, Islamic Azad University, Shiraz, Iran.

³Department of Mathematics, Sepidan Branch, Islamic Azad University, Sepidan, Iran.

ABSTRACT

We study the direct correlation function (DCF) of a classical fluid of non-spherical molecules. The components of the fluid are hard spherocylinder (SC) molecules. The required homogeneous DCF is obtained by solving Ornstein-Zernike (OZ) integral equation numerically, using the Percus-Yevich (PY) approximation and the procedure proposed by Ram and co-workers. We also obtained the closest approach between two spherocylinders by using two different methods, first extending the algorithm proposed by Vega and Lago by introducing a new geometry and second use the finite element procedure. Results are in agreement in two methods. The calculation is performed for various values of packing fractions of the fluid and for the aspect ratio, $L/D=5.0, 10.0$. The coefficient expansions of DCF are obtained. The results are in agreement with the other recent works.

Keywords: Molecular fluid, direct correlation function, hard spherocylinder.

1. INTRODUCTION

The problem is the prediction of the physical properties of simple and multi-component molecular fluids based on the information about molecular shapes and intermolecular interactions. Considerable progress, however, has been made over the last two decades in understanding the behavior of these fluids at the molecular level by considering simplified models.

For these fluids we can consider models in which molecules are treated as hard particles with a given shape [1]. The main reasons to study systems of hard convex bodies is that they provide us with simple reference systems which are used for the investigation of more realistic systems such as liquid crystals [2].

Among simple geometric forms, the spherocylinder would seem to approximate best the shape of the most nematogenic molecules. For this reason, a fluid of rigid SC is useful model for investigating the fundamental nature of the nematic-isotropic phase transition in liquid crystals.

In the recent years, there have been many investigations of the structural properties of interface and confined molecular fluids using the liquid state such as integral equation [3,4] density functional theory [5-8] and computer simulations [9,10]. Since the integral equation theory has been found to be quite successful in describing molecular fluids, this theory is used in order to study the fluid.

The direct correlation function (DCF) plays an important role in describing thermodynamical properties and the structure of simple [11-13], molecular [14-16] and multicomponent fluids

* Tel.: +989371322746.

E-mail address: tooraj.gh@gmail.com

[17-20]. The DCF can be used to calculate the equation of state [21], free energy [22], phase transition [23, 24], elastic constants [25-27], etc.

In our work, we consider a system composed of hard SC particles and obtain the DCF of this system. We use the Ornstein-Zernike (OZ) integral equation [28] by the Percus-Yevick (PY) approximation [29] to study this system.

This article organized as follows: In Sec.2, we introduce the SC model and explain how to find the DCF of this model by OZ equation, in Sec.3 the methods of calculating the closest approach between SC molecules are described. In Sec.4, we report and discuss the results and finally in Sec.5 our conclusions are summarized.

2. DCF OF SC

DCF of spherical fluids is only dependent on distance between the centers of the molecules. In non spherical fluids, in addition the distance, the orientation of the molecules is also effective. For solving this problem, expansion of the DCF according to spherical harmonics is suggested [4]. Because the spherical harmonics include polar and azimuthal angles, that are useful tools to describe the orientation axis of each molecule.

In the SC model, system formed by cylinders of length L and diameter D terminating in hemispheres at the two bases. These spherocylinders interact according to the potential:

$$u(\vec{r}, \hat{\Omega}_1, \hat{\Omega}_2) = \begin{cases} \infty & \sigma(\vec{r}, \hat{\Omega}_1, \hat{\Omega}_2) < D \\ 0 & \sigma(\vec{r}, \hat{\Omega}_1, \hat{\Omega}_2) > D \end{cases} \quad (1)$$

where $\hat{\Omega}_i = (\theta_i, \phi_i)$ describes the orientation of particle i and $\vec{r}$ is a vector along the line connecting the centers of the two particles and $\sigma(\vec{r}, \hat{\Omega}_1, \hat{\Omega}_2)$ is the closest approach between the segments constituting the axis of the cylinder.

The DCF of fluid with non-spherical molecules can be defined through the OZ equation [28]

$$h(r_i, \hat{\Omega}_1, r_2, \hat{\Omega}_2) = c(r_i, \hat{\Omega}_1, r_2, \hat{\Omega}_2) + \frac{\rho}{4\pi} \int d\Omega_3 \int dr_3 c(r_i, \hat{\Omega}_1, r_3, \hat{\Omega}_3) h(r_3, \hat{\Omega}_3, r_2, \hat{\Omega}_2) \quad (2)$$

where r_i is the position of the center of mass and ρ shows the number density of the molecules. The symbols h and c are the well-known total and direct correlation functions.

To solve OZ equation, we first introduce the expansion of DCF. There are two common choices for this expansion. In one expansion, the molecular orientations are referred to an intermolecular reference frame in which the polar axis is along the intermolecular vector $\vec{r}_{12}$, body frame (BF). In the other one, the molecular orientations are defined in a space-fixed or laboratory frame (LF) of reference. We expand the angular dependence of the correlation functions, h or c, using orthogonal basis set of harmonics.

Expansions in BF is

$$c(\vec{r}, \hat{\Omega}_1, \hat{\Omega}_2) = \sum_{l_1, l_2, m} c(l_1, l_2, m; r) Y_{l_1}^m(\hat{\Omega}_1) Y_{l_2}^{-m}(\hat{\Omega}_2) \quad (3)$$

and in LF can be defined by

$$c(\vec{r}, \hat{\Omega}_1, \hat{\Omega}_2) = \sum_{l_1, l_2, l} c(l_1, l_2, l; r) \phi_{l_1, l_2, l}(\hat{\Omega}_1, \hat{\Omega}_2, \hat{\Omega}_r). \quad (4)$$

Where $\hat{\Omega}_r$ is the unit vector along $\vec{r}$, $c(l_1, l_2, m; r)$ and $c(l_1, l_2, l; r)$ are the expansion coefficients in body and laboratory frames. $\phi_{l_1, l_2, l}(\hat{\Omega}_1, \hat{\Omega}_2, \hat{\Omega}_r)$ is rotational invariant given by

$$\phi_{l_1, l_2, l}(\hat{\Omega}_1, \hat{\Omega}_2, \hat{\Omega}_r) = \sum_{m_1, m_2, m} C(l_1, l_2, l; m_1, m_2, m) Y_{l_1}^{m_1}(\hat{\Omega}_1) Y_{l_2}^{m_2}(\hat{\Omega}_2) Y_l^{m*}(\hat{\Omega}_r) \quad (5)$$

82

83 Where $C(l_1, l_2, l; m_1, m_2, m)$ is a Clebsch-Gordan coefficient and Y_l^m is the spherical
 84 harmonics, and * indicates the complex conjugate. In BF expansion, the z-axis of the
 85 coordinate system was chosen along the axis connecting the two molecules that are
 86 correlated. Therefore, we only need to deal with one index m. To find the direct correlation
 87 function the following procedure is required:

88 (1) Guess an initial value for $c(l_1, l_2, m; r)$.

89 (2) Calculate the expansion coefficient in LF by using the first step

$$c(l_1, l_2, l; r) = \sum_m \sqrt{\frac{4\pi}{2l+1}} c(l_1, l_2, m; r) C(l_1, l_2, l; m, -m, 0) \quad (6)$$

91 (3) Use Fourier-Hankel transformation to obtain the coefficients in the Fourier space

$$c(l_1, l_2, l; k) = 4\pi(-i)^l \int_0^\infty dr r^2 j_l(kr) c(l_1, l_2, l; r) \quad (7)$$

93 (4) The coefficients $c(l_1, l_2, l; k)$ are used to obtain

$$c(l_1, l_2, m; k) = \sum_l \sqrt{\frac{2l+1}{4\pi}} c(l_1, l_2, l; k) C(l_1, l_2, l; m, -m, 0) \quad (8)$$

95 These three equations transform a two particle correlation function from r frame in real space
 96 into a function in k space; within the complete set of the spherical harmonics we can
 97 transform the OZ equation as

$$h(l_1, l_2, m; k) = c(l_1, l_2, m; k) + \frac{\rho}{4\pi} \sum_l c(l_1, l, m; k) h(l, l_2, m; k) \quad (9)$$

99 This can be written in matrix form for each m and k value

$$\underline{\underline{h}}(m; k) = \underline{\underline{c}}(m; k) + \frac{\rho}{4\pi} \underline{\underline{c}}(m; k) \underline{\underline{h}}(m; k) \quad (10)$$

101 where $\underline{\underline{c}}$ and $\underline{\underline{h}}$ are symmetry matrices with indices l_1 and l_2 .

102 Here we introduce an auxiliary function $\underline{\underline{y}}(m; k)$

$$\underline{\underline{y}}(m; k) = \underline{\underline{h}}(m; k) - \underline{\underline{c}}(m; k) \quad (11)$$

104 (5) The OZ equation is written as

$$\underline{\underline{y}}(m; k) = \frac{\rho}{4\pi} \left[1 - \frac{\rho}{4\pi} \underline{\underline{c}}(m; k) \right]^{-1} [\underline{\underline{c}}(m; k)]^2 \quad (12)$$

106 and it is solved for $y(l_1, l_2, m; k)$.

107 (6) Converting $y(l_1, l_2, m; k)$ to $y(l_1, l_2, l; k)$ yields

$$y(l_1, l_2, l; k) = \sum_m \sqrt{\frac{4\pi}{2l+1}} y(l_1, l_2, m; k) C(l_1, l_2, l; m, -m, 0) \quad (13)$$

109 (7) By using the inverse Fourier-Hankel transform we can get $y(l_1, l_2, l; r)$

$$y(l_1, l_2, l; r) = \frac{i^l}{2\pi^2} \int_0^\infty dk k^2 j_l(kr) y(l_1, l_2, l; k) \quad (14)$$

(8) These expansions are used to obtain $y(l_1, l_2, m; r)$

$$y(l_1, l_2, m; r) = \sum_l \sqrt{\frac{2l+1}{4\pi}} y(l_1, l_2, l; r) C(l_1, l_2, l; m, -m, 0) \quad (15)$$

(9) To solve the OZ equation, we require an appropriate closure relation. Here we apply the PY closure relation [29]

$$c(l, 2) = f(l, 2) [y(l, 2) + l] \quad (16)$$

where $f(l, 2)$ is the Mayer function and is defined by

$$f(l, 2) = \exp(-\beta u(l, 2)) - l \quad (17)$$

and for the pair potential energy of interaction between particles, which defined in Eq.(1) is

$$f(l, 2) = f(\vec{r}, \hat{\Omega}_1, \hat{\Omega}_2) = \begin{cases} -l & \sigma(\vec{r}, \hat{\Omega}_1, \hat{\Omega}_2) < D \\ 0 & \sigma(\vec{r}, \hat{\Omega}_1, \hat{\Omega}_2) > D \end{cases} \quad (18)$$

(10) We can determine $c(l_1, l_2, m; r)$ with the help of the PY equation

$$c(l_1, l_2, m; r) = \frac{1}{4\pi} \sum_{l'_1 l'_2} \sqrt{\frac{(2l'_1+1)(2l'_2+1)(2l''_1+1)(2l''_2+1)}{(2l_1+1)(2l_2+1)}} \quad (19)$$

$$\times C(l'_1, l'_2, l_1; 0, 0, 0) C(l'_2, l'_2, l_2; 0, 0, 0) \sum_{m' m''} C(l'_1, l'_2, l_1; m', m'', m) \\ \times C(l'_2, l'_2, l_2; -m', -m'', -m) f(l'_1, l'_2, m'; r) [y(l''_1, l''_2, m''; r) + 4\pi \delta_{000}^{l'_1 l'_2 m''}]$$

Where $f(l_1, l_2, m; r)$ is the Mayer expansion coefficient and given by

$$f(l_1, l_2, m; r) = \int d\hat{\Omega}_1 \int d\hat{\Omega}_2 f(r, \hat{\Omega}_1, \hat{\Omega}_2) Y_{l_1}^m(\hat{\Omega}_1) Y_{l_2}^{m*}(\hat{\Omega}_2) \quad (20)$$

With these new values of $c(l_1, l_2, m; r)$, we return to step (2) and repeat the next steps again. This iteration is continued until a required convergency is achieved. By using these coefficients, we can obtain the DCF of SC fluid.

3. CLOSEST APPROACH BETWEEN HARD SC MOLECULES

In molecular liquid theory calculating the closest approach between two molecules is very important. In order to calculate the DCF of the fluid it is necessary to calculate the closet approach between the molecules, therefore, we calculate the closest approach of two SCs by using two different methods.

In the first method we use a fast algorithm proposed by Vega and Lago [30]. They actually applied their method for calculating the closest approach between two rods and we extend this to find the closest approach between two SCs. If the shortest distance between two rods obtained and subtract this value from the half diameter of the each SC that value is the closest approach between SCs.

In the second method we use the finite element procedure to find the closest approach. In this method, the main axis of the SCs (cylinders axis) are divided into equal parts.


141 With having direction ($\hat{\Omega}_l$), diameter, length of SC and $\vec{r}_{l2}$ the vector that connects the
 142 centers of SCs, the coordinate of the begin and end of the main axis of the cylinder can be
 143 determined and given by
 144

$$145 \quad \vec{A} = \vec{r}_{l2} + \frac{L_l}{2} \hat{\Omega}_l ; \vec{B} = \vec{r}_{l2} - \frac{L_l}{2} \hat{\Omega}_l \quad (21)$$

146 To divide this segment (axis of cylinder) into n equal parts from the following equation is
 147 used

$$148 \quad F_i = \frac{(i-1)B + (n-i+1)A}{n} \quad (22)$$

149 where F_i is the coordinate of the i^{th} node.
 150



151
 152 **Fig. 1. Coordinates and orientation of two SCs**
 153

154 The main axis of the other cylinder is divided into m equal parts similarly. Now, the distances
 155 between the pairs of points listed from each cylinder which is the combination of 2 from m+n
 156 are calculated. The minimum value from the calculated distances is chosen. Finally, we
 157 subtract this value from the half diameter of the each SC that value is the closest approach
 158 between SCs. These methods are more general and can also be used for the mixtures.
 159

160 3. RESULTS AND DISCUSSION

162 We solve the OZ integral equation numerically to find the DCF of hard SC fluid. For
 163 calculating the DCF of the fluid, it is required to calculate the closet approach between the
 164 molecules, therefore, we calculate the closest approach of two SCs by using two different
 165 methods which described. According to the results shown in the Table 1, for the aspect ratio,
 166 $L/D=5.0$, these methods are in good agreement. Here L and D are the length and diameter
 167 of SC, respectively. In order To make the calculation more accurate, we use the average of
 168 these two methods. All values are randomly selected.
 169
 170
 171
 172
 173
 174
 175
 176


177 **Table 1. Calculated closest approach of SCs with L/D=5.0, the angles are in radian.**
178

r	θ	φ	θ_1	φ_1	θ_2	φ_2	Vega & Lago Method	Finite Element Method
2.8591	1.2419	5.1896	2.7181	2.6025	2.4872	5.1296	0	0
4.2640	1.3555	1.8970	1.3601	1.5154	1.9937	2.3071	0.2196	0.2177
11.7971	2.5181	1.2959	0.6010	4.5526	3.0588	4.4264	6.5927	6.5851
1.4388	2.3084	3.1228	2.7033	1.6170	0.8722	1.8106	0.1033	0.1016
5.5697	3.0763	2.9350	0.6200	3.2049	0.6594	3.6442	2.9350	2.5141
3.7358	1.3938	5.2771	3.0850	2.5906	1.0920	4.1983	2.2485	2.5141
6	0	0	0	0	0	0	0	0
2.8719	1.5483	6.0874	2.5982	0.8067	2.6460	4.9604	1.2461	1.2437
3.7071	0	0	0.785	1.57	0.785	1.57	0	0

179
180 In Table1 r , $\Omega_i = (\theta_i, \varphi_i)$ and $\Omega = (\theta, \varphi)$ are the distance between the centers of
181 SCs, orientation of i^{th} molecule and orientation of the vector which connects the center of
182 SCs respectively.
183 Now, we can calculate the Mayer function with Eq. (18). The packing fraction of SC can be
184 given by

$$185 \quad \eta = \rho \left[\frac{4}{3} \pi \left(\frac{D}{2} \right)^3 \right] + \left[\pi \left(\frac{D}{2} \right)^2 L \right] \quad (23)$$

186
187 In Figure 2, the selected DCF expansion coefficients for packing fraction $\eta = 0.5$ with
188 aspect ratio L/D=5.0 are plotted.
189
190
191
192
193
194
195
196
197
198
199



200

201
202

203

204


205

206

207

Fig. 2. Selected expansion coefficients of the DCF of hard SCs as a function of the reduced distance, $r^* = \frac{r}{D}$, for $\eta = 0.5$.

In Figure 3 the selected DCF expansion coefficients of DCF for packing fraction $\eta = 0.441$ with aspect ratio $L/D=5.0$ are plotted.



208

209
210

211

212

213

214

215

216

Fig. 3. Selected expansion coefficients of the DCF of hard SCs as a function of the reduced distance, $r^* = \frac{r}{D}$, for $\eta = 0.441$.

In the following, the expansion coefficients are compared with computer simulation [10] in Figure 4. These coefficients are calculated in LF for aspect ratio $L/D=10$ and $\eta = 0.1592$. The obtained expansion coefficients of the DCF using the PY integral equation are in good agreement with the coefficients provided by computer simulation.




Fig. 4. Selected expansion coefficients of the DCF as a function of reduced distance $r^* = \frac{r}{D}$, for $\eta = 0.1592$ and $L/D=10.0$ in LF. The solid and dotted curves are due to our results and simulation [10], respectively.

4. CONCLUSION

For studying a system containing spherical and non-spherical members knowing the DCF is required. We consider a system composed of hard SC particles and obtain the DCF of this package. We use the integral equation by the PY closure. First, we should define how to calculate closest approach between two molecules and, then, compare these methods. As shown in Table 1 these methods are in fairly agreement. With calculating closest approach, the expansion coefficients of DCF are obtained. These coefficients are plotted for different

packing fraction and aspect ratio. Finally, we compared the expansion coefficients with computer simulation, which are in good agreement.

REFERENCES

1. Allen MP, Evans GT, Frenkel D, Mulder BM. Hard convex body fluids. *Adv.Chem. Phys.*1993; 86: 1- 166.
2. Colot JL, Wu XG, Xu H,Baus M. Density-functional, Landau, and Onsager theories of the isotropic-nematic transition of hard ellipsoids. *Phy. Rev. A.*1988; 38, 2022- 2036.
3. Ward DA, Lado F. Structure, thermodynamics, and orientational correlations of the nematogenic hard ellipse fluid from the Percus-Yevick equation. *Mol. Phys.*1988; 63, 623-638.
4. Hansen JP, McDonald IR. *Theory of Simple Liquids*: Academic Press, London; 1986.
5. Rickayzen G. Phase transitions in molecular fluids. *Mol. Phys.* 1995; 80: 1093-1103.
6. Calleja M, Rickayzen G. A model for homogeneous and inhomogeneous hard molecular fluids: ellipsoidal fluids. *J. Phys. Condens.Matter.* 1995; 7: 8839-8856.
7. Chrzanowska A, Teixeira PIC,Ehrenttraut H, Cleaver DJ. Ordering of hard particles between hard walls. *J. Phys.Condens.Matter.* 2001; 13: 4715-4726.
8. Velasco E, Mederes L, Sullivan DE. Density-functional study of the nematic-isotropic interface of hard spherocylinders. *Phys. Rev. E.* 2002; 66: 021708-021714.
9. Allen MP, Frenkel D. Observation of dynamical precursors of the isotropic-nematic transition by computer simulation. *Phys. Rev. Lett.* 1987; 58: 1748-1750.
10. Savenko SV, Marjolein Dijkstra. Asymptotic decay of the pair correlation function in molecular fluids: Application to hard rods. *Phys. Rev. E.* 2005; 72: 021202-021212.
11. Henderson D, Chen KY, Degreve L. The direct correlation functions and bridge functions for hard spheres near a large hard sphere. *J. Chem. Phys.* 1994; 101: 6975-6978.
12. Stelzer J, Longa L, Terbin HR. Molecular dynamics simulations of a Gay-Berne nematic liquid crystal – elastic properties from direct correlation functions. *J. Chem. Phys.* 1995; 103: 3098-3117.
13. Moradi M, Shahri H. Equation of state and freezing of gmsa hard spheres. *Int. J. Mod. Phys. B.* 2003; 17: 6057-6065.
14. Ram J, Singh RC, Singh Y. Solution of the Percus-Yevick equation for pair-correlation functions of molecular fluids. *Phys. Rev. E.* 1994; 49: 5117-5126.
15. Letz M, Latz A. Fluids of hard ellipsoids: Phase diagram including a nematic instability from Percus-Yevick theory. *Phys. Rev. E.* 1999; 60: 5865-5871.
16. Singh RC, Ram J, Singh Y. Thermodynamically self-consistent integral-equation theory for pair-correlation functions of a molecular fluid. *Phys. Rev. E.* 1996; 54: 977-980.

- 284 17. Patra CN, Ghosh SK. Structure of nonuniform three-component fluid mixtures: A density-
285 functional approach. *J. Chem. Phys.* 2003; 118: 3668-3677.
286
- 287 18. Moradi M, Khordad R. Direct correlation functions of binary mixtures of hard Gaussian
288 overlap molecules. *J. Chem. Phys.* 2006; 125: 214504-214510.
289
- 290 19. DuBois S, Perera A. Entropy driven demixing in fluids of rigidly ordered particles.
291 *J. Chem. Phys.* 2002; 116: 6354-6367.
292
- 293 20. Biben T, Hansen JP. Phase separation of asymmetric binary hard-sphere fluids. *Phys.*
294 *Rev. Lett.* 1991; 66: 2215-2218.
295
- 296 21. Barrio C, Solana J R. Contact pair correlation functions and equation of state for additive
297 hard disk fluid mixtures. *J. Chem. Phys.* 2001; 115: 7123-7129.
298
- 299 22. Tang Y, Wu J. Modeling inhomogeneous van der Waals fluids using an analytical direct
300 correlation function. *Phys. Rev. E.* 2004; 70: 011201-011208.
301
- 302 23. Marko J F. First-order phase transitions in the hard-ellipsoid fluid from variationally
303 optimized direct pair correlations. *Phys. Rev. A.* 1989; 39: 2050-2062.
304
- 305 24. Singh UP, Singh Y. Molecular theory for freezing of a system of hard ellipsoids:
306 Properties of isotropic-plastic and isotropic-nematic transitions. *Phys. Rev. A.* 1986; 33:
307 2725-2734.
308
- 309 25. Poniewierski A, Stecki J. Statistical theory of elastic constants of nematic liquid crystals.
310 *Mol. Phys.* 1979; 38: 1931-1940.
311
- 312 26. Phuong NH, Germano G, Schmid F. Elastic constants from direct correlation functions in
313 nematic liquid crystals: a computer simulation study. *J. Chem. Phys.* 2001; 115: 7227-7234.
314
- 315 27. Avazpour A, Mahdavi V, Avazpour L. Effect of direct correlation functions on elastic
316 constants of molecular liquids. *Phys. Rev. E.* 2010; 82: 041701-041709.
317
- 318 28. Ornstein LS, Zernike F. Accidental deviations of density and opalescence at the critical
319 point of a single substance. *Proc. Acad.Sci. Amsterdam*, 1914; 17: 793-806.
320
- 321 29. Percus JK, Yevick G. Analysis of Classical Statistical Mechanics by Means of Collective
322 Coordinates. *Phys. Rev.* 1958; 110: 1-13.
323
- 324 30. Vega C, Lago S. A fast algorithm to evaluate the shortest distance between rods.
325 *Comput. Chem.* 1994; 18: 55-59.