

# New $(G'/G)$ -expansion method and its applications to nonlinear PDE

Xu Lanlan <sup>\*1,2</sup> Chen Huaitang <sup>1,2</sup>

<sup>1</sup> School of Sciences, Linyi University, Linyi, Shandong, 276005, China.

Department of Mathematics, Shandong Normal University, Jinan, Shandong, 250014, China.

Received: XX December 20XX

Accepted: XX December 20XX

Online Ready: XX December 20XX

## Abstract

In this paper, the new  $(G'/G)$ -expansion method is proposed for constructing more general exact solutions of nonlinear evolution equation with the aid of symbolic computation. By using this method many new and more general exact solutions have been obtained. To illustrate the novelty and advantage of the proposed method, we solve the Zakharov-Kuznetsov-Benjamin-Bona-Mahony (ZKBBM) equation. Abundant exact travelling wave solutions of these equations are obtained, which include the exponential function solutions, the hyperbolic function solutions and the trigonometric function solutions. Also it is shown that the proposed method is efficient for solving nonlinear evolution equations in mathematical physics and in engineering.

**Keywords:**  $(G'/G)$ -expansion method; ZKBBM equation; exact solutions;

MSC-AMS: 35Q53, 35Q51

## 1 Introduction

In recent years, because of the wide applications of soliton theory in natural science, it is important to seek explicit exact solutions of nonlinear partial differential equations (NLPDEs). Many powerful methods for constructing exact solutions of nonlinear evolution equations have been established and developed, such as the Backlund transform [1], the Hirota's bilinear operators [2], the tanh-coth function expansion [3][4], the Jacobi elliptic function expansion [5], the F-expansion [6], the sub-ODE method [7], the homogeneous balance method [8], the sine-cosine method and so on [9]. But there is no unified method that can be used to deal with all types of nonlinear evolution equations.

Recently, the  $(G'/G)$ -expansion method, firstly introduced by wang et al. [10] has become widely used to search for various exact solutions of NLEEs. The value of the  $(G'/G)$ -expansion method is that one treats nonlinear problems by essentially linear methods. Very lately to enhance the  $(G'/G)$ -expansion method and expand the range of its applicability, further research has been carried out by several authors. Anand Malik improved the method to deal with ten nonlinear equations of physical importance viz [11][12], and Ghodrati Ebadi used the method to carry out the solutions of

\*Corresponding author: E-mail: [xulanlan777@163.com](mailto:xulanlan777@163.com); [chenhuaitang@sina.com](mailto:chenhuaitang@sina.com)

the integro-differential equation[13]. Zhang et al. [14] improved the method to deal with the (2+1)-dimensional Broer-Kaup equation with variable coefficients. Blismail Aslan explored a new application of this method to Toda type lattice differential equations[15]. Junchao Chen used the extended multiple  $(G'/G)$ -expansion method to obtain six nonlinear equations of physical importance viz[16].

The present paper is motivated by the desire to use the improved  $(G'/G)$ -expansion method to construct a series of some types of exact solutions. We will get more interaction solutions of the nonlinear Zakharov-Kuznetsov- Benjamin-Bona-Mahony (ZKBBM) equation, which are very important nonlinear evolution equations in the mathematical physics and have been paid attention by many researchers.

## 2 Summary of the expansion method

The new auxiliary ordinary differential equation is expressed as follows:

$$GG'' = AG^2 + BGG' + C(G')^2 \quad (2.1)$$

where the prime denotes derivative with respect to  $\xi$ .  $A, B, C$  are real parameters.  $F(\xi)$  is

$$F(\xi) = \frac{G'(\xi)}{G(\xi)} \quad (2.2)$$

Using the general solutions of Eq. (2.1), with the help of Maple we have the following four solutions of Eq. (2.2):

(i). when  $B \neq 0$  and  $\Delta_1 = B^2 + 4A - 4AC \geq 0$ , then

$$F(\xi) = \frac{B}{2(1-C)} + \frac{B\sqrt{\Delta_1}}{2(1-C)} \frac{c_1 \exp^{\frac{\sqrt{\Delta_1}}{2}\xi} + c_2 \exp^{-\frac{\sqrt{\Delta_1}}{2}\xi}}{c_1 \exp^{\frac{\sqrt{\Delta_1}}{2}\xi} - c_2 \exp^{-\frac{\sqrt{\Delta_1}}{2}\xi}} \quad (2.3)$$

(ii). when  $B \neq 0$  and  $\Delta_1 = B^2 + 4A - 4AC < 0$ , then

$$F(\xi) = \frac{B}{2(1-C)} + \frac{B\sqrt{-\Delta_1}}{2(1-C)} \frac{ic_1 \cos(\frac{\sqrt{-\Delta_1}}{2}\xi) - c_2 \sin(-\frac{\sqrt{-\Delta_1}}{2}\xi)}{ic_1 \sin(\frac{\sqrt{-\Delta_1}}{2}\xi) + c_2 \cos(-\frac{\sqrt{-\Delta_1}}{2}\xi)} \quad (2.4)$$

(iii). when  $B = 0$  and  $\Delta_2 = A(C-1) \geq 0$ , then

$$F(\xi) = \frac{\sqrt{\Delta_2}}{(1-C)} \frac{c_1 \cos(\sqrt{\Delta_2}\xi) + c_2 \sin(\sqrt{\Delta_2}\xi)}{c_1 \sin(\sqrt{\Delta_2}\xi) - c_2 \cos(\sqrt{\Delta_2}\xi)} \quad (2.5)$$

(iv). when  $B = 0$  and  $\Delta_2 = A(C-1) < 0$ , then

$$F(\xi) = \frac{\sqrt{-\Delta_2}}{(1-C)} \frac{ic_1 \cosh(\sqrt{-\Delta_2}\xi) - c_2 \sinh(\sqrt{-\Delta_2}\xi)}{ic_1 \sinh(\sqrt{-\Delta_2}\xi) - c_2 \cosh(\sqrt{-\Delta_2}\xi)} \quad (2.6)$$

where  $\xi = x - \omega t$ ,  $\omega$  is wave velocity,  $A, B, C$  and  $c_1, c_2$  are real parameters.

Suppose that we have a NLEE for  $u(x, t)$  in the form

$$P(u, u_t, u_x, u_{tt}, u_{xt}, u_{xx}, \dots) = 0 \quad (2.7)$$

where  $P$  is a polynomial in its arguments, which includes nonlinear terms and the highest order derivatives.

Next, the main steps of this method are given as follows:

**Step 1.** The transformation  $u(x, t) = u(\xi)$ ,  $\xi = x - \omega t$  reduces Eq. (2.7) to the ordinary differential equation (ODE)

$$H(u, u_\xi, u_{\xi\xi}, u_{\xi\xi\xi}, \dots) = 0 \quad (2.8)$$

**Step 2.** We assume that the solution of Eq. (2.8) is of the form

$$u(\xi) = \sum_{i=-m}^m a_i (d + F(\xi))^i \quad (2.9)$$

where  $F(\xi)$  satisfy the new auxiliary ordinary differential Eq.(2.1), and  $\omega, d, a_i (i = -m, \dots, m)$  can be determined later. We can determine the positive integer  $m$  by balancing the highest nonlinear terms and the highest partial derivative terms in the given system equations.

**Step 3.** Substituting Eq. (2.9) along with (2.1) and (2.2) into Eq. (2.8) and using Maple yields a system of equations of  $F^i(\xi)$ , setting the coefficients of  $F^i(\xi)$  in the obtained system of equations to zero, we can deduce the following set of algebraic polynomials with the respect unknowns  $\omega, d, a_i (i = -m, \dots, m)$  namely.

**Step 4.** Solving the over-determined system of differential equations by using the symbolic computation as Maple, we obtain expressions for  $\omega, d, a_i (i = -m, \dots, m)$ .

**Step 5.** Since the general solutions of (2.1) have been well known for us, then substituting  $\omega, d, a_i (i = -m, \dots, m)$  and the general solutions (2.3)-(2.6) into (2.2) we have more exact solutions of the non-linear partial differential Eq. (2.7).

### 3 Travelling wave solitons for ZKBBM equation

In this section, we apply this method to construct the exact interaction soliton solutions of the ZKBBM equation

$$u_t + u_x - 2auu_x - bu_{xxx} = 0 \quad (3.1)$$

The transformation  $u(x, t) = u(\xi), \xi = x + Vt$  reduces Eq. (3.1) to the ordinary differential equation (ODE):

$$(1 + V)u' - 2auu' - bVu''' = 0 \quad (3.2)$$

We can determine the positive integer  $n$  by balancing  $uu'$  and  $u'''$  in the given system equations. So we can suppose that Eq. (3.2) has the following ansatz:

$$u(\xi) = \frac{a_{-2}}{(d + F(\xi))^2} + \frac{a_{-1}}{d + F(\xi)} + a_0 + a_1(d + F(\xi)) + a_2(d + F(\xi))^2 \quad (3.3)$$

Substituting Eq. (3.3) along with (2.1) and (2.2) into Eq. (3.2) and using Maple yields a system of equations of  $F^i(\xi)$ , setting the coefficients of  $F^i(\xi) (i = 0, 1, 2, \dots)$  in the obtained system of equations to zero, we can deduce the set of algebraic polynomials with the respect unknowns  $V, d, a_i (i = -m, \dots, m)$  namely. Solving the over-determined system of differential equations by using the symbolic computation as Maple, we obtain expressions for  $V, d, a_i (i = -m, \dots, m)$ .

**Case 1.**

$$\begin{aligned} d &= \frac{1}{2} \frac{B}{C-1}, V = V, a_1 = 0, a_2 = 0 \\ a_{-1} &= 0, a_0 = -\frac{1}{2} \frac{(8bVAC - 8bVA - 2bVB^2 - 1 - V)}{a} \\ a_{-2} &= -\frac{3}{8} \frac{bV(-8AB^2C + 8AB^2 + B^4 + 16A^2C^2 - 32A^2C + 16A^2)}{a(C^2 - 2C + 1)} \end{aligned}$$

where  $A, B, C, V, a, b$  are arbitrary constants, and  $a \neq 0, C \neq 1$ .

**Case 2.**

$$\begin{aligned} d &= \frac{2A}{B}, V = V, a_1 = 0, a_2 = 0 \\ a_{-2} &= -\frac{6bVA^2(-8AB^2C + 8AB^2 + B^4 + 16A^2C^2 - 32A^2C + 16A^2)}{aB^4} \end{aligned}$$

---


$$a_{-1} = \frac{6bVA(-8AB^2C + 8AB^2 + B^4 + 16A^2C^2 - 32A^2C + 16A^2)}{aB^3}$$

$$a_0 = -\frac{1}{2} \frac{48bVA^2C^2 - 16bVACB^2 - 96bVA^2C + 16bVAB^2 - B^2 - VB^2 + 48bVA^2 + bVB^4}{B^2a}$$

where  $A, B, C, V, a, b$  are arbitrary constants, and  $B \neq 0, a \neq 0$ .

**Case 3.**

$$d = d, V = V, a_{-2} = 0, a_{-1} = 0$$

$$a_1 = -\frac{6bV(-2dC^2 + 4Cd + BC - 2d - B)}{a}, a_2 = -\frac{6bV(C^2 - 2C + 1)}{a}$$

$$a_0 = -\frac{1}{2} \frac{12bVC^2d^2 + 8bVAC - 24bVCd^2 - 12bVBCd - 1 + 12bVd^2 + bVB^2 + 12bVBd - V - 8bVA}{a}$$

where  $A, B, C, V, d, a, b$  are arbitrary constants, and  $a \neq 0$ .

**Case 4.**

$$d = \frac{1}{2} \frac{B}{C-1}, V = V, a_{-1} = 0, a_1 = 0$$

$$a_0 = -\frac{1}{2} \frac{8bVAC - 8bVA - 2bVB^2 - 1 - V}{a}, a_2 = -\frac{6bV(C^2 - 2C + 1)}{a}$$

$$a_{-2} = -\frac{3}{8} \frac{bV(-8AB^2C + 8AB^2 + B^4 + 16A^2C^2 - 32A^2C + 16A^2)}{a(C^2 - 2C + 1)}$$

where  $A, B, C, V, a, b$  are arbitrary constants, and  $a \neq 0, C \neq 1$ .

Substituting those cases in (3.3), we obtain the following solutions of Eq. (3.2). These solutions are:

$$u_1(\xi) = -\frac{3}{8} \frac{bV(-8AB^2C + 8AB^2 + B^4 + 16A^2C^2 - 32A^2C + 16A^2)}{a(C^2 - 2C + 1)} \frac{1}{(\frac{1}{2} \frac{B}{C-1} + F(\xi))^2}$$

$$- \frac{1}{2} \frac{(8bVAC - 8bVA - 2bVB^2 - 1 - V)}{a}$$

$$u_2(\xi) = -\frac{6bVA^2(-8AB^2C + 8AB^2 + B^4 + 16A^2C^2 - 32A^2C + 16A^2)}{aB^4} \frac{1}{(\frac{2A}{B} + F(\xi))^2}$$

$$+ \frac{6bVA(-8AB^2C + 8AB^2 + B^4 + 16A^2C^2 - 32A^2C + 16A^2)}{aB^3} \frac{1}{\frac{2A}{B} + F(\xi)} - \frac{1}{2}$$

$$\frac{48bVA^2C^2 - 16bVACB^2 - 96bVA^2C + 16bVAB^2 - B^2 - VB^2 + 48bVA^2 + bVB^4}{B^2a}$$

$$u_3(\xi) = -\frac{1}{2} \frac{12bVC^2d^2 + 8bVAC - 24bVCd^2 - 12bVBCd - 1 + 12bVd^2 + bVB^2 + 12bVBd - V - 8bVA}{a}$$

$$- \frac{6bV(-2dC^2 + 4Cd + BC - 2d - B)}{a} (d + F(\xi)) - \frac{6bV(C^2 - 2C + 1)}{a} (d + F(\xi))^2$$

$$u_4(\xi) = -\frac{3}{8} \frac{bV(-8AB^2C + 8AB^2 + B^4 + 16A^2C^2 - 32A^2C + 16A^2)}{a(C^2 - 2C + 1)} \frac{1}{(\frac{1}{2} \frac{B}{C-1} + F(\xi))^2} - \frac{1}{2}$$

$$\frac{8bVAC - 8bVA - 2bVB^2 - 1 - V}{a} - \frac{6bV(C^2 - 2C + 1)}{a} (\frac{1}{2} \frac{B}{C-1} + F(\xi))^2$$

where  $\xi = x + Vt$ .

According to (2.3)-(2.6), we obtain the following exponential function solutions, hyperbolic function solutions and triangular function solutions of Eq. (3.1). For example

(1). When  $B \neq 0$  and  $\Delta_1 = B^2 + 4A - 4AC \geq 0$ , then

$$u_{41}(\xi) = -\frac{3}{8} \frac{bV(-8AB^2C + 8AB^2 + B^4 + 16A^2C^2 - 32A^2C + 16A^2)}{a(C^2 - 2C + 1)} \\ - \frac{1}{2} \frac{8bVAC - 8bVA - 2bVB^2 - 1 - V}{a} - \frac{6bV(C^2 - 2C + 1)}{a} \\ \frac{1}{\left( \frac{B\sqrt{\Delta_1}}{2(1-C)} \frac{c_1 \exp \frac{\sqrt{\Delta_1}}{2} \xi + c_2 \exp -\frac{\sqrt{\Delta_1}}{2} \xi}{c_1 \exp \frac{\sqrt{\Delta_1}}{2} \xi - c_2 \exp -\frac{\sqrt{\Delta_1}}{2} \xi} \right)^2} \\ \left( \frac{B\sqrt{\Delta_1}}{2(1-C)} \frac{c_1 \exp \frac{\sqrt{\Delta_1}}{2} \xi + c_2 \exp -\frac{\sqrt{\Delta_1}}{2} \xi}{c_1 \exp \frac{\sqrt{\Delta_1}}{2} \xi - c_2 \exp -\frac{\sqrt{\Delta_1}}{2} \xi} \right)^2$$

where  $\xi = x + Vt$ .

(2). When  $B \neq 0$  and  $\Delta_1 = B^2 + 4A - 4AC < 0$ , then

$$u_{42}(\xi) = -\frac{3}{8} \frac{bV(-8AB^2C + 8AB^2 + B^4 + 16A^2C^2 - 32A^2C + 16A^2)}{a(C^2 - 2C + 1)} \\ - \frac{1}{2} \frac{8bVAC - 8bVA - 2bVB^2 - 1 - V}{a} - \frac{6bV(C^2 - 2C + 1)}{a} \\ \frac{1}{\left( \frac{B\sqrt{-\Delta_1}}{2(1-C)} \frac{ic_1 \cos(\frac{\sqrt{\Delta_1}}{2} \xi) - c_2 \sin(-\frac{\sqrt{\Delta_1}}{2} \xi)}{ic_1 \sin(\frac{\sqrt{\Delta_1}}{2} \xi) + c_2 \cos(-\frac{\sqrt{\Delta_1}}{2} \xi)} \right)^2} \\ \left( \frac{B\sqrt{-\Delta_1}}{2(1-C)} \frac{ic_1 \cos(\frac{\sqrt{\Delta_1}}{2} \xi) - c_2 \sin(-\frac{\sqrt{\Delta_1}}{2} \xi)}{ic_1 \sin(\frac{\sqrt{\Delta_1}}{2} \xi) + c_2 \cos(-\frac{\sqrt{\Delta_1}}{2} \xi)} \right)^2$$

where  $\xi = x + Vt$ .

(3). When  $B = 0$  and  $\Delta_2 = A(C - 1) \geq 0$ , then

$$u_{43}(\xi) = -\frac{3}{8} \frac{bV(16A^2C^2 - 32A^2C + 16A^2)}{a(C^2 - 2C + 1)} \frac{1}{\left( \frac{\sqrt{\Delta_2}}{(1-C)} \frac{c_1 \cos(\sqrt{\Delta_2} \xi) + c_2 \sin(\sqrt{\Delta_2} \xi)}{c_1 \sin(\sqrt{\Delta_2} \xi) - c_2 \cos(\sqrt{\Delta_2} \xi)} \right)^2} \\ - \frac{1}{2} \frac{8bVAC - 8bVA - 1 - V}{a} - \frac{6bV(C^2 - 2C + 1)}{a} \left( \frac{\sqrt{\Delta_2}}{(1-C)} \frac{c_1 \cos(\sqrt{\Delta_2} \xi) + c_2 \sin(\sqrt{\Delta_2} \xi)}{c_1 \sin(\sqrt{\Delta_2} \xi) - c_2 \cos(\sqrt{\Delta_2} \xi)} \right)^2$$

where  $\xi = x + Vt$ .

(4). When  $B = 0$  and  $\Delta_2 = A(C - 1) < 0$ , then

$$u_{44}(\xi) = -\frac{3}{8} \frac{bV(16A^2C^2 - 32A^2C + 16A^2)}{a(C^2 - 2C + 1)} \frac{1}{\left( \frac{\sqrt{-\Delta_2}}{(1-C)} \frac{ic_1 \cosh(\sqrt{-\Delta_2} \xi) - c_2 \sinh(\sqrt{-\Delta_2} \xi)}{ic_1 \sinh(\sqrt{-\Delta_2} \xi) - c_2 \cosh(\sqrt{-\Delta_2} \xi)} \right)^2} \\ - \frac{1}{2} \frac{8bVAC - 8bVA - 1 - V}{a} - \frac{6bV(C^2 - 2C + 1)}{a} \left( \frac{\sqrt{-\Delta_2}}{(1-C)} \frac{ic_1 \cosh(\sqrt{-\Delta_2} \xi) - c_2 \sinh(\sqrt{-\Delta_2} \xi)}{ic_1 \sinh(\sqrt{-\Delta_2} \xi) - c_2 \cosh(\sqrt{-\Delta_2} \xi)} \right)^2$$

where  $\xi = x + Vt$

If  $c_1 = -c_2, u_{41}(\xi)$  can be rewritten :

---


$$u'_{41}(\xi) = -\frac{3}{8} \frac{bV(-8AB^2C + 8AB^2 + B^4 + 16A^2C^2 - 32A^2C + 16A^2)}{a(C^2 - 2C + 1)}$$

$$-\frac{1}{\left(\frac{B\sqrt{\Delta_1}}{2(1-C)} \tanh\left(\frac{\sqrt{\Delta_1}}{2}\xi\right)\right)^2}$$

$$-\frac{1}{2} \frac{8bVAC - 8bVA - 2bVB^2 - 1 - V}{a} - \frac{6bV(C^2 - 2C + 1)}{a}$$

$$\left(\frac{B\sqrt{\Delta_1}}{2(1-C)} \tanh\left(\frac{\sqrt{\Delta_1}}{2}\xi\right)\right)^2$$

where  $\xi = x + Vt$ . Because of  $\tanh^2 y = 1 - \text{sech}^2 y$ ,  $u'_{41}(\xi)$  becomes

$$u''_{41}(\xi) = -\frac{3}{8} \frac{bV(-8AB^2C + 8AB^2 + B^4 + 16A^2C^2 - 32A^2C + 16A^2)}{a(C^2 - 2C + 1)}$$

$$-\frac{1}{\frac{B^2\Delta_1}{4(1-C)^2} (1 - \text{sech}^2(\frac{\sqrt{\Delta_1}}{2}\xi))}$$

$$-\frac{1}{2} \frac{8bVAC - 8bVA - 2bVB^2 - 1 - V}{a} - \frac{6bV(C^2 - 2C + 1)}{a}$$

$$\left(\frac{B^2\Delta_1}{4(1-C)^2} (1 - \text{sech}^2(\frac{\sqrt{\Delta_1}}{2}\xi))\right)$$

Again, if  $c_1 = c_2$ ,  $u_{41}(\xi)$  can be rewritten :

$$u'''_{41}(\xi) = -\frac{3}{8} \frac{bV(-8AB^2C + 8AB^2 + B^4 + 16A^2C^2 - 32A^2C + 16A^2)}{a(C^2 - 2C + 1)}$$

$$-\frac{1}{\left(\frac{B\sqrt{\Delta_1}}{2(1-C)} \coth\left(\frac{\sqrt{\Delta_1}}{2}\xi\right)\right)^2}$$

$$-\frac{1}{2} \frac{8bVAC - 8bVA - 2bVB^2 - 1 - V}{a} - \frac{6bV(C^2 - 2C + 1)}{a}$$


$$\left(\frac{B\sqrt{\Delta_1}}{2(1-C)} \coth\left(\frac{\sqrt{\Delta_1}}{2}\xi\right)\right)^2$$

where  $\xi = x + Vt$ .


By the improved  $(G'/G)$ -expansion method, Zhang et al. [18] obtained seven solutions of the ZKBBM equations, but by means of the proposed expansion method we obtained solutions are different to Zhang et al. [18] solutions . Furthermore, we obtain solutions  $u_{41}(\xi)$ ,  $u_{42}(\xi)$ ,  $u_{43}(\xi)$ ,  $u_{44}(\xi)$ . These solutions are new and were not obtained by Zhang et al.[18]. On the other hand, the auxiliary equation used in this paper is different, so the solutions obtained is also different.

## 4 Summary and conclusion

In summary, the improved  $(G'/G)$ -expansion method with symbolic computation is developed to deal with the nonlinear ZKBBM equation. When applying the proposed method to construct the exact interaction soliton solutions of the nonlinear ZKBBM equation, we get a rich variety of exact solutions which include exponential function solutions, hyperbolic function solutions and triangular function



1.jpg



2.jpg

Fig 1:  $A = 0, B = C = 2, a = b = V = 1, c_1 = -c_2$ ;

Fig 2:  $A = 0, B = C = 2, a = b = V = 1, c_1 = c_2$

solutions. We make graphs of obtained solutions, so that they can depict the importance of each obtained solution and physically interpret the importance of parameters too. Further more, our method can obtain more types of travelling solutions mentioned above. We also see that our method is different from the old  $(G'/G)$ -expansion method. We use the new auxiliary ordinary differential equations to construct more types of travelling solutions. Our method is more powerful and much easier to solve nonlinear evolution equations. We believe that this method should play an important role in finding exact solutions of NLPDEs.

Note that the nonlinear evolution equations proposed in the present paper are difficult and more general. Therefore, the solutions of the proposed nonlinear evolution equation in this paper have many potential applications in physics.

## Acknowledgment

This research was supported by the Natural Science Foundation of Shandong Province of China under Grant (ZR2010AM014) and the Scientific Research Fund from Shandong Provincial Education Department (J07WH04).

## References

- [1] Ablowitz, P.A. Clarkson, Solitons, Nonlinear evolution equations and inverse scattering transform. Cambridge Univ. Press, Cambridge. (1991)
- [2] V.A. Matveev, M.A. Salle, Darboux transformation and solitons, [M]. Springer, Berlin. (1991)
- [3] Houria TRIKI, Ahmet YILDIRIM, T. HAYAT, Omar M. ALDOSSAR, Anjan BISWAS, TOPOLOGICAL AND NON-TOPOLOGICAL SOLITON SOLUTIONS OF THE BRETHERTON EQUATION. PROCEEDINGS OF THE ROMANIAN ACADEMY, Series A, Volume 13, Number 2/2012, pp. 103-108
- [4] HOURIA TRIKI, SIHON CRUTCHER, AHMET YILDIRIM, BRIGHT AND DARK SOLITONS OF THE MODIFIED COMPLEX GINZBURG LANDAU EQUATION WITH PARABOLIC AND DUAL-POWER LAW NONLINEARITY, Romanian Reports in Physics, Vol. 64, No. 2, P. 367-380, 2012.

- 
- [5] Huiqun Zhang, Extended Jacobi elliptic function expansion method and its applications, [J]. Communications in Nonlinear Science and Numerical Simulation, Issue 5, (2007) pp. 627-635.
  - [6] M.A. Abdou, The extended F-expansion method and its application for a class of nonlinear evolution equations, [J]. Chaos, Solitons and Fractals, 31, Issue 1, January, (2007) pp. 95-104.
  - [7] Houria Triki, Abdul-Majid Wazwaz, Sub-ODE method and soliton solutions for the variable-coefficient mKdV equation, [J]. Applied Mathematics and Computation, Volume 214, Issue 2, 15 August, (2009) pp. 370-373.
  - [8] S.A. El-Wakil, E.M. Abulwafa, A. Elhanbaly, M.A. Abdou, The extended homogeneous balance method and its applications for a class of nonlinear evolution equations, [J]. Chaos, Solitons and Fractals, Volume 33, Issue 5, pp. 1512-1522, August 2007.
  - [9] E. Yusufolu, A. Bekir, M. Alp, Periodic and solitary wave solutions of Kawahara and modified Kawahara equations by using Sine-Cosine method, [J]. Chaos, Solitons and Fractals, Volume 37, Issue 4, August, (2008) pp. 1193-1197.
  - [10] Mingliang Wang, Xiangzheng Li, Jinliang Zhang, The  $(G'/G)$ -expansion method and travelling wave solutions of nonlinear evolution equations in mathematical physics. Physics Letters A, 372, 2008 pp. 417-423.
  - [11] Anand Malik, Fakir Chand \*, S.C. Mishra, Exact travelling wave solutions of some nonlinear equations by  $(G'/G)$ -expansion method. Applied Mathematics and Computation, 216 (2010) 2596-2612
  - [12] ANAND MALIK, FAKIR CHAND, HITENDER KUMAR and S.C. MISHRA, Exact solutions of some physical models using the  $(G'/G)$ -expansion method [J]. Indian Academy of Sciences, Vol. 78, No. 4, 2012, pp. 513-529.
  - [13] Ghodrat EBADI, A.H. KARA, Marko D. PETKOVIC, Ahmet YILDIRIM, Anjan BISWAS, SOLITONS AND CONSERVED QUANTITIES OF THE ITO EQUATION, [J]. PROCEEDINGS OF THE ROMANIAN ACADEMY, Series A, 13, Number 3, 2012, pp. 215-224.
  - [14] Sheng Zhang \*, Wei Wang, Jing-Lin Tong, A generalized  $(G'/G)$ -expansion method and its application to the  $(2+1)$ -dimensional Broer-Kaup equations Applied Mathematics and Computation, 209 (2009) 399-404.
  - [15] Ismail Aslan, Some exact solutions for Toda type lattice differential equations using the improved  $(G'/G)$ -expansion method, [J]. Mathematical Methods in the Applied Sciences, 2010, pp. 1579.
  - [16] JUNCHAO CHEN. and BIAO LI, Multiple  $(G'/G)$ -expansion method and its applications to nonlinear evolution equations in mathematical physics, [J]. Indian Academy of Sciences, 78, No. 3 2012, pp. 375-388.
  - [17] Xiaohua Liu, Weiguo Zhang and Zhengming Li, Application of improved  $(G'/G)$ -expansion method to traveling wave solutions of two nonlinear evolution equations, [J/OL]. Advances in Applied Mathematics and Mechanics, 4, No. 1, 2012, pp. 122-130.
  - [18] Jiao Zhang, Fengli Jiang, Xiaoying Zhao, An improved  $(G'/G)$ -expansion method for solving nonlinear evolution equations. International Journal of Computer Mathematics, 55, 2010, pp. 1716-1725.