

Electromagnetic fields of self-modes in spherical resonators

ABSTRACT

In this article a physical theory of self-modes of electromagnetic resonators is presented. It is known, that Maxwell equations predict non-physical singular behavior of self-modes in spherical resonators. This shows that Maxwell theory is incomplete. For the improvement of the theory this problem is investigated with the help of Maxwell-Einstein theory. Maxwell-Einstein equations take into account space-time curvature. Regular implementation of this approach permits to avoid the influence of singularity. Another result consists of that modes with large values of orbital angular moment are not observable. An analogy with CMB in the Universe is made.

Keywords: resonator, self-mode, singularity, cosmic microwave background

1. INTRODUCTION

Field in the electromagnetic resonators is usually described using Maxwell's equations, which are superimposed on the solutions appropriate boundary conditions. So solutions are obtained in the form of standing waves corresponding to the eigenmodes of resonators. Upon excitation of the resonator by external sources it creates a field that can be represented as an expansion in eigenmodes which forms a complete orthogonal system. Below we are interested only in eigenmodes. In an empty cavity they are excited by radiation emitted by atoms of the cavity walls. Consider radiation of atoms located on one of the walls of resonator. For a qualitative analysis of the field of the resonator eigenmode we use the Huygens-Fresnel principle [1]. Suppose that each atom emits independently of the other, and thus the total radiation is a combination of waves of incoherent sources. Front of such a wave in no way corresponds to the shape of the cavity walls and the wave reaching the opposite wall, and reflected from it, will come to the original wall with random phase, which does not correspond to phase of the emitted wave. Thus, the reflected wave, having interacted with the original one, destroy it. This will not happen if the atoms radiate in phase. Then, the wave front shape corresponds to the shape of cavity wall, the reflected wave coming from the emitting panel having at each point the same phase shift, and if it is a multiple 2π ¹, the resultant wave doesn't destroy and will comply with eigenmode of the resonator. In general, this situation is typical for the formation of eigenmodes for cavities of any shape. Of course, the condition for the survival of mode can't be considered as a reason

¹ For the eigenmodes with a sufficiently large number (spherical cavity)

for causing the wall atoms radiate coherently. The essential reason may be the synchronization atoms by eigenmode itself.

The above picture is consistent with the definition of the eigenmode field using Maxwell's equations for rectangular resonators, when their solutions have no singularities and have simple interpretation. For resonators of spherical shape solutions of Maxwell equations have a singularity at $r = 0$, what requires assumptions about the nonphysical infinite energy density at the origin, which is located in the center of the cavity. When one tries to give a physically meaningful interpretation of the eigenmodes of spherical resonators this fact must be taken into account and requires going beyond the Maxwell theory. First physically reasonable solution to this problem was proposed in the paper [2], devoted to the definition of the metric of space-time, curves by spherical electromagnetic waves (SEMW). This requires along with Maxwell's equations also use the Einstein equations for the Riemann tensor, which describes the curvature of space-time, which right side contains energy-momentum tensor of SEMW. This is justified, at least by two reasons:

1. The metric tensor of the problem [2, 3] contains a component that is independent of the amplitude of the electric wave and significant at distances of the order of the wavelength.
2. In general solutions of Einstein's equations have singularities, which can prove as an example of specific solutions (Schwarzschild metric), and with the help of the theorems on the global structure of space-time [4].

An attempts was made to interpret the singular solutions using the Maxwell equations alone (or methods of geometrical optics) for the fields in the cavities or open optical systems, focusing the incident field at the point (focus), but did not give conclusive results [5]². These failures can be considered as a third reason justifying the use Maxwell-Einstein equations to solve the problem.

Solutions of the Maxwell equation which was used in [2]³ obey degeneration, connecting with arbitrariness of z – axis' direction of co-ordinate system. If direction of z – axis is fixed the initial spherical symmetry of problem is lowered.

2. ANGLE DISTRIBUTION OF THE SELF-MODES OF SPHERICAL RESONATORS

In quantum mechanics recovery of breaking symmetry is due to so-called zero modes [7]. In our problem all directions of z – axis are equivalent: all solutions corresponding to its different directions are possible and have the same energy. In order to eliminate zero modes, one must explicitly take into account the transitions between degenerate states. For the simplicity one can do this in quantum description. A simplification of problem is connected with fact that angular behavior of photon wave function is just the same as for classical SEMW. Let us calculate the probability of transition from the state with orbital quantum number l , which angular behavior is described by $P_l(\cos(\theta))$ in co-ordinate system with given axis z , to the state with the same quantum number in co-ordinate system with axis z' deviating from z on angle $\Delta\theta$. In this latter co-ordinate system angular behavior of

² In [5] a notion of an effective sources for divergent SEMW so on as sinks for convergent ones is introduced.

³ So as all similar solutions, which can be found in scientific literature (see [6], for example). As a consequence, field distribution in spheroidal electromagnetic resonator has axial symmetry.

78 wave function describes as $P_l(\cos(\theta+\Delta\theta))^4$. Let us expand $P_l(\cos(\theta+\Delta\theta))$ in series on
 79 associated Legendre polynomials $P_l^k(\cos(\theta))$:

$$P_l(\cos(\theta + \Delta\theta)) = \sum_{k=-l}^l I_k(\Delta\theta) P_l^k(\cos(\theta))$$

(1)

82 The amplitude of transition probability $I_k(\Delta\theta)$ one can find with the help of adding theorem for
 83 spherical functions [5]:

$$P_l(\cos(\theta + \Delta\theta)) = P_l(\cos(\theta))P_l(\cos(\Delta\theta)) + 2 \sum_{k=1}^l \frac{(l-k)!}{(l+k)!} P_l^k(\cos(\theta))P_l^k(\cos(\Delta\theta))$$

(2)

86 Due to orthogonality of associated Legendre polynomials [8], we receive:

$$I_k(\Delta\theta) = P_k(\cos(\Delta\theta)) \cdot \delta_{k0} + 2 \frac{(l-k)!}{(l+k)!} P_l^k(\cos(\Delta\theta))$$

(3)


89 where δ_{kl} – Kroneker's symbol. Interested probabilities look as follows:

$$w_l = \frac{[P_l(\cos(\Delta\theta))]^2}{\sum_{k=0}^l [I_k(\cos(\Delta\theta))]^2}$$


(4)

92 Denominator in (4) arises because associated Legendre polynomial are orthogonal, but not-
 93 orthonormal set of functions. Quantity w_l is a "fraction" of function $P_l(\cos(\theta+\Delta\theta))$ in initial
 94 function $P_l(\cos(\theta))$. Fig.1 represents results of calculation $w_l(\Delta\theta)$ for different l :


⁴ P_l are Legendre polynomials, P_l^k - are associated Legendre polynomials.




a)



b)



c)



d)

95
96
97

98
99

100 **Fig. 1.** Plot of $w_l(\Delta\theta)$
101 Ordinata: points – $(P_l(\cos(\theta)))^2$, solid curve – $w_l(\Delta\theta)$; Abscissa: angles θ and $\Delta\theta$ from 0 to π ;
102 a) $l = 1$ (curves coincide), b) $l = 2$, c) $l = 3$, d) $l = 4$.

103

104 These results show that angle region $\Delta\theta_c$, where fraction of “shifted” harmonic $P_l(\cos(\theta+\Delta\theta))$
105 in the “basic” one $P_l(\cos(\theta))$ is significant, is comparable with scale θ_c of angular dependence
106 of $P_l(\cos(\theta))$, which has order of value $1/l$. This can be assigned to effect of zero modes,
107 because both abovementioned harmonics have the same energy. Of course, this effect

108 vanishes when direction of z axis is fixed physically, for instance, with the help of external
109 field.

110 Recall that the field of electrical oscillations in a spherical cavity is defined by the function U ,
111 which has the form [6]

$$112 \quad U = A\Psi_l(kr)P_l^m(\cos\theta)\begin{cases} \cos m\varphi \\ \sin m\varphi \end{cases} \quad (5)$$

113
114 A – is a constant, P_l^m – associated Legendre polynomial, r, θ, φ – spherical coordinates,
115 $\Psi_l(kr)$ – radial part of the field, k – wavenumber. Equations for U are given in [6]. We have
116 already mentioned that the expression (5) is valid only for the modes excited by an external
117 source (antenna), which specifies the direction of the OZ axis coordinate system. Question
118 about the arbitrariness of the choice of direction of the axis OZ is also addressed in [6], but
119 the answer seems unconvincing.

120 In the spirit of this approach the correct expression for the eigenmode field U of a spherical
121 cavity must take into account the degeneracy of the directions the axis OZ . Simplify the
122 problem by putting $m = 0$. This means that we fix a plane in which lies the axis OZ . As
123 mentioned above, all directions $\theta_{0n} = \pi n/l, 0 \leq n \leq l-1$ in this plane, measured from some
124 arbitrary reference direction $\theta_{00} = 0$, may be taken from the same ground as the orientation
125 axis OZ .

126 Desired expression for U must be of the form (in the general case $m \neq 0$)

$$127 \quad U' = B\Psi_l(kr)\sum_{n=0}^{l-1} P_l(\cos(\theta - \theta_{0n}))\begin{cases} \cos m\varphi \\ \sin m\varphi \end{cases} \quad (6)$$

128
129 B – is a constant defined as A in (5) by the normalization condition. Due to linearity of
130 Maxwell's equations (6) is a mode of a spherical cavity, but in contrast to (5), it allows for
131 the maximum degree of symmetry of the problem.

132 Expression (6) was used in [3] when recording energy-momentum tensor of SEMW before
133 averaging it over the angle θ . Such a procedure is always applied when considering the free-
134 oriented systems [9].

135 3. ELECTROMAGNETIC FIELDS OF THE SELF-MODES OF 136 SPHERICAL RESONATORS

137 In the spherical cavity eigenmodes also are excited by atoms of wall which are synchronized
138 by radiated wave. When one traditionally considers the spherical cavity eigenmodes within
139 Maxwell's theory he will receive for the radial parts of the complex field amplitudes well-
140 known expressions of the form of standing waves $\sim J_{n+1/2}(kr)/(kr)^{3/2}e^{i\omega t}$, containing the above-
141 mention singularity [6] (J - Bessel function, r - radial coordinate, k - wave number, ω - the
142 angular frequency, n - integer). Physically reasonable to present them as the sum of a
143 convergent ($\sim e^{i(kr+\pi n/2+\omega t)}$) and divergent ($\sim e^{i(-kr+\pi n/2+\omega t)}$) waves⁵. The radiation of the atoms of
144 walls excites convergent wave, which converges to the point $r = 0$, passes it some way, then
145 is transformed into a divergent one, reaches the walls of the cavity and, in the case of a
146 phase shift multiple to 2π creates a stable eigenmode. The latter condition determines the

⁵ Given expressions are valid for $kr \gg 1$.

mode spectrum, i.e. a set of allowed values $\omega=\omega_n$ ⁶. This is reminiscent of the argument given above for the rectangular cavity. There is, however, a subtle place associated with the passage of the convergent wave the point $r = 0$. As was shown in [10], convergent wave is partially captured by the curvature of the metric at $r = 0$ in the domain which size is of the order of the wavelength λ and can't conventionally, i.e. classically be transformed into divergent one. For this to happen, it is necessary to involve solutions of the M-E equations of other, non-wave type, the existence of which is proved in [2]⁷. This reminds the tunneling process in quantum mechanics: a convergent electromagnetic wave is converted into an instanton, and from it - in the divergent wave. This process occurs with probability $w \sim \exp(-\Lambda_0/\hbar)$, where $\hbar = h/2\pi$, h - is Plank constant, Λ_0 - finite Euclidean action of the instanton [2, 10]. Thus, each eigenmode of spherical cavity has a probability $w = w(\omega)$ ⁸. Electromagnetic field of the instanton and the magnitude Λ_0 were calculated in [2] and [10]. The results of both papers agree qualitatively. In [2], the action of the instanton Λ_0 was determined from the equations for the electromagnetic field produced by a variation of the action S of the field on the independent components of the field tensor F_{jk} . In [10] action Λ was recorded taking into account ties imposed on components F_{jk} , arising out of the field equations, and then the variation $\delta\Lambda$ was calculated and action Λ_0 was calculated from the condition $\delta\Lambda = 0$.

According to the results of [10] the instanton field is exponentially small at the vicinity of $r = 0$, that it corresponds to the nature of the tunnelling, and solves the problem of singularity of field of spherical electromagnetic wave at the point $r = 0$, although the metric is singular at this point.

168

169 3. ISOTROPISATION OF SELF-MODES IN SPHERICAL RESONATORS

Space-time metric, curved by the presence of a SEMW is found in [10] and looks as follows

$$\begin{aligned}
 ds^2 &= e^{-\alpha} c^2 dt^2 - e^{\alpha} dr^2 - r^2 (d\theta^2 + \sin^2 \theta \cdot d\varphi^2) \\
 e^{-\alpha} &= g_{00} = 1 - \frac{r_c}{r} + \left(\frac{r_s}{r} \right)^2 \\
 r_c &= \frac{l(l+1)c}{\omega}, r_s^2 = \frac{K}{2c^4} |G|^2
 \end{aligned}
 \tag{7}$$

G - the amplitude of the electromagnetic wave, l - an integer specifying the orbital angular momentum of the SEMW, K - the gravitational constant. Eigenmodes of the spherical cavity, as shown in [10], can be divided into scattered by curvature of metrics and captured by it. Scattered modes in terms of geometrical optics are associated with rays, which are corresponding to the areas of the front of the SEMW satisfying conditions $\theta < \theta_*$ or $\pi > \theta > \pi - \theta_*$, where θ_* - polar angle, and

⁶ In electromagnetic theory eigenmode spectrum is obtained from boundary condition on the wall of the resonator at $r = R$, R - is the radius of resonator which leads to an equation $J_{n+1/2}(kR)=0$.

⁷ In [2] they were called as instanton-like solutions

⁸ Coincidence of mode's frequency distribution with Plank one permits to connect instanton parameters with temperature of equilibrium radiation in cavity.

$$\sin \theta_* = \frac{r_c}{\rho_*} \frac{m}{l(l+1)} \quad (8)$$

180
181
182
183
184
185
186

ρ_* - is impact distance of the ray, at which capture takes place for the first time, m – integer which defines projection of orbital kinetic moment on axis OZ, $-l < m < l$ [10]. All other modes are captured by the curvature of the metric. Here we consider the scattered modes and clarify their role in shaping the field of eigenfields of the spherical electromagnetic resonators.

187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212

As is known from electromagnetic theory, electromagnetic fields in spherical resonators have axial symmetry [3]. This is due to the fixed direction of axis OZ, from which the angle θ is measured. This is true for forced oscillations in resonators excited by an external source, such as an antenna, which sets the preferred direction. However, for eigenmodes none of the preferred directions among others can be selected as the orientation of the axis OZ. The only thing that can be observed in the experiment - is the angular distribution of the eigenmodes. It doesn't permit to determine unequivocally the direction axis OZ. For example, for $l = 1$ (dipole mode) directions corresponding to $\theta=0$ and $\theta=\pi$ are equivalent and both may be selected as the orientation of axis OZ. For $l > 1$ the situation becomes even more ambiguous: all directions $\theta=m\pi/l$ are equivalent [8]⁹. The situation is exacerbated when one considers the scattering modes by curved metric. To summarize, for $l \gg 1$ any direction can be selected with an equal basis as the axis OZ, because the angular distribution of the higher modes becomes completely isotropic and gives no basis for choosing a particular direction as the axis OZ. This can be illustrated with the following considerations. Mode of the order l has an angular distribution (in the angle θ), which is characterized by the maximums with width $1/l$. Near each maximum scattered modes are concentrated, the maximum deviation angle¹⁰ of which is determined by the formula $\delta\vartheta \approx 2r_c / \rho_*$ (if one neglects the amplitude of SEMW¹¹) [9]. For the impact distance ρ_* , from which the capture begins, one can take a value $\rho_* = \sqrt{27}r_c / 2$ which is corresponding to SEMW of small amplitude [9, 15]. Then, one receives $\delta\vartheta_{\max} = 4/\sqrt{27} \approx 0.77$. Overlapping of neighboring peaks occur when the inequality $1/l + \delta\vartheta_{\max} \geq \pi/l$ will be valid, what takes place for $l \geq 3$. Thus, the observation of non-uniform angular distribution of the eigenfields of the spherical resonator is possible only for small $l=1, 2$. This corresponds to values of j , defining full kinetic moment of SEMW $j = l \pm 1 = 1, 2, 3$ (value $j = 0$ is forbidden). For the eigenfields of higher order electromagnetic fields are isotropic. Recall that we are talking about the oscillation amplitude, phase retains the dependence on the azimuthal angle φ .

213

4. APPLICATION TO COSMIC MICROWAVE BACKGROUND

214
215
216
217
218
219
220

It is interesting that these results are applicable in cosmology. Indeed, the cosmic microwave background (CMB) shows features characteristic of eigenmodes of spherical resonators: a high degree of isotropy and Planck frequency distribution [11]. To reinforce the analogy, we note two facts. First, there is a model of the universe¹², representing it as a spherical cavity with a radius increasing with time [12, 13]. The role of the walls of that cavity plays so-called last scattering surface. Cosmic microwave background radiation in this model is represented as a standing electromagnetic waves - the eigenmodes of the cavity. The model

⁹ Up to a small error, which decreases with increasing l

¹⁰ Defined by the angle of deflection of the ray corresponding to a small part of the front of the SEMW

¹¹ What can be done for the entire observable universe.

¹² Closed model of the universe

221 predicts the correct dependence of the radiation frequency of the radius of the Universe¹³
 222 [12]. It should be noted that, despite the different nature of the sources of the eigenmodes of
 223 the resonator and the relict radiation of the universe, the analogy between them is
 224 permissible, because the received radiation is likely not the primary born as a result of
 225 annihilation processes in lepton-baryon plasma that filled the universe immediately after Big
 226 Bang. Between the birth of the primary photons of the CMB and their detection by devices
 227 considerable time has passed, during which in the "universe – resonator" could finish
 228 transients formed the standing waves, taken as a relict by devices.

229 Secondly, it follows from the experimental data, the CMB in the long end of the range can be
 230 described as classical electromagnetic waves. According to generally accepted ideas CMB –
 231 is a photon gas which is formed in the Big Bang and are currently in thermal equilibrium at
 232 temperature $\sim 2,7^\circ K$ [11]. This gas fills the universe, which is described by one of the
 233 cosmological models, which are based on Einstein's equations. CMB is observed in the
 234 range from $0.33 Sm$ to $73.5 Sm$ [11]. In long wave diapason of the CMB quantum numbers
 235 of filling of photonic levels $N_k = \langle E^2 \rangle c^3 / \hbar \omega^4 \gg 1$ [14], $\langle E^2 \rangle$ - the average energy density of
 236 the microwave radiation, which is equal to $4 \cdot 10^{-13} \text{ эрг/см}^3$ [15]. This allows one to use
 237 classical equations for its description. Shortwave portion for which $N_k \ll 1$, by contrast,
 238 allows one to apply the concepts of geometrical optics.

239 5. CONCLUSIONS

240
 241 The results of this article concerning the distribution of the electromagnetic fields in cavities
 242 associated with the elimination of unphysical singularities allow for a fresh look at the role of
 243 space-time curvature in applications. The results relating to cosmology, give reason to
 244 assume that the observed anisotropy of CMB associated with harmonics with low values of
 245 the orbital angular momentum and attributed Intergalactic movements may actually be the
 246 property of the CMB caused by the interaction of electromagnetic waves with a static
 247 component of the gravitational field, or more precisely, the influence of the curvature of
 248 space-time metric, created by them. Studies conducted earlier, consider the interaction of
 249 the CMB only with gravitational waves [13].

250 Based on these results we can conclude that the light can be not only a carrier of
 251 information, but also act as its source.

252 Another finding concerns the focusing of rays in the lens system. We have already
 253 mentioned about trying to solve this problem using fictitious sinks and sources [5].
 254 Consideration of this problem in the curved space-time allows us to give another solution. It
 255 is known that the minimum area of a sphere of radius r in the space-time possessing a
 256 Schwarzschild metric is $S_{min} = 4\pi r_g^2$, r_g – gravitational radius [16]. In our problem with the
 257 metric (7), the role of gravitational radius plays the r_c ¹⁴. Instanton allows sphere (spherical
 258 front of the SEMW) after reaching the minimum area to expand in the same region I , from
 259 which it began its contraction, and not in the unphysical region I' [16]¹⁵.

260
 261
 262
 263

¹³ Strictly speaking, in these arguments the role of the radius of the universe should play radius of the sphere of last scattering.

¹⁴ At distances $r \sim r_c$ last term in (7) can be neglected, so there is a complete analogy with the Schwarzschild problem [14].

¹⁵ The latter is a figure of speech [16], because there is no time- like geodesic going from I to I' .

264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299

REFERENCES

1. V.V. Nikolsky, T.I. Nikolskaya. Electrodynamics and radio-wave propagation. Nauka, Moscow, (1989), Russian
2. Y. N. Zayko. Explicit Solutions of the Maxwell-Einstein Equations, International Journal of Theoretical and Mathematical Physics, 2011, 1(1), 1-6.
3. Zayko Y.N. Explicit Solution of Maxwell-Einstein Equations. II. International Journal of Theoretical and Mathematical Physics – IJTMP, 2011, V. 1, # 1, pp 17 – 21.
4. S. Hawking, R. Penrose. The Nature of Space and Time. Princeton Univ. Press, Princeton, New Jersey, 1999.
5. Kuligin V.A. Wave behavior in the vicinity of focus. In Problems of scattering and optimal receiving of radio waves. Proc. of Voronezh state univ, Voronezh, 1975, Russian
6. Vainshtein L.A. Electromagnetic waves. Moscow, Radio I Svyaz, 1988, Russian.
7. R. Rajaraman. An Introduction to Solitons and Instantons in Quantum Field Theory,. North-Holland Publishing Company, Amsterdam-NY-Oxford (1982).
8. G. Arfken. Mathematical Methods for Physicists, Academic Press, NY and London.
9. Berestetsky V.B., Lifshits E.M., Pitaevsky L.P. Course of theoretical physics in 10 volumes, V. 4, ch. 1, Relativistic quantum theory, Moscow, Nauka, 1968, Russian
10. Zayko Y.N. Completeness problems of being transmitted information. Proc. of the Saratov State University, 2013, V. 13, # 2, pp. 15-21, Russian
11. G.L. Trigg. Landmark EXPERIMENTS IN Twentieth Century. Physics. Crane, Russak & Company, Inc, New York, 1975
12. Berkov A.B., Kobzarev I.Yu. Gravitation Einstein theory application to astrophysics and cosmology. Tutorial, Moscow, MIFI, 1990, Russian.
13. Zeldovich Ya. B., Novikov I.D. Structure and evolution of the Universe, Moscow, Nauka, 1975, Russian.
14. Landau L.D., Lifshits E.M. Theoretical Physics. In 10 volumes, V. 2, Field Theory, 5-th Ed, Moscow, Fizmatlit, 1967, Russian
15. Ginzburg V.L. About physics and astrophysics, 2-nd Ed, Moscow, Nauka, 1992, Russian.
16. S.W. Hawking, G.F.R. Ellis. The Large Scale Structure of Space-Time. Cambridge Univ. Press, 1973.