

Original Research Article**Effect of High Voltage on Texture, Color, and Growth of Aloe Vera Leaves**

ABSTRACT

With the growth in transmission lines all over the world an evident observation comes in front regarding the effect of high voltage on the plants. High-voltage transmission lines are used to transmit electric power over long distances. High voltage may have positive or negative effects on plants and other living beings. The objective of this study is to investigate the effect of high voltage on aloe vera plants with respect to changes in texture, colour, and growth of leaves after exposure to high voltage for different durations. Images of exposed as well as unexposed aloe vera leaves were taken and analysed over a period of ten days. Other control variables like temperature, humidity, sun light were maintained almost the same for all observations.

Keywords: Aloe Vera, effect of high voltage, Tesla coil.

1. INTRODUCTION

Revolutionary and ever growing field of electrical engineering have enhanced the high voltage transmissions in day to day life. Since last few decades high voltage transmission is becoming a vital part of our life. The high voltage has much effect on living organisms as well as human beings. Plants are also multicellular, complex organisms like human beings and some studies have reported the effect of high voltage on them as well [1]. The growth and development of the plant can be affected by mechanical stimulations like wind, vibrations, and touch. This process is called thigmomorphogenesis [2]. In plants, there are various natural and artificial impacts along with large variation of potential. These days, huge amount of information regarding electrical effects in plants is available. Plants play a significant role in the living world as main producers of food and oxygen; therefore it would be advantageous to investigate their relations with today's increased exposure to high voltage electrical fields. Ksoy et al. [3] in 2010 investigated the growth of allium cepa bulbs in pots for three days on treatment area on which 380 KV high voltage power lines were passing. Ten bulbs were set up for each treatment area. Triticum baeoticum boiss. subsp. baeoticum seeds were collected at same distance from power lines on planted field. Ten seeds from each area were germinated in petri dishes for three days in laboratory. The treatment groups were compared with the control group for mitotic index and chromosome aberrations. Data obtained from this revealed that the electromagnetic fields from high voltage power lines improved the mitotic index and chromosome aberrations. Maziah et al. [4] investigated that electromagnetic field could be used as a tool to promote mustard growth by means of photosynthesis once the right EMF strength is exposed. The effects of electromagnetic fields at high voltage transmission line on biochemical and antioxidant system changes in mustard leaf (*brassica chinensis*) were investigated under field condition. Mustard leaves were exposed to EMF from power lines at various distances. The effects of

EMF on leaf mustard planted at different distances from the power line showed that the leaf mustard planted within 20 m from the power line had considerably higher protein, soluble protein, soluble nitrogen, and chlorophyll contents due to the higher EMF strength which decreased with increasing distance from the line. Higher EMF strength closer to the 275 KV power line resulted in higher peroxidase enzymatic activity, and chlorophyll content. Thus the new and energetic area of scientific research is the studies associated to the high voltage effects. In India, high voltage exposure is very common as shown in Fig. 1.



Fig. 1 Plants under high voltage agricultural fields in roop nagar area of Jammu

Agricultural science takes interest not only in the general and valued crop-forming factors, but also in those which are inexpensive and generally underestimated, though more pro-ecological ones which include ionizing, laser, ultraviolet radiation, electric, and magnetic fields [5]. The objective of this paper is to investigate the effect of high voltage on texture, colour as well as on the growth of the Aloe Vera plants. The Aloe Vera has been chosen because of its numerous applications. The Aloe Vera and its characteristics are described in Section 2. The methodology is presented in Section 3 and results and discussions in Section 4 along with the conclusions in Section 5.

2. ALOE VERA

Aloe Vera is a perennial liliaceous plant which has juicy green leaves attached at the stem in a whorled pattern [6]. It is a hardy, perennial, tropical, drought-resistant, succulent plant and has been used for a variety of medicinal purposes. Aloe Vera is a stem less or very short-stemmed plant growing upto 60– 90 cm in height. The stem has dormant root buds that develop to form new roots. The leaves are thick and fleshy, green to grey-green, with some varieties showing white flecks on the upper and lower stem surfaces [7]. Mature plants can be grown as tall as four feet with average height around 30 inches. Each plant has an average of 15 leaves usually and weighs up to 2-3 kg when fully matured. The Aloe Vera plant can be harvested following every 6 to 8 weeks by removing 3-4 leaves per plant [8]. The peripheral bundle sheath cells of Aloe Vera produces extremely bitter, yellow latex, which is commonly known as aloe juice or sap [9]. Aloe Vera has also been used for medicinal purposes in several countries like India, Mexico, Greece, Egypt, China and Japan [10]. Fig. 2 shows a Aloe Vera tree grown in a house.



Fig. 2 Aloe Vera

Aloe Vera has extensive applications in Ayurveda for its anti-burn effect. It also acts as a tonic, antiseptic, antibiotic, anti-diarrheal, anti-fungal, anti-viral, and also a good hair conditioner [11]. In Ayurveda, which is also the traditional medicine of India, Aloe Vera has manifold uses such as it is used as a laxative, in haemorrhoid remedy, and as a uterine stimulant. It is also use to treat eczema or psoriasis [12]. Recently, Aloe Vera has been used to treat canker sores [13]. The most important part of Aloe Vera is Aloe gel. Aloe Vera gel is 99% water. It has a pH of 4.5 and is a common ingredient in many non-prescription skin salves. The gel also consists of the bulk of the leaf substance, which serves as the water storage space organ for the plant. In addition to this, the Aloe gel consist of polysaccharide, glucomannan, etc. [14].

3. METHODOLOGY

The investigations were carried out on Aloe Vera plants to study the effect of high voltage on the leaves with respect to texture, colour, and growth. In this research, the photograph of Aloe Vera leaf before high voltage exposure was taken and it was studied. After that the Aloe Vera plant was exposed to high voltage. The Tesla coil shown Fig. 3 was used to produce high voltage of order 50 KV.



Fig. 3 Tesla coil

103

104

105

106

107

108


109

110

111

112

Tesla coil is an electrical resonant transformer which is used to produce high voltage low current, high frequency alternating current electricity. It is essentially a high frequency air core transformer. Tesla coil can produce comparatively higher voltages than other artificial sources of high-voltage discharges or electrostatic machines. The voltage across the Tesla coil was 50 kV. After that two Aloe Vera plants were taken and were kept at a distance of 15 cm and 50 cm respectively from the Tesla coil respectively. Three leaves from each Aloe Vera plant were taken to study the effect of high voltage on it. On each of the leaves two surface electrodes were attached at a distance of 6.5 cm in between them. The high voltage experimental setup can be explained in the Fig. 4.



113

114

115

Fig. 4 Block diagram of the experiment

116

117

118

119

120

121

Then these Plants were kept in Faraday cage along with the Tesla coil to study the effect of high voltage. Faraday cage is a metallic enclosure that prevents the entry or escape of an electromagnetic field. It is an enclosure formed by conducting materials which blocks external electromagnetic fields. For best performance, the cage should be directly connected to an earth ground. A heavy duty Faraday cage can shield against direct lightning strikes.



122

123

124

Fig. 5 Faraday's cage

125 The exposed Aloe Vera leaf was kept under natural environmental condition for ten days i.e.
 126 the control variables like temperature, humidity, sun light were maintained almost same for
 127 all observations. The plant was watered every day. The Aloe Vera leaf was observed with
 128 respect to change in physical properties like texture, colour and growth. The texture and
 129 colour was observed by taking images of the Aloe Vera leaf and the growth was recorded by
 130 a thick thread measured from the base of the leaf just above the root to the tip of the leaf.
 131 The setup to record the growth of the Aloe Vera leaf is shown in Fig. 6.
 132



Fig. 6 Setup to record growth of Aloe Vera leaves

133
 134
 135
 136

4. RESULTS AND DISCUSSIONS

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

5. CONCLUSIONS

155

156

157

158

159

160

161

162

163

164

The investigations were carried out to study the effect of high voltage on Aloe Vera leaves. The analysis of the results showed that high voltage has a definite adverse effect on the Aloe Vera leaves and growth. With passage of time, the recovery process of the plant tries to minimize the damage. If the level of the exposure is very high, the damage is irreversible. Some colour changes were also observed in the unexposed leaves. This may be due to the wrapping of the leaves, which stops the sun light in the wrapped portion of the leaves. This type of damage is easily recovered within 2-3 days.

165

166 **Table I. Texture, colour, and growth of Aloe Vera leaves before and after exposure to**
 167 **high voltages.**

168

DAY	PARAMETER	PLANT 1 L1	PLANT 1 L2	PLANT 1 L3 (UNEXPOSED)	PLANT 2 L1	PLANT 2 L2	PLANT 2 L3 (UNEXPOSED)
D0	TEXTURE	HARD	HARD	HARD	HARD	HARD	HARD
	COLOUR	GREEN	GREEN	GREEN	GREEN	GREEN	GREEN
	LENGTH	30 CM	31 CM	28 CM	29 CM	30.5 CM	28.5 CM
	DAMAGE	NIL	NIL	NIL	NIL	NIL	NIL
D1	TEXTURE	HARD	HARD	HARD	HARD	HARD	HARD
	COLOUR	GREEN	GREEN	GREEN	GREEN	GREEN	GREEN
	GROWTH	29CM	31 CM	28 CM	28 CM	30.5 CM	28.5 CM
	DAMAGE	LIGHT BLACK SPOTS	NIL	NIL	ONE BIG AND ONE SMALL BLACK SPOT	TIP OF LEAF STARTED SHRINKING	NIL
D3	TEXTURE	HARD	HARD	HARD	HARD	HARD	HARD
	COLOUR	YELLOW GREEN	GREEN	GREEN	YELLOW GREEN	GREEN	GREEN
	GROWTH	29 CM	31 CM	28 CM	28 CM	30 CM	28.5 CM
	DAMAGE	DARK BLACK SPOTS	NIL	NIL	PROMINENT BLACK SPOTS	NIL	NIL
D5	TEXTURE	HARD	HARD	HARD	HARD	HARD	HARD
	COLOUR	YELLOW GREEN	YELLOW GREEN (STARTED FADING)	GREEN	YELLOW GREEN	YELLOW GREEN	YELLOW GREEN
	GROWTH	28 CM	31 CM	28 CM	29 CM	30 CM	28.5 CM
	DAMAGE	DARK BLACK SPOTS	VERY SMALL BLACK SPOTS	NIL	DARK BLACK SPOTS	TIP OF THE LEAF SHRINKED FURTHER	NIL
D7	TEXTURE	HARD	HARD	HARD	HARD	HARD	HARD
	COLOUR	YELLOW (FADED)	YELLOW	STARTED TO FADE YELLOW.		YELLOW	YELLOW
	GROWTH	15 CM	31 CM	28 CM	29 CM	30 CM	28.5 CM
	DAMAGE	TOTALLY DAMAGED AND LEAF FALLEN OFF	SMALL BLACK SPOTS	NIL	NIL		

169




Fig. 7 Images of Aloe Vera leaves taken on 1st, 3rd, 5th and 7th day.

REFERENCES

1. Kulkarni G, Gandhare WZ. Proximity Effects of High Voltage Transmission Lines on Humans. International Journal on Electrical and Power Engineering. 2012; 3:28-32.

2. Uchida A, Yamamoto KT. Effect of mechanical vibration on seed germination of Arabidopsis thaliana (L.) Heynh, Plant Cell Physiology. 2002; 43:647-651.

- 179 3. Aksoy H, Unal G, Ozcan S. Genotoxic Effects of Electromagnetic Fields from High
180 Voltage Power Lines on Some Plants. International Journal of Environmental
181 Research. 2010; 4:595-606.
182
- 183 4. Maziah M, Ooi BB, Tengku M, and Sreeramanan S. Effects of electromagnetic field
184 of 33 and 275 kV influences on physiological, biochemical and antioxidant system
185 changes of leaf mustard (*Brassica chinensis*). African Journal of Biotechnology.
186 2012; 11:13016-13029.
187
- 188 5. Kordas L. The Effect of Magnetic Field on Growth, Development and the Yield of
189 Spring Wheat, Polish Journal of Environmental Studies. 2002;11: 527-530.
190
- 191 6. Sheikh F.A, Singh R.P.P, Singh J.B., Lehana P. Effect of microwaves on the
192 resistance of Aloe Vera leaves. International Journal of Engineering Research and
193 Applications. 2013; 3:242-247.
194
- 195 7. Available online: www.wikipedia.org/Aloe_vera.
196
- 197 8. Nandal U, Bhardwaj RL, Aloe Vera: A valuable wonder plant for food, medicine and
198 cosmetic use - A review, International Journal of Pharmaceutical Sciences Review
199 and Research, 2012; 13:59-67.
200
- 201 9. Vogler BK, Ernest E. Aloe Vera: a systematic review of its clinical effectiveness.
202 British Journal of General Practice. 1999; 49:823-828.
203
- 204 10. Marshall JM, Aloe Vera gel: What is the evidence? The Pharmaceutical
205 Journal, 1990; 360-36.
206
- 207 11. Prajapati M, Patel PS, Vyas PJ. Phytochemical Analysis of Aloe Vera And Study of
208 Mixing Antibiotic with Aloe Vera and Its Antibacterial Activity, Asian Journal of
209 Biochemical and Pharmaceutical Research. 2011; 1:473-479.
210
- 211 12. Saeed MA, Ahmad I, Yaqub U, Akbar, Waheed MA, Saleem A, and Nasir-ud-Din.
212 Aloe vera a plant of vital significance, Science vision. 2004; 9:1-2.
213
- 214 13. Cera LM, Hegggers JP, Robson MC, and Hagstrom WJ. The Therapeutical Efficacy of
215 Aloe Vera Cream (Dermaid Aloe) in Thermal Injuries: Two Case Reports, Journal of
216 the American Animal Hospital Association. 1988; 16:786-772.
217
- 218 14. Sharma SK, Singh R, Lehana P. To Investigate the Electrical Impedance of the Aloe
219 Barbadensis Miller Leaves, International Journal of Soft Computing and
220 Engineering, 2012; 2:234-238.