

The Modified Simple Equation Method and Its Application to Solve NLEEs Associated with Engineering Problem

Md. Ashrafuzzaman Khan and M. Ali Akbar

Department of Applied Mathematics, University of Rajshahi, Rajshahi-6205, Bangladesh

ABSTRACT

The modified simple equation (MSE) method is an important mathematical tool for searching closed-form solutions to nonlinear evolution equations (NLEEs). In the present paper, by using the MSE method, we derive some impressive solitary wave solutions to NLEEs via the strain wave equation in microstructured solids which is a very important equation in the field of engineering. The solutions contain some free parameters and for particular values of the parameters some known solutions are derived. The solutions exhibit necessity and reliability of the MSE method.

Keywords: Modified simple equation method; balance number; solitary wave solutions; strain wave equation; microstructured solids.

Mathematics Subject Classification: 35C07, 35C08, 35P99.

1. INTRODUCTION

Physical systems are in general explained with nonlinear partial differential equations. The mathematical modeling of microstructured solid materials that change over time depends closely on the study of a variety of systems of ordinary and partial differential equations. Similar models are developed in diverse fields of study, ranging from the natural and physical sciences, population ecology to economics, infectious disease epidemiology, neural

E-mail address: ali_math74@yahoo.com.

24 networks, biology, mechanics etc. In spite of the eclectic nature of the fields wherein these
25 models are formulated, different groups of them contribute adequate common attributes that
26 make it possible to examine them within a unified theoretical structure. Such study is an area
27 of functional analysis, usually called the theory of evolution equations. Therefore, the
28 investigation of solutions to NLEEs plays a very important role to uncover the obscurity of
29 many phenomena and processes throughout the natural sciences. However, one of the
30 essential problems is to obtain their closed-form solutions. For that reason, diverse groups
31 of engineers, physicists, and mathematicians have been working tirelessly to investigate
32 closed-form solutions to NLEEs. Accordingly, in the recent years, they establish several
33 methods to search exact solutions, for instance, the Darboux transformation method [1], the
34 Jacobi elliptic function method [2, 3], the He's homotopy perturbation method [4, 5], the tanh-
35 function method [6, 7], the extended tanh-function method [8, 9], the Lie group symmetry
36 method [10], the variational iteration method [11], the Hirota's bilinear method [12], the
37 Backlund transformation method [13, 14], the inverse scattering transformation method [15],
38 the sine-cosine method [16, 17], the Painleve expansion method [18], the Adomian
39 decomposition method [19, 20], the (G'/G) -expansion method [21-26], the first integration
40 method [27], the F-expansion method [28], the auxiliary equation method [29], the ansatz
41 method [30, 31], the Exp-function method [32, 33], the homogeneous balance method [34],
42 the modified simple equation method [35-47], the $\exp(-\phi(\eta))$ -expansion method [48, 49], the
43 Miura transformation method [50], and others.

44 Microstructured materials like crystallites, alloys, ceramics, and functionally graded materials
45 have gained broad application. The modeling of wave propagation in such materials should
46 be able to account for various scales of microstructure [51]. In the past years, many authors
47 have studied the strain wave equation in microstructured solids, such as, Alam et al. [51]
48 solved this equation by using the new generalized (G'/G) -expansion method. Pastrone et
49 al. [52], Porubov and Pastrone [53] examined bell-shaped and kink-shaped solutions of this
50 engineering problem. Akbar et al. [54] constructed traveling wave solutions of this equation
E-mail address: ali_math74@yahoo.com.

by using the generalized and improved (G'/G) -expansion method. The above analysis shows that several methods to achieve exact solutions to this equation have been accomplished in the recent years. But, the equation has not been studied by means of the MSE method. In this article, our aim is, we will apply the MSE method following the technique derived in the Ref. [55] to examine some new and impressive solitary wave solutions to this equation.

The structure of this article is as follows: In section 2, we describe the method. In section 3, we apply the MSE method to the strain wave equation in microstructured solids. In section 4, we provide the physical interpretations of the obtained solutions. Finally, in section 5, conclusions are given.

2. DESCRIPTION OF THE METHOD

Assume the nonlinear evolution equation has the following form

$$P(u, u_t, u_x, u_y, u_z, u_{xx}, u_{tt}, \dots) = 0, \quad (2.1)$$

where $u = u(x, y, z, t)$ is an unidentified function, P is a polynomial function in $u = u(x, y, z, t)$ and its partial derivatives, wherein nonlinear term of the highest order and the highest order linear terms exist and subscripts indicate partial derivatives. To solve (2.1) by using the MSE method [35-47], we need to perform the subsequent steps:

Step 1: Now, we combine the real variable x and t by a compound variable ξ as follows:

$$u(x, y, z, t) = U(\xi), \quad \xi = x + y + z \pm \omega t. \quad (2.2)$$

Here ξ is called the wave variable it allows us to switch Eq. (2.1) into an ordinary differential equation (ODE):

$$Q(U, U', U'', U''', \dots), \quad (2.3)$$

where Q is a polynomial in $U(\xi)$ and its derivatives, where $U'(\xi) = \frac{dU}{d\xi}$.

Step 2: We assume that Eq. (2.3) has the traveling wave solution in the following form,

$$U(\xi) = \sum_{i=0}^N a_i \frac{\psi'(\xi)}{\psi(\xi)}^i, \quad (2.4)$$

where a_i , ($i=0,1,2,\dots,N$) are arbitrary constants, such that $a_N \neq 0$, and $\psi(\xi)$ is an unidentified function which is to be determined later. In (G'/G) -expansion method, Exp-function method, tanh-function method, sine-cosine method, Jacobi elliptic function method etc., the solutions are initiated through several auxiliary functions which are previously known, but in the MSE method, $\psi(\xi)$ is neither a pre-defined function nor a solution of any pre-defined differential equation. Therefore, it is not possible to speculate from formerly, what kind of solution can be found by this method.

Step 3: We determine the positive integer N , come out in Eq. (2.4) by taking into account the homogeneous balance between the highest order nonlinear terms and the derivatives of the highest order occurring in Eq. (2.3).

Step 4: We calculate the necessary derivatives U' , U'' , U''' , etc., then insert them into Eq. (2.3) and then taken into consideration the function $\psi(\xi)$. As a result of this insertion, we obtain a polynomial in $(\psi'(\xi)/\psi(\xi))$. We equate all the coefficients of $(\psi'(\xi)/\psi(\xi))^i$, ($i=0,1,2,\dots,N$) to this polynomial to zero. This procedure yields a system of algebraic and differential equations whichever can be solved for getting a_i ($i=0,1,2,\dots,N$), $\psi(\xi)$ and the value of the other parameters.

3. APPLICATION OF THE METHOD

In this section, we will execute the application of the MSE method to extract solitary wave solutions to the strain wave equation in microstructured solids which is a very important equation in the field of engineering. Let us consider the strain wave equation in microstructured solids:

$$u_{tt} - u_{xx} - \varepsilon \alpha_1 (u^2)_{xx} - \gamma \alpha_2 u_{xxt} + \delta \alpha_3 u_{xxxx} - (\delta \alpha_4 - \gamma^2 \alpha_7) u_{xxtt} + \gamma \delta (\alpha_5 u_{xxxxt} + \alpha_6 u_{xxtt}) = 0. \quad (3.1)$$

97

98 **3.1. THE NON-DISSIPATIVE CASE**

99 The system is non-dissipative, if $\gamma=0$ and determined by the double dispersive equation
100 (see [52], [53], [56], [57] for details)

$$u_{tt} - u_{xx} - \varepsilon \alpha_1 (u^2)_{xx} + \delta \alpha_3 u_{xxxx} - \delta \alpha_4 u_{xxtt} = 0. \quad (3.2)$$

101 The balance between dispersion and nonlinearities happen when $\delta = O(\varepsilon)$. Therefore, (3.2)
102 becomes

$$u_{tt} - u_{xx} - \varepsilon \{ \alpha_1 (u^2)_{xx} - \alpha_3 u_{xxxx} + \alpha_4 u_{xxtt} \} = 0. \quad (3.3)$$

103 In order to extract solitary wave solutions of the strain wave equation in microstructured
104 solids by using the MSE method, we use the traveling wave variable

$$u(x, t) = U(\xi), \quad \xi = x - \omega t. \quad (3.4)$$

105 The wave transformation (3.4) reduces Eq. (3.3) into the ODE in the following form:

$$(\omega^2 - 1) U'' - \varepsilon \{ \alpha_1 (U^2)'' - (\alpha_3 - \omega^2 \alpha_4) U^{(iv)} \} = 0. \quad (3.5)$$

106 where primes indicate differential coefficients with respect to ξ . Eq. (3.5) is integrable,
107 therefore, integration (3.5) as many time as possible, we obtain the following ODE:

$$(\omega^2 - 1) U - \varepsilon \{ \alpha_1 U^2 - (\alpha_3 - \omega^2 \alpha_4) U'' \} = 0. \quad (3.6)$$

108 where the integration constants are set zero, as we are seeking solitary wave solutions.

109 Taking homogeneous balance between the terms U'' and U^2 appearing in Eq. (3.6), we
110 obtain $N=2$. Therefore, the shape of the solution of Eq. (3.6) becomes

$$U(\xi) = a_0 + \frac{a_1 \psi'}{\psi} + \frac{a_2 (\psi')^2}{\psi^2}. \quad (3.7)$$

111 wherein a_0 , a_1 and a_2 are constants to be find out afterward such that $a_2 \neq 0$, and $\psi(\xi)$ is
 112 an unknown function. The derivatives of U are given in the following:

$$U' = -\frac{a_1(\psi')^2}{\psi^2} - \frac{2a_2(\psi')^3}{\psi^3} + \frac{a_1\psi''}{\psi} + \frac{2a_2\psi'\psi''}{\psi^2}. \quad (3.8)$$

$$U'' = a_1 \left\{ \frac{2(\psi')^3}{\psi^3} - \frac{3\psi'\psi''}{\psi^2} + \frac{\psi'''}{\psi} \right\} + 2a_2 \left\{ \frac{(\psi'')^2}{\psi^2} + \frac{\psi'\psi'''}{\psi^2} - \frac{5(\psi')^2\psi''}{\psi^3} + \frac{3(\psi')^4}{\psi^4} \right\}. \quad (3.9)$$

113 Inserting the values of U , U' and U'' into Eq. (3.6), and setting each coefficient of ψ^j , $j =$
 114 $0, 1, 2, \dots$ to zero, we derive, successively

$$a_0(-1 + \omega^2 - \varepsilon a_0\alpha_1) = 0. \quad (3.10)$$

$$a_1\{(-1 + \omega^2 - 2\varepsilon a_0\alpha_1)\psi' + \varepsilon(\alpha_3 - \omega^2\alpha_4)\psi'''\} = 0. \quad (3.11)$$

$$\begin{aligned} -\varepsilon a_1\psi'\{a_1\alpha_1\psi' + 3(\alpha_3 - \omega^2\alpha_4)\psi''\} + 2a_2\varepsilon(\alpha_3 - \omega^2\alpha_4)\psi'\psi''' \\ + a_2\{(-1 + \omega^2 - 2\varepsilon a_0\alpha_1)(\psi')^2 + 2\varepsilon(\alpha_3 - \omega^2\alpha_4)(\psi'')^2\} = 0. \end{aligned} \quad (3.12)$$

$$-2\varepsilon(\psi')^2\{a_1(a_2\alpha_1 - \alpha_3 + \omega^2\alpha_4)\psi' + 5a_2(\alpha_3 - \omega^2\alpha_4)\psi''\} = 0. \quad (3.13)$$

$$-\varepsilon a_2(a_2\alpha_1 - 6\alpha_3 + 6\omega^2\alpha_4)(\psi')^4 = 0. \quad (3.14)$$

115 From Eq. (3.10) and Eq. (3.14), we obtain

$$a_0 = 0, \quad \frac{-1 + \omega^2}{\varepsilon\alpha_1} \quad \text{and} \quad a_2 = \frac{6(\alpha_3 - \omega^2\alpha_4)}{\alpha_1}, \quad \text{since } a_2 \neq 0.$$

116 Therefore, for the values of a_0 , there arise the following cases:

117 **Case 1:** When $a_0 = 0$, from Eqs. (3.11)-(3.13), we obtain

$$a_1 = \pm \frac{6\sqrt{1 - \omega^2}\sqrt{\alpha_3 - \omega^2\alpha_4}}{\sqrt{\varepsilon}\alpha_1}$$

118 and

$$\psi(\xi) = c_2 + \frac{\varepsilon c_1(-\alpha_3 + \omega^2\alpha_4)}{-1 + \omega^2} e^{\mp \frac{\xi\sqrt{1 - \omega^2}}{\sqrt{\varepsilon}\sqrt{\alpha_3 - \omega^2\alpha_4}}},$$

119 where c_1 and c_2 are integration constants.

120 Substituting the values of a_0, a_1, a_2 and $\psi(\xi)$ into Eq. (3.7), we obtain the following
 121 exponential form solution:

$$U(\xi) = \frac{6e^{\pm \frac{\xi\sqrt{1-\omega^2}}{\sqrt{\varepsilon}\sqrt{\alpha_3-\omega^2\alpha_4}}}(-1+\omega^2)^2c_1c_2(-\alpha_3+\omega^2\alpha_4)}{\alpha_1 \left((-1+\omega^2)c_2 e^{\pm \frac{i\xi\sqrt{-1+\omega^2}}{\sqrt{\varepsilon}\sqrt{\alpha_3-\omega^2\alpha_4}}} + \varepsilon c_1(-\alpha_3+\omega^2\alpha_4) \right)^2}. \quad (3.15)$$

122 Simplifying the required solution (3.15), we derive the following close-form solution of the
 123 strain wave equation in microstructured solids (3.3):

$$u(x, t) = \{6(-1+\omega^2)^2c_1c_2(-\alpha_3+\omega^2\alpha_4)\} \\ / \left[\alpha_1 \left\{ \pm i \sin((x-t\omega)\beta) \{(-1+\omega^2)c_2 + \varepsilon c_1(\alpha_3 - \omega^2\alpha_4)\} \right. \right. \\ \left. \left. + \cos((x-t\omega)\beta) \{(-1+\omega^2)c_2 + \varepsilon c_1(-\alpha_3 + \omega^2\alpha_4)\} \right\}^2 \right] \quad (3.16)$$

124 where $\beta = \frac{\sqrt{-1+\omega^2}}{2\sqrt{\varepsilon}\sqrt{\alpha_3-\omega^2\alpha_4}}$. Solution (3.16) is the generalized solitary wave solution of the strain
 125 wave equation in microstructured solids. Since c_1 and c_2 are arbitrary constants, one might
 126 arbitrarily choose their values. Therefore, if we choose $c_1 = (-1+\omega^2)$ and $c_2 = \varepsilon(-\alpha_3 +$
 127 $\omega^2\alpha_4)$ then from (3.16), we obtain the following bell shaped soliton solution:

$$u_1(x, t) = \frac{3(-1+\omega^2)}{2\varepsilon\alpha_1} \operatorname{sech}^2 \left(\frac{(x-t\omega)\sqrt{-1+\omega^2}}{2\sqrt{\varepsilon}\sqrt{-\alpha_3+\omega^2\alpha_4}} \right). \quad (3.17)$$

128 Again, if we choose $c_1 = (-1+\omega^2)$ and $c_2 = -\varepsilon(-\alpha_3 + \omega^2\alpha_4)$, then from (3.16), we obtain
 129 the following singular soliton:

$$u_2(x, t) = -\frac{3(-1+\omega^2)}{2\varepsilon\alpha_1} \operatorname{csch}^2 \left(\frac{(x-t\omega)\sqrt{-1+\omega^2}}{2\sqrt{\varepsilon}\sqrt{-\alpha_3+\omega^2\alpha_4}} \right). \quad (3.18)$$

130 On the other hand, when $c_1 = (-1+\omega^2)$ and $c_2 = \pm i \varepsilon(-\alpha_3 + \omega^2\alpha_4)$, from solution (3.16),
 131 we obtain the following trigonometric solution:

$$u_3(x, t) = \frac{3(-1+\omega^2)}{2\varepsilon\alpha_1} \sec^2 \left[\frac{1}{4} \left\{ \pi + \frac{2(x-t\omega)\sqrt{-1+\omega^2}}{\sqrt{\varepsilon}\sqrt{\alpha_3-\omega^2\alpha_4}} \right\} \right]. \quad (3.19)$$

E-mail address: ali_math74@yahoo.com.

132 Again when $c_1 = (-1 + \omega^2)$ and $c_2 = \mp i \varepsilon(-\alpha_3 + \omega^2 \alpha_4)$, then the generalized solitary wave
 133 solution (3.16) can be simplified as:

$$u_4(x, t) = \frac{3(-1 + \omega^2)}{2\varepsilon\alpha_1} \csc^2 \left[\frac{1}{4} \left\{ \pi + \frac{2(-x + t\omega)\sqrt{-1 + \omega^2}}{\sqrt{\varepsilon}\sqrt{\alpha_3 - \omega^2\alpha_4}} \right\} \right]. \quad (3.20)$$

134 If we choose more different values of c_1 and c_2 , we may derive a lot of general solitary
 135 wave solutions to the Eq. (3.3) through the MSE method. For succinctness, other solutions
 136 have been overlooked.

137 **Case 2:** When $a_0 = \frac{-1 + \omega^2}{\varepsilon\alpha_1}$, then Eqs. (3.11)-(3.13) yield

$$a_1 = \pm \frac{6\sqrt{-1 + \omega^2}\sqrt{\alpha_3 - \omega^2\alpha_4}}{\sqrt{\varepsilon}\alpha_1}$$

138 And

$$\psi(\xi) = c_2 + \frac{\varepsilon c_1(\alpha_3 - \omega^2 \alpha_4)}{-1 + \omega^2} e^{\mp \frac{\xi\sqrt{-1 + \omega^2}}{\sqrt{\varepsilon}\sqrt{\alpha_3 - \omega^2\alpha_4}}},$$

139 where c_1 and c_2 are constants of integration.

140 Now, by means of the values of a_0 , a_1 , a_2 and $\psi(\xi)$, from Eq. (3.7), we obtain the
 141 subsequent solution:

$$U(\xi) = \frac{-1 + \omega^2}{\varepsilon\alpha_1} + \frac{6(-1 + \omega^2)^2 c_1 c_2 (-\alpha_3 + \omega^2 \alpha_4) e^{\pm \frac{\xi\sqrt{-1 + \omega^2}}{\sqrt{\varepsilon}\sqrt{\alpha_3 - \omega^2\alpha_4}}}}{\alpha_1 \left\{ (-1 + \omega^2) c_2 e^{\pm \frac{\xi\sqrt{-1 + \omega^2}}{\sqrt{\varepsilon}\sqrt{\alpha_3 - \omega^2\alpha_4}}} + \varepsilon c_1 (\alpha_3 - \omega^2 \alpha_4) \right\}^2}. \quad (3.21)$$

142 Now, transforming the required exponential function solution (3.21) into hyperbolic function,
 143 we obtain the following solution to the strain wave equation in the microstructured solids:

$$\begin{aligned}
u(x, t) = & (-1 + \omega^2) [(-1 + \omega^2)^2 \{ \cosh(2\rho(x - t\omega)) + \sinh(2\rho(x - t\omega)) \} c_2^2 \\
& + \varepsilon^2 \{ \cosh(2\rho(x - t\omega)) - \sinh(2\rho(x - t\omega)) \} c_1^2 (\alpha_3 - \omega^2 \alpha_4)^2 \\
& + 4\varepsilon(-1 + \omega^2) c_1 c_2 (-\alpha_3 + \omega^2 \alpha_4)] \\
& / \left(\varepsilon \alpha_1 [(-1 + \omega^2) \{ \cosh(\rho(x - t\omega)) + \sinh(\rho(x - t\omega)) \} c_2 \right. \\
& \left. + \varepsilon \{ \cosh(\rho(x - t\omega)) - \sinh(\rho(x - t\omega)) \} c_1 (\alpha_3 - \omega^2 \alpha_4) \right]^2. \quad (3.22)
\end{aligned}$$

144 Thus, we acquire the generalized solitary wave solution (3.22) to the strain wave equation in
145 microstructured solids, where $\rho = \frac{\sqrt{-1+\omega^2}}{2\sqrt{\varepsilon}\sqrt{\alpha_3-\omega^2\alpha_4}}$. Since c_1 and c_2 are integration constants,
146 therefore, somebody might randomly pick their values. So, if we pick $c_1 = (-1 + \omega^2)$ and
147 $c_2 = -\varepsilon(\alpha_3 - \omega^2 \alpha_4)$, then from (3.22), we obtain the subsequent solitary wave solution:

$$u_5(x, t) = \frac{(-1 + \omega^2)}{2\varepsilon\alpha_1} \left\{ 2 + 3 \operatorname{csch}^2 \left(\frac{(x - t\omega)\sqrt{-1 + \omega^2}}{2\sqrt{\varepsilon}\sqrt{\alpha_3 - \omega^2 \alpha_4}} \right) \right\}. \quad (3.23)$$

148 Again, if we pick $c_1 = (-1 + \omega^2)$ and $c_2 = \varepsilon(\alpha_3 - \omega^2 \alpha_4)$, then the solitary wave solution
149 (3.22) reduces to:

$$u_6(x, t) = -\frac{(-1 + \omega^2)}{2\varepsilon\alpha_1} \left\{ -2 + 3 \operatorname{sech}^2 \left(\frac{(x - t\omega)\sqrt{-1 + \omega^2}}{2\sqrt{\varepsilon}\sqrt{\alpha_3 - \omega^2 \alpha_4}} \right) \right\}. \quad (3.24)$$

150 Moreover, if we pick $c_1 = (-1 + \omega^2)$ and $c_2 = \mp i \varepsilon(\alpha_3 - \omega^2 \alpha_4)$, then from (3.22), we derive
151 the following solution:

$$u_7(x, t) = \frac{(-1 + \omega^2)}{\varepsilon\alpha_1} \left\{ 1 - \frac{3}{2} \operatorname{csc}^2 \left(\frac{\pi}{4} - \frac{1}{2} \frac{(x - t\omega)\sqrt{-1 + \omega^2}}{\sqrt{\varepsilon}\sqrt{-\alpha_3 + \omega^2 \alpha_4}} \right) \right\}. \quad (3.25)$$

152 Again, if we pick $c_1 = (-1 + \omega^2)$ and $c_2 = \pm i \varepsilon(\alpha_3 - \omega^2 \alpha_4)$, then from (3.22), we obtain the
153 following solution:

$$u_8(x, t) = \frac{(-1 + \omega^2)}{\varepsilon\alpha_1} \left\{ 1 - \frac{3}{2} \operatorname{csc}^2 \left(\frac{\pi}{4} + \frac{1}{2} \frac{(x - t\omega)\sqrt{-1 + \omega^2}}{\sqrt{\varepsilon}\sqrt{-\alpha_3 + \omega^2 \alpha_4}} \right) \right\}. \quad (3.26)$$

Forasmuch as, c_1 and c_2 are arbitrary constants, if we choose more different values of them, we may derive a lot of general solitary wave solutions to the Eq. (3.3) through the MSE method easily. But, we did not write down the other solutions for minimalism.

Remark 1: Solutions (3.17)-(3.20) and (3.23)-(3.26) have been confirmed by inserting them into the main equation and found accurate.

3.2. THE DISSIPATIVE CASE

If $\gamma \neq 0$, then the system is dissipative. Therefore, for $\delta = \gamma = O(\varepsilon)$, the balance should be between nonlinearity, dispersion and dissipation, perturbed by the higher order dissipative terms to the strain wave equation in microstructured solids (see [52], [53], [56], [57] for details)

$$u_{tt} - u_{xx} - \varepsilon \{ \alpha_1 (u^2)_{xx} + \alpha_2 u_{xxt} - \alpha_3 u_{xxxx} + \alpha_4 u_{xxtt} \} = 0. \quad (3.27)$$

where $\varepsilon \rightarrow 0$, so the higher order term are omitted.

The traveling wave transformation (3.4) reduces Eq. (3.27) to the following ODE:

$$(\omega^2 - 1) U'' - \varepsilon \{ \alpha_1 (U^2)'' - \omega \alpha_2 U''' - (\alpha_3 - \omega^2 \alpha_4) U^{(iv)} \} = 0. \quad (3.28)$$

where prime stands for the differential coefficient. Integrating Eq. (3.28) with respect to ξ , we get

$$(\omega^2 - 1) U - \varepsilon \{ \alpha_1 U^2 - \omega \alpha_2 U' - (\alpha_3 - \omega^2 \alpha_4) U'' \} = 0. \quad (3.29)$$

The homogeneous between the highest order nonlinear term and the linear terms of the highest order, we obtain $N = 2$. Thus, the structure of the solution of Eq. (3.29) is one and the same to the form of the solution (3.7).

Inserting the values of U , U' and U'' into Eq. (3.29) and then setting each coefficient of ψ^{-j} , $j = 0, 1, 2, \dots$ to zero, we successively obtain

$$a_0(-1 + \omega^2 - \varepsilon a_0 \alpha_1) = 0. \quad (3.30)$$

$$a_1 \{ (-1 + \omega^2 - 2\varepsilon a_0 \alpha_1) \psi' + \varepsilon \omega \alpha_2 \psi'' + \varepsilon (\alpha_3 - \omega^2 \alpha_4) \psi''' \} = 0. \quad (3.31)$$

E-mail address: ali_math74@yahoo.com.

$$\begin{aligned}
& -\varepsilon a_1 \psi' \{ (a_1 \alpha_1 + \omega \alpha_2) \psi' + 3(\alpha_3 - \omega^2 \alpha_4) \psi'' \} + 2\varepsilon a_2 \psi' \{ \omega \alpha_2 \psi'' + (\alpha_3 - \omega^2 \alpha_4) \psi''' \} \\
& + a_2 [(-1 + \omega^2 - 2\varepsilon a_0 \alpha_1) (\psi')^2 + 2\varepsilon (\alpha_3 - \omega^2 \alpha_4) (\psi'')^2] = 0. \quad (3.32)
\end{aligned}$$

$$-2\varepsilon a_1 (a_2 \alpha_1 - \alpha_3 + \omega^2 \alpha_4) (\psi')^3 - 2\varepsilon a_2 \{ \omega \alpha_2 \psi' + 5(\alpha_3 - \omega^2 \alpha_4) \psi'' \} (\psi')^2 = 0. \quad (3.33)$$

$$-\varepsilon a_2 (a_2 \alpha_1 - 6\alpha_3 + 6\omega^2 \alpha_4) (\psi')^4 = 0. \quad (3.34)$$

173 From Eqs. (3.30) and (3.34), we obtain

$$a_0 = 0, \quad \frac{-1 + \omega^2}{\varepsilon \alpha_1} \quad \text{and} \quad a_2 = \frac{6(\alpha_3 - \omega^2 \alpha_4)}{\alpha_1}, \quad \text{since } a_2 \neq 0.$$

174 Therefore, depending on the values of a_0 , the following different cases arise:

175 **Case 1:** When $a_0 = 0$, from Eqs. (3.31) - (3.33), we get

$$\begin{aligned}
\psi(\xi) &= c_2 + \frac{30c_1(\alpha_3 - \omega^2 \alpha_4)}{-5a_1 \alpha_1 - 6\omega \alpha_2} e^{\frac{\xi(-5a_1 \alpha_1 - 6\omega \alpha_2)}{30(\alpha_3 - \omega^2 \alpha_4)}}, \\
a_1 &= 0, \quad \omega = \pm \frac{\sqrt{\frac{6\varepsilon \alpha_2^2 - 25(\alpha_3 + \alpha_4) + \sqrt{\{6\varepsilon \alpha_2^2 - 25(\alpha_3 + \alpha_4)\}^2 - 2500\alpha_3 \alpha_4}}{-\alpha_4}}}{5\sqrt{2}} = \pm \theta,
\end{aligned}$$

176 and

$$\begin{aligned}
a_1 &= \frac{3 \left[3\varepsilon \omega \alpha_1 \alpha_2 + 5\sqrt{\varepsilon \alpha_1^2 \{ \varepsilon \omega^2 \alpha_2^2 + 4(-1 + \omega^2)(-\alpha_3 + \omega^2 \alpha_4) \}} \right]}{5\varepsilon \alpha_1^2}, \\
\omega &= -\frac{\sqrt{25 + \frac{6\varepsilon \alpha_2^2}{\alpha_4} + \frac{25\alpha_3}{\alpha_4} \pm \frac{\sqrt{(-6\varepsilon \alpha_2^2 - 25\alpha_3 - 25\alpha_4)^2 - 2500\alpha_3 \alpha_4}}{\alpha_4}}}{5\sqrt{2}},
\end{aligned}$$

177 where c_1 and c_2 are integration constants.

178 Hence for the values of a_1 and ω , there also arise three cases. But when $a_1 \neq 0$ then the
179 shape of the solutions for dissipative case is distorted and the solution size is very long. So
180 we have omitted the other value of a_1 and discussed only for $a_1 = 0$.

181 When $a_1 = 0$ then we get also the solutions to the above mentioned equation depends for
182 the values of ω . Thus,

$$\psi(\xi) = c_2 - \frac{5c_1(\alpha_3 - \omega^2 \alpha_4)}{\omega \alpha_2} e^{-\frac{\xi \omega \alpha_2}{5(\alpha_3 - \omega^2 \alpha_4)}}$$

E-mail address: ali_math74@yahoo.com.

183 Now, by means of the values of a_0 , a_1 , a_2 and $\psi(\xi)$ from Eq. (3.7), we achieve the
 184 subsequent solution:

$$U(\xi) = - \frac{6\omega^2 c_1^2 \alpha_2^2 (-\alpha_3 + \omega^2 \alpha_4)}{\alpha_1 \left\{ \omega c_2 \alpha_2 e^{\frac{\xi \omega \alpha_2}{5\alpha_3 - 5\omega^2 \alpha_4}} - 5c_1 (\alpha_3 - \omega^2 \alpha_4) \right\}^2}. \quad (3.35)$$

185 Simplifying the required solution (3.35), we derive the following close-form solution of the
 186 strain wave equation in microstructured solids for dissipative case (3.27):

$$u(x, t) = [6\omega^2 \{-\cosh(2\sigma(x - t\omega)) + \sinh(2\sigma(x - t\omega))\} c_1^2 \alpha_2^2 (-\alpha_3 + \omega^2 \alpha_4)] \\ / \left(\alpha_1 [\omega \{\cosh(\sigma(x - t\omega)) + \sinh(\sigma(x - t\omega))\} c_2 \alpha_2 \right. \\ \left. + 5\{-\cosh(\sigma(x - t\omega)) + \sinh(\sigma(x - t\omega))\} c_1 (\alpha_3 - \omega^2 \alpha_4)]^2 \right). \quad (3.36)$$

187 where $\sigma = \frac{\omega \alpha_2}{10(\alpha_3 - \omega^2 \alpha_4)}$, $\omega = \pm \theta$ or and c_1 , c_2 are integrating constants. Since c_1 and c_2 are
 188 integration constants, one might arbitrarily select their values. If we choose $c_1 = \alpha_2 \omega$ and
 189 $c_2 = -5(\alpha_3 - \omega^2 \alpha_4)$, then from (3.36), we obtain

$$u_9(x, t) = \frac{3\omega^2 \alpha_2^2}{50\alpha_1 (\alpha_3 - \omega^2 \alpha_4)} \left\{ 1 + \tanh \left(\frac{\omega(-x + t\omega)\alpha_2}{10(\alpha_3 - \omega^2 \alpha_4)} \right) \right\}^2. \quad (3.37)$$

190 Again if we choose $c_1 = \alpha_2 \omega$ and $c_2 = 5(\alpha_3 - \omega^2 \alpha_4)$, then from (3.36), we attain the
 191 subsequent soliton solution:

$$u_{10}(x, t) = \frac{3\omega^2 \alpha_2^2}{50\alpha_1 (\alpha_3 - \omega^2 \alpha_4)} \left\{ 1 + \coth \left(\frac{\omega(-x + t\omega)\alpha_2}{10(\alpha_3 - \omega^2 \alpha_4)} \right) \right\}^2. \quad (3.38)$$

192 **Case 2:** When $a_0 = \frac{-1 + \omega^2}{\varepsilon \alpha_1}$, from Eq.(3.31)-(3.33), we obtain

$$\psi(\xi) = c_2 + \frac{30c_1(\alpha_3 - \omega^2 \alpha_4)}{-5a_1 \alpha_1 - 6\omega \alpha_2} e^{\frac{\xi(-5a_1 \alpha_1 - 6\omega \alpha_2)}{30(\alpha_3 - \omega^2 \alpha_4)}},$$

193 where c_1 and c_2 are integration constants and

$$\left\{ a_1 = 0, \omega = \left[\begin{array}{l} \pm \frac{\sqrt{\frac{6\varepsilon \alpha_2^2 + 25\alpha_3 + 25\alpha_4 - \sqrt{\{6\varepsilon \alpha_2^2 + 25(\alpha_3 + \alpha_4)\}^2 - 2500\alpha_3 \alpha_4}}{\alpha_4}}}{5\sqrt{2}} = \pm \vartheta_1 (\text{say}) \\ \pm \frac{\sqrt{\frac{6\varepsilon \alpha_2^2 + 25\alpha_3 + 25\alpha_4 + \sqrt{\{6\varepsilon \alpha_2^2 + 25(\alpha_3 + \alpha_4)\}^2 - 2500\alpha_3 \alpha_4}}{\alpha_4}}}{5\sqrt{2}} = \pm \vartheta_2 (\text{say}) \end{array} \right] \right\};$$

E-mail address: ali_math74@yahoo.com.

$$\left\{ \begin{aligned} a_1 &= \frac{3 \left[3\varepsilon\omega\alpha_1\alpha_2 + 5\sqrt{\varepsilon\alpha_1^2 \{ \varepsilon\omega^2\alpha_2^2 + 4(-1 + \omega^2)(\alpha_3 - \omega^2\alpha_4) \}} \right]}{5\varepsilon\alpha_1^2}, \\ \omega &= -\sqrt{\frac{-6\varepsilon\alpha_2^2 + 25\alpha_3 + 25\alpha_4 \pm \sqrt{\{6\varepsilon\alpha_2^2 - 25(\alpha_3 + \alpha_4)\}^2 - 2500\alpha_3\alpha_4}}{\alpha_4}} \Bigg/ 5\sqrt{2} \end{aligned} \right\};$$

$$\left\{ \begin{aligned} a_1 &= \frac{3 \left[3\varepsilon\omega\alpha_1\alpha_2 - 5\sqrt{\varepsilon\alpha_1^2 \{ \varepsilon\omega^2\alpha_2^2 + 4(-1 + \omega^2)(\alpha_3 - \omega^2\alpha_4) \}} \right]}{5\varepsilon\alpha_1^2}, \\ \omega &= \sqrt{\frac{-6\varepsilon\alpha_2^2 + 25\alpha_3 + 25\alpha_4 \pm \sqrt{\{6\varepsilon\alpha_2^2 - 25(\alpha_3 + \alpha_4)\}^2 - 2500\alpha_3\alpha_4}}{\alpha_4}} \Bigg/ 5\sqrt{2} \end{aligned} \right\}.$$

194 Hence for the values of a_1 and ω , there arises also three cases. When $a_1 \neq 0$, then the form
 195 of solutions to the strain wave equation in microstructured solids for dissipative case (3.24)
 196 indistinct and the solution size is very lengthy. So we omitted the extra value of a_1 and only
 197 discuss for $a_1 = 0$.

198 When $a_1 = 0$ then we find also the solutions to the above revealed equation depends for the
 199 values of ω , i.e. $\omega = \pm\vartheta_1$ and $\omega = \pm\vartheta_2$. Therefore,

$$\psi(\xi) = c_2 - \frac{5c_1(\alpha_3 - \omega^2\alpha_4)}{\omega\alpha_2} e^{-\frac{\xi\omega\alpha_2}{5(\alpha_3 - \omega^2\alpha_4)}}$$

200 where . $\omega = \pm\vartheta_1$ or $\omega = \pm\vartheta_2$, c_1 and c_2 are constants of integration.

201 Substituting the values of a_0 , a_1 , a_2 and $\psi(\xi)$ into Eq. (3.7), we accomplish the following
 202 solution:

$$U(\xi) = \frac{-1 + \omega^2}{\varepsilon \alpha_1} - \frac{6\omega^2 c_1^2 \alpha_2^2 (-\alpha_3 + \omega^2 \alpha_4)}{\alpha_1 \left\{ \omega c_2 \alpha_2 e^{\frac{\xi \omega \alpha_2}{5\alpha_3 - 5\omega^2 \alpha_4}} - 5c_1 (\alpha_3 - \omega^2 \alpha_4) \right\}^2}. \quad (3.39)$$

203 Simplifying the required exponential function solution (3.39) into trigonometric function
 204 solution, we derive the solution of Eq. (3.27) as follows:

$$\begin{aligned} u(x, t) = & \left[\omega^2 (-1 + \omega^2) \{ \cosh(2\varphi(x - t\omega)) + \sinh(2\varphi(x - t\omega)) \} c_2^2 \alpha_2^2 \right. \\ & + \{ \cosh(2\varphi(x - t\omega)) - \sinh(2\varphi(x - t\omega)) \} c_1^2 (\alpha_3 - \omega^2 \alpha_4) \{ 6\varepsilon \omega^2 \alpha_2^2 \\ & - 25(-1 + \omega^2)(-\alpha_3 + \omega^2 \alpha_4) \} + 10\omega(-1 + \omega^2) c_1 c_2 \alpha_2 (-\alpha_3 + \omega^2 \alpha_4)] \\ & / \left(\varepsilon \alpha_1 [\omega \{ \cosh(\varphi(x - t\omega)) + \sinh(\varphi(x - t\omega)) \} c_2 \alpha_2 \right. \\ & \left. \left. + 5 \{ -\cosh(\varphi(x - t\omega)) + \sinh(\varphi(x - t\omega)) \} c_1 (\alpha_3 - \omega^2 \alpha_4) \right]^2 \right). \quad (3.40) \end{aligned}$$

205 Therefore, we obtain the generalized soliton solution (3.40) to the strain wave equation in
 206 microstructured solids for dissipative case, where $\varphi = \frac{\omega \alpha_2}{10(\alpha_3 - \omega^2 \alpha_4)}$ and $\omega = \pm \vartheta_1$ or $\omega = \pm \vartheta_2$.
 207 But, since c_1 and c_2 are arbitrary constants, someone may arbitrarily choose their values.
 208 So, if we choose $c_1 = \alpha_2 \omega$ and $c_2 = 5(\alpha_3 - \omega^2 \alpha_4)$, from (3.20), we get the subsequent
 209 soliton solutions:

$$u_{11}(x, t) = \frac{(-1 + \omega^2)}{\alpha_1 \varepsilon} - \frac{3\omega^2 \alpha_2^2}{50\alpha_1 (-\alpha_3 + \omega^2 \alpha_4)} \left\{ -1 + \coth \left(\frac{\omega(x - t\omega) \alpha_2}{10(\alpha_3 - \omega^2 \alpha_4)} \right) \right\}^2. \quad (3.41)$$

210 Again, if we choose $c_1 = \alpha_2 \omega$ and $c_2 = -5(\alpha_3 - \omega^2 \alpha_4)$, the solitary wave solution (3.40)
 211 becomes

$$u_{12}(x, t) = \frac{(-1 + \omega^2)}{\varepsilon \alpha_1} + \frac{3\varepsilon \omega^2 \alpha_2^2}{50\varepsilon \alpha_1 (\alpha_3 - \omega^2 \alpha_4)} \left\{ -1 + \tanh \left(\frac{\omega(x - t\omega) \alpha_2}{10(\alpha_3 - \omega^2 \alpha_4)} \right) \right\}^2. \quad (3.42)$$

212 As c_1 and c_2 are arbitrary constants, one may pick many other values of them and each of
 213 this selection construct new solution. But for minimalism, we have not recorded these
 214 solutions.

E-mail address: ali_math74@yahoo.com.

215 **Remark 2:** The solutions (3.37)-(3.38), where $\omega = \pm\theta_1$ or $\omega = \pm\theta_2$ and the solutions (3.41)-
 216 (3.42) $\omega = \pm\vartheta_1$ or $\omega = \pm\vartheta_2$ have been confirmed by satisfying the original equation.

217 4. PHYSICAL INTERPRETATIONS OF THE SOLUTIONS

218 In this sub-section, we draw the graph of the derived solutions and explain the effect of the
 219 parameters on the solutions for both non-dissipative and dissipative cases. The solution u_1
 220 in (3.17) depends on the physical parameters $\alpha_1, \alpha_3, \alpha_4, \varepsilon$ and the group velocity ω . Now,
 221 we will discuss all the possible physical significances for $-2 \leq \alpha_1, \alpha_3, \alpha_4, \varepsilon \leq 2$, and soliton
 222 exists for $|\omega| > 1$ and $|\omega| < 1$. For the value of parameters $\alpha_1, \alpha_3, \alpha_4, \varepsilon < 0$ and $|\omega| > 1$, the
 223 solution u_1 in (3.17) represents the bell shape soliton and when $|\omega| < 1$ then the solution u_1
 224 represents the anti-bell shape soliton. It is shown in Fig. 1. Also if the values of the
 225 parameters are $\alpha_1 > 0, \alpha_3, \alpha_4, \varepsilon < 0$ and $|\omega| > 1$, then the solution u_1 represents the anti-
 226 bell shape soliton and when $|\omega| < 1$, then the solution u_1 represents the bell shape soliton. It
 227 is shown the Fig. 2. Again, for $\alpha_1, \alpha_3, \alpha_4 < 0, \varepsilon > 0$ and $|\omega| < 1$, the solution u_1 in (3.17)
 228 represents the multi-soliton and when $|\omega| > 1$, the solution u_1 represents the anti-bell shape
 229 soliton. It is plotted in Fig. 3. Again, if the values of the physical parameters are
 230 $\alpha_1 > 0, \alpha_3, \alpha_4 < 0, \varepsilon > 0$ and $|\omega| > 1$, then the solution u_1 represents the anti-bell shape
 231 soliton and when $|\omega| < 1$ then the solution u_1 represents the bell shape soliton. It is shown in
 232 Fig. 4. We can sketch the other figures of the solution u_1 for different values of the
 233 parameters. But for page limitation in this article we have omitted these figures. So, for other
 234 cases we do not draw the figures but we discuss for other cases with the following table:

$\varepsilon > 0$	$ \omega > 1$	$\alpha_1 < 0, \alpha_3 < 0, \alpha_4 < 0$	Anti-bell shape soliton
		$\alpha_1 > 0, \alpha_3 < 0, \alpha_4 < 0$	Bell shape soliton
		$\alpha_1 > 0, \alpha_3 > 0, \alpha_4 < 0$	Bell shape soliton
		$\alpha_1 > 0, \alpha_3 > 0, \alpha_4 > 0$	Bell shape soliton
		$\alpha_1 > 0, \alpha_3 < 0, \alpha_4 > 0$	Bell shape soliton

E-mail address: ali_math74@yahoo.com.

		$\alpha_1 < 0, \alpha_3 > 0, \alpha_4 < 0$	Anti-bell shape soliton
		$\alpha_1 < 0, \alpha_3 > 0, \alpha_4 > 0$	Anti-bell shape soliton
		$\alpha_1 < 0, \alpha_3 < 0, \alpha_4 > 0$	Anti-bell shape soliton
	$ \omega < 1$	$\alpha_1 < 0, \alpha_3 < 0, \alpha_4 < 0$	Bell shape soliton
		$\alpha_1 > 0, \alpha_3 < 0, \alpha_4 < 0$	Anti-bell shape soliton
		$\alpha_1 > 0, \alpha_3 > 0, \alpha_4 < 0$	Anti-bell shape soliton
		$\alpha_1 > 0, \alpha_3 > 0, \alpha_4 > 0$	Anti-bell shape soliton
		$\alpha_1 > 0, \alpha_3 < 0, \alpha_4 > 0$	Anti-bell shape soliton
		$\alpha_1 < 0, \alpha_3 > 0, \alpha_4 < 0$	Bell shape soliton
		$\alpha_1 < 0, \alpha_3 > 0, \alpha_4 > 0$	Bell shape soliton
		$\alpha_1 < 0, \alpha_3 < 0, \alpha_4 > 0$	Periodic bell shape solution
$\varepsilon < 0$	$ \omega > 1$	$\alpha_1 < 0, \alpha_3 < 0, \alpha_4 < 0$	Bell shape or Periodic bell shape solution
		$\alpha_1 > 0, \alpha_3 < 0, \alpha_4 < 0$	Anti-bell shape soliton or Periodic anti-bell shape solution
		$\alpha_1 > 0, \alpha_3 > 0, \alpha_4 < 0$	Anti-bell shape soliton
		$\alpha_1 > 0, \alpha_3 > 0, \alpha_4 > 0$	Periodic anti-bell shape solution
		$\alpha_1 > 0, \alpha_3 < 0, \alpha_4 > 0$	Periodic anti-bell shape solution
		$\alpha_1 < 0, \alpha_3 > 0, \alpha_4 < 0$	Bell shape soliton
		$\alpha_1 < 0, \alpha_3 > 0, \alpha_4 > 0$	Periodic bell shape solution
		$\alpha_1 < 0, \alpha_3 < 0, \alpha_4 > 0$	Periodic bell shape solution
	$ \omega < 1$	$\alpha_1 < 0, \alpha_3 < 0, \alpha_4 < 0$	Anti-bell shape soliton or Periodic anti-bell shape solution
		$\alpha_1 > 0, \alpha_3 < 0, \alpha_4 < 0$	Bell shape or Periodic bell shape solution
		$\alpha_1 > 0, \alpha_3 > 0, \alpha_4 < 0$	Periodic bell shape solution
		$\alpha_1 > 0, \alpha_3 > 0, \alpha_4 > 0$	Bell shape or Periodic bell shape solution
		$\alpha_1 > 0, \alpha_3 < 0, \alpha_4 > 0$	Bell shape soliton
		$\alpha_1 < 0, \alpha_3 > 0, \alpha_4 < 0$	Periodic anti-bell shape solution
		$\alpha_1 < 0, \alpha_3 > 0, \alpha_4 > 0$	Anti-bell shape soliton or Periodic anti-bell shape solution
		$\alpha_1 < 0, \alpha_3 < 0, \alpha_4 > 0$	Anti-bell shape soliton

235

236

237

238

239

240

Also the soliton u_2 in (3.18) depends on the parameters $\alpha_1, \alpha_3, \alpha_4, \varepsilon$ and ω . Now, we will discuss all the possible physical significances for $-2 \leq \alpha_1, \alpha_3, \alpha_4, \varepsilon \leq 2$, and soliton exists for $|\omega| > 1$ and $|\omega| < 1$. For the value of parameters contains $\alpha_1, \alpha_3, \alpha_4, \varepsilon > 0$ and $|\omega| > 1$, then the solution u_2 in (3.18) represents the singular anti-bell shape soliton and when $|\omega| < 1$ then the solution u_2 represents the singular bell shape soliton. It is shown in Fig. 5. Also, for

E-mail address: ali_math74@yahoo.com.

241 $\alpha_1, \alpha_3, \alpha_4 < 0, \varepsilon > 0$ and $|\omega| > 1$, then the solution u_2 in (3.18) represents the periodic
 242 singular anti-bell shape solution and when $|\omega| < 1$ then the solution u_2 represents the
 243 periodic singular bell shape solution. It is plotted of the Fig. 6. On the other hand, the
 244 solutions u_3 in (3.19) and u_4 in (3.20) exist for $(\alpha_3 - \alpha_4 \omega^2) > 0, \varepsilon < 0$ or
 245 $(\alpha_3 - \alpha_4 \omega^2) < 0, \varepsilon > 0$ when $|\omega| > 1$ or $|\omega| < 1$. For the value of the parameters are
 246 $\alpha_1 = -1.25, \alpha_3 = -0.1, \alpha_4 = -2, \varepsilon = -1$, when $\omega = 0.96$, the solution u_3 in (3.19) represents
 247 the anti-bell shape soliton and $\alpha_1 = -1.5, \alpha_3 = -0.1, \alpha_4 = 2, \varepsilon = -1$, when $\omega = 1.5$, the
 248 solution u_4 represents the periodic solution. It is shown in Fig. 7. Again, the travelling wave
 249 solution u_5 in (3.23) represents the bell shape singular solitons for $\alpha_1 = -1 = \alpha_3, \alpha_4 = 1$,
 250 $\varepsilon = 0.5, \omega = -1.5$ and $\omega = 0.5$ respectively, in Fig. 8 and Fig. 9 from u_6 in (3.24)
 251 represents the bell shape soliton, when $\omega = 1.5$ and the anti-bell shape soliton, when
 252 $\omega = -0.75$. In Fig. 10, we have plotted of the periodic bell shape and anti-bell shape solution
 253 for $\alpha_1 = \alpha_3 = -1.25, \alpha_4 = 1, \varepsilon = 0.7, \omega = -1.2$ and $\alpha_1 = \alpha_3 = -1.25, \alpha_4 = 1, \varepsilon = -0.7$,
 254 $\omega = 0.25$ respectively to the solution of u_7 in (3.25) and Fig. 11 plotted the periodic anti-bell
 255 shape solution and bell shape solution for $\alpha_1 = 1.25, \alpha_3 = -1.25, \alpha_4 = 1, \varepsilon = 0.7, \omega = -1.2$
 256 and $\alpha_1 = \alpha_3 = -1.25, \alpha_4 = 1, \varepsilon = -0.7, \omega = -0.25$ respectively to the solution of u_8 in
 257 (3.26). Fig. 12 and 13 represent the kink shape solutions u_9 given in (3.37) are respectively,
 258 for $\alpha_1 = 1, \alpha_2 = 1, \alpha_3 = -1.5, \alpha_4 = -1$ and $\alpha_1 = -1, \alpha_2 = 1, \alpha_3 = -1.5, \alpha_4 = -1$
 259 respectively, when $\omega = \pm\mu_1$ and for $\alpha_1 = 1, \alpha_2 = 1, \alpha_3 = -1.5, \alpha_4 = -1$ and $\alpha_1 = -1$,
 260 $\alpha_2 = 1, \alpha_3 = -1.5, \alpha_4 = -1$ respectively, when $\omega = \pm\mu_2$. Also sketch the figures 14 and 15,
 261 singular bell shape solutions u_{10} in (3.38) for $\alpha_1 = 1, \alpha_2 = 1, \alpha_3 = -1.5, \alpha_4 = -1$ and
 262 $\alpha_1 = -1, \alpha_2 = 1, \alpha_3 = -1.5, \alpha_4 = -1$ respectively, when $\omega = \pm\mu_1$ and for $\alpha_1 = 1, \alpha_2 = 1$,
 263 $\alpha_3 = -1.5, \alpha_4 = -1$ and $\alpha_1 = -1, \alpha_2 = 1, \alpha_3 = -1.5, \alpha_4 = -1$ respectively, when $\omega = \pm\mu_2$.
 264 On the other hand, Fig. 16 and 17 are singular bell and singular anti-bell shape soliton
 E-mail address: ali_math74@yahoo.com.

265 solitons u_{11} in (3.41) for $\alpha_1=1, \alpha_2=1, \alpha_3=1, \alpha_4=1, \varepsilon=0.5$ and $\alpha_1=-1, \alpha_2=1, \alpha_3=1,$
 266 $\alpha_4=1, \varepsilon=0.5$ respectively, when $\omega=\pm\theta_1$ and for $\alpha_1=1, \alpha_2=1, \alpha_3=1, \alpha_4=1, \varepsilon=0.5$
 267 and $\alpha_1=-1, \alpha_2=1, \alpha_3=1, \alpha_4=1, \varepsilon=0.5$ respectively, when $\omega=\pm\theta_2$. Also, draw the
 268 Figures 18 and 19 are kink shape solitons u_{12} in (3.42) for $\alpha_1=1, \alpha_2=1, \alpha_3=1, \alpha_4=1,$
 269 $\varepsilon=0.5$ and $\alpha_1=-1, \alpha_2=1, \alpha_3=1, \alpha_4=1, \varepsilon=0.5$ respectively, when $\omega=\pm\theta_1$ and for
 270 $\alpha_1=1, \alpha_2=1, \alpha_3=1, \alpha_4=1, \varepsilon=0.5$ and $\alpha_1=-1, \alpha_2=1, \alpha_3=1, \alpha_4=1, \varepsilon=0.5$
 271 respectively, when $\omega=\pm\theta_2$. All figures are drawn within $-10 \leq x, t \leq 10$.

272 There is another kind of solution which is not a kink, anti-kink, dark or bell-shape soliton,
 273 known as Love wave [58, 59]. A Love wave is define to be a surface wave having a
 274 horizontal motion that is transverse or perpendicular to the direction the wave is traveling.
 275 We can discuss the solutions u_2 to u_{12} for other values of the parameters. But for page
 276 limitation in this article we have omitted these figures in details.

277




Fig. 1: Sketch of the solution u_1 for $\alpha_1 = -0.001, \alpha_3 = \alpha_4 = \varepsilon = \omega = -1.5$ and
 $\alpha_1 = -0.001, \alpha_3 = \alpha_4 = \varepsilon = \omega = -0.75$ respectively.

278

279

280




Fig. 2: Plot of the solution u_1 for $\alpha_1 = 0.001$, $\alpha_3 = \alpha_4 = \varepsilon = \omega = -1.5$ and $\alpha_1 = 0.001$, $\alpha_3 = \alpha_4 = \varepsilon = \omega = -0.75$ respectively.




Fig. 3: Sketch of the solution u_1 for $\alpha_1 = \alpha_3 = \alpha_4 = -1.2$, $\varepsilon = \omega = 0.5$ and $\alpha_1 = \alpha_3 = \alpha_4 = -1.2$, $\varepsilon = 0.5$, $\omega = 1.25$ respectively.




Fig. 4: Sketch of the solution u_1 for $\alpha_1 = 0.75$, $\alpha_3 = \alpha_4 = -1.2$, $\varepsilon = 0.5$, $\omega = 1.25$ and $\alpha_1 = 0.75$, $\alpha_3 = \alpha_4 = -1.2$, $\varepsilon = 0.5$, $\omega = 0.5$ respectively.




Fig. 5: Sketch of the singular dark and singular bell shape soliton u_2 for $\alpha_1 = \alpha_3 = \alpha_4 = 0.5$, $\varepsilon = 0.75$, $\omega = -1.5$ and $\alpha_1 = \alpha_3 = \alpha_4 = 0.5$, $\varepsilon = 0.75$, $\omega = -0.25$ respectively.




Fig. 6: Sketch of the periodic singular solution u_2 for $\alpha_1 = \alpha_3 = \alpha_4 = -1.5$, $\varepsilon = 0.75$, $\omega = -1.5$ and $\alpha_1 = \alpha_3 = \alpha_4 = -1.5$, $\varepsilon = 0.75$, $\omega = -0.25$ respectively.






Fig. 7: Sketch of the solution u_3 and the solution u_4 for $\alpha_1 = -1.25$, $\alpha_3 = -0.1$, $\alpha_4 = -2$, $\varepsilon = -1$, $\omega = 0.96$ and $\alpha_1 = -1.5$, $\alpha_3 = -0.1$, $\alpha_4 = 2$, $\varepsilon = -1$, $\omega = 1.5$ respectively.




306
307

Fig. 8: Sketch of the solutions u_5 for $\alpha_1 = -1 = \alpha_3$, $\alpha_4 = 1$, $\varepsilon = 0.5$, $\omega = -1.5$ and $\omega = 0.5$ respectively.



308
309
310

Fig. 9: Sketch of the bell shape soliton and anti-bell shape soliton u_6 for $\alpha_1 = \alpha_3 = \alpha_4 = -1$, $\varepsilon = 0.5$, $\omega = 1.5$ and $\omega = -0.75$ respectively.



311
312
313

Fig. 10: Sketch of the solutions u_7 for $\alpha_1 = \alpha_3 = -1.25$, $\alpha_4 = 1$, $\varepsilon = 0.7$, $\omega = -1.2$ and $\alpha_1 = \alpha_3 = -1.25$, $\alpha_4 = 1$, $\varepsilon = -0.7$, $\omega = 0.25$ respectively.




Fig. 11: Sketch of the solutions u_8 for $\alpha_1 = 1.25$, $\alpha_3 = -1.25$, $\alpha_4 = 1$, $\varepsilon = 0.7$, $\omega = -1.2$ and $\alpha_1 = \alpha_3 = -1.25$, $\alpha_4 = 1$, $\varepsilon = -0.7$, $\omega = -0.25$ respectively.




Fig. 12: Kink shape soliton obtained from u_9 for $\alpha_1 = 1$, $\alpha_2 = 1$, $\alpha_3 = -1.5$, $\alpha_4 = -1$, $\varepsilon = 0.5$ and $\alpha_1 = -1$, $\alpha_2 = 1$, $\alpha_3 = -1.5$, $\alpha_4 = -1$, $\varepsilon = 0.5$ respectively, when $\omega = \pm\mu_1$.




Fig. 13: Kink shape soliton obtained from u_9 for $\alpha_1 = 1$, $\alpha_2 = 1$, $\alpha_3 = -1.5$, $\alpha_4 = -1$, $\varepsilon = 0.5$ and $\alpha_1 = -1$, $\alpha_2 = 1$, $\alpha_3 = -1.5$, $\alpha_4 = -1$, $\varepsilon = 0.5$ respectively, when $\omega = \pm\mu_2$.




Fig. 14: Singular bell shape and anti-bell shape soliton u_{10} for $\alpha_1 = 1, \alpha_2 = 1, \alpha_3 = -1.5, \alpha_4 = -1, \varepsilon = 0.5$ and $\alpha_1 = -1, \alpha_2 = 1, \alpha_3 = -1.5, \alpha_4 = -1, \varepsilon = 0.5$ respectively, when $\omega = \pm\mu_1$.




Fig. 15: Singular anti-bell shape and bell shape soliton u_{10} in (3.38) for $\alpha_1 = 1, \alpha_2 = 1, \alpha_3 = -1.5, \alpha_4 = -1, \varepsilon = 0.5$ and $\alpha_1 = -1, \alpha_2 = 1, \alpha_3 = -1.5, \alpha_4 = -1, \varepsilon = 0.5$ respectively, when $\omega = \pm\mu_2$.




Fig. 16: Sketch the singular bell type and anti-bell soliton u_{11} for $\alpha_1 = 1, \alpha_2 = 1, \alpha_3 = 1, \alpha_4 = 1, \varepsilon = 0.5$ and $\alpha_1 = -1, \alpha_2 = 1, \alpha_3 = 1, \alpha_4 = 1, \varepsilon = 0.5$ respectively, when $\omega = \pm\theta_1$.




Fig. 17: Singular anti-bell shape and bell shape soliton u_{11} for $\alpha_1 = 1, \alpha_2 = 1, \alpha_3 = 1, \alpha_4 = 1, \varepsilon = 0.5$ and $\alpha_1 = -1, \alpha_2 = 1, \alpha_3 = 1, \alpha_4 = 1, \varepsilon = 0.5$ respectively, when $\omega = \pm\theta_2$.




Fig. 18: Kink shape soliton u_{12} for $\alpha_1 = 1, \alpha_2 = 1, \alpha_3 = 1, \alpha_4 = 1, \varepsilon = 0.5$ and $\alpha_1 = -1, \alpha_2 = 1, \alpha_3 = 1, \alpha_4 = 1, \varepsilon = 0.5$ respectively, when $\omega = \pm\theta_1$.




Fig. 19: Kink shape soliton u_{12} for $\alpha_1 = 1, \alpha_2 = 1, \alpha_3 = 1, \alpha_4 = 1, \varepsilon = 0.5$ and $\alpha_1 = -1, \alpha_2 = 1, \alpha_3 = 1, \alpha_4 = 1, \varepsilon = 0.5$ respectively, when $\omega = \pm\theta_2$.

5. CONCLUSION

E-mail address: ali_math74@yahoo.com.

343 In this article, we have implemented the MSE method to obtain soliton solutions to the strain
344 wave equation in microstructured solids for both non-dissipative and dissipative cases. In
345 fact, we have derived general solitary wave solutions to this equation associated with
346 arbitrary constants, and for particular values of these constants solitons are originated from
347 the general solitary wave solutions. We have illustrated the solitary wave properties of the
348 solutions for various values of the free parameters by means of the graphs. This work shows
349 that the MSE method is competent and more powerful and can be used for many other
350 equations NLEEs applied mathematics and engineering.

351 **ACKNOWLEDGEMENTS**

352 The authors wish to take this opportunity to express their gratitude to the referees for their
353 valuable comments and suggestions which enhanced the quality of this article.

354 **REFERENCES**

- 355 1. V.B. Matveev and M.A. Salle, Darboux transformation and solitons, Springer, Berlin,
356 1991.
- 357 2. G. Xu, An elliptic equation method and its applications in nonlinear evolution equations,
358 Chaos, Solitons Fract., 29 (2006) 942-947.
- 359 3. E. Yusufoglu and A. Bekir, Exact solution of coupled nonlinear evolution equations,
360 Chaos, solitons Fract., 37 (2008) 842-848.
- 361 4. D.D. Ganji, The application of He's homotopy perturbation method to nonlinear
362 equations arising in heat transfer, Phys. Lett. A, 355 (2006) 137-141.
- 363 5. D.D. Ganji, G.A. Afrouzi and R.A. Talarposhti, Application of variational iteration method
364 and homotopy perturbation method for nonlinear heat diffusion and heat transfer
365 equations, Phys. Lett. A, 368 (2007) 450-457.
- 366 6. W. Malfliet and W. Hereman, The tanh method II: Perturbation technique for
367 conservative systems, Phys. Scr., 54 (1996) 563-569.

E-mail address: ali_mam174@yahoo.com.

- 368 7. H.A. Nassar, M.A. Abdel-Razek and A.K. Seddeek, Expanding the tanh-function method
369 for solving nonlinear equations, *Appl. Math.*, 2 (2011) 1096-1104.
- 370 8. A.J.M. Jawad, M.D. Petkovic, P. Laketa and A. Biswas, Dynamics of shallow water
371 waves with Boussinesq equation, *Scientia Iranica, Trans. B: Mech. Engr.*, 20(1) (2013)
372 179-184.
- 373 9. M.A. Abdou, The extended tanh method and its applications for solving nonlinear
374 physical models, *Appl. Math. Comput.*, 190 (1) (2007) 988-996.
- 375 10. A.L. Guo and J. Lin, Exact solutions of (2+1)-dimensional HNLS equation, *Commun.*
376 *Theor. Phys.*, 54 (2010) 401-406.
- 377 11. S.T. Mohyud-Din, , M.A. Noor and K.I. Noor, Modified Variational Iteration Method for
378 Solving Sine-Gordon Equations, *World Appl. Sci. J.*, 6 (7) (2009) 999-1004.
- 379 12. R. Hirota, The direct method in soliton theory, Cambridge University Press, Cambridge,
380 2004.
- 381 13. C. Rogers and W.F. Shadwick, Backlund transformations and their applications, Vol. 161
382 of *Mathematics in Science and Engineering*, Academic Press, New York, USA, 1982.
- 383 14. L. Jianming, D. Jie and Y. Wenjun, Backlund transformation and new exact solutions of
384 the Sharma-Tasso-Olver equation, *Abstract and Appl. Analysis*, 2011 (2011) Article ID
385 935710, 8 pages.
- 386 15. M.J. Ablowitz and P.A. Clarkson, Soliton, nonlinear evolution equations and inverse
387 scattering, Cambridge University Press, New York, 1991.
- 388 16. A.M. Wazwaz, A sine-cosine method for handle nonlinear wave equations, *Appl. Math.*
389 *Comput. Modeling*, 40 (2004) 499-508.
- 390 17. E. Yusufoglu, and A. Bekir, Solitons and periodic solutions of coupled nonlinear
391 evolution equations by using Sine-Cosine method, *Int. J. Comput. Math.*, 83 (12) (2006)
392 915-924.

E-mail address: ali_math74@yahoo.com.

- 393 18. J. Weiss, M. Tabor and G. Carnevale, The Painlevé property for partial differential
394 equations, J. Math. Phys., 24 (1982) 522-526.
- 395 19. A.M. Wazwaz, Partial Differential equations: Method and Applications, Taylor and
396 Francis, 2002.
- 397 20. M.A. Helal and M.S. Mehana, A comparison between two different methods for solving
398 Boussinesq-Burgers equation, Chaos, Solitons Fract., 28 (2006) 320-326.
- 399 21. M. Wang, X. Li and J. Zhang, The (G'/G) -expansion method and traveling wave
400 solutions of nonlinear evolution equations in mathematical physics, Phys. Lett. A, 372
401 (2008) 417-423.
- 402 22. J. Zhang, F. Jiang and X. Zhao, An improved (G'/G) -expansion method for solving
403 nonlinear evolution equations, Inter. J. Comput. Math., 87 (8) (2010) 1716-1725.
- 404 23. J. Feng, W. Li and Q. Wan, Using (G'/G) -expansion method to seek the traveling
405 wave solution of Kolmogorov-Petrovskii-Piskunov equation, Appl. Math. Comput., 217
406 (2011) 5860-5865.
- 407 24. M.A. Akbar, N.H.M. Ali and E.M.E. Zayed, Abundant exact traveling wave solutions of
408 the generalized Bretherton equation via (G'/G) -expansion method, Commun. Theor.
409 Phys., 57 (2012) 173-178.
- 410 25. R. Abazari, The (G'/G) -expansion method for Tziteica type nonlinear evolution
411 equations, Math. Comput. Modelling, 52 (2010) 1834-1845.
- 412 26. M.A. Akbar, N.H.M. Ali and S.T. Mohyud-Din, Further exact traveling wave solutions to
413 the (2+1)-dimensional Boussinesq and Kadomtsev-Petviashvili equation, J. Comput.
414 Analysis Appl., 15 (3) (2013) 557-571.
- 415 27. N. Taghizadeh and M. Mirzazadeh, The first integral method to some complex nonlinear
416 partial differential equations, J. Comput. Appl. Math., 235 (2011) 4871-4877.

E-mail address: ali_math74@yahoo.com.

- 417 28. M.L. Wang and X.Z. Li, Extended F-expansion method and periodic wave solutions for
418 the generalized Zakharov equations, Phys. Lett. A, 343 (2005) 48-54.
- 419 29. Sirendaoreji, Auxiliary equation method and new solutions of Klein-Gordon equations,
420 Chaos, Solitons Fract., 31 (2007) 943-950.
- 421 30. H. Triki, A. Chowdhury and A. Biswas, Solitary wave and shock wave solutions of the
422 variants of Boussinesq equation, U.P.B. Sci. Bull., Series A, 75(4) (2013) 39-52.
- 423 31. H. Triki, A.H. Kara and A. Biswas, Domain walls to Boussinesq type equations in (2+1)-
424 dimensions, Indian J. Phys., 88(7) (2014) 751-755.
- 425 32. J.H. He and X.H. Wu, Exp-function method for nonlinear wave equations, Chaos,
426 Solitons Fract., 30 (2006) 700-708.
- 427 33. H. Naher, A.F. Abdullah and M.A. Akbar, New traveling wave solutions of the higher
428 dimensional nonlinear partial differential equation by the Exp-function method, J. Appl.
429 Math., 2012 (2012) Article ID 575387, 14 pages.
- 430 34. M. Wang, Solitary wave solutions for variant Boussinesq equations, Phy. Lett. A, 199
431 (1995) 169-172.
- 432 35. A.J.M. Jawad, M.D. Petkovic and A. Biswas, Modified simple equation method for
433 nonlinear evolution equations, Appl. Math. Comput., 217 (2010) 869-877.
- 434 36. E.M.E. Zayed and S.A.H. Ibrahim, Exact solutions of nonlinear evolution equations in
435 mathematical physics using the modified simple equation method, Chin. Phys. Lett.,
436 29(6) (2012) 060201.
- 437 37. K. Khan, M.A. Akbar and M.N. Alam, Traveling wave solutions of the nonlinear Drinfel'd-
438 Sokolov-Wilson equation and modified Benjamin-Bona-Mahony equations, J. Egyptian
439 Math. Soc., 21 (2013) 233-240.

- 440 38. K. Khan and M.A. Akbar, Exact and solitary wave solutions for the Tzitzeica-Dodd-
441 Bullough and the modified Boussinesq-Zakharov-Kuznetsov equations using the
442 modified simple equation method, Ain Shams Engr. J., 4 (2013) 903-909.
- 443 39. E.M.E. Zayed and S.A.H. Ibrahim, Exact Solutions of Kolmogorov-Petrovskii-Piskunov
444 Equation Using the Modified Simple Equation Method, Acta Mathematicae Applicatae
445 Sinica, 30(3) (2014) 749-754.
- 446 40. E.M.E. Zayed and A.H. Arnous, The enhanced modified simple equation method for
447 solving nonlinear evolution equations with variable coefficients, AIP Conf. Proceedings
448 of ICNAAM 2013, 1558 (2013) 1999-2005.
- 449 41. E.M.E. Zayed and A.H. Arnous, Exact Solutions for a Nonlinear Dynamical System in a
450 New Double-Chain Model of DNA Using the Modified Simple Equation Method," Inform.
451 Sci. Comp., 2013(1), Article ID ISC080713, 08 pages
- 452 42. E.M.E. Zayed, The modified simple equation method for two nonlinear PDEs with power
453 law and kerr law nonlinearity, PanAmerican Math. J., 24(1) (2014) 65-74.
- 454 43. E.M.E. Zayed, The modified simple equation method applied to nonlinear two models
455 of diffusion-reaction equations", J. Math. Res. Applications, 2(2) (2014) 5-13.
- 456 44. E.M.E. Zayed and A.H. Arnous, The modified simple equation method with applications
457 to (2+1)-dimensional systems of nonlinear evolution equations in mathematical physics,
458 Sci. Res. Essays, 8(40) (2013) 1973-1982.
- 459 45. E.M.E. Zayed, The modified simple equation method and its applications for solving
460 nonlinear evolution equations in mathematical physics, Commun. Appl. Nonlinear
461 Analysis, 20(3) (2013) 95-104.
- 462 46. E.M.E. Zayed and A.H. Arnous, Exact traveling wave solutions of nonlinear PDEs in
463 mathematical physics using the modified simple equation method, Appl. Appl. Math.: An
464 Int. J., 8(2) (2013) 553-572.

- 465 47. E.M.E. Zayed and S.A.H. Ibrahim, Modified simple equation method and its applications
466 for some nonlinear evolution equations in mathematical physics, Int. J. Computer Appl.,
467 67(6) (2013) 39-44.
- 468 48. K. Khan and M.A. Akbar, Application of $\exp(-\phi(\xi))$ -expansion method to find the exact
469 solutions of modified Benjamin-Bona-Mahony equation, World Appl. Sci. J., 24(10)
470 (2013) 1373-1377.
- 471 49. M.G. Hafez, M.N. Alam and M.A. Akbar, Traveling wave solutions for some important
472 coupled nonlinear physical models via the coupled Higgs equation and the Maccari
473 system, J. King Saud Univ.-Sci., 27(2) (2015) 105-112.
- 474 50. T.L. Bock and M.D. Kruskal, A two-parameter Miura transformation of the Benjamin-One
475 equation, Phys. Lett. A, 74 (1979) 173-176.
- 476 51. M.N. Alam, M.A. Akbar, S. T. Mohyud-Din, General traveling wave solutions of the strain
477 wave equation in microstructured solids via the new approach of generalized (G'/G) -
478 expansion method, Alexandria Engr. J., 53 (2014) 233-241.
- 479 52. F. Pastrone, P. Cermelli and A. Porubov, Nonlinear waves in 1-D solids with
480 microstructure, Mater. Phys. Mech., 7 (2004) 9-16
- 481 53. A.V. Porubov and F. Pastrone, Non-linear bell-shaped and kink-shaped strain waves in
482 microstructured solids, Int. J. Nonlinear Mech., 39(8) (2004) 1289-1299.
- 483 54. M.A. Akbar, N.H.M. Ali and E.M.E. Zayed, A generalized and improved (G'/G) -
484 expansion method for nonlinear evolution equations, Math. Prob. Engr., Vol. 2012,
485 Article ID 459879, 22 pages, DOI:10.1155/2012/459879.
- 486 55. M.A. Khan and M.A. Akbar, Exact and solitary wave solutions to the generalized fifth-
487 order KdV equation by using the modified simple equation method, Appl. Comput. Math.,
488 4(3) (2015) 122-129.

- 489 56. A.M. Samsonov, Strain Solitons and How to Construct Them, Chapman and Hall/CRC,
490 Boca Raton, Fla, USA, 2001.
- 491 57. F. Pastrone, F. Hierarchy of Nonlinear waves in complex microstructured solids, Int.
492 Con. Com. Of nonlinear waves, (2009) 5-7.
- 493 58. A. A. Zakharenko, Analytical studying the group velocity of three-partial Love (type)
494 waves in both isotropic and anisotropic media, Nondestructive Testing and Evaluation,
495 20(4), Dec. 2005, 237-254.
- 496 59. A. A. Zakharenko, Slow acoustic waves with the anti-plane polarization in layered
497 systems, Int. J. of Modern Phys. B, 24(4) (2010) 515-536.