

# The Modified Simple Equation Method and Its Application to Solve NLEEs Associated with Engineering Problem

Md. Ashrafuzzaman Khan and M. Ali Akbar

Department of Applied Mathematics, University of Rajshahi, Rajshahi-6205, Bangladesh

## ABSTRACT

The modified simple equation (MSE) method is an important mathematical tool for searching closed-form solutions to nonlinear evolution equations (NLEEs). In the present paper, by using the MSE method, we derive some impressive solitary wave solutions to NLEES via the strain wave equation in microstructured solids which is a very important equation in the field of engineering. The solutions contain some free parameters and for particular values of the parameters some known solutions are derived. The solutions exhibit necessity and reliability of the MSE method.

**Keywords:** Modified simple equation method; balance number; solitary wave solutions; strain wave equation; microstructured solids.

**Mathematics Subject Classification:** 35C07, 35C08, 35P99.

## 1. INTRODUCTION

Physical systems are in general explained with nonlinear partial differential equations. The mathematical modeling of microstructured solid materials that change over time depends closely on the study of a variety of systems of ordinary and partial differential equations. Similar models are developed in diverse fields of study, ranging from the natural and physical sciences, population ecology to economics, infectious disease epidemiology, neural

E-mail address: ali\_math74@yahoo.com.

24 networks, biology, mechanics etc. In spite of the eclectic nature of the fields wherein these  
25 models are formulated, different groups of them contribute adequate common attributes that  
26 make it possible to examine them within a unified theoretical structure. Such study is an area  
27 of functional analysis, usually called the theory of evolution equations. Therefore, the  
28 investigation of solutions to NLEEs plays a very important role to uncover the obscurity of  
29 many phenomena and processes throughout the natural sciences. However, one of the  
30 essential problems is to obtain their closed-form solutions. For that reason, diverse groups  
31 of engineers, physicists, and mathematicians have been working tirelessly to investigate  
32 closed-form solutions to NLEEs. Accordingly, in the recent years, they establish several  
33 methods to search exact solutions, for instance, the Darboux transformation method [1], the  
34 Jacobi elliptic function method [2, 3], the He's homotopy perturbation method [4, 5], the tanh-  
35 function method [6, 7], the extended tanh-function method [8, 9], the Lie group symmetry  
36 method [10], the variational iteration method [11], the Hirota's bilinear method [12], the  
37 Backlund transformation method [13, 14], the inverse scattering transformation method [15],  
38 the sine-cosine method [16, 17], the Painleve expansion method [18], the Adomian  
39 decomposition method [19, 20], the  $(G'/G)$ -expansion method [21-26], the first integration  
40 method [27], the F-expansion method [28], the auxiliary equation method [29], the ansatz  
41 method [30, 31], the Exp-function method [32, 33], the homogeneous balance method [34],  
42 the modified simple equation method [35-47], the  $\exp(-\phi(\eta))$ -expansion method [48, 49], the  
43 Miura transformation method [50], and others.

44 Microstructured materials like crystallites, alloys, ceramics, and functionally graded materials  
45 have gained broad application. The modeling of wave propagation in such materials should  
46 be able to account for various scales of microstructure [51]. In the past years, many authors  
47 have studied the strain wave equation in microstructured solids, such as, Alam et al. [51]  
48 solved this equation by using the new generalized  $(G'/G)$ -expansion method. Pastrone et al.  
49 [52], Porubov and Pastrone [53] examined bell-shaped and kink-shaped solutions of this  
50 engineering problem. Akbar et al. [54] constructed traveling wave solutions of this equation  
E-mail address: ali\_math74@yahoo.com.

51 by using the generalized and improved  $(G'/G)$ -expansion method. The above analysis  
 52 shows that several methods to achieve exact solutions to this equation have been  
 53 accomplished in the recent years. But, the equation has not been studied by means of the  
 54 MSE method. In this article, our aim is, we will apply the MSE method following the  
 55 technique derived in the Ref. [55] to examine some new and impressive solitary wave  
 56 solutions to this equation.

57 The structure of this article is as follows: In section 2, we describe the method. In section 3,  
 58 we apply the MSE method to the strain wave equation in microstructured solids. In section 4,  
 59 we provide the physical interpretations of the obtained solutions. Finally, in section 5,  
 60 conclusions are given.

## 61 2. DESCRIPTION OF THE METHOD

62 Assume the nonlinear evolution equation has the following form

$$P(u, u_t, u_x, u_y, u_z, u_{tt}, u_{xx}, u_{yy}, u_{zz}, \dots) = 0, \quad (2.1)$$

63 where  $u = u(x, y, z, t)$  is an unidentified function,  $P$  is a polynomial function in  $u =$   
 64  $u(x, y, z, t)$  and its partial derivatives, wherein nonlinear term of the highest order and the  
 65 highest order linear terms exist and subscripts indicate partial derivatives. To solve (2.1) by  
 66 using the MSE method [35-47], we need to perform the subsequent steps:

67 **Step 1:** Now, we combine the real variable  $x$  and  $t$  by a compound variable  $\xi$  as follows:

$$\xi = x + y + z \pm \omega t, \quad u(x, y, z, t) = U(\xi), \quad (2.2)$$

68 Here  $\xi$  is called the wave variable it allows us to switch Eq. (2.1) into an ordinary differential  
 69 equation (ODE):

$$Q(U, U', U'', U''', \dots) = 0, \quad (2.3)$$

70 where  $Q$  is a polynomial in  $U(\xi)$  and its derivatives, where  $U'(\xi) = \frac{dU}{d\xi}$ .

71 **Step 2:** We assume that Eq. (2.3) has the traveling wave solution in the following form,

$$U(\xi) = \sum_{i=0}^N a_i \left\{ \frac{\psi'(\xi)}{\psi(\xi)} \right\}^i, \quad (2.4)$$

E-mail address: ali\_math74@yahoo.com.

72 where  $a_i$  ( $i = 0, 1, 2, \dots, N$ ) are arbitrary constants, such that  $a_N \neq 0$ , and  $\psi(\xi)$  is an  
 73 unidentified function which is to be determined later. In  $(G'/G)$ -expansion method, Exp-  
 74 function method, tanh-function method, sine-cosine method, Jacobi elliptic function method  
 75 etc., the solutions are initiated through several auxiliary functions which are previously  
 76 known, but in the MSE method,  $\psi(\xi)$  is neither a pre-defined function nor a solution of any  
 77 pre-defined differential equation. Therefore, it is not possible to speculate from formerly,  
 78 what kind of solution can be found by this method.

79 **Step 3:** We determine the positive integer  $N$ , come out in Eq. (2.4) by taking into account  
 80 the homogeneous balance between the highest order nonlinear terms and the derivatives of  
 81 the highest order occurring in Eq. (2.3).

82 **Step 4:** We calculate the necessary derivatives  $U', U'', U'''$  etc., then insert them into Eq.  
 83 (2.3) and then taken into consideration the function  $\psi(\xi)$ . As a result of this insertion, we  
 84 obtain a polynomial in  $(\psi'(\xi)/\psi(\xi))$ . We equate all the coefficients of  $(\psi(\xi))^{-i}$ , ( $i =$   
 85  $0, 1, 2, \dots, N$ ) to this polynomial to zero. This procedure yields a system of algebraic and  
 86 differential equations whichever can be solved for getting  $a_i$  ( $i = 0, 1, 2, \dots, N$ ),  $\psi(\xi)$  and the  
 87 value of the other parameters.

### 88 3. APPLICATION OF THE METHOD

89 In this section, we will execute the application of the MSE method to extract solitary wave  
 90 solutions to the strain wave equation in microstructured solids which is a very important  
 91 equation in the field of engineering. Let us consider the strain wave equation in  
 92 microstructured solids:

$$u_{tt} - u_{xx} - \varepsilon \alpha_1 (u^2)_{xx} - \gamma \alpha_2 u_{xxt} + \delta \alpha_3 u_{xxxx} - (\delta \alpha_4 - \gamma^2 \alpha_7) u_{xxtt} \\ + \gamma \delta (\alpha_5 u_{xxxxt} + \alpha_6 u_{xxtt}) = 0. \quad (3.1)$$

93

#### 94 3.1. THE NON-DISSIPATIVE CASE

E-mail address: ali\_math74@yahoo.com.

95 The system is non-dissipative, if  $\gamma = 0$  and determined by the double dispersive equation  
 96 (see [52], [53], [56], [57] for details)

$$u_{tt} - u_{xx} - \varepsilon \alpha_1 (u^2)_{xx} + \delta \alpha_3 u_{xxxx} - \delta \alpha_4 u_{xxtt} = 0. \quad (3.2)$$

97 The balance between dispersion and nonlinearities happen when  $\delta = O(\varepsilon)$  Therefore, (3.2)  
 98 becomes

$$u_{tt} - u_{xx} - \varepsilon \{ \alpha_1 (u^2)_{xx} - \alpha_3 u_{xxxx} + \alpha_4 u_{xxtt} \} = 0. \quad (3.3)$$

99 In order to extract solitary wave solutions of the strain wave equation in microstructured  
 100 solids by using the MSE method, we use the traveling wave variable

$$u(x, t) = U(\xi), \quad \xi = x - \omega t. \quad (3.4)$$

101 The wave transformation (3.4) reduces Eq. (3.3) into the ODE in the following form:

$$(\omega^2 - 1) U'' - \varepsilon \{ \alpha_1 (U^2)'' - (\alpha_3 - \omega^2 \alpha_4) U^{(iv)} \} = 0. \quad (3.5)$$

102 where primes indicate differential coefficients with respect to  $\xi$ . Eq. (3.5) is integrable,  
 103 therefore, integration (3.5) as many time as possible, we obtain the following ODE:

$$(\omega^2 - 1) U - \varepsilon \{ \alpha_1 U^2 - (\alpha_3 - \omega^2 \alpha_4) U'' \} = 0. \quad (3.6)$$

104 where the integration constants are set zero, as we are seeking solitary wave solutions.  
 105 Taking homogeneous balance between the terms  $U''$  and  $U^2$  appearing in Eq. (3.6), we  
 106 obtain  $N = 2$ . Therefore, the shape of the solution of Eq. (3.6) becomes

$$U(\xi) = a_0 + \frac{a_1 \psi'}{\psi} + \frac{a_2 (\psi')^2}{\psi^2}. \quad (3.7)$$

107 wherein  $a_0$ ,  $a_1$  and  $a_2$  are constants to be find out afterward such that  $a_2 \neq 0$ , and  $\psi(\xi)$  is an  
 108 unknown function. The derivatives of  $U$  are given in the following:

$$U' = -\frac{a_1 (\psi')^2}{\psi^2} - \frac{2a_2 (\psi')^3}{\psi^3} + \frac{a_1 \psi''}{\psi} + \frac{2a_2 \psi' \psi''}{\psi^2}. \quad (3.8)$$

$$U'' = a_1 \left\{ \frac{2(\psi')^3}{\psi^3} - \frac{3\psi' \psi''}{\psi^2} + \frac{\psi'''}{\psi} \right\} + 2a_2 \left\{ \frac{(\psi'')^2}{\psi^2} + \frac{\psi' \psi'''}{\psi^2} - \frac{5(\psi')^2 \psi''}{\psi^3} + \frac{3(\psi')^4}{\psi^4} \right\}. \quad (3.9)$$

E-mail address: ali\_math74@yahoo.com.

109 Inserting the values of  $U$ ,  $U'$  and  $U''$  into Eq. (3.6), and setting each coefficient of  $\psi^{-i}$ ,  $i =$   
 110  $0, 1, 2, \dots$  to zero, we derive, successively

$$a_0(-1 + \omega^2 - \varepsilon a_0 \alpha_1) = 0. \quad (3.10)$$

$$a_1\{(-1 + \omega^2 - 2\varepsilon a_0 \alpha_1)\psi' + \varepsilon(\alpha_3 - \omega^2 \alpha_4)\psi'''\} = 0. \quad (3.11)$$

$$\begin{aligned} & -\varepsilon a_1 \psi' \{a_1 \alpha_1 \psi' + 3(\alpha_3 - \omega^2 \alpha_4)\psi''\} + 2a_2 \varepsilon (\alpha_3 - \omega^2 \alpha_4) \psi' \psi''' \\ & + a_2 \{(-1 + \omega^2 - 2\varepsilon a_0 \alpha_1)(\psi')^2 + 2\varepsilon(\alpha_3 - \omega^2 \alpha_4)(\psi'')^2\} = 0. \end{aligned} \quad (3.12)$$

$$-2\varepsilon(\psi')^2 \{a_1(a_2 \alpha_1 - \alpha_3 + \omega^2 \alpha_4)\psi' + 5a_2(\alpha_3 - \omega^2 \alpha_4)\psi''\} = 0. \quad (3.13)$$

$$-\varepsilon a_2(a_2 \alpha_1 - 6\alpha_3 + 6\omega^2 \alpha_4)(\psi')^4 = 0. \quad (3.14)$$

111 From Eq. (3.10) and Eq. (3.14), we obtain

$$a_0 = 0, \quad \frac{-1 + \omega^2}{\varepsilon \alpha_1} \quad \text{and} \quad a_2 = \frac{6(\alpha_3 - \omega^2 \alpha_4)}{\alpha_1}, \quad \text{since } a_2 \neq 0.$$

112 Therefore, for the values of  $a_0$ , there arise the following cases:

113 **Case 1:** When  $a_0 = 0$ , from Eqs. (3.11)-(3.13), we obtain

$$a_1 = \pm \frac{6\sqrt{1 - \omega^2} \sqrt{\alpha_3 - \omega^2 \alpha_4}}{\sqrt{\varepsilon} \alpha_1}$$

114 and

$$\psi(\xi) = c_2 + \frac{\varepsilon c_1(-\alpha_3 + \omega^2 \alpha_4)}{-1 + \omega^2} e^{\mp \frac{\xi \sqrt{1 - \omega^2}}{\sqrt{\varepsilon} \sqrt{\alpha_3 - \omega^2 \alpha_4}}},$$

115 where  $c_1$  and  $c_2$  are integration constants.

116 Substituting the values of  $a_0$ ,  $a_1$ ,  $a_2$  and  $\psi(\xi)$  into Eq. (3.7), we obtain the following

117 exponential form solution:

$$U(\xi) = \frac{6e^{\pm \frac{\xi \sqrt{1 - \omega^2}}{\sqrt{\varepsilon} \sqrt{\alpha_3 - \omega^2 \alpha_4}}} (-1 + \omega^2)^2 c_1 c_2 (-\alpha_3 + \omega^2 \alpha_4)}{\alpha_1 \left( (-1 + \omega^2) c_2 e^{\pm \frac{i \xi \sqrt{-1 + \omega^2}}{\sqrt{\varepsilon} \sqrt{\alpha_3 - \omega^2 \alpha_4}}} + \varepsilon c_1 (-\alpha_3 + \omega^2 \alpha_4) \right)^2}. \quad (3.15)$$

118 Simplifying the required solution (3.15), we derive the following close-form solution of the  
 119 strain wave equation in microstructured solids (3.3):

$$u(x, t) = \{6(-1 + \omega^2)^2 c_1 c_2 (-\alpha_3 + \omega^2 \alpha_4)\} \\ / \left[ \alpha_1 \left\{ \pm i \sin((x - t\omega)\beta) \{(-1 + \omega^2)c_2 + \varepsilon c_1 (\alpha_3 - \omega^2 \alpha_4)\} \right. \right. \\ \left. \left. + \cos((x - t\omega)\beta) \{(-1 + \omega^2)c_2 + \varepsilon c_1 (-\alpha_3 + \omega^2 \alpha_4)\} \right\}^2 \right] \quad (3.16)$$

120 where  $\beta = \frac{\sqrt{-1+\omega^2}}{2\sqrt{\varepsilon}\sqrt{\alpha_3-\omega^2\alpha_4}}$ . Solution (3.16) is the generalized solitary wave solution of the strain  
 121 wave equation in microstructured solids. Since  $c_1$  and  $c_2$  are arbitrary constants, one might  
 122 arbitrarily choose their values. Therefore, if we choose  $c_1 = (-1 + \omega^2)$  and  $c_2 = \varepsilon(-\alpha_3 +$   
 123  $\omega^2 \alpha_4)$  then from (3.16), we obtain the following bell shaped soliton solution:

$$u_1(x, t) = \frac{3(-1 + \omega^2)}{2\varepsilon\alpha_1} \operatorname{sech}^2 \left( \frac{(x - t\omega)\sqrt{-1 + \omega^2}}{2\sqrt{\varepsilon}\sqrt{-\alpha_3 + \omega^2 \alpha_4}} \right). \quad (3.17)$$

124 Again, if we choose  $c_1 = (-1 + \omega^2)$  and  $c_2 = -\varepsilon(-\alpha_3 + \omega^2 \alpha_4)$ , then from (3.16), we obtain  
 125 the following singular soliton:

$$u_2(x, t) = -\frac{3(-1 + \omega^2)}{2\varepsilon\alpha_1} \operatorname{csch}^2 \left( \frac{(x - t\omega)\sqrt{-1 + \omega^2}}{2\sqrt{\varepsilon}\sqrt{-\alpha_3 + \omega^2 \alpha_4}} \right). \quad (3.18)$$

126 On the other hand, when  $c_1 = (-1 + \omega^2)$  and  $c_2 = \pm i \varepsilon(-\alpha_3 + \omega^2 \alpha_4)$ , from solution (3.16),  
 127 we obtain the following trigonometric solution:

$$u_3(x, t) = \frac{3(-1 + \omega^2)}{2\varepsilon\alpha_1} \sec^2 \left[ \frac{1}{4} \left\{ \pi + \frac{2(x - t\omega)\sqrt{-1 + \omega^2}}{\sqrt{\varepsilon}\sqrt{\alpha_3 - \omega^2 \alpha_4}} \right\} \right]. \quad (3.19)$$

128 Again when  $c_1 = (-1 + \omega^2)$  and  $c_2 = \mp i \varepsilon(-\alpha_3 + \omega^2 \alpha_4)$ , then the generalized solitary wave  
 129 solution (3.16) can be simplified as:

$$u_4(x, t) = \frac{3(-1 + \omega^2)}{2\varepsilon\alpha_1} \csc^2 \left[ \frac{1}{4} \left\{ \pi + \frac{2(-x + t\omega)\sqrt{-1 + \omega^2}}{\sqrt{\varepsilon}\sqrt{\alpha_3 - \omega^2 \alpha_4}} \right\} \right]. \quad (3.20)$$

130 If we choose more different values of  $c_1$  and  $c_2$ , we may derive a lot of general solitary wave  
 131 solutions to the Eq. (3.3) through the MSE method. For succinctness, other solutions have  
 132 been overlooked.

E-mail address: ali\_math74@yahoo.com.

133 **Case 2:** When  $a_0 = \frac{-1 + \omega^2}{\varepsilon \alpha_1}$ , then Eqs. (3.11)-(3.13) yield

$$a_1 = \pm \frac{6\sqrt{-1 + \omega^2}\sqrt{\alpha_3 - \omega^2\alpha_4}}{\sqrt{\varepsilon}\alpha_1}$$

134 And

$$\psi(\xi) = c_2 + \frac{\varepsilon c_1(\alpha_3 - \omega^2\alpha_4)}{-1 + \omega^2} e^{\mp \frac{\xi\sqrt{-1 + \omega^2}}{\sqrt{\varepsilon}\sqrt{\alpha_3 - \omega^2\alpha_4}}},$$

135 where  $c_1$  and  $c_2$  are constants of integration.

136 Now, by means of the values of  $a_0$ ,  $a_1$ ,  $a_2$  and  $\psi(\xi)$ , from Eq. (3.7), we obtain the  
137 subsequent solution:

$$U(\xi) = \frac{-1 + \omega^2}{\varepsilon \alpha_1} + \frac{6(-1 + \omega^2)^2 c_1 c_2 (-\alpha_3 + \omega^2 \alpha_4) e^{\pm \frac{\xi\sqrt{-1 + \omega^2}}{\sqrt{\varepsilon}\sqrt{\alpha_3 - \omega^2\alpha_4}}}}{\alpha_1 \left\{ (-1 + \omega^2) c_2 e^{\pm \frac{\xi\sqrt{-1 + \omega^2}}{\sqrt{\varepsilon}\sqrt{\alpha_3 - \omega^2\alpha_4}}} + \varepsilon c_1 (\alpha_3 - \omega^2 \alpha_4) \right\}^2}. \quad (3.21)$$

138 Now, transforming the required exponential function solution (3.21) into hyperbolic function,  
139 we obtain the following solution to the strain wave equation in the microstructured solids:

$$\begin{aligned} u(x, t) = & (-1 + \omega^2) [(-1 + \omega^2)^2 \{ \cosh(2\rho(x - t\omega)) + \sinh(2\rho(x - t\omega)) \} c_2^2 \\ & + \varepsilon^2 \{ \cosh(2\rho(x - t\omega)) - \sinh(2\rho(x - t\omega)) \} c_1^2 (\alpha_3 - \omega^2 \alpha_4)^2 \\ & + 4\varepsilon(-1 + \omega^2) c_1 c_2 (-\alpha_3 + \omega^2 \alpha_4)] \\ & / \left( \varepsilon \alpha_1 [(-1 + \omega^2) \{ \cosh(\rho(x - t\omega)) + \sinh(\rho(x - t\omega)) \} c_2 \right. \\ & \left. + \varepsilon \{ \cosh(\rho(x - t\omega)) - \sinh(\rho(x - t\omega)) \} c_1 (\alpha_3 - \omega^2 \alpha_4) \right]^2. \end{aligned} \quad (3.22)$$

140 Thus, we acquire the generalized solitary wave solution (3.22) to the strain wave equation in  
141 microstructured solids, where  $\rho = \frac{\sqrt{-1 + \omega^2}}{2\sqrt{\varepsilon}\sqrt{\alpha_3 - \omega^2\alpha_4}}$ . Since  $c_1$  and  $c_2$  are integration constants,  
142 therefore, somebody might randomly pick their values. So, if we pick  $c_1 = (-1 + \omega^2)$  and  
143  $c_2 = -\varepsilon(\alpha_3 - \omega^2\alpha_4)$ , then from (3.22), we obtain the subsequent solitary wave solution:

$$u_5(x, t) = \frac{(-1 + \omega^2)}{2\varepsilon\alpha_1} \left\{ 2 + 3 \operatorname{csch}^2 \left( \frac{(x - t\omega)\sqrt{-1 + \omega^2}}{2\sqrt{\varepsilon}\sqrt{\alpha_3 - \omega^2\alpha_4}} \right) \right\}. \quad (3.23)$$

144 Again, if we pick  $c_1 = (-1 + \omega^2)$  and  $c_2 = \varepsilon(\alpha_3 - \omega^2\alpha_4)$ , then the solitary wave solution  
 145 (3.22) reduces to:

$$u_6(x, t) = -\frac{(-1 + \omega^2)}{2\varepsilon\alpha_1} \left\{ -2 + 3 \operatorname{sech}^2 \left( \frac{(x - t\omega)\sqrt{-1 + \omega^2}}{2\sqrt{\varepsilon}\sqrt{\alpha_3 - \omega^2\alpha_4}} \right) \right\}. \quad (3.24)$$

146 Moreover, if we pick  $c_1 = (-1 + \omega^2)$  and  $c_2 = \mp i \varepsilon(\alpha_3 - \omega^2\alpha_4)$ , then from (3.22), we derive  
 147 the following solution:

$$u_7(x, t) = \frac{(-1 + \omega^2)}{\varepsilon\alpha_1} \left\{ 1 - \frac{3}{2} \operatorname{csc}^2 \left( \frac{\pi}{4} - \frac{1}{2} \frac{(x - t\omega)\sqrt{-1 + \omega^2}}{\sqrt{\varepsilon}\sqrt{-\alpha_3 + \omega^2\alpha_4}} \right) \right\}. \quad (3.25)$$

148 Again, if we pick  $c_1 = (-1 + \omega^2)$  and  $c_2 = \pm i \varepsilon(\alpha_3 - \omega^2\alpha_4)$ , then from (3.22), we obtain the  
 149 following solution:

$$u_8(x, t) = \frac{(-1 + \omega^2)}{\varepsilon\alpha_1} \left\{ 1 - \frac{3}{2} \operatorname{csc}^2 \left( \frac{\pi}{4} + \frac{1}{2} \frac{(x - t\omega)\sqrt{-1 + \omega^2}}{\sqrt{\varepsilon}\sqrt{-\alpha_3 + \omega^2\alpha_4}} \right) \right\}. \quad (3.26)$$

150 Forasmuch as,  $c_1$  and  $c_2$  are arbitrary constants, if we choose more different values of them,  
 151 we may derive a lot of general solitary wave solutions to the Eq. (3.3) through the MSE  
 152 method easily. But, we did not write down the other solutions for minimalism.

153 **Remark 1:** Solutions (3.17)-(3.20) and (3.23)-(3.26) have been confirmed by inserting them  
 154 into the main equation and found accurate.

155

### 156 3.2. THE DISSIPATIVE CASE

157 If  $\gamma \neq 0$ , then the system is dissipative. Therefore, for  $\delta = \gamma = O(\varepsilon)$ , the balance should be  
 158 between nonlinearity, dispersion and dissipation, perturbed by the higher order dissipative  
 159 terms to the strain wave equation in microstructured solids (see [52], [53], [56], [57] for  
 160 details)

E-mail address: ali\_math74@yahoo.com.

$$u_{tt} - u_{xx} - \varepsilon \{ \alpha_1 (u^2)_{xx} + \alpha_2 u_{xxt} - \alpha_3 u_{xxxx} + \alpha_4 u_{xxtt} \} = 0. \quad (3.27)$$

161 where  $\varepsilon \rightarrow 0$ , so the higher order term are omitted.

162 The traveling wave transformation (3.4) reduces Eq. (3.27) to the following ODE:

$$(\omega^2 - 1) U'' - \varepsilon \{ \alpha_1 (U^2)'' - \omega \alpha_2 U''' - (\alpha_3 - \omega^2 \alpha_4) U^{(iv)} \} = 0. \quad (3.28)$$

163 where prime stands for the differential coefficient. Integrating Eq. (3.28) with respect to  $\xi$ ,

164 we get

$$(\omega^2 - 1) U - \varepsilon \{ \alpha_1 U^2 - \omega \alpha_2 U' - (\alpha_3 - \omega^2 \alpha_4) U'' \} = 0. \quad (3.29)$$

165 The homogeneous between the highest order nonlinear term and the linear terms of the  
166 highest order, we obtain  $N = 2$ . Thus, the structure of the solution of Eq. (3.29) is one and  
167 the same to the form of the solution (3.7).

168 Inserting the values of  $U$ ,  $U'$  and  $U''$  into Eq. (3.29) and then setting each coefficient of

169  $\psi^{-j}$ ,  $j = 0, 1, 2, \dots$  to zero, we successively obtain

$$a_0(-1 + \omega^2 - \varepsilon a_0 \alpha_1) = 0. \quad (3.30)$$

$$a_1 \{ (-1 + \omega^2 - 2\varepsilon a_0 \alpha_1) \psi' + \varepsilon \omega \alpha_2 \psi'' + \varepsilon (\alpha_3 - \omega^2 \alpha_4) \psi''' \} = 0. \quad (3.31)$$

$$\begin{aligned} & -\varepsilon a_1 \psi' \{ (a_1 \alpha_1 + \omega \alpha_2) \psi' + 3(\alpha_3 - \omega^2 \alpha_4) \psi'' \} + 2\varepsilon a_2 \psi' \{ \omega \alpha_2 \psi'' + (\alpha_3 - \omega^2 \alpha_4) \psi''' \} \\ & + a_2 [ (-1 + \omega^2 - 2\varepsilon a_0 \alpha_1) (\psi')^2 + 2\varepsilon (\alpha_3 - \omega^2 \alpha_4) (\psi'')^2 ] = 0. \end{aligned} \quad (3.32)$$

$$-2\varepsilon a_1 (a_2 \alpha_1 - \alpha_3 + \omega^2 \alpha_4) (\psi')^3 - 2\varepsilon a_2 \{ \omega \alpha_2 \psi' + 5(\alpha_3 - \omega^2 \alpha_4) \psi'' \} (\psi')^2 = 0. \quad (3.33)$$

$$-\varepsilon a_2 (a_2 \alpha_1 - 6\alpha_3 + 6\omega^2 \alpha_4) (\psi')^4 = 0. \quad (3.34)$$

170

171 From Eqs. (3.30) and (3.34), we obtain

$$a_0 = 0, \quad \frac{-1 + \omega^2}{\varepsilon \alpha_1} \quad \text{and} \quad a_2 = \frac{6(\alpha_3 - \omega^2 \alpha_4)}{\alpha_1}, \quad \text{since } a_2 \neq 0.$$

172 Therefore, depending on the values of  $a_0$ , the following different cases arise:

173 **Case 1:** When  $a_0 = 0$ , from Eqs. (3.31) - (3.33), we get

E-mail address: ali\_math74@yahoo.com.

$$\psi(\xi) = c_2 + \frac{30c_1(\alpha_3 - \omega^2\alpha_4)}{-5a_1\alpha_1 - 6\omega\alpha_2} e^{\frac{\xi(-5a_1\alpha_1 - 6\omega\alpha_2)}{30(\alpha_3 - \omega^2\alpha_4)}},$$

$$a_1 = 0, \quad \omega = \pm \frac{\sqrt{\frac{6\varepsilon\alpha_2^2 - 25(\alpha_3 + \alpha_4) + \sqrt{\{6\varepsilon\alpha_2^2 - 25(\alpha_3 + \alpha_4)\}^2 - 2500\alpha_3\alpha_4}}{-\alpha_4}}}{5\sqrt{2}} = \pm\theta,$$

174 and

$$a_1 = \frac{3 \left[ 3\varepsilon\omega\alpha_1\alpha_2 + 5\sqrt{\varepsilon\alpha_1^2\{\varepsilon\omega^2\alpha_2^2 + 4(-1 + \omega^2)(-\alpha_3 + \omega^2\alpha_4)\}} \right]}{5\varepsilon\alpha_1^2},$$

$$\omega = -\frac{\sqrt{25 + \frac{6\varepsilon\alpha_2^2}{\alpha_4} + \frac{25\alpha_3}{\alpha_4} \pm \frac{\sqrt{(-6\varepsilon\alpha_2^2 - 25\alpha_3 - 25\alpha_4)^2 - 2500\alpha_3\alpha_4}}{\alpha_4}}}{5\sqrt{2}},$$

175 where  $c_1$  and  $c_2$  are integration constants.

176 Hence for the values of  $a_1$  and  $\omega$ , there also arise three cases. But when  $a_1 \neq 0$  then the  
177 shape of the solutions for dissipative case is distorted and the solution size is very long. So  
178 we have omitted the other value of  $a_1$  and discussed only for  $a_1 = 0$ .

179 When  $a_1 = 0$  then we get also the solutions to the above mentioned equation depends for  
180 the values of  $\omega$ . Thus,

$$\psi(\xi) = c_2 - \frac{5c_1(\alpha_3 - \omega^2\alpha_4)}{\omega\alpha_2} e^{-\frac{\xi\omega\alpha_2}{5(\alpha_3 - \omega^2\alpha_4)}}$$

181 Now, by means of the values of  $a_0$ ,  $a_1$ ,  $a_2$  and  $\psi(\xi)$  from Eq. (3.7), we achieve the  
182 subsequent solution:

$$U(\xi) = -\frac{6\omega^2c_1^2\alpha_2^2(-\alpha_3 + \omega^2\alpha_4)}{\alpha_1 \left\{ \omega c_2 \alpha_2 e^{\frac{\xi\omega\alpha_2}{5\alpha_3 - 5\omega^2\alpha_4}} - 5c_1(\alpha_3 - \omega^2\alpha_4) \right\}^2}. \quad (3.35)$$

183 Simplifying the required solution (3.35), we derive the following close-form solution of the  
184 strain wave equation in microstructured solids for dissipative case (3.27):

$$\begin{aligned} u(x, t) = & \left[ 6\omega^2 \{ -\cosh(2\sigma(x - t\omega)) + \sinh(2\sigma(x - t\omega)) \} c_1^2 \alpha_2^2 (-\alpha_3 + \omega^2 \alpha_4) \right] \\ & / \left( \alpha_1 [ \omega \{ \cosh(\sigma(x - t\omega)) + \sinh(\sigma(x - t\omega)) \} c_2 \alpha_2 \right. \\ & \left. + 5 \{ -\cosh(\sigma(x - t\omega)) + \sinh(\sigma(x - t\omega)) \} c_1 (\alpha_3 - \omega^2 \alpha_4) \}^2 \right). \end{aligned} \quad (3.36)$$

E-mail address: ali\_math74@yahoo.com.

185 where  $\sigma = \frac{\omega \alpha_2}{10(\alpha_3 - \omega^2 \alpha_4)}$ ,  $\omega = \pm \theta$  or and  $c_1, c_2$  are integrating constants. Since  $c_1$  and  $c_2$  are  
 186 integration constants, one might arbitrarily select their values. If we choose  $c_1 = \alpha_2 \omega$  and  
 187  $c_2 = -5(\alpha_3 - \omega^2 \alpha_4)$ , then from (3.36), we obtain

$$u_9(x, t) = \frac{3\omega^2 \alpha_2^2}{50\alpha_1(\alpha_3 - \omega^2 \alpha_4)} \left\{ 1 + \tanh \left( \frac{\omega(-x + t\omega)\alpha_2}{10(\alpha_3 - \omega^2 \alpha_4)} \right) \right\}^2. \quad (3.37)$$

188 Again if we choose  $c_1 = \alpha_2 \omega$  and  $c_2 = 5(\alpha_3 - \omega^2 \alpha_4)$ , then from (3.36), we attain the  
 189 subsequent soliton solution:

$$u_{10}(x, t) = \frac{3\omega^2 \alpha_2^2}{50\alpha_1(\alpha_3 - \omega^2 \alpha_4)} \left\{ 1 + \coth \left( \frac{\omega(-x + t\omega)\alpha_2}{10(\alpha_3 - \omega^2 \alpha_4)} \right) \right\}^2. \quad (3.38)$$

190 **Case 2:** When  $a_0 = \frac{-1 + \omega^2}{\varepsilon \alpha_1}$ , from Eq.(3.31)-(3.33), we obtain

$$\psi(\xi) = c_2 + \frac{30c_1(\alpha_3 - \omega^2 \alpha_4)}{-5a_1\alpha_1 - 6\omega\alpha_2} e^{\frac{\xi(-5a_1\alpha_1 - 6\omega\alpha_2)}{30(\alpha_3 - \omega^2 \alpha_4)}},$$

191 where  $c_1$  and  $c_2$  are integration constants and

$$a_1 = 0, \quad \omega = \begin{cases} \pm \sqrt{\frac{6\varepsilon\alpha_2^2 + 25\alpha_3 + 25\alpha_4 - \sqrt{\{6\varepsilon\alpha_2^2 + 25(\alpha_3 + \alpha_4)\}^2 - 2500\alpha_3\alpha_4}}{\alpha_4}}}{5\sqrt{2}} = \pm \vartheta_1(\text{say}) \\ \pm \sqrt{\frac{6\varepsilon\alpha_2^2 + 25\alpha_3 + 25\alpha_4 + \sqrt{\{6\varepsilon\alpha_2^2 + 25(\alpha_3 + \alpha_4)\}^2 - 2500\alpha_3\alpha_4}}{\alpha_4}}}{5\sqrt{2}} = \pm \vartheta_2(\text{say}); \end{cases}$$

$$a_1 = \frac{3 \left[ 3\varepsilon\omega\alpha_1\alpha_2 + 5\sqrt{\varepsilon\alpha_1^2\{\varepsilon\omega^2\alpha_2^2 + 4(-1 + \omega^2)(\alpha_3 - \omega^2\alpha_4)\}} \right]}{5\varepsilon\alpha_1^2},$$

$$\omega = -\sqrt{\frac{-6\varepsilon\alpha_2^2 + 25\alpha_3 + 25\alpha_4 \pm \sqrt{\{6\varepsilon\alpha_2^2 - 25(\alpha_3 + \alpha_4)\}^2 - 2500\alpha_3\alpha_4}}{\alpha_4}}}{5\sqrt{2}};$$

$$a_1 = \frac{3 \left[ 3\varepsilon\omega\alpha_1\alpha_2 - 5\sqrt{\varepsilon\alpha_1^2\{\varepsilon\omega^2\alpha_2^2 + 4(-1 + \omega^2)(\alpha_3 - \omega^2\alpha_4)\}} \right]}{5\varepsilon\alpha_1^2},$$

$$\omega = \sqrt{\frac{-6\varepsilon\alpha_2^2 + 25\alpha_3 + 25\alpha_4 \pm \sqrt{\{6\varepsilon\alpha_2^2 - 25(\alpha_3 + \alpha_4)\}^2 - 2500\alpha_3\alpha_4}}{\alpha_4}}}{5\sqrt{2}}.$$

192 Hence for the values of  $a_1$  and  $\omega$ , there arises also three cases. When  $a_1 \neq 0$ , then the form  
 193 of solutions to the strain wave equation in microstructured solids for dissipative case (3.24)  
 194 indistinct and the solution size is very lengthy. So we omitted the extra value of  $a_1$  and only  
 195 discuss for  $a_1 = 0$ .

196 When  $a_1 = 0$  then we find also the solutions to the above revealed equation depends for the  
 197 values of  $\omega$ , i.e.  $\omega = \pm\vartheta_1$  and  $\omega = \pm\vartheta_2$ . Therefore,

$$\psi(\xi) = c_2 - \frac{5c_1(\alpha_3 - \omega^2\alpha_4)}{\omega\alpha_2} e^{-\frac{\xi\omega\alpha_2}{5(\alpha_3 - \omega^2\alpha_4)}}$$

198 where  $\omega = \pm\vartheta_1$  or  $\omega = \pm\vartheta_2$ ,  $c_1$  and  $c_2$  are constants of integration.

199 Substituting the values of  $a_0$ ,  $a_1$ ,  $a_2$  and  $\psi(\xi)$  into Eq. (3.7), we accomplish the following  
 200 solution:

$$U(\xi) = \frac{-1 + \omega^2}{\varepsilon\alpha_1} - \frac{6\omega^2 c_1^2 \alpha_2^2 (-\alpha_3 + \omega^2 \alpha_4)}{\alpha_1 \left\{ \omega c_2 \alpha_2 e^{\frac{\xi\omega\alpha_2}{5\alpha_3 - 5\omega^2\alpha_4}} - 5c_1(\alpha_3 - \omega^2\alpha_4) \right\}^2}. \quad (3.39)$$

201 Simplifying the required exponential function solution (3.39) into trigonometric function  
 202 solution, we derive the solution of Eq. (3.27) as follows:

$$\begin{aligned} u(x, t) = & \left[ \omega^2(-1 + \omega^2) \{ \cosh(2\varphi(x - t\omega)) + \sinh(2\varphi(x - t\omega)) \} c_2^2 \alpha_2^2 \right. \\ & + \{ \cosh(2\varphi(x - t\omega)) - \sinh(2\varphi(x - t\omega)) \} c_1^2 (\alpha_3 - \omega^2 \alpha_4) \{ 6\varepsilon\omega^2 \alpha_2^2 \\ & - 25(-1 + \omega^2)(-\alpha_3 + \omega^2 \alpha_4) \} + 10\omega(-1 + \omega^2) c_1 c_2 \alpha_2 (-\alpha_3 + \omega^2 \alpha_4) ] \\ & / \left( \varepsilon\alpha_1 [ \omega \{ \cosh(\varphi(x - t\omega)) + \sinh(\varphi(x - t\omega)) \} c_2 \alpha_2 \right. \\ & \left. \left. + 5 \{ -\cosh(\varphi(x - t\omega)) + \sinh(\varphi(x - t\omega)) \} c_1 (\alpha_3 - \omega^2 \alpha_4) \right]^2 \right). \quad (3.40) \end{aligned}$$

203 Therefore, we obtain the generalized soliton solution (3.40) to the strain wave equation in  
 204 microstructured solids for dissipative case, where  $\varphi = \frac{\omega \alpha_2}{10(\alpha_3 - \omega^2 \alpha_4)}$  and  $\omega = \pm\vartheta_1$  or  $\omega = \pm\vartheta_2$ .  
 205 But, since  $c_1$  and  $c_2$  are arbitrary constants, someone may arbitrarily choose their values. So,

206 if we choose  $c_1 = \alpha_2 \omega$  and  $c_2 = 5(\alpha_3 - \omega^2 \alpha_4)$ , from (3.20), we get the subsequent soliton  
 207 solutions:

$$u_{11}(x, t) = \frac{(-1 + \omega^2)}{\alpha_1 \varepsilon} - \frac{3\omega^2 \alpha_2^2}{50\alpha_1(-\alpha_3 + \omega^2 \alpha_4)} \left\{ -1 + \coth \left( \frac{\omega(x - t\omega)\alpha_2}{10(\alpha_3 - \omega^2 \alpha_4)} \right) \right\}^2. \quad (3.41)$$

208 Again, if we choose  $c_1 = \alpha_2 \omega$  and  $c_2 = -5(\alpha_3 - \omega^2 \alpha_4)$ , the solitary wave solution (3.40)  
 209 becomes

$$u_{12}(x, t) = \frac{(-1 + \omega^2)}{\varepsilon \alpha_1} + \frac{3\varepsilon \omega^2 \alpha_2^2}{50\varepsilon \alpha_1(\alpha_3 - \omega^2 \alpha_4)} \left\{ -1 + \tanh \left( \frac{\omega(x - t\omega)\alpha_2}{10(\alpha_3 - \omega^2 \alpha_4)} \right) \right\}^2. \quad (3.42)$$

210 As  $c_1$  and  $c_2$  are arbitrary constants, one may pick many other values of them and each of  
 211 this selection construct new solution. But for minimalism, we have not recorded these  
 212 solutions.

213 **Remark 2:** The solutions (3.37)-(3.38), where  $\omega = \pm\theta_1$  or  $\omega = \pm\theta_2$  and the solutions (3.41)-  
 214 (3.42)  $\omega = \pm\theta_1$  or  $\omega = \pm\theta_2$  have been confirmed by satisfying the original equation.

215

216

#### 217 4. PHYSICAL INTERPRETATIONS OF THE SOLUTIONS

218 In this sub-section, we draw the graph of the derived solutions and explain the effect of the  
 219 parameters on the solutions for both non-dissipative and dissipative cases. The solution  $u_1$   
 220 in (3.17) depends on the physical parameters  $\alpha_1, \alpha_3, \alpha_4, \varepsilon$  and the group velocity  $\omega$ . Now,  
 221 we will discuss all the possible physical significances for  $-2 \leq \alpha_1, \alpha_3, \alpha_4, \varepsilon \leq 2$ , and soliton  
 222 exists for  $|\omega| > 1$  and  $|\omega| < 1$ . For the value of parameters  $\alpha_1, \alpha_3, \alpha_4, \varepsilon < 0$  and  $|\omega| > 1$ , the  
 223 solution  $u_1$  in (3.17) represents the bell shape soliton and when  $|\omega| < 1$  then the solution  $u_1$   
 224 represents the anti-bell shape soliton. It is shown in Fig. 1. Also if the values of the  
 225 parameters are  $\alpha_1 > 0, \alpha_3, \alpha_4, \varepsilon < 0$  and  $|\omega| > 1$ , then the solution  $u_1$  represents the anti-

226 bell shape soliton and when  $|\omega| < 1$ , then the solution  $u_1$  represents the bell shape soliton. It  
 227 is shown the Fig. 2. Again, for  $\alpha_1, \alpha_3, \alpha_4 < 0, \varepsilon > 0$  and  $|\omega| < 1$ , the solution  $u_1$  in (3.17)  
 228 represents the multi-soliton and when  $|\omega| > 1$ , the solution  $u_1$  represents the anti-bell shape  
 229 soliton. It is plotted in Fig. 3. Again, if the values of the physical parameters are  
 230  $\alpha_1 > 0, \alpha_3, \alpha_4 < 0, \varepsilon > 0$  and  $|\omega| > 1$ , then the solution  $u_1$  represents the anti-bell shape  
 231 soliton and when  $|\omega| < 1$  then the solution  $u_1$  represents the bell shape soliton. It is shown in  
 232 Fig. 4. We can sketch the other figures of the solution  $u_1$  for different values of the  
 233 parameters. But for page limitation in this article we have omitted these figures. So, for other  
 234 cases we do not draw the figures but we discuss for other cases with the following table:

$\varepsilon > 0$	$ \omega  > 1$	$\alpha_1 < 0, \alpha_3 < 0, \alpha_4 < 0$	Anti-bell shape soliton
		$\alpha_1 > 0, \alpha_3 < 0, \alpha_4 < 0$	Bell shape soliton
		$\alpha_1 > 0, \alpha_3 > 0, \alpha_4 < 0$	Bell shape soliton
		$\alpha_1 > 0, \alpha_3 > 0, \alpha_4 > 0$	Bell shape soliton
		$\alpha_1 > 0, \alpha_3 < 0, \alpha_4 > 0$	Bell shape soliton
		$\alpha_1 < 0, \alpha_3 > 0, \alpha_4 < 0$	Anti-bell shape soliton
		$\alpha_1 < 0, \alpha_3 > 0, \alpha_4 > 0$	Anti-bell shape soliton
		$\alpha_1 < 0, \alpha_3 < 0, \alpha_4 > 0$	Anti-bell shape soliton
	$ \omega  < 1$	$\alpha_1 < 0, \alpha_3 < 0, \alpha_4 < 0$	Bell shape soliton
		$\alpha_1 > 0, \alpha_3 < 0, \alpha_4 < 0$	Anti-bell shape soliton
		$\alpha_1 > 0, \alpha_3 > 0, \alpha_4 < 0$	Anti-bell shape soliton
		$\alpha_1 > 0, \alpha_3 > 0, \alpha_4 > 0$	Anti-bell shape soliton
		$\alpha_1 > 0, \alpha_3 < 0, \alpha_4 > 0$	Anti-bell shape soliton
		$\alpha_1 < 0, \alpha_3 > 0, \alpha_4 < 0$	Bell shape soliton
		$\alpha_1 < 0, \alpha_3 > 0, \alpha_4 > 0$	Bell shape soliton
		$\alpha_1 < 0, \alpha_3 < 0, \alpha_4 > 0$	Periodic bell shape solution
$\varepsilon < 0$	$ \omega  > 1$	$\alpha_1 < 0, \alpha_3 < 0, \alpha_4 < 0$	Bell shape or Periodic bell shape solution
		$\alpha_1 > 0, \alpha_3 < 0, \alpha_4 < 0$	Anti-bell shape soliton or Periodic anti-bell shape solution
		$\alpha_1 > 0, \alpha_3 > 0, \alpha_4 < 0$	Anti-bell shape soliton
		$\alpha_1 > 0, \alpha_3 > 0, \alpha_4 > 0$	Periodic anti-bell shape solution
		$\alpha_1 > 0, \alpha_3 < 0, \alpha_4 > 0$	Periodic anti-bell shape solution
		$\alpha_1 < 0, \alpha_3 > 0, \alpha_4 < 0$	Bell shape soliton
		$\alpha_1 < 0, \alpha_3 > 0, \alpha_4 > 0$	Periodic bell shape solution
		$\alpha_1 < 0, \alpha_3 < 0, \alpha_4 > 0$	Periodic bell shape solution

E-mail address: ali\_math74@yahoo.com.


	$ \omega  < 1$	$\alpha_1 < 0, \alpha_3 < 0, \alpha_4 < 0$	Anti-bell shape soliton or Periodic anti-bell shape solution
		$\alpha_1 > 0, \alpha_3 < 0, \alpha_4 < 0$	Bell shape or Periodic bell shape solution
		$\alpha_1 > 0, \alpha_3 > 0, \alpha_4 < 0$	Periodic bell shape solution
		$\alpha_1 > 0, \alpha_3 > 0, \alpha_4 > 0$	Bell shape or Periodic bell shape solution
		$\alpha_1 > 0, \alpha_3 < 0, \alpha_4 > 0$	Bell shape soliton
		$\alpha_1 < 0, \alpha_3 > 0, \alpha_4 < 0$	Periodic anti-bell shape solution
		$\alpha_1 < 0, \alpha_3 > 0, \alpha_4 > 0$	Anti-bell shape soliton or Periodic anti-bell shape solution
		$\alpha_1 < 0, \alpha_3 < 0, \alpha_4 > 0$	Anti-bell shape soliton

235  
 236 Also the soliton  $u_2$  in (3.18) depends on the parameters  $\alpha_1, \alpha_3, \alpha_4, \varepsilon$  and  $\omega$ . Now, we will  
 237 discuss all the possible physical significances for  $-2 \leq \alpha_1, \alpha_3, \alpha_4, \varepsilon \leq 2$ , and soliton exists  
 238 for  $|\omega| > 1$  and  $|\omega| < 1$ . For the value of parameters contains  $\alpha_1, \alpha_3, \alpha_4, \varepsilon > 0$  and  $|\omega| > 1$ ,  
 239 then the solution  $u_2$  in (3.18) represents the singular anti-bell shape soliton and when  $|\omega| < 1$   
 240 then the solution  $u_2$  represents the singular bell shape soliton. It is shown in Fig. 5. Also, for  
 241  $\alpha_1, \alpha_3, \alpha_4 < 0, \varepsilon > 0$  and  $|\omega| > 1$ , then the solution  $u_2$  in (3.18) represents the periodic  
 242 singular anti-bell shape solution and when  $|\omega| < 1$  then the solution  $u_2$  represents the  
 243 periodic singular bell shape solution. It is plotted of the Fig. 6. On the other hand, the  
 244 solutions  $u_3$  in (3.19) and  $u_4$  in (3.20) exist for  $(\alpha_3 - \alpha_4 \omega^2) > 0, \varepsilon < 0$  or  
 245  $(\alpha_3 - \alpha_4 \omega^2) < 0, \varepsilon > 0$  when  $|\omega| > 1$  or  $|\omega| < 1$ . For the value of the parameters are  
 246  $\alpha_1 = -1.25, \alpha_3 = -0.1, \alpha_4 = -2, \varepsilon = -1$ , when  $\omega = 0.96$ , the solution  $u_3$  in (3.19) represents  
 247 the anti-bell shape soliton and  $\alpha_1 = -1.5, \alpha_3 = -0.1, \alpha_4 = 2, \varepsilon = -1$ , when  $\omega = 1.5$ , the  
 248 solution  $u_4$  represents the periodic solution. It is shown in Fig. 7. Again, the travelling wave  
 249 solution  $u_5$  in (3.23) represents the bell shape singular solitons for  $\alpha_1 = -1 = \alpha_3, \alpha_4 = 1$ ,  
 250  $\varepsilon = 0.5, \omega = -1.5$  and  $\omega = 0.5$  respectively, in Fig. 8 and Fig. 9 from  $u_6$  in (3.24)  
 251 represents the bell shape soliton, when  $\omega = 1.5$  and the anti-bell shape soliton, when  
 252  $\omega = -0.75$ . In Fig. 10, we have plotted of the periodic bell shape and anti-bell shape solution

253 for  $\alpha_1 = \alpha_3 = -1.25$ ,  $\alpha_4 = 1$ ,  $\varepsilon = 0.7$ ,  $\omega = -1.2$  and  $\alpha_1 = \alpha_3 = -1.25$ ,  $\alpha_4 = 1$ ,  $\varepsilon = -0.7$ ,  
 254  $\omega = 0.25$  respectively to the solution of  $u_7$  in (3.25) and Fig. 11 plotted the periodic anti-bell  
 255 shape solution and bell shape solution for  $\alpha_1 = 1.25$ ,  $\alpha_3 = -1.25$ ,  $\alpha_4 = 1$ ,  $\varepsilon = 0.7$ ,  $\omega = -1.2$   
 256 and  $\alpha_1 = \alpha_3 = -1.25$ ,  $\alpha_4 = 1$ ,  $\varepsilon = -0.7$ ,  $\omega = -0.25$  respectively to the solution of  $u_8$  in  
 257 (3.26). Fig. 12 and 13 represent the kink shape solutions  $u_9$  given in (3.37) are respectively,  
 258 for  $\alpha_1 = 1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = -1.5$ ,  $\alpha_4 = -1$  and  $\alpha_1 = -1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = -1.5$ ,  $\alpha_4 = -1$   
 259 respectively, when  $\omega = \pm\mu_1$  and for  $\alpha_1 = 1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = -1.5$ ,  $\alpha_4 = -1$  and  $\alpha_1 = -1$ ,  
 260  $\alpha_2 = 1$ ,  $\alpha_3 = -1.5$ ,  $\alpha_4 = -1$  respectively, when  $\omega = \pm\mu_2$ . Also sketch the figures 14 and 15,  
 261 singular bell shape solutions  $u_{10}$  in (3.38) for  $\alpha_1 = 1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = -1.5$ ,  $\alpha_4 = -1$  and  
 262  $\alpha_1 = -1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = -1.5$ ,  $\alpha_4 = -1$  respectively, when  $\omega = \pm\mu_1$  and for  $\alpha_1 = 1$ ,  $\alpha_2 = 1$ ,  
 263  $\alpha_3 = -1.5$ ,  $\alpha_4 = -1$  and  $\alpha_1 = -1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = -1.5$ ,  $\alpha_4 = -1$  respectively, when  $\omega = \pm\mu_2$ .  
 264 On the other hand, Fig. 16 and 17 are singular bell and singular anti-bell shape soliton  
 265 solitons  $u_{11}$  in (3.41) for  $\alpha_1 = 1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = 1$ ,  $\alpha_4 = 1$ ,  $\varepsilon = 0.5$  and  $\alpha_1 = -1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = 1$ ,  
 266  $\alpha_4 = 1$ ,  $\varepsilon = 0.5$  respectively, when  $\omega = \pm\theta_1$  and for  $\alpha_1 = 1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = 1$ ,  $\alpha_4 = 1$ ,  $\varepsilon = 0.5$   
 267 and  $\alpha_1 = -1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = 1$ ,  $\alpha_4 = 1$ ,  $\varepsilon = 0.5$  respectively, when  $\omega = \pm\theta_2$ . Also, draw the  
 268 Figures 18 and 19 are kink shape solitons  $u_{12}$  in (3.42) for  $\alpha_1 = 1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = 1$ ,  $\alpha_4 = 1$ ,  
 269  $\varepsilon = 0.5$  and  $\alpha_1 = -1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = 1$ ,  $\alpha_4 = 1$ ,  $\varepsilon = 0.5$  respectively, when  $\omega = \pm\theta_1$  and for  
 270  $\alpha_1 = 1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = 1$ ,  $\alpha_4 = 1$ ,  $\varepsilon = 0.5$  and  $\alpha_1 = -1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = 1$ ,  $\alpha_4 = 1$ ,  $\varepsilon = 0.5$   
 271 respectively, when  $\omega = \pm\theta_2$ . All figures are drawn within  $-10 \leq x, t \leq 10$ .  
 272 There is another kind of solution which is not a kink, anti-kink, dark or bell-shape soliton,  
 273 known as Love wave [58, 59]. A Love wave is define to be a surface wave having a  
 274 horizontal motion that is transverse or perpendicular to the direction the wave is traveling.  
 275 We can discuss the solutions  $u_2$  to  $u_{12}$  for other values of the parameters. But for page  
 276 limitation in this article we have omitted these figures in details.

E-mail address: ali\_math74@yahoo.com.

277




278

279

280

Fig. 1: Sketch of the solution  $u_1$  for  $\alpha_1 = -0.001$ ,  $\alpha_3 = \alpha_4 = \varepsilon = \omega = -1.5$  and  $\alpha_1 = -0.001$ ,  $\alpha_3 = \alpha_4 = \varepsilon = \omega = -0.75$  respectively.




281

282

283

Fig. 2: Plot of the solution  $u_1$  for  $\alpha_1 = 0.001$ ,  $\alpha_3 = \alpha_4 = \varepsilon = \omega = -1.5$  and  $\alpha_1 = 0.001$ ,  $\alpha_3 = \alpha_4 = \varepsilon = \omega = -0.75$  respectively.



284

285

286

287

288

Fig. 3: Sketch of the solution  $u_1$  for  $\alpha_1 = \alpha_3 = \alpha_4 = -1.2$ ,  $\varepsilon = \omega = 0.5$  and  $\alpha_1 = \alpha_3 = \alpha_4 = -1.2$ ,  $\varepsilon = 0.5$ ,  $\omega = 1.25$  respectively.




Fig. 4: Sketch of the solution  $u_1$  for  $\alpha_1 = 0.75$ ,  $\alpha_3 = \alpha_4 = -1.2$ ,  $\varepsilon = 0.5$ ,  $\omega = 1.25$  and  $\alpha_1 = 0.75$ ,  $\alpha_3 = \alpha_4 = -1.2$ ,  $\varepsilon = 0.5$ ,  $\omega = 0.5$  respectively.




Fig. 5: Sketch of the singular dark and singular bell shape soliton  $u_2$  for  $\alpha_1 = \alpha_3 = \alpha_4 = 0.5$ ,  $\varepsilon = 0.75$ ,  $\omega = -1.5$  and  $\alpha_1 = \alpha_3 = \alpha_4 = 0.5$ ,  $\varepsilon = 0.75$ ,  $\omega = -0.25$  respectively.




Fig. 6: Sketch of the periodic singular solution  $u_2$  for  $\alpha_1 = \alpha_3 = \alpha_4 = -1.5$ ,  $\varepsilon = 0.75$ ,  $\omega = -1.5$  and  $\alpha_1 = \alpha_3 = \alpha_4 = -1.5$ ,  $\varepsilon = 0.75$ ,  $\omega = -0.25$  respectively.




Fig. 7: Sketch of the solution  $u_3$  and the solution  $u_4$  for  $\alpha_1 = -1.25$ ,  $\alpha_3 = -0.1$ ,  $\alpha_4 = -2$ ,  $\varepsilon = -1$ ,  $\omega = 0.96$  and  $\alpha_1 = -1.5$ ,  $\alpha_3 = -0.1$ ,  $\alpha_4 = 2$ ,  $\varepsilon = -1$ ,  $\omega = 1.5$  respectively.




Fig. 8: Sketch of the solutions  $u_5$  for  $\alpha_1 = -1 = \alpha_3$ ,  $\alpha_4 = 1$ ,  $\varepsilon = 0.5$ ,  $\omega = -1.5$  and  $\omega = 0.5$  respectively.




Fig. 9: Sketch of the bell shape soliton and anti-bell shape soliton  $u_6$  for  $\alpha_1 = \alpha_3 = \alpha_4 = -1$ ,  $\varepsilon = 0.5$ ,  $\omega = 1.5$  and  $\omega = -0.75$  respectively.




Fig. 10: Sketch of the solutions  $u_7$  for  $\alpha_1 = \alpha_3 = -1.25$ ,  $\alpha_4 = 1$ ,  $\varepsilon = 0.7$ ,  $\omega = -1.2$  and  $\alpha_1 = \alpha_3 = -1.25$ ,  $\alpha_4 = 1$ ,  $\varepsilon = -0.7$ ,  $\omega = 0.25$  respectively.




Fig. 11: Sketch of the solutions  $u_8$  for  $\alpha_1 = 1.25$ ,  $\alpha_3 = -1.25$ ,  $\alpha_4 = 1$ ,  $\varepsilon = 0.7$ ,  $\omega = -1.2$  and  $\alpha_1 = \alpha_3 = -1.25$ ,  $\alpha_4 = 1$ ,  $\varepsilon = -0.7$ ,  $\omega = -0.25$  respectively.




Fig. 12: Kink shape soliton obtained from  $u_9$  for  $\alpha_1 = 1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = -1.5$ ,  $\alpha_4 = -1$ ,  $\varepsilon = 0.5$  and  $\alpha_1 = -1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = -1.5$ ,  $\alpha_4 = -1$ ,  $\varepsilon = 0.5$  respectively, when  $\omega = \pm\mu_1$ .




Fig. 13: Kink shape soliton obtained from  $u_9$  for  $\alpha_1 = 1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = -1.5$ ,  $\alpha_4 = -1$ ,  $\varepsilon = 0.5$  and  $\alpha_1 = -1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = -1.5$ ,  $\alpha_4 = -1$ ,  $\varepsilon = 0.5$  respectively, when  $\omega = \pm\mu_2$ .




Fig. 14: Singular bell shape and anti-bell shape soliton  $u_{10}$  for  $\alpha_1 = 1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = -1.5$ ,  $\alpha_4 = -1$ ,  $\varepsilon = 0.5$  and  $\alpha_1 = -1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = -1.5$ ,  $\alpha_4 = -1$ ,  $\varepsilon = 0.5$  respectively, when  $\omega = \pm\mu_1$ .







Fig. 15: Singular anti-bell shape and bell shape soliton  $u_{10}$  in (3.38) for  $\alpha_1 = 1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = -1.5$ ,  $\alpha_4 = -1$ ,  $\varepsilon = 0.5$  and  $\alpha_1 = -1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = -1.5$ ,  $\alpha_4 = -1$ ,  $\varepsilon = 0.5$  respectively, when  $\omega = \pm\mu_2$ .



330 Fig. 16: Sketch the singular bell type and anti-bell soliton  $u_{11}$  for  $\alpha_1 = 1, \alpha_2 = 1, \alpha_3 = 1, \alpha_4 = 1, \varepsilon = 0.5$   
 331 and  $\alpha_1 = -1, \alpha_2 = 1, \alpha_3 = 1, \alpha_4 = 1, \varepsilon = 0.5$  respectively, when  $\omega = \pm\theta_1$ .  
 332



333 Fig. 17: Singular anti-bell shape and bell shape soliton  $u_{11}$  for  $\alpha_1 = 1, \alpha_2 = 1, \alpha_3 = 1, \alpha_4 = 1, \varepsilon = 0.5$  and  
 334  $\alpha_1 = -1, \alpha_2 = 1, \alpha_3 = 1, \alpha_4 = 1, \varepsilon = 0.5$  respectively, when  $\omega = \pm\theta_2$ .  
 335



336 Fig. 18: Kink shape soliton  $u_{12}$  for  $\alpha_1 = 1, \alpha_2 = 1, \alpha_3 = 1, \alpha_4 = 1, \varepsilon = 0.5$  and  $\alpha_1 = -1, \alpha_2 = 1,$   
 337  $\alpha_3 = 1, \alpha_4 = 1, \varepsilon = 0.5$  respectively, when  $\omega = \pm\theta_1$ .  
 338




Fig. 19: Kink shape soliton  $u_{12}$  for  $\alpha_1 = 1$ ,  $\alpha_2 = 1$ ,  $\alpha_3 = 1$ ,  $\alpha_4 = 1$ ,  $\varepsilon = 0.5$  and  $\alpha_1 = -1$ ,

$\alpha_2 = 1$ ,  $\alpha_3 = 1$ ,  $\alpha_4 = 1$ ,  $\varepsilon = 0.5$  respectively, when  $\omega = \pm\theta_2$ .

## 5. CONCLUSION

In this article, we have implemented the MSE method to obtain soliton solutions to the strain wave equation in microstructured solids for both non-dissipative and dissipative cases. In fact, we have derived general solitary wave solutions to this equation associated with arbitrary constants, and for particular values of these constants solitons are originated from the general solitary wave solutions. We have illustrated the solitary wave properties of the solutions for various values of the free parameters by means of the graphs. This work shows that the MSE method is competent and more powerful and can be used for many other equations NLEEs applied mathematics and engineering.

## ACKNOWLEDGEMENTS

The authors wish to take this opportunity to express their gratitude to the referees for their valuable comments and suggestions which enhanced the quality of this article.

## REFERENCES

1. V.B. Matveev and M.A. Salle, Darboux transformation and solitons, Springer, Berlin, 1991.

E-mail address: ali\_math74@yahoo.com.

- 358 2. G. Xu, An elliptic equation method and its applications in nonlinear evolution equations,  
359 Chaos, Solitons Fract., 29 (2006) 942-947.
- 360 3. E. Yusufoglu and A. Bekir, Exact solution of coupled nonlinear evolution equations,  
361 Chaos, solitons Fract., 37 (2008) 842-848.
- 362 4. D.D. Ganji, The application of He's homotopy perturbation method to nonlinear  
363 equations arising in heat transfer, Phys. Lett. A, 355 (2006) 137-141.
- 364 5. D.D. Ganji, G.A. Afrouzi and R.A. Talarposhti, Application of variational iteration method  
365 and homotopy perturbation method for nonlinear heat diffusion and heat transfer  
366 equations, Phys. Lett. A, 368 (2007) 450-457.
- 367 6. W. Malfliet and W. Hereman, The tanh method II: Perturbation technique for  
368 conservative systems, Phys. Scr., 54 (1996) 563-569.
- 369 7. H.A. Nassar, M.A. Abdel-Razek and A.K. Seddeek, Expanding the tanh-function method  
370 for solving nonlinear equations, Appl. Math., 2 (2011) 1096-1104.
- 371 8. A.J.M. Jawad, M.D. Petkovic, P. Laketa and A. Biswas, Dynamics of shallow water  
372 waves with Boussinesq equation, Scientia Iranica, Trans. B: Mech. Engr., 20(1) (2013)  
373 179-184.
- 374 9. M.A. Abdou, The extended tanh method and its applications for solving nonlinear  
375 physical models, Appl. Math. Comput., 190 (1) (2007) 988-996.
- 376 10. A.L. Guo and J. Lin, Exact solutions of (2+1)-dimensional HNLS equation, Commun.  
377 Theor. Phys., 54 (2010) 401-406.
- 378 11. S.T. Mohyud-Din, , M.A. Noor and K.I. Noor, Modified Variational Iteration Method for  
379 Solving Sine-Gordon Equations, World Appl. Sci. J., 6 (7) (2009) 999-1004.
- 380 12. R. Hirota, The direct method in soliton theory, Cambridge University Press, Cambridge,  
381 2004.

- 382 13. C. Rogers and W.F. Shadwick, Backlund transformations and their applications, Vol. 161  
383 of Mathematics in Science and Engineering, Academic Press, New York, USA, 1982.
- 384 14. L. Jianming, D. Jie and Y. Wenjun, Backlund transformation and new exact solutions of  
385 the Sharma-Tasso-Olver equation, Abstract and Appl. Analysis, 2011 (2011) Article ID  
386 935710, 8 pages.
- 387 15. M.J. Ablowitz and P.A. Clarkson, Soliton, nonlinear evolution equations and inverse  
388 scattering, Cambridge University Press, New York, 1991.
- 389 16. A.M. Wazwaz, A sine-cosine method for handle nonlinear wave equations, Appl. Math.  
390 Comput. Modeling, 40 (2004) 499-508.
- 391 17. E. Yusufoglu, and A. Bekir, Solitons and periodic solutions of coupled nonlinear  
392 evolution equations by using Sine-Cosine method, Int. J. Comput. Math., 83 (12) (2006)  
393 915-924.
- 394 18. J. Weiss, M. Tabor and G. Carnevale, The Painlevé property for partial differential  
395 equations, J. Math. Phys., 24 (1982) 522-526.
- 396 19. A.M. Wazwaz, Partial Differential equations: Method and Applications, Taylor and  
397 Francis, 2002.
- 398 20. M.A. Helal and M.S. Mehana, A comparison between two different methods for solving  
399 Boussinesq-Burgers equation, Chaos, Solitons Fract., 28 (2006) 320-326.
- 400 21. M. Wang, X. Li and J. Zhang, The  $(G'/G)$ -expansion method and traveling wave  
401 solutions of nonlinear evolution equations in mathematical physics, Phys. Lett. A, 372  
402 (2008) 417-423.
- 403 22. J. Zhang, F. Jiang and X. Zhao, An improved  $(G'/G)$ -expansion method for solving  
404 nonlinear evolution equations, Inter. J. Comput. Math., 87 (8) (2010) 1716-1725.

- 405 23. J. Feng, W. Li and Q. Wan, Using  $(G'/G)$ -expansion method to seek the traveling  
406 wave solution of Kolmogorov-Petrovskii-Piskunov equation, Appl. Math. Comput., 217  
407 (2011) 5860-5865.
- 408 24. M.A. Akbar, N.H.M. Ali and E.M.E. Zayed, Abundant exact traveling wave solutions of  
409 the generalized Bretherton equation via  $(G'/G)$ -expansion method, Commun. Theor.  
410 Phys., 57 (2012) 173-178.
- 411 25. R. Abazari, The  $(G'/G)$ -expansion method for Tziteica type nonlinear evolution  
412 equations, Math. Comput. Modelling, 52 (2010) 1834-1845.
- 413 26. M.A. Akbar, N.H.M. Ali and S.T. Mohyud-Din, Further exact traveling wave solutions to  
414 the (2+1)-dimensional Boussinesq and Kadomtsev-Petviashvili equation, J. Comput.  
415 Analysis Appl., 15 (3) (2013) 557-571.
- 416 27. N. Taghizadeh and M. Mirzazadeh, The first integral method to some complex nonlinear  
417 partial differential equations, J. Comput. Appl. Math., 235 (2011) 4871-4877.
- 418 28. M.L. Wang and X.Z. Li, Extended F-expansion method and periodic wave solutions for  
419 the generalized Zakharov equations, Phys. Lett. A, 343 (2005) 48-54.
- 420 29. Sirendaoreji, Auxiliary equation method and new solutions of Klein-Gordon equations,  
421 Chaos, Solitons Fract., 31 (2007) 943-950.
- 422 30. H. Triki, A. Chowdhury and A. Biswas, Solitary wave and shock wave solutions of the  
423 variants of Boussinesq equation, U.P.B. Sci. Bull., Series A, 75(4) (2013) 39-52.
- 424 31. H. Triki, A.H. Kara and A. Biswas, Domain walls to Boussinesq type equations in (2+1)-  
425 dimensions, Indian J. Phys., 88(7) (2014) 751-755.
- 426 32. J.H. He and X.H. Wu, Exp-function method for nonlinear wave equations, Chaos,  
427 Solitons Fract., 30 (2006) 700-708.

- 428 33. H. Naher, A.F. Abdullah and M.A. Akbar, New traveling wave solutions of the higher  
429 dimensional nonlinear partial differential equation by the Exp-function method, J. Appl.  
430 Math., 2012 (2012) Article ID 575387, 14 pages.
- 431 34. M. Wang, Solitary wave solutions for variant Boussinesq equations, Phy. Lett. A, 199  
432 (1995) 169-172.
- 433 35. A.J.M. Jawad, M.D. Petkovic and A. Biswas, Modified simple equation method for  
434 nonlinear evolution equations, Appl. Math. Comput., 217 (2010) 869-877.
- 435 36. E.M.E. Zayed and S.A.H. Ibrahim, Exact solutions of nonlinear evolution equations in  
436 mathematical physics using the modified simple equation method, Chin. Phys. Lett.,  
437 29(6) (2012) 060201.
- 438 37. K. Khan, M.A. Akbar and M.N. Alam, Traveling wave solutions of the nonlinear Drinfel'd-  
439 Sokolov-Wilson equation and modified Benjamin-Bona-Mahony equations, J. Egyptian  
440 Math. Soc., 21 (2013) 233-240.
- 441 38. K. Khan and M.A. Akbar, Exact and solitary wave solutions for the Tzitzeica-Dodd-  
442 Bullough and the modified Boussinesq-Zakharov-Kuznetsov equations using the  
443 modified simple equation method, Ain Shams Engr. J., 4 (2013) 903-909.
- 444 39. E.M.E. Zayed and S.A.H. Ibrahim, Exact Solutions of Kolmogorov-Petrovskii-Piskunov  
445 Equation Using the Modified Simple Equation Method, Acta Mathematicae Applicatae  
446 Sinica, 30(3) (2014) 749-754.
- 447 40. E.M.E. Zayed and A.H. Arnous, The enhanced modified simple equation method for  
448 solving nonlinear evolution equations with variable coefficients, AIP Conf. Proceedings  
449 of ICNAAM 2013, 1558 (2013) 1999-2005.
- 450 41. E.M.E. Zayed and A.H. Arnous, Exact Solutions for a Nonlinear Dynamical System in a  
451 New Double-Chain Model of DNA Using the Modified Simple Equation Method," Inform.  
452 Sci. Comp., 2013(1), Article ID ISC080713, 08 pages

- 453 42. E.M.E. Zayed, The modified simple equation method for two nonlinear PDEs with power  
454 law and kerr law nonlinearity, PanAmerican Math. J., 24(1) (2014) 65–74.
- 455 43. E.M.E. Zayed, The modified simple equation method applied to nonlinear two models  
456 of diffusion-reaction equations”, J. Math. Res. Applications, 2(2) (2014) 5-13.
- 457 44. E.M.E. Zayed and A.H. Arnous, The modified simple equation method with applications  
458 to (2+1)-dimensional systems of nonlinear evolution equations in mathematical physics,  
459 Sci. Res. Essays, 8(40) (2013) 1973-1982.
- 460 45. E.M.E. Zayed, The modified simple equation method and its applications for solving  
461 nonlinear evolution equations in mathematical physics, Commun. Appl. Nonlinear  
462 Analysis, 20(3) (2013) 95-104.
- 463 46. E.M.E. Zayed and A.H. Arnous, Exact traveling wave solutions of nonlinear PDEs in  
464 mathematical physics using the modified simple equation method, Appl. Appl. Math.: An  
465 Int. J., 8(2) (2013) 553-572.
- 466 47. E.M.E. Zayed and S.A.H. Ibrahim, Modified simple equation method and its applications  
467 for some nonlinear evolution equations in mathematical physics, Int. J. Computer Appl.,  
468 67(6) (2013) 39-44.
- 469 48. K. Khan and M.A. Akbar, Application of  $\exp(-\phi(\xi))$ -expansion method to find the exact  
470 solutions of modified Benjamin-Bona-Mahony equation, World Appl. Sci. J., 24(10)  
471 (2013) 1373-1377.
- 472 49. M.G. Hafez, M.N. Alam and M.A. Akbar, Traveling wave solutions for some important  
473 coupled nonlinear physical models via the coupled Higgs equation and the Maccari  
474 system, J. King Saud Univ.-Sci., 27(2) (2015) 105-112.
- 475 50. T.L. Bock and M.D. Kruskal, A two-parameter Miura transformation of the Benjamin-One  
476 equation, Phys. Lett. A, 74 (1979) 173-176.

- 477 51. M.N. Alam, M.A. Akbar, S. T. Mohyud-Din, General traveling wave solutions of the strain  
478 wave equation in microstructured solids via the new approach of generalized  $(G'/G)$ -  
479 expansion method, Alexandria Engr. J., 53 (2014) 233-241.
- 480 52. F. Pastrone, P. Cermelli and A. Porubov, Nonlinear waves in 1-D solids with  
481 microstructure, Mater. Phys. Mech., 7 (2004) 9-16
- 482 53. A.V. Porubov and F. Pastrone, Non-linear bell-shaped and kink-shaped strain waves in  
483 microstructured solids, Int. J. Nonlinear Mech., 39(8) (2004) 1289-1299.
- 484 54. M.A. Akbar, N.H.M. Ali and E.M.E. Zayed, A generalized and improved  $(G'/G)$ -  
485 expansion method for nonlinear evolution equations, Math. Prob. Engr., Vol. 2012,  
486 Article ID 459879, 22 pages, DOI:10.1155/2012/459879.
- 487 55. M.A. Khan and M.A. Akbar, Exact and solitary wave solutions to the generalized fifth-  
488 order KdV equation by using the modified simple equation method, Appl. Comput. Math.,  
489 4(3) (2015) 122-129.
- 490 56. A.M. Samsonov, Strain Solitons and How to Construct Them, Chapman and Hall/CRC,  
491 Boca Raton, Fla, USA, 2001.
- 492 57. F. Pastrone, F. Hierarchy of Nonlinear waves in complex microstructured solids, Int.  
493 Con. Com. Of nonlinear waves, (2009) 5-7.
- 494 58. A. A. Zakharenko, Analytical studying the group velocity of three-partial Love (type)  
495 waves in both isotropic and anisotropic media, Nondestructive Testing and Evaluation,  
496 20(4), Dec. 2005, 237-254.
- 497 59. A. A. Zakharenko, Slow acoustic waves with the anti-plane polarization in layered  
498 systems, Int. J. of Modern Phys. B, 24(4) (2010) 515-536.